

SInstitute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham, New Hampshire 03824, USA

1. Bai, Y., Han, X., Wu, J., Chen, Z. & Li, L. *Nature* **431**, 181–184 (2004).
2. Doak, D. F. *et al.* *Am. Nat.* **151**, 264–276 (1998).
3. Tilman, D. *Ecology* **77**, 350–363 (1996).
4. McNaughton, S. J. *Am. Nat.* **111**, 515–525 (1977).
5. McNaughton, S. J. *Ecol. Monogr.* **55**, 259–294 (1985).
6. Huston, M. C. & McBride, A. C. in *Biodiversity and Ecosystem*

Functioning: Synthesis and Perspectives (eds Loreau, M., Naeem, S. & Inchausti, P.) 47–60 (Oxford Univ. Press, Oxford, 2002).

7. Xiao, X. M., Wang, Y., Jiang, S., Ojima, D. S. & Bonham, C. D. *J. Arid Environ.* **31**, 283–299 (1995).
8. Xiao, X. M., Jiang, S., Wang, Y., Ojima, D. S. & Bonham, C. D. *Vegetatio* **123**, 1–12 (1996).
9. Chen, Z. Z. & Wang, S. P. (eds) *Typical Steppe Ecosystem of China* (Science, Beijing, 2000).

doi:10.1038/nature03862

PLANT COMMUNITIES

Ecosystem maturity and performance

Arising from: Bai, Y., Han, X., Wu, J., Chen, Z. & Li, L. *Nature* **431**, 181–184 (2004)

The effect of maturity, or successional stage, on ecosystem performance (measured as productivity or stability, for example) is important for both basic ecology and ecosystem management. On the basis of the results of a long-term study of two different plant communities at two sites in the Inner Mongolia grassland¹, Bai *et al.* claim that these communities simultaneously achieve high species richness, productivity and ecosystem stability at the late successional stage¹. However, I question their interpretation of the data and suggest that this claim is undermined by evidence from other empirical and theoretical studies.

Bai *et al.*¹ present data on above-ground biomass of the total community and on various plant groups, but they provide no time series to indicate how diversity, productivity and stability might have changed during succession.




Figure 1 | Changes during succession. Model of temporal changes in diversity (species richness; green curve), productivity (blue), biomass (turquoise) and stability (red) during succession, based on an extensive literature review^{5,6}. In the transitional stage, both early (short-lived) and late (long-lived) species coexist, leading to high species diversity. In this stage, biomass is relatively low and resource level is still high, promoting higher productivity. In the late stage, stability may be high but diversity and productivity may be low because of competitive exclusions⁷ and sequestration of limited resources by accumulated litter and biomass^{5,12}.

The two communities compared at their site A represent two extreme degrees of disturbance: one undisturbed and the other heavily degraded. Although the undisturbed community did support relatively higher diversity and above-ground biomass than the highly degraded community, intermediate amounts of disturbance or transitional stages, which may well support even higher diversity and productivity, were not accounted for^{2–6} (Fig. 1).

Over space (that is, within one habitat), the ‘intermediate disturbance’ hypothesis⁷ predicts that diversity will be highest at an intermediate level of disturbance^{2,3}. Extensive evidence^{2–12} in support of this hypothesis is not in agreement with the conclusions of Bai *et al.*

Finally, over time at one locality, successional studies from various ecosystems — particularly those covering entire successional cycles — reveal that biodiversity and productivity are highest in the mid- or transitional-stage of succession, when both early- and late-stage species coexist. The high diversity and productivity then gradually decline owing to accumulated biomass and litter and therefore to increasing competition^{2–12} (Fig. 1).

Although each particular case would be expected to show some deviation from the general patterns in Fig. 1, because of the life history of dominant species, less destructive

disturbance, or variation in resources available over time¹³, for example, it is not clear why the Inner Mongolia grassland should be so different¹. If community stability, whose estimate depends on how it is measured, and biomass both increase with succession and are really high in undisturbed mature ecosystems, then the contrasting patterns between diversity and stability call their relationship into question.

The high stability in mature, or late-stage succession, grassland may be at least in part caused by the longer lifespan of the remaining, competitive perennial species (unlike annuals or short-lived plants in early succession) and by the high accumulated biomass, rather than by species diversity. For example, if community stability is measured as the coefficient of variation in biomass (CV, variance/mean)¹, then the CV, which is not independent of mean biomass¹⁴, will be lower when the biomass is higher, so the stability should be higher. Long-term, simultaneous monitoring of these variables in both above- and below-ground communities over the entire successional cycles of the grassland would help to clarify this point.

Qinfeng Guo

US Geological Survey, NPWRC, Jamestown, Nevada 58401, USA

e-mail: qguo@usgs.gov

1. Bai, Y., Han, X., Wu, J., Chen, Z. & Li, L. *Nature* **431**, 181–184 (2004).
2. Connell, J. H. *Science* **199**, 1302–1310 (1978).
3. Huston, M. *Biological Diversity* (Cambridge Univ. Press, Cambridge, 1994).
4. Gibson, D. J. & Hulbert, L. C. *Vegetatio* **72**, 175–185 (1987).
5. Guo, Q. *J. Veg. Sci.* **14**, 121–128 (2003).
6. *J. Veg. Sci.* http://www.opuluspress.se/pub/archives/JVS/archive_J.html
7. Grime, J. P. *Nature* **242**, 344–347 (1973).
8. Guo, Q. *Ambio* **32**, 428–430 (2003).
9. Lichter, J. *Ecol. Monogr.* **68**, 487–510 (1998).
10. Odum, E. P. *Science* **164**, 262–270 (1969).
11. Whittaker, R. H. *Communities and Ecosystems* 2nd edn (Macmillan, New York, 1975).
12. Ryan, M. G., Binkley, D., Fownes, J. H., Giardina, C. P. & Senock, R. S. *Ecol. Monogr.* **74**, 393–414 (2004).
13. Yang, L. H. *Science* **306**, 1565–1567 (2004).
14. Collins, S. L. *Trends Ecol. Evol.* **10**, 95–96 (1995).

doi:10.1038/nature03583

PLANT COMMUNITIES

Wu *et al.* reply

Reply to: Wang, S. *et al.* *Nature* doi:10.1038/nature03862 (2005) and Guo, Q. *Nature* doi:10.1038/nature03583 (2005)

Some of our findings¹ are questioned by Wang *et al.*², on the basis that we use inconsistent data and fail to distinguish spatial heterogeneity effects. Here we show that both claims are unfounded. We also address the questions raised by Guo³ concerning how the steppe communities vary as they mature.

In 1998 (not in 1999, as Wang *et al.* indicate), the sampling transect in our site A (the

Leymus chinensis community) was moved about 60 m northward within the same fenced plant community in a fairly uniform environmental setting to allow for more replicates and other sampling activities. To assure data consistency, we chose 5 quadrats in the new location at exactly the same distance interval as in the old transect. Our reanalysis of the 1980–97 data confirms our previous result¹ that live