

NEUROSCIENCE

Rewiring the adult brain

Arising from: S. M. Smirnakis *et al.* *Nature* 435, 300–307 (2005)

Any analysis of plastic reorganization at a neuronal locus needs a veridical measure of changes in the functional output — that is, spiking responses of the neurons in question. In a study of the effect of retinal lesions on adult primary visual cortex (V1), Smirnakis *et al.*¹ propose that there is no cortical reorganization. Their results are based, however, on BOLD (blood-oxygen-level-dependent) fMRI (functional magnetic resonance imaging), which provides an unreliable gauge of spiking activity. We therefore question their criterion for lack of plasticity, particularly in the light of the large body of earlier work that demonstrates cortical plasticity.

Plasticity in adult V1 has been demonstrated by multiple, independent lines of evidence from more than twenty studies in three species (see refs 2–6, for example). Physiologically, the evidence derives from measurements of lesion-induced shifts in the locations of V1 neuronal receptive fields. By plotting receptive fields before and at various points after making retinal lesions, it was shown that the affected cortex — with receptive fields originally inside the lesion — develops new, shifted receptive field positions after recovery. These shifts are cortically mediated because the lateral geniculate nucleus, the source of thalamic input to V1, shows limited reorganization⁷. All these measurements were made using suprathreshold spiking neuronal responses: this is an important point as the neuronal output from V1 to subsequent cortical stages is carried entirely by spikes. Any measure of V1 reorganization — and consequent functional remapping of visual information — therefore needs to assess the effect on spiking activity. These physiological results are buttressed by anatomical findings showing a selective increase in the density of axon collaterals in reorganized cortex⁸, and the sequential expression of biochemical markers^{9,10}.

By contrast, the primary evidence for lack of plasticity offered by Smirnakis *et al.* is the observation that the V1 'silent zone', mapped with BOLD fMRI immediately following a retinal lesion, did not change over time. There are plausible reasons why fMRI maps may fail to change, despite re-emergent neuronal activity. The reorganization of cortex is believed to be mediated by long-range horizontal connections within V1 (refs 8, 11, 12). In normal V1, these connections mediate subthreshold modulation. Following retinal lesions, horizontal connections stretching from 'normal' cortex into the lesion projection zone (LPZ) are believed to strengthen their synapses — but not to change anatomical extent. They there-

fore induce re-emergent spiking activity, but only in neurons lying within their target zone in the silenced cortex.

Such re-emergent activity could involve reduction of inhibition as much as an increase in excitation. As the BOLD signal probably reflects synaptic input into a region rather than spiking output, the 'silent zone' observed by Smirnakis *et al.* immediately following a lesion may mark not the edge of the real LPZ but the inner edge of subthreshold activation spreading into the LPZ through horizontal connections. In subsequent measurements, the BOLD signal would continue to show the unchanging position of this inner boundary while being blind to synaptic reorganization, which would lead to re-emergent spiking activity over the extent of the horizontal connections. The single set of electrode recordings by Smirnakis *et al.* after months of recovery might simply show the extent of largely completed recovery and, not surprisingly, produce a border in register with the edge of the BOLD signal.

Furthermore, BOLD gives a local measure of the total cortical activity, a significant component of which comes from thalamocortical inputs, the contribution of which is probably further accentuated by the disproportionately high vascularization of layer 4, the cortical input layer. However, the neurons showing recovery may reside primarily in the superficial layers¹², which receive the long-range horizontal connections, as opposed to layer 4.

Owing to these uncertainties about the validity of BOLD fMRI as a yardstick of functional reorganization in V1, we believe that Smirnakis *et al.* do not present a convincing contradiction to the body of earlier evidence indicating substantial receptive field plasticity in adult animals following retinal lesion. The recovered activity demonstrated in the earlier studies has a likely corollary in the recovery of visual perception: human subjects suffering

from macular degeneration, or with artificially induced retinal lesions, show improved perceptual fill-in over time after the lesions^{13–15}.

Michael B. Calford*, Yu-ze M. Chino†, Aniruddha Das‡, Ulf T. Eysel§, Charles D. Gilbert||, Stephen J. Heinen¶, Jon H. Kaas#, Shimon Ullman☆

*School of Biomedical Sciences, The University of Newcastle, New South Wales 2308, Australia

†College of Optometry, University of Houston, Houston, Texas 77004, USA

‡Columbia University Medical School, New York, New York 10032, USA

§Department of Neurophysiology, Faculty of Medicine, Ruhr-University Bochum, 44780 Bochum, Germany

||The Rockefeller University, New York, New York 10021, USA

¶The Smith-Kettlewell Eye Research Institute, San Francisco, California 94115, USA

#Vanderbilt University, Nashville, Tennessee 37240, USA

☆Weizmann Institute, Rehovot 76100, Israel

- Smirnakis, S. M. *et al.* *Nature* 435, 300–307 (2005).
- Heinen, S. J. & Skavenski, A. A. *Exp. Brain Res.* 83, 670–674 (1991).
- Calford, M. B. *et al.* *J. Physiol. Lond.* 524, 587–602 (2000).
- Gilbert, C. D., Hirsch, J. A. & Wiesel, T. N. *Cold Spring Harbor Symp. Quant. Biol.* 55, 663–677 (1990).
- Kaas, J. H. *et al.* *Science* 248, 229–231 (1990).
- Chino, Y. M., Smith, E. L. III, Kaas, J. H., Sasaki, Y. & Cheng, H. *J. Neurosci.* 15, 2417–2433 (1995).
- Eysel, U. T. *Nature* 299, 442–444 (1982).
- Darian-Smith, C. & Gilbert, C. D. *Nature* 368, 737–740 (1994).
- Obata, S., Obata, J., Das, A. & Gilbert, C. D. *Cereb. Cortex* 9, 238–248 (1999).
- Arckens, L. *et al.* *Eur. J. Neurosci.* 12, 4222–4232 (2000).
- Das, A. & Gilbert, C. D. *Nature* 375, 780–784 (1995).
- Calford, M. B., Wright, L. L., Metha, A. B. & Tagliavetti, V. *J. Neurosci.* 23, 6434–6442 (2003).
- Craik, K. J. W. in *The Nature of Psychology* (ed. Sherwood, S. L.) 98–103 (Cambridge Univ. Press, Cambridge, 1966).
- Gerrits, H. J. & Timmerman, G. J. *Vision Res.* 9, 439–442 (1969).
- Zur, D. & Ullman, S. *Vision Res.* 43, 971–982 (2003).

doi:10.1038/nature04359

NEUROSCIENCE

Smirnakis *et al.* reply

Replying to: M. B. Calford *et al.* *Nature* 438, doi:10.1038/04359 (2005)

We disagree with Calford *et al.*¹ that there is a consensus on adult plasticity in primate V1 cortex: for example, macaque area V1 cytochrome oxidase levels remained depressed for several months after binocular retinal lesions²; no reorganization in macaque V1 after monocular retinal lesions was found³; and no area

V1 reorganization in a patient with macular degeneration was detected⁴.

Calford *et al.*¹ agree that subthreshold activity shows no long-term reorganization. They propose that plasticity comprises only an increase in the likelihood of transforming subthreshold signals into action