

Nitrogen saturation and net ecosystem production

Arising from: F. Magnani *et al.* *Nature* **447**, 848–850 (2007)

Magnani *et al.*¹ found that net carbon (C) sequestration of temperate and boreal forests is clearly driven by nitrogen (N) deposition. From the positive relationship between average net ecosystem production (NEP) and wet N deposition, the authors further conclude that “no signs of N saturation were apparent” in the studied forests and that this is “casting doubts on the risk of widespread ecosystem nitrogen saturation”. Nitrogen additions can clearly alter net ecosystem production, but net ecosystem production cannot be used as an indicator of N saturation.

Nitrogen saturation implies a change in N cycling from a closed internal cycle to an open cycle² where excess N is leached and/or emitted from the forest ecosystem. These changes in forest ecosystem functioning have been extensively documented^{3,4}. Examples from literature lead us to suggest that some of the forest ecosystems discussed in the concerned article might be N saturated, irrespective of the increased net ecosystem production. Evidence for N saturation has been observed in forest ecosystems subject to N deposition levels similar to the relatively low deposition range reported in the concerned article¹ (that is, less than 10 kg N ha⁻¹ yr⁻¹ wet deposition), including considerable nitrate loss (up to 10 kg N ha⁻¹ yr⁻¹) through runoff or seepage water^{5–7} and elevated emissions of NO and N₂O⁸. Because Magnani *et al.*¹ did not measure any of these pathways of N loss, they cannot rule out N saturation in the studied forest ecosystems.

The demonstrated relationship between N deposition and C sequestration is an important finding, consistent with other literature⁹. In our opinion, however, Magnani *et al.*¹ demonstrate an incorrect view on the phenomenon of N saturation and, in doing so, greatly ignore the effect of N deposition and saturation on soil acidification, groundwater and surface water quality, biodiversity, and ecosystem services other than C sequestration. Because the data presented by Magnani *et al.*¹ do not allow an evaluation of the N saturation status of the studied forests, the expressed “doubts on the risk of widespread ecosystem nitrogen saturation” are not substantiated. At the moment, N saturation of forest ecosystems is probably not yet a widespread problem on a global scale, but it is surely a widespread problem in densely populated and more industrialized regions^{4,7}. From the Europe-covering IFEF (Indicators of Forest Ecosystem Functioning) and Level-II (UN-ECE/EC intensive monitoring plots) databases^{10,11}, it can be deduced that more than 25% of the European forests included in these databases are N saturated (considering the

nitrate seepage flux as indicator and 5 kg (357 mol) N ha⁻¹ yr⁻¹ as a threshold value of N saturation). Furthermore, from a large number of sites in the northeastern United States⁴, N saturation was indicated as a frequently occurring phenomenon. As substantial increases in global N emissions are predicted for the coming 50 yr¹², the potential risk of widespread N saturation of forest ecosystems in the long term cannot be denied.

An De Schrijver¹, Kris Verheyen¹, Jan Mertens¹, Jeroen Staelens¹, Karen Wuyts¹ & Bart Muys²

¹Laboratory of Forestry, Ghent University, Geraardsbergse Steenweg 267, 9090 Gontrode, Belgium.

e-mail: An.Deschrijver@Ugent.be

²Division Forest, Nature and Landscape, Katholieke Universiteit Leuven, Celestijnenlaan 200E, 3001 Leuven, Belgium.

Received 20 July; accepted 30 October 2007.

- Magnani, F. *et al.* The human footprint in the carbon cycle of temperate and boreal forests. *Nature* **447**, 848–850 (2007).
- Matson, P., Lohse, K. A. & Hall, S. J. The globalization of nitrogen deposition: Consequences for terrestrial ecosystems. *Ambio* **31**, 113–119 (2002).
- Aber, J. D. *et al.* Nitrogen saturation in temperate forest ecosystems—Hypotheses revisited. *Bioscience* **48**, 921–934 (1998).
- Aber, J. D. *et al.* Is nitrogen deposition altering the nitrogen status of northeastern forests? *Bioscience* **53**, 375–389 (2003).
- Dise, N. B., Matzner, E. & Gundersen, P. Synthesis of nitrogen pools and fluxes from European forest ecosystems. *Wat. Air Soil Pollut.* **105**, 143–154 (1998).
- Macdonald, J. A. *et al.* Nitrogen input together with ecosystem nitrogen enrichment predict nitrate leaching from European forests. *Glob. Change Biol.* **8**, 1028–1033 (2002).
- Gundersen, P. K., Schmidt, I. K. & Raulund-Rasmussen, K. Leaching of nitrate from temperate forests—effects of air pollution and forest management. *Environ. Rev.* **14**, 1–57 (2006).
- Pilegaard, K. *et al.* Factors controlling regional differences in forest soil emission of nitrogen oxides (NO and N₂O). *Biogeosciences* **3**, 651–661 (2006).
- Karjalainen, T. *et al.* Causes and Consequences of Forest Growth Trends in Europe—Results of the RECOGNITION Project. European Forest Institute Research Report (Brill, Leiden, in the press).
- Lorenz, M. *et al.* Nitrogen retention and release in European forests: Deriving indicators from large databases. In *Forest Condition in Europe: 2005 Technical Report* (Federal Research Centre for Forestry and Forest Products (BFH), UNECE Geneva, 2005).
- Van der Salm, C., de Vries, W., Reinds, G. J. & Dise, N. B. N leaching across European forests: Derivation and validation of empirical relationships using data from intensive monitoring plots. *For. Ecol. Manage.* **238**, 81–91 (2007).
- Galloway, J. N. *et al.* Nitrogen cycles: past, present, and future. *Biogeochemistry* **70**, 153–226 (2004).

doi:10.1038/nature06578

Ecologically implausible carbon response?

Arising from: F. Magnani *et al.* *Nature* **447**, 848–850 (2007)

Magnani *et al.*¹ present a very strong correlation between mean lifetime net ecosystem production (NEP, defined as the net rate of carbon (C) accumulation in ecosystems²) and wet nitrogen (N) deposition. For their data in the range 4.9–9.8 kg N ha⁻¹ yr⁻¹, on which the correlation largely depends, the response is approximately 725 kg C per kg N in wet deposition. According to the authors, the maximum N wet deposition level of 9.8 kg N ha⁻¹ yr⁻¹ is equivalent to a total deposition of 15 kg N ha⁻¹ yr⁻¹, implying a net sequestration near 470 kg C per kg N of total deposition. We question the ecological plausibility of the relationship and show, from a

multi-factor analysis of European forest measurements, how interactions with site productivity and environment imply a much smaller NEP response to N deposition.

The C response to N deposition is restricted by the C:N stoichiometry of the forest ecosystem compartments. The implied NEP response of 470 kg C per kg N would require that the fate of the deposited nitrogen was exclusively in stem wood, which is the only carbon sink with a C:N ratio of this magnitude. This is unreasonable because N-limited forest stands, as suggested¹, invest primarily in roots³, with C:N ratios near 50–100. The impossibility of near-total

Table 1 | Multivariate regression results at stand level and individual tree level**a** Results at stand level for N sensitive plots with parameter estimates

Tree species	Site productivity*	Age†	SDI‡	N deposition§	Drought	Temperature change¶
All plots						
Norway spruce	0.054	-0.005	-	0.020#	-	0.524
Scots pine	-	-0.017	-	0.010	-0.0032	-
Sensitive plots						
Norway spruce	0.039	-0.004	-	0.022	-	0.32
Scots pine	-	-0.017	0.001	0.013	-0.002	-

b Results at individual tree level with parameter estimates for the main influencing factors included

Tree species	BAL☆	SDI‡	C:N _{soil} **	N deposition§	Temperature ††	Temperature change¶
Norway spruce	-0.39	-0.00056	-0.023	0.013	-	-
Scots pine	-0.29	-0.00066	-	0.015	0.053	-
Common beech	-0.16	-	-	0.012	-	0.064
Oak	-0.38	-0.00062	-	0.013	0.080	-

Multivariate regression results indicating the relative change in stem volume growth per unit change in influencing factor (for example, a value of 0.013 for N deposition implies an increase in stem growth of 1.3% for each additional 1 kg ha⁻¹ yr⁻¹ of N deposition). Note that a dash (-) implies that the effect was insignificant ($P > 0.05$). Sensitive plots are plots with a C:N ratio above 25.

* Site productivity is a variable (m³ ha⁻¹ yr⁻¹) derived from selected European site index curves, with input variables being age and top height.

† Stand age (yr).

‡ SDI, stand density index (number of trees per ha).

§ N deposition is total N deposition (unit change: kg ha⁻¹ yr⁻¹).

|| Drought is a variable describing drought given as a relative value (unit change: %) to the normal (30-yr mean) drought stress at each site.

¶ The temperature difference during the growing period compared with the 30-yr average temperature (unit change: °C).

Results from a linear regression; in the multivariate analysis the coefficient was just not significant at $P < 0.05$.

☆ BAL is basal area of larger trees, which affects tree competition (m² ha⁻¹).

** C:N_{soil} is the C:N ratio of the mineral topsoil (0–30 cm).

†† Temperature is average yearly temperature during the investigation period 1993–2000 (°C).

storage of deposited N in stem wood follows also from the expected N leaching rates, varying between 10–50% of the N input in a range of 10–25 kg N ha⁻¹ yr⁻¹ (refs 4, 5), which is the likely range for total N deposition at the plots of Magnani *et al.*¹ where the high C:N response was found (that is, their European sites where wet N deposition was estimated at 4.9–9.8 kg N ha⁻¹ yr⁻¹). This total N deposition range follows from an application of the EMEP (European Monitoring and Evaluation Programme of the Long-range Transmission of Air Pollutants) model, used in the NITROEUROPE project, for the year 2000 to the sites of Magnani *et al.*¹. Using this model, the estimates of which are in close agreement with measured atmospheric N deposition at forest sites⁶, we found that total N deposition at the sites of Magnani *et al.*¹ is 2–7 times greater than wet deposition. This analysis implies a relationship of approximately 175 kg C per kg N in the mentioned total N deposition range.

However, even this lower response is unlikely. ¹⁵N-labelled tracer experiments in temperate forests indicate that N retention hardly occurs in stem wood but mainly in the soil⁷. Considering the fate of N and the ranges in C:N ratios in forest ecosystem compartments, this implies a carbon response near 50 kg C per kg N in forest ecosystems^{7,8}. Even though the above-ground C sequestration may be underestimated by Nadelhoffer *et al.*⁷, owing to neglecting the effect of direct foliar uptake^{9,10}, this effect is likely to be small, as above-ground foliar N uptake is generally less than 5 kg N ha⁻¹ yr⁻¹ (ref. 11), whereas below-ground uptake is generally more than 50 kg N ha⁻¹ yr⁻¹. Furthermore, similar results are found in long-term (15–30 yr) nitrogen-fertilizer trials at rates of nitrogen addition below 50 kg N ha⁻¹ yr⁻¹ (refs 12, 13) and in process-based model simulations¹⁴. The reason for the extremely high influence of N deposition on NEP suggested by Magnani *et al.*¹ is probably due to the contribution of other factors—which co-vary with wet N deposition—to the derived relationship. The authors filtered out the effects of age and investigated the separate effect of temperature but they aggregated all tree species and site characteristics, such as site fertility and stand density, into one relationship.

We carried out a multi-factor analysis of measured forest growth data at nearly 400 intensively monitored forest plots in Europe, including Norway spruce, Scots pine, common beech and oak. The influence of nitrogen and acid deposition was considered by using values during the growth period (1993–2000), whereas the impacts of temperature, precipitation and drought were addressed by taking the

deviation of these climatic parameters in the growth period (1993–2000) from the 30-yr mean. We simultaneously accounted for site factors influencing measured tree growth, including site productivity, stand age and stand density. We also applied a multi-factor analysis with measured basal-area increment of each individual tree as responding factor. The influencing factors used in this study and the results of the multivariate analyses at stand and individual tree level are shown in Table 1. The approach at tree level indicated a 1.2–1.5% increase in basal-area increment, depending on tree species, in response to 1 kg of N ha⁻¹ yr⁻¹. The approach at stand level indicated only a significant response of Norway spruce and Scots pine to N with roughly a 1–2% increase in volume growth in response to 1 kg of N ha⁻¹ yr⁻¹, depending on the C:N ratio of the plots. We recalculated these responses in terms of C sequestration by multiplying the mean measured volume growth at each stand with the estimated growth increase and the mean wood density of each tree species, assuming a C content of 50%. The results of our analyses at both tree and stand level indicate a response of trees between approximately 20–40 kg C per kg N. Additionally, results of long-term nitrogen addition experiments indicate soil responses of 10–30 kg C per kg N^{12,13,15}. Thus, the total NEP response would be about 30–70 kg C per kg N, which is much smaller than that estimated by Magnani *et al.*¹.

METHODS

The multivariate regression at stand level was carried out by a backward stepwise method, where the model was reduced step-by-step by removing nonsignificant effects. The results of the hypothesis testing of the effects are based on partial *F*-tests.

The multivariate regression analysis at tree level was carried out by using tree size and tree competition variables on tree level and site factors and environmental factors on plot level, including plot as a random effect and applying the restricted maximum likelihood (REML) method for parameter estimation. Parameters in the model, which exhibited significant ($P < 0.05$) coefficients and behaved according to their known impacts, are included in Table 1b. We accounted for correlations, such as those between climatic parameters and N deposition (for example, we found a quadratic relation between N deposition and temperature; $R^2 = 0.62$).

Wim de Vries¹, Svein Solberg², Matthias Dobbertin³, Hubert Sterba⁴, Daniel Laubhahn⁴, Gert Jan Reinds¹, Gert-Jan Nabuurs¹, Per Gundersen⁵ & Mark A. Sutton⁶

¹Alterra, Wageningen University and Research Centre, PO Box 47, 6700 AA Wageningen, The Netherlands.

e-mail: wim.devries@wur.nl

²Norwegian Forest and Landscape Institute, PO Box 115, N-1431 Ås, Norway.

³Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland.

⁴University of Natural Resources and Applied Life Sciences, Peter Jordan-Straße 82, A-1190 Vienna, Austria.

⁵Danish Centre for Forest, Landscape and Planning, University of Copenhagen, Hoersholm, Denmark.

⁶Centre for Ecology and Hydrology, Edinburgh Research Station, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK.

Received 23 July; accepted 16 November 2007.

- Magnani, F. *et al.* The human footprint in the carbon cycle of temperate and boreal forests. *Nature* **447**, 848–850 (2007).
- Woodwell, G. M. & Whittaker, R. H. Primary production in terrestrial ecosystems. *Am. Zool.* **8**, 19–30 (1968).
- Brouwer, R. Functional equilibrium: sense or nonsense. *Neth. J. Agric. Sci.* **31**, 335–348 (1983).
- Gundersen, P., Schmidt, I. K. & Raulund-Rasmussen, K. Leaching of nitrate from temperate forests – effects of air pollution and forest management. *Environ. Rev.* **14**, 1–57 (2006).
- De Vries, W., van der Salm, C., Reinds, G. J. & Erisman, J. W. Element fluxes through European forest ecosystems and their relationships with stand and site characteristics. *Environ. Pollut.* **148**, 501–513 (2007).

- Simpson, D., Fagerli, H., Hellsten, S., Knulst, J. C. & Westling, O. Deposition and emissions of reactive nitrogen over European forests: A modelling study. *Biogeosciences* **3**, 337–355 (2006).
- Nadelhoffer, K. J. *et al.* Nitrogen deposition makes a minor contribution to carbon sequestration in temperate forests. *Nature* **398**, 145–148 (1999).
- De Vries, W., Reinds, G. J., Gundersen, P. & Sterba, H. The impact of nitrogen deposition on carbon sequestration in European forests and forest soils. *Glob. Change Biol.* **12**, 1151–1173 (2006).
- Jenkinson, D. S., Goulding, K. & Powlson, D. S. Nitrogen deposition and carbon sequestration. *Nature* **400**, 629 (1999).
- Sievering, H. Nitrogen deposition and carbon sequestration. *Nature* **400**, 629–630 (1999).
- Sievering, H., Fernandez, I., Lee, J., Hom, J. & Rustad, L. Forest canopy uptake of atmospheric nitrogen deposition at eastern US conifer sites: Carbon storage implications? *Glob. Biogeochem. Cycles* **14**, 1153–1160 (2000).
- Hyvönen, R. *et al.* Impact of long-term nitrogen addition on carbon stocks in trees and soils in northern Europe. *Biogeochemistry* doi:10.1007/s10533-007-9121-3 (2007).
- Högberg, P., Fan, H., Quist, M., Binkley, D. & Tamm, C. O. Tree growth and soil acidification in response to 30 years of experimental nitrogen loading on boreal forest. *Glob. Change Biol.* **12**, 489–499 (2006).
- Levy, P. E., Wendler, R., Van Oijen, M., Cannell, M. G. R. & Millard, P. The effect of nitrogen enrichment on the carbon sink in coniferous forests: uncertainty and sensitivity analyses of three ecosystem models. *Water Air Soil Pollut. Focus* **4**, 67–74 (2004).
- Evans, C. D. *et al.* Modelling nitrogen saturation and carbon accumulation in heathland soils under elevated nitrogen deposition. *Environ. Pollut.* **143**, 468–478 (2006).

doi:10.1038/nature06579

Magnani *et al.* reply

Replying to: A. De Schrijver *et al.* *Nature* **451**, doi:10.1038/nature06578; W. de Vries *et al.* *Nature* **451**, doi:10.1038/nature06579 (2008)

Nitrogen (N) deposition alters ecosystem function in several ways, with important effects on N leaching and water quality, as well as on interspecific competition and biodiversity. These changes have been attributed to ecosystem N saturation, defined as the alleviation of N limitations on rates of biological function¹. After an initial fertilization effect, N saturation has also been suggested to reduce plant function and growth², eventually leading to forest dieback. Although our observation of a substantial positive effect of N deposition on forest carbon (C) sequestration³ does not imply the absence of nitrate losses or other negative effects, as rightly stressed by De Schrijver *et al.*⁴, the sustained response observed demonstrates that the fear of a generalized forest decline in response to N fertilization could be overstated, at least within the rather broad N deposition range explored in our analysis. The nature of the observed response of forest C sequestration to N deposition, however, has been questioned outright by de Vries *et al.*⁵, who suggested that it could be an artefact resulting from the covariation between N deposition and other environmental variables. The arguments proposed against an overwhelming N effect, however, do not seem to stand up to close scrutiny.

We agree that ecosystem gross primary production (GPP) and plant growth are, to a large extent, controlled by local climate, drought and fertility (that is, N mineralization associated with soil organic matter decomposition), although fertility could be itself influenced by current and past N deposition⁶. However, the same environmental factors would modulate in parallel ecosystem respiration, and as a result do not seem to affect net ecosystem production (NEP), which is the difference between GPP and ecosystem respiration and is the subject of our analysis³. Both components of NEP seem to be also affected by N deposition, but in opposite directions: apart from the positive effects on plant growth considered by de Vries *et al.*⁵, respiration is known to be significantly reduced by N fertilization, as demonstrated by manipulation experiments^{7,8} as well as regional transect studies⁹. The combined effect at the ecosystem level is largely missed when focusing on tree growth alone.

The question remains of the magnitude of the observed response to N deposition. Assuming a linear relationship between NEP and N

deposition, a slope of 445 ± 38 kg C per kg N of wet N deposition can be inferred from our entire data set ($n = 20$, rather than the subsample of 8 data points in the analysis by de Vries *et al.*⁵). If we assume, rather conservatively, that wet deposition constitutes 40–50% of total N deposition¹⁰, this would imply a NEP sensitivity to total N deposition of approximately 175–225 kg C per kg N, which is consistent with the stoichiometry of plant tissues and soil organic matter. Although it is true that fine roots account for a significant fraction of forest growth, it should be noted that one of the main effects of increased N availability is an increased allocation to woody tissues (with a high C:N ratio of up to 500:1) away from fine roots¹¹. This mechanism could indeed represent an important component of the observed response to N deposition.

Far from implausible, a 200:1 sensitivity is nevertheless higher than suggested by long-term forest fertilization experiments¹². Potential problems with N manipulation studies have already been discussed¹³. In particular, they overlook the role of canopy N uptake, which enables plants to absorb a relevant fraction of incoming N without any competition from soil microbes. Canopy N uptake amounts to up to 70% of N deposition, providing as much as one-third of tree N requirements^{9,14,15}. The critical comparison of results from ecosystem manipulation and observational studies could be providing a rare, unforeseen insight into the key factors controlling C–N relations in forest ecosystems.

Federico Magnani¹, Maurizio Mencuccini², Marco Borghetti³, Frank Berninger⁴, Sylvain Delzon⁵, Achim Grelle⁶, Pertti Hari⁷, Paul G. Jarvis², Pasi Kolari⁷, Andrew S. Kowalski⁸, Harry Lankreijer⁹, Beverly E. Law¹⁰, Anders Lindroth⁹, Denis Loustau¹¹, Giovanni Manca¹², John B. Moncrieff², Vanessa Tedeschi³, Riccardo Valentini¹³ & John Grace²

¹Department of Fruit Tree and Woody Plant Science, University of Bologna, Bologna I-40127, Italy.

e-mail: federico.magnani@unibo.it

²School of GeoSciences, University of Edinburgh, Edinburgh EH93JU, UK.

³Department of Crop Systems, Forestry and Environmental Sciences, University of Basilicata, Potenza I-85100, Italy.

⁴Departement des Sciences Biologiques, University of Québec à Montréal, Montréal, Quebec H3C 3P8, Canada.

⁵UMR BIOGECO, INRA—Université Bordeaux 1, F-33400 Talence Cedex, France.

⁶Department of Ecology and Environmental Research, Swedish University of Agricultural Sciences, SE-75007 Uppsala, Sweden.

⁷Department of Forest Ecology, University of Helsinki, FIN-00014 Helsinki, Finland.

⁸Department of Applied Physics, University of Granada, E-18071 Granada, Spain.

⁹Department of Physical Geography and Ecosystems Analysis, Lund University, S-223 62 Lund, Sweden.

¹⁰College of Forestry, Oregon State University, Corvallis, Oregon 97331, USA.

¹¹INRA, UR1263 EPHYSE, F-33883 Villenave d'Ornon, France.

¹²Institute for Environment and Sustainability—Climate Change Unit, Joint Research Center, European Commission, I-21020 Ispra, Italy.

¹³Department of Forest Resources and Environment, University of Tuscia, Viterbo I-01100, Italy.

1. Aber, J. D. *et al.* Nitrogen saturation in temperate forest ecosystems. *Bioscience* **48**, 921–934 (1998).
2. Aber, J. D. *et al.* Is nitrogen deposition altering the nitrogen status of northeastern forests? *Bioscience* **53**, 375–389 (2003).
3. Magnani, F. *et al.* The human footprint in the carbon cycle of established temperate and boreal forests. *Nature* **447**, 848–850 (2007).
4. De Schrijver, A. *et al.* Nitrogen saturation and net ecosystem production. *Nature* **451**, doi:10.1038/nature06578 (2008).
5. de Vries, W. *et al.* Ecologically implausible carbon response? *Nature* **451**, doi:10.1038/nature06579 (2008).
6. Spiecker, H. Overview of recent growth trends in European forests. *Wat. Air Soil Pollut.* **116**, 33–46 (1999).
7. Bowden, R. D., Davidson, E., Savage, K., Arabia, C. & Steudler, P. Chronic nitrogen additions reduce total soil respiration and microbial respiration in temperate forest soils at the Harvard Forest. *For. Ecol. Manage.* **196**, 43–56 (2004).
8. Olsson, P., Linder, S., Giesler, R. & Högberg, P. Fertilization of boreal forest reduces both autotrophic and heterotrophic soil respiration. *Glob. Change Biol.* **11**, 1745–1753 (2005).
9. Schulze, E.-D. (ed.) *Carbon and Nitrogen Cycling in European Forest Ecosystems* (Springer, Berlin, 2000).
10. Dentener, F. *et al.* Nitrogen and sulfur deposition on regional and global scales. A multimodel evaluation. *Glob. Biogeochem. Cycles* **20**, GB4003 (2006).
11. Litton, C. M., Raich, J. W. & Ryan, M. G. Carbon allocation in forest ecosystems. *Glob. Change Biol.* **13**, 2089–2109 (2007).
12. Hyvönen, R. *et al.* Impact of long-term nitrogen addition on carbon stocks in trees and soils in northern Europe. *Biogeochemistry* doi:10.1007/s10533-007-9121-3 (2007).
13. Sievering, H. Nitrogen deposition and carbon sequestration. *Nature* **400**, 629–630 (1999).
14. Sievering, H., Tomaszewski, T. & Torizzo, J. Canopy uptake of atmospheric N deposition at a conifer forest. Part I—Canopy N budget, photosynthetic efficiency and net ecosystem exchange. *Tellus* **59B**, 483–492 (2007).
15. Gaige, E. *et al.* Changes in canopy processes following whole-forest canopy nitrogen fertilization of a mature spruce-hemlock forest. *Ecosystems* **10**, 1133–1147 (2007).

doi:10.1038/nature06580