


Corrigendum: Rethinking 'secondary' metabolism: physiological roles for phenazine antibiotics

Alexa Price-Whelan, Lars E P Dietrich & Dianne K Newman

Nature Chemical Biology 2, 71–78 (2006); published online 18 January 2006; corrected after print 7 March 2006

In the version of this article initially published, some of the values in Table 1 were incorrect and three chemical names were misspelled. The correct table is below. The errors have been corrected in the PDF version of the article.



No	Name	R ₁	R ₂	R ₃	R ₄	K _{OW} ^{ox}	K _{OW} ^{red}	E ^{o'} (mV)
1	Aeruginosin A	COOH		CH ₃	NH ₂	-0.71	0.46	NA
2	Phenazine-1-carboxylic acid (PCA)	COOH				2.17	3.72	-177
3	Pyocyanin (PYO)	OH		CH ₃		1.60	2.89	-34
4	2-Hydroxyphenazine-1-carboxylic acid (2-OHPCA)	COOH	OH			2.54	3.32	NA
5	Phenazine-1-carboxamide (PCN)	CONH ₂				1.04	2.19	-115
6	1-Hydroxyphenazine (1-OHPHZ)	OH				1.81	2.35	-172

Table 1 Characteristics of some phenazines excreted by pseudomonads^{90–92}. Octanol-water partition coefficients (K_{OW}) were calculated using the KOWWIN demo program available at http://www.syrres.com/esc/est_kowdemo.htm. E^{o'}, standard redox potential, pH 7; NA, not available.

Erratum: Secramine inhibits Cdc42-dependent functions in cells and Cdc42 activation *in vitro*

Henry E Pelish, Jeffrey R Peterson, Susana B Salvarezza, Enrique Rodriguez-Boulan, Ji-Long Chen, Mark Stamnes, Eric Macia, Yan Feng, Matthew D Shair & Tomas Kirchhausen

Nature Chemical Biology 2, 39–46 (2006); published online 20 November 2005; corrected after print 7 March 2006.

In the version of this article initially published online, two labels in Figure 3a are incorrect. The label in the square below the arrow for step 2 should read 'Cdc42 GTP' instead of 'Cdc42 GDP', and 'P_i' in step 4 should be 'P_i⁻'. These errors have been corrected in the PDF version of the article. The corrected figure is below.

