

ARTICLE

Received 8 Jul 2014 | Accepted 11 Nov 2014 | Published 18 Dec 2014

DOI: 10.1038/ncomms6820

OPEN

Slow unloading leads to DNA-bound β_2 -sliding clamp accumulation in live *Escherichia coli* cells

M. Charl Moolman¹, Sriram Tiruvadi Krishnan¹, Jacob W.J. Kerssemakers¹, Aafke van den Berg¹, Pawel Tulinski¹, Martin Depken¹, Rodrigo Reyes-Lamothe², David J. Sherratt³ & Nynke H. Dekker¹

The ubiquitous sliding clamp facilitates processivity of the replicative polymerase and acts as a platform to recruit proteins involved in replication, recombination and repair. While the dynamics of the *E. coli* β_2 -sliding clamp have been characterized *in vitro*, its *in vivo* stoichiometry and dynamics remain unclear. To probe both β_2 -clamp dynamics and stoichiometry in live *E. coli* cells, we use custom-built microfluidics in combination with single-molecule fluorescence microscopy and photoactivated fluorescence microscopy. We quantify the recruitment, binding and turnover of β_2 -sliding clamps on DNA during replication. These quantitative *in vivo* results demonstrate that numerous β_2 -clamps in *E. coli* remain on the DNA behind the replication fork for a protracted period of time, allowing them to form a docking platform for other enzymes involved in DNA metabolism.

¹Department of Bionanoscience, Kavli Institute of Nanoscience, Faculty of Applied Sciences, Delft University of Technology, Lorentzweg 1, 2628 CJ Delft, The Netherlands. ²Department of Biology, McGill University, Montreal, Quebec, Canada H3G 0B1. ³Department of Biochemistry, University of Oxford, Oxford OX1 3QU, UK. Correspondence and requests for materials should be addressed to N.H.D. (email: N.H.Dekker@tudelft.nl).

The multi-protein replisome complex (replisome, Fig. 1a) is responsible for the accurate and timely duplication of the genome before cell division. The sliding clamp protein complex is a key subunit of the replisome and is vital for protein–DNA interactions related to DNA metabolism in all three domains of life^{1–3}. Through their interaction with polymerases, DNA ligase, replication initiation protein DnaA, the dynamine-like protein CrfC, as well as different mismatch-repair proteins, sliding clamps have important roles in replication and repair^{4–15}. In *E. coli*, the β_2 -sliding clamp (β_2 -clamp) is a homo-dimer¹⁶ (Fig. 1a inset) that encircles double-stranded DNA (dsDNA) and tethers DNA Polymerase III (DNA Pol III) to the template, thereby ensuring sufficiently high processivity during synthesis^{17,18}.

The β_2 -sliding clamp is actively assembled and disassembled onto DNA during the synthesis of the two complementary DNA strands (Fig. 1b). The loading reaction of a β_2 -clamp onto each new primer-template junction¹⁹ is catalysed by an ATP-dependent heteropentameric clamp-loader complex (clamp-loader), also known as the γ -complex²⁰. The clamp-loader pries open the β_2 -clamp, recognizes the primer-template junction²¹ and closes the β_2 -clamp around the dsDNA before release²². The clamp-loader is also thought to chaperone DNA Pol III onto a newly loaded β_2 -clamp²³ and to unload inactive DNA-bound β_2 -clamps via the δ -subunit²⁴. During all of these reactions, the loader complex and the various clamp binding proteins compete for the carboxy (C)-terminal face of the clamp. In accordance with the proposed model in which the replisome includes three core DNA polymerase IIIs^{25–27}, three β_2 -clamps can be active at the replication fork, one for each of the three polymerases (Fig. 1a). While leading-strand replication is thought to be continuous, utilizing only a single β_2 -clamp, the lagging-strand template is copied in discrete 1–2 kb Okazaki fragments²⁸, each utilizing a separate β_2 -clamp. These fragments are initiated by the continuous formation of 10–12 nt RNA primers by the primase (DnaG), which, together with the helicase (DnaB), sets the replication fork clock²⁹. Since the number of Okazaki fragments (2,000–4,000) for the 4.6 Mbp genome is roughly an order of magnitude greater than the average number of β_2 -clamps per cell in a nutrient-rich culture^{24,30}, continuous recycling of β_2 -clamps is necessary for total genome replication to occur.

Despite numerous *in vitro* and *in vivo* studies, it still remains unclear whether recycling of the *E. coli* β_2 -clamps takes place immediately following the completion of an Okazaki fragment, or at a later time. A slow recycling could permit a β_2 -clamp to fulfil additional functions, while remaining bound to the newly synthesized DNA. Quantitative *in vitro* unloading assays^{24,31} indicate that in the absence of the clamp-loader, a loaded β_2 -clamp has a long half-life of $t_{\text{unload}} > 1$ h on the DNA. Although this is decreased by more than an order of magnitude to $t_{\text{unload}} \sim 127$ s per β_2 in the presence of clamp-loader, this unloading time still remains long compared with the typical time required to complete an Okazaki fragment (on the order of seconds). Such a slow unloading time suggests that many β_2 -clamps are left behind in the wake of the replication fork³². However, a recent *in vitro* single-molecule study indicates that lagging-strand synthesis can persist *in vitro* in the absence of excess β_2 -clamps in solution, implying that a β_2 -clamp can be directly reused at a successive primer-template junction³³. Two *in vivo* studies, one in *Bacillus subtilis* (*B. subtilis*)³⁴ and the other in *E. coli*²⁶, provided contrasting results. Hence, to understand the regulatory mechanism that underlies the recycling of β_2 -clamps in *E. coli*, further insights into their *in vivo* dynamics are required.

To gain detailed insight into the *in vivo* recruitment and turnover of the β_2 -clamp, we investigate its dynamics in




Figure 1 | The *E. coli* replisome and β_2 -clamp assembly during replication.

(a) The position of the β_2 -sliding clamp within the *E. coli* replisome complex. The helicase (DnaB) unwinds dsDNA ahead of the replicative polymerase (DNA Pol III), which subsequently duplicates the template strands. Different configurations of Pol III are potentially possible. Primase (DnaG) synthesizes short RNA primers on the lagging strand for Okazaki fragment initiation. Single-strand binding proteins (SSB) remove the secondary structure of ssDNA and protect it from digestion. To ensure sufficient processivity during replication, Pol III is tethered to the DNA by the β_2 -sliding clamp. β_2 is assembled onto primer-template junctions by the multi-protein ($(\tau/\gamma)_3\delta\delta'\psi\chi$) clamp-loader complex. (Inset) A ribbon representation of the DNA-bound β_2 -sliding clamp (generated using the Protein Data Bank (PDB) file, 2POL¹⁶). The β_2 -sliding clamp is a homo-dimer that consists of six globular domains¹⁶. The monomers are arranged in a ring that encircles the DNA⁷⁰ and can slide freely along it. Different proteins can bind to the two hydrophobic pockets of the β_2 -clamp via a conserved sequence motif¹⁰. (b) The life cycle of the β_2 -clamp during replication. (top) The β_2 -clamp is actively loaded by the clamp-loader, which opens the closed clamp and places it onto dsDNA before release²². (middle) The β_2 -clamp remains DNA-bound as long as an Okazaki fragment is being synthesized. (bottom) After the β_2 -clamp has reached the end of an Okazaki fragment, DNA Pol III is signalled to release⁵². The β_2 -clamp is believed to be disassembled by the clamp-loader.

individual live cells with single-molecule sensitivity. We use both conventional fluorescence microscopy and Photoactivated Localization Microscopy (PALM)^{35,36}, in combination with custom-built microfluidics. Single-molecule techniques have provided us with insights into the dynamics of processes—such as replication, transcription and translation—that are not readily accessible with conventional ensemble-averaging techniques^{37,38}. *In vivo* single-molecule fluorescence imaging, in particular, has provided detailed insights into the behaviour of individual molecules in live cells^{39–41}. Combining single-molecule fluorescence microscopy with microfluidics allows us to image individual molecules in live cells over multiple cell cycles, without chemical fixation that could potentially perturb the dynamic behaviour of the protein under investigation⁴².

By using this experimental approach, we have measured the number of DNA-bound β_2 -clamps during chromosomal replication over the entire course of the cell cycle. In addition, we have determined the time required to unload an individual DNA-bound β_2 -clamp during replication, as well as the effective time required to load a new β_2 -clamp during replication. Our data reveal that the number of DNA-bound β_2 -clamps accumulates on the DNA after initiation, and then levels off to a constant steady-state number of DNA-bound β_2 -clamps on the order of minutes. This steady state is maintained throughout the rest of the replication process, until termination occurs and a concomitant decrease of DNA-bound β_2 -clamps is observed. The number of DNA-bound β_2 -clamps in steady state exceeds the estimates of a previously published *in vivo* study²⁶ by an order of magnitude. The measured values for the effective loading time and unloading time during replication, in the context of the live cell, are in good agreement with previous biochemical *in vitro* experiments²⁴. Taken together, our data indicate that a β_2 -clamp remains on the DNA for a protracted period of time following the completion of an Okazaki fragment. DNA-bound β_2 -clamps that are left behind during fork progression may facilitate the recruitment of

additional proteins active during the cell cycle for different processes such as DNA repair.

Results

The *in vivo* dynamics of β_2 -clamps measured in single cells. To study the dynamics of β_2 -clamps by wide-field fluorescence microscopy, we perform long time-lapse imaging of labelled β_2 -clamps over multiple replication cycles. During such experiments, we ensure healthy cell physiology by implementing a custom-built microfluidic device (Fig. 2a; see Methods)^{43,44} in which cells growing in steady state are immobilized in micron-sized growth channels. Through a neighbouring central trench, growth medium is continuously supplied throughout an experiment (see Methods). In such a microfluidic device, cells experience minimal perturbation over the course of the time-lapse experiment, as stable growth conditions remain continuously present. This contrasts with long time-lapse




Figure 2 | Long time-lapse fluorescence microscopy of the β_2 -sliding clamp at the single-cell level utilizing microfluidics. (a) The microfluidic device used for performing long time-lapse fluorescence microscopy. *E. coli* cells are immobilized in growth channels perpendicular to a main trench through which growth medium is actively pumped. (inset) A brightfield image and corresponding YPet- β_2 fluorescence image (80 ms laser light exposure) are acquired every 2.5 min for the duration of the time-lapse experiment. Scale bar, 3 μm . (b) YPet- β_2 molecules that are either DNA-bound or freely diffusing are studied using wide-field fluorescence microscopy. (left) Freely diffusing YPet- β_2 molecules in the cytoplasm of a cell. This signal is representative to YPet- β_2 dynamics before and after replication. (middle) A clear focus is observed due to DNA-bound YPet- β_2 molecules. The observation of a single focus, instead of two distinct foci, shortly after initiation results from the overlap of diffraction limited spots. (right) Two distinct foci are visible, indicative of two individual replisomes. Scale bar, 800 nm. (c) Kymograph of a single growth channel during an overnight time-lapse experiment. The cells first grow the growth channel full, and maintain a steady state growth rate as can be observed from the curved shape of the fluorescence signal. The shape of the fluorescence signal is due to the individual cells growing and pushing each other in the direction of the main trench. Clear observable diffuse patterns occur at regular intervals, indicative of no DNA-bound β_2 -clamps before initiation or after termination. This repeating pattern is due to the multiple cycles of replication (indicated with repeating white dashed lines). (d) A kymograph of an individual replication cycle indicated in c. The blue lines are the cell boundaries detected from the brightfield images. The illustrations on the right-hand side indicate the different stages of replication that can be observed during the cell cycle.

experiments performed on agarose pads in which nutrients and water may become depleted, leading to non-steady state cell populations as a result. Additional benefits of such a device are that daughter cells ultimately grow out of the growth channels, preventing the accumulation of cells, and that the cells are always aligned, which facilitates data analysis.

Labelling of the β_2 -clamp was accomplished by using a functional amino (N)-terminal YPet⁴⁵ fusion²⁶ (Supplementary Fig. 1; Supplementary Table 1; see Methods) expressed from (and replacing) the endogenous *E. coli dnaN* gene locus. Fluorescence images are acquired under shuttered 515 nm laser excitation (see Methods; Fig. 2a (inset)). Fluorescence images of YPet- β_2 within individual cells either yielded no (Fig. 2b, left), a single (Fig. 2b, middle) or two cellular foci (Fig. 2b, right), depending on the stage of replication, in agreement with previous reports of fluorescently labelled replisome components⁴⁶. Before each fluorescence image, a brightfield image is acquired to provide details of the cell periphery (Fig. 2a (inset)). This alternating imaging sequence has a sufficiently long period to avoid giving rise to any notable deleterious growth effects, as assessed by comparing the doubling time of cell growth in a shake flask with cells grown in the microfluidic device (Supplementary Figs 1 and 2).

Using this approach, we are able to observe numerous consecutive replication (and corresponding division) cycles of cells in the different growth channels. We examine the global replication dynamics of multiple cells within a growth channel by converting the time-lapse images into a kymograph (Fig. 2c; see Methods). A distinct reoccurring pattern indicative of multiple replication cycles in the generations of cells is clearly noticeable (indicated by repeating dashed lines in Fig. 2c). Under these experimental conditions (see Methods), the analysis of individual cells ($n = 137$) in our microfluidic system yields an average replication time of $t_{\text{rep}} = 68 \pm 10$ min, and a doubling time of $t_{\text{double}} = 84 \pm 17$ min (Supplementary Fig. 2). In both cases, the error is \pm s.d. We further analyse these kymographs to investigate the subcellular dynamics of the YPet- β_2 molecules within the individual cells from cell birth till cell division (Fig. 2d). One can clearly observe the dynamics of the two β_2 -clamp foci associated with the two independent replisomes.

The assembly and accumulation of β_2 -clamps on DNA. We use the fluorescence intensity from the YPet- β_2 fusion to determine the number of β_2 -clamps that are DNA-bound as well as in the total number in the cell during the life cycle of a cell. A sample montage of the YPet- β_2 fluorescence signal from cell birth till after cell division (Fig. 3a) illustrates that there is a distinct increase in the foci following the B-period⁴⁷ of the cell cycle (represented by a diffuse signal after birth) (Fig. 3a (inset)) and a similar decrease before cell division. The fraction of fluorescence that originates from DNA-bound YPet- β_2 (foci) provides clear evidence that $>50\%$ of β_2 -clamps are DNA-bound shortly after the initiation of replication (Fig. 3b). The steady decline in the fraction of DNA-bound β_2 -clamps that commences roughly 10 min after initiation, results from the increase of total number of β_2 -clamps in the cell as the cell grows. In assessing this intensity fraction, we verified that very little out-of-focus fluorescence escapes detection (Supplementary Note 1; Supplementary Fig. 3).

A constant number of DNA-bound β_2 -clamps is maintained.

To precisely quantify the number of DNA-bound β_2 -clamps as a function of the replication cycle, we exploited a single-molecule *in vitro* calibration method²⁶ that allows us to reliably convert the detected YPet- β_2 signal into an absolute number of molecules

(Supplementary Note 2). We immobilize single purified YPet molecules on a cover glass and determine the average intensity of a single YPet fluorescent protein under these conditions (Supplementary Fig. 4). Using this calibration, we perform control stoichiometry experiments of previously studied DNA-bound replisome components²⁶, specifically the ϵ -subunit of Pol III and the τ -subunit of the clamp-loader to verify that our *in vitro* single-molecule calibration remains reliable *in vivo*⁴⁸ (Supplementary Fig. 5). For both the proteins, we reproduced the stoichiometry for the pair of sister replisomes as previously published²⁶, namely 5.74 ± 0.04 molecules in total for the ϵ -subunit ($n = 64$) and 6.12 ± 0.03 molecules in total for the τ -subunit ($n = 66$). Here the error is \pm s.e.m. We also verified that a YPet- β intensity standard provides the same mean intensity value under our experimental conditions (Supplementary Fig. 4g). Therefore, we subsequently use this average intensity value to estimate the number of β_2 -clamps in our experiments. In calculating the number of β_2 -molecules, we correct for photobleaching (Supplementary Note 3; Supplementary Fig. 3) and verify that the fraction of immature, dark YPet proteins is negligible (Supplementary Note 4, Supplementary Fig. 3). In our conversion from intensity to molecules, we also take into account that β_2 -clamps are dimers by dividing the measured YPet signal by two. This is a realistic assumption as it is believed that β_2 -clamps are in closed conformation even when they are not DNA-bound²².

Using this calibration standard, we quantify the absolute number of DNA-bound YPet- β_2 molecules for individual traces of DNA replication. Representative individual time-traces of single cells clearly demonstrate that following the initiation of replication, a gradual increase of the number of DNA-bound β_2 occurs until a steady state plateau is reached (Fig. 3c). This plateau is maintained until decrease is observed shortly after or during termination (Fig. 3c). From the individual traces, one can observe that there is significant cell-to-cell variability in the absolute number of DNA-bound β_2 -clamps, but that the overall trend in which the number of DNA-bound β_2 -clamps is constant for a significant fraction of the cell cycle is the same for all cells. We compared this temporal behaviour with that of a different replisome component, the τ -subunit of the clamp-loader in a strain in which both the β_2 -clamp and the τ -subunit are fluorescently labelled (Supplementary Note 5). The τ -YPet fluorescence signal fluctuates strongly and does not yield a stable plateau, in contrast to the mCherry- β_2 fluorescent signal (Supplementary Fig. 6).

To obtain statistically significant values for both the total number of β_2 -clamps in the cell and the number of DNA-bound β_2 -clamps, we extracted the average behaviour from analysis over numerous cells ($n = 137$; Fig. 3d,e). Figure 3d clearly depicts that for an average cell, the fraction of DNA-bound β_2 -clamps is more than half of the total content in the cell, which decreases to roughly zero after termination. During the cell cycle, an average cell doubles its YPet- β_2 content from ~ 60 to 120 molecules. This number of β_2 -clamps in the cell is in good agreement with ensemble western estimates we performed under the same growth conditions (Supplementary Note 6; Supplementary Fig. 7). Remarkably, the number of DNA-bound YPet- β_2 is held at a stable value of $N_{\beta_2} = 46$ (s.d. = 12, s.e.m. = 1; Fig. 3e (inset)). We also observe that the number of DNA-bound β_2 -clamps are close to zero before initiation and after termination. We ruled out the presence of an ectopic *dnaN* gene, by verifying that only a single copy of the *dnaN* gene is present in the strain that we used for these experiments (Supplementary Note 7; Supplementary Fig. 7). The experiment was successfully reproduced with a different fluorescent protein fusion (mCherry- β_2), which strengthens the argument that accumulation is unlikely the result of fluorophore




Figure 3 | Quantification of the *in vivo* β_2 -sliding clamp stoichiometry during replication. (a) A representative temporal montage of the YPet- β_2 fluorescence signal from before initiation until after cell division. A clear intensity increase is observed at the focus formation following initiation (indicated with white arrows in inset) Scale bar, 1.6 μm . (b,c) Traces of focus formation in individual cells. In (b), we plot the fraction of YPet- β_2 molecules that are DNA-bound compared with the total number in the cell. More than 50% of the total YPet- β_2 molecules are DNA-bound. The gradual decline in this fraction results from the increase of β_2 during the cell cycle. The inset indicates how the DNA-bound YPet- β_2 molecules and the total YPet- β_2 in the cell are defined. In (c), we plot the absolute number of DNA-bound YPet- β_2 molecules. Here, the gradual increase, steady state and gradual decrease of the DNA-bound YPet- β_2 molecules can clearly be seen. In both (b) and (c), the traces have been aligned with respect to initiation. (d,e) The average behaviour of individual YPet- β_2 molecules measured in individual cells. (d) The fraction of DNA-bound YPet- β_2 molecules is on average >50% half way through the replication cycle. (e) The YPet- β_2 molecules in the whole cell (blue curve) approximately doubles during the cell cycle, from 60 to 120 YPet- β_2 molecules. The DNA-bound YPet- β_2 molecules (red curve) remarkably increases to a mean steady state value of 46 YPet- β_2 molecules (s.d. = 12, s.e.m. = 1) following initiation. This value is maintained throughout the replication process until a concomitant decrease is observed after or during termination. Individual traces have been normalized with respect to initiation and termination to make averaging possible. (inset) A histogram of the distribution of number of DNA-bound YPet- β_2 molecules during steady state. ($n = 137$).

aggregation^{49,50}, but rather due to physiological build-up of DNA-bound clamps (Supplementary Note 8; Supplementary Fig. 8). The slightly lower mean number of DNA-bound clamps ($N_{\beta_2} = 34$, s.d. = 12, s.e.m. = 1.5) as measured using the mCherry- β_2 protein fusion, in combination with the mCherry intensity calibration, possibly results from the less ideal photophysical properties of mCherry, which make it less suitable than YPet for rigorous quantitative fluorescence microscopy.

Single β_2 -clamps are not rapidly unloaded *in vivo*. To study the *in vivo* unloading time of an individual β_2 -clamp, we utilized single-molecule PALM (Fig. 4a). The endogenous *dnaN* gene was replaced with a functional N-terminal PAMCherry⁵¹ fusion (see Methods). Fluorescence images are acquired under shuttered 561 nm excitation (see Methods), while activation is performed once with low 405 nm laser illumination, such that on average less than one DNA-bound PAMCherry- β per cell is activated.




Figure 4 | Direct measurement of the *in vivo* unloading time of the β_2 -sliding clamp during replication. (a) Illustration of the measurement sequence to image a single β_2 -clamp unloading event. First a phase-contrast (PH) and pre-activation snapshot are taken, after which molecules are activated only once, and subsequently imaged until foci are no longer visible (b,c) Single PAmCherry- β molecules are visualized by PALM. The sample PH image together with the respective pre-activation and post-activation fluorescence images illustrate that a single PAmCherry- β molecule can successfully be photoactivated. The corresponding pre-activation (red) and post-activation (blue) line profile plots of the single DNA-bound PAmCherry- β molecule. Scale bar, 1.6 μm . (d) A representative example of a montage showing the fluorescence intensity of a PAmCherry- β molecule over time and the corresponding intensity trace of the signal. The single-step disappearance is indicative of a single molecule. (e) On-time distribution for individual PAmCherry- β molecules ($n = 84$) fitted with an exponential (red line), and the distribution for the unloading times corrected for photobleaching (dashed green line). The inset shows the distribution of the fitted unloading time constants over the 10^6 bootstrapped data sets from which the confidence interval for the unloading time is determined.

Before fluorescence activation, a phase-contrast (PH) image is acquired to determine the cell's position and its periphery. Sample pre- and post-activation images (Fig. 4b), together with the corresponding line-profile intensity plots (Fig. 4c), demonstrate successful activation of individual DNA-bound PAmCherry- β molecules in our strain. The advantage of PALM over more conventional techniques like Fluorescence Recovery After Photobleaching and Fluorescence Loss In Photobleaching for measuring protein turnover is that it allows one to directly image a single unloading event, as shown in the sample temporal montage and the corresponding integrated intensity trace (Fig. 4d). We image a different field of view of cells

for each complete PALM measurement sequence (Fig. 4a) to ensure that the cell physiology and β_2 -clamp behaviour are not influenced by excessive 405 nm light exposure. Using the individual analysed traces from different cells, we are able to build up a distribution for the on-time events of single PAmCherry- β molecules (Fig. 4d). To visualize a single unloading event, we only image one out of the two PAmCherry- β_2 dimer subunits. After correcting for photobleaching (Supplementary Note 9; Supplementary Fig. 9), we estimate the *in vivo* unloading time to be $t_{\text{unload}} = 195 \pm 58$ s per β_2 (Fig. 4e). This result is in good agreement with previous *in vitro* experiments (127 s per β_2 ; ref. 24).

The effective *in vivo* loading rate of β_2 -clamps. The *in vivo* loading time of a β_2 -clamp during chromosomal replication provides us with insight into how frequently a new clamp, β_2 , is needed for processive genome duplication. We utilize both the long time lapse and the single-molecule PALM data to compute the effective loading time *in vivo* of a new β_2 -clamp ($t_{\text{load}}^{\text{eff}}$). The designation ‘effective’ is added as we do not directly measure the loading of an individual β_2 -clamp, but rather the total loading rate of β_2 -clamps onto DNA. We have shown in the preceding section that the number of DNA-bound β_2 -clamps remains essentially constant ($N_{\beta_2} = 46$) during 2/3 of the replication process. We independently determined the *in vivo* unloading time via PALM to be $t_{\text{unload}} = 195$ s per β_2 during the replication. In the steady-state regime, the total unloading rate of β_2 -clamps is balanced by the effective loading rate of β_2 -clamps ($t_{\text{load}}^{\text{eff}}$) onto newly formed primer for Okazaki fragment synthesis:

$$\frac{1}{t_{\text{load}}^{\text{eff}}} = N_{\beta_2} \cdot \frac{1}{t_{\text{unload}}} \quad (1)$$

Using equation (1), we compute the *in vivo* effective loading time for a β_2 -clamp during the replication to be $t_{\text{load}}^{\text{eff}} = 4 \pm 1$ s per β_2 . The reader is referred to Supplementary Note 10 and Supplementary Fig. 10 for a more detailed discussion of equation (1).

Discussion

DNA replication, orchestrated by the multi-protein replisome-complex, is a process essential to cell viability. By using *in vivo* single-molecule fluorescence microscopy in combination with microfluidics, we were able to investigate the detailed dynamics of an essential component of the replisome, the β_2 -clamp, during DNA replication in live *E. coli* cells. Lagging-strand synthesis is a complex and highly dynamic process, and the sliding clamp is one of the key proteins involved. Each new primer-template junction requires a loaded β_2 -clamp to ensure processive replication by DNA Pol III, which is signalled to cycle from one Okazaki fragment to the next⁵² as the replication fork progresses at ~ 600 bp s⁻¹. Given the average replication fork speed and the typical size of an Okazaki fragment (1–2 kb), one would expect a β_2 -clamp to be necessary every ~ 1.5 –3 s. Leading-strand synthesis might be less processive than commonly believed, which would imply that a new β_2 -clamp would also need to be loaded on the leading strand. In what follows, however, we have assumed that during normal replication, the exchange of β_2 -clamps on the leading strand is a much less frequent occurrence than β_2 -clamp exchange for the lagging strand. Until now, it was not demonstrated *in vivo* whether these loaded β_2 -clamps are predominantly recycled (that is, immediately unloaded and reloaded) between successive Okazaki fragments, or whether β_2 -clamps remain bound to a completed fragment for a prolonged period of time.

Our results indicate that the number of DNA-bound β_2 -sliding clamps increases during the course of the cell cycle, peaking at more than 20 behind an individual fork. Following the initiation of replication, we observe that the number of DNA-bound β_2 -clamps gradually increases until a steady-state plateau is reached. This plateau, whose magnitude is such that $\sim 50\%$ of the total β_2 -clamps in the cell are DNA-bound, is maintained throughout the remainder of the cell cycle. We determined the number of β_2 -clamps in the cell (60–120) during replication, as well as the total number of DNA-bound β_2 -clamps ($N_{\beta_2} = 46$). After termination, β_2 -clamps are presumably no longer being loaded, and the fraction of DNA-bound β_2 -clamps decays accordingly.

Notably, our measurements for the number of DNA-bound β_2 -clamps differ from the value measured previously by some of us in a comprehensive *in vivo* study of the whole *E. coli* replisome

complex²⁶. In this study, the number of β_2 -clamps was estimated to be three for each of the two independent replisomes⁴⁶, for a total of six DNA-bound β_2 -clamps present during replication. We note that stoichiometries for most of the other proteins have been duplicated independently, and, therefore, the difference of the number of DNA-bound β_2 -clamps appears an isolated case²⁷. Although we cannot fully explain the difference in the number of DNA-bound β_2 -clamps, we can nonetheless delineate some possible contributions. The difference may result from the cell physiology due to the immobilization method, lower statistics due to the challenging nature of the ‘slim-field’ experiments at the time, or due to inadvertent changes of the imaging system since measurements spanned across months in the earlier study. It is thus crucial to maintain healthy cell physiology and cell cycle synchronization during experiments, which highlights the utility of microfluidics in live cell single-molecule fluorescence measurements.

The substantial number of DNA-bound β_2 -clamps behind each replication fork suggests that β_2 -clamps are not rapidly recycled during replication. To corroborate this view, we have utilized PALM to directly measure the *in vivo* unloading rate of a single β_2 -clamp ($t_{\text{unload}} = 195$ s per β_2). Together with the number of DNA-bound β_2 -clamps in steady state, this allows us to calculate the effective time of loading (equation (1)) a β_2 -clamp during replication as $t_{\text{load}}^{\text{eff}} = 4$ s per β_2 . This result is in good agreement with our previously calculated average estimate of the primer formation time using the Okazaki fragment size range and the typical size of the *E. coli* genome. Also, this effective loading rate is in accordance with the model that DnaG sets the fork speed²⁹. DnaG is thought to synthesize RNA primers at a rate of approximately one primer every one to two seconds⁵³, which is in good agreement with our calculated *in vivo* effective loading time. We suggest that an individual β_2 -clamp remains on the DNA for a protracted period of time during chromosomal replication, as has been proposed on this basis of *in vitro* experiments^{54,24} and plasmid replication⁵⁵. Our results are in agreement with the behaviour of the sliding clamp for the Gram-positive bacterium *B. subtilis*³⁴. In this bacterium, the number of DNA-bound β_2 -clamps was estimated at ~ 200 during replication, indicative of clamps being left behind during fork progression. There is a slight possibility that the loading and unloading reaction could be sterically hampered by the fusion protein. However, we have no reason to believe that this is the case since our results are in very good agreement with previous *in vitro*²⁴ and *in vivo* work³⁴. Our study shows that rapid recycling of β_2 -clamps for subsequent lagging-strand synthesis³³, though observed in *in vitro* experiments in the absence of excess β_2 -clamps in solution, is not the predominant mode *in vivo*. Although our data does not exclude that β_2 -clamps are rapidly recycled at the replication fork, the fact that the loading rate from solution matches the estimated primer formation rate strongly suggests that direct recycling is not the dominant mode of clamp loading.

To illustrate the overall β_2 -clamp dynamics during replication, we perform a Monte Carlo simulation (see Methods) that takes our experimentally determined values for N_{β_2} , t_{unload} , $t_{\text{load}}^{\text{eff}}$ and t_{rep} as input parameters (Fig. 5a). As the approximate rate of clamp removal during termination (Fig. 3e) agrees with the value measured by PALM during steady-state replication (~ 195 s per β_2), we simply input the latter (likely more accurate) value into the simulations. The simulation starts at $t = 0$ with no DNA-bound β_2 -clamps, after which it takes < 10 min to reach a stable steady-state number of β_2 bound to DNA (Fig. 5b, left). This value is maintained for ~ 60 min (Fig. 5b, middle), after which termination occurs and the clamps are unloaded in < 5 min (Fig. 5b, right). The number of β_2 -clamps in steady state as well as the rise and fall times underline our measured results, and are depicted schematically in Fig. 5c.




Figure 5 | Describing the β_2 -sliding clamp recycling process during replication. (a) A Monte Carlo simulation of the β_2 -clamp assembly and disassembly reaction. For illustrative purposes, we perform a Monte Carlo simulation of the proposed model, utilizing the experimentally determined primer formation and unloading rate, as well as the replication time, and under the assumption that primer formation is rate limiting. We show the simulation results for five individual traces (coloured lines). The black curve is the analytical solution for the average number of loaded β_2 -clamps. Here we divide the total trace into three time regions, namely initiation (red), steady state (green) and termination (blue). (b) A zoom of the different sections from a. (left) A build-up of loaded β_2 -clamps on the DNA proceeds for ~ 10 min. (middle) After the gradual accumulation of loaded β_2 -clamps, a steady state plateau of 46 DNA-bound β_2 -clamps is maintained for $\sim 2/3$ of the replication process. (right) After termination, all DNA-bound β_2 -clamps are unloaded in ~ 5 min. (c) A cartoon illustrating the DNA-bound β_2 -clamp build-up during replication. As the rate at which β_2 -clamps are loaded (one every 4 s) is much faster than the unloading rate of individual β_2 -clamps (once every 195 s) during replication, there will be a dynamic reservoir of β_2 -clamps that have not yet been unloaded left on the lagging strand.

The steady-state build-up of DNA-bound β_2 -sliding clamps forms a β_2 -landing pad³⁴ for different proteins to dock themselves to DNA during the life cycle of the cell. Numerous different proteins utilize the β_2 -clamp via the same binding pocket⁵⁶ to perform their respective biological function. These other β_2 -clamp-binding proteins range from DNA ligase for Okazaki fragment maturation, inactivation of DnaA through the β_2 -Hda₂ interaction^{11–13,57,58}, potential screening of DNA damage due to sliding capability of the β_2 -clamps⁵⁶, the tethering of the necessary polymerases for repair^{4–8}, overcoming replication barriers⁵⁹, as well as coupling mismatch-detection and replication by positioning MutS at newly replicated DNA⁶⁰. It is still unclear which of the above-mentioned (or other) proteins are the main users of the DNA-bound clamps that are not directly situated at

the replication fork. As Okazaki fragment maturation seems to be relatively fast as assessed via Ligase and Pol I dynamics⁶¹, it is likely not these proteins that predominantly occupy the DNA-bound clamps. The extent to which DNA-bound β_2 -clamps are utilized while being docked to the Okazaki fragment will most likely be dependent on the physiological state of the cell at a particular time in its cell cycle. Under stress conditions, MutS and the different repair polymerases might predominantly make use of DNA-bound β_2 -clamps, whereas under minimal stress conditions, ligase CrfC and Hda₂ are the likely candidates. A thorough *in vivo* investigation of the stoichiometry and dynamic of different β_2 -associated proteins over the course of the cell cycle would provide the quantitative underpinning required to provide further insight into these biological processes.

Methods

Strains and strain construction. All strains are derivatives of *E. coli* K12 AB1157. Strains were constructed either by P1 transduction⁶² or by λ -Red recombination⁶³.

The *YPet-dnaN:tetR-mCerulean* was constructed using P1 transduction by transducing the *YPet-dnaN* fusion²⁶ together with the adjacent *kanR* gene into a strain that contains a *tetO* array (50 kb clockwise from the *dit*-site), as well as the chromosomal integrated chimeric gene *tetR-mCerulean*⁴⁶. The presence of the *YPet-dnaN* gene fusion was verified using the oligonucleotides: 5'-CGTTGGCA CCTACCAGAAAG-3' and 5'-ATGCCTGCCGTAAGATCGAG-3'. The sequence of the *YPet-dnaN* fusion was confirmed by DNA sequencing.

A chromosomal fusion of the gene encoding for the photoactivatable fluorescent protein (PAmCherry1)⁵¹ to the N terminus of the *dnaN* gene was created using λ -Red recombination⁶³. The gene encoding for PAmCherry1 was amplified by PCR. The forward primer used contains an *XmaI* restriction site (5'-CGGGCCCCGGGATGGTAGCAAGGGCGAGGAG-3'). The reverse primer used contains a sequence coding for an 11 amino-acid linker and a *SacI* site (5'-CGATCGGAGCTCCGCGCTGCCAGAACCCGCGGCGGAGCCTGCC GACTTGTACAGCTCGTCCATGCC-3'). The PCR product was cloned into the backbone of pROD44 (ref. 26) containing a kanamycin-resistance cassette, flanked by *frt* sites, resulting in the template plasmid PAmCherry1.

This plasmid was then used as a template plasmid for generating the insert sequence used during λ -Red recombination to create the *PAmCherry-dnaN* strain. The primer sequences used were: forward 5'-ACGATATCAAAGAAGATTTTTC AATTTAATCAGAACATTGTCTATCGTAACTGTAGGCTGGAGCTGCTTC-3'; reverse 5'-ACCTGTTGTAGCGGTTTTAATAAATGCTCAGCTTCTACGGTAA ATTTATCGCGCTGCCAGAACCCGCGG-3'. The DNA fragment was gel purified and ~700 ng of the linear DNA was used for electroporation of AB1157 cells overexpressing λ -Red proteins from pKD46 (ref. 63). The correct insertion of the fragment into the chromosome of the resulting strain was assayed by PCR. The oligonucleotides used were 5'-CGTTGGCACCTACCAGAAAG-3' and 5'-ATGC CTGCCGTAAGATCGAG-3'. The sequence of the fusion gene in this strain was confirmed by DNA sequencing.

Construction of the *mCherry-dnaN* strain. The *mCherry* gene was amplified by PCR. The forward primer used for this contains an *XmaI* restriction site (5'-TAG GTCCTCCGGGATGAGCAAGGGCGAGGATAAC-3'). The reverse primer used contains a *SacI* site and sequence coding for an 11 amino-acid linker (5'-AAGG AGCTCGGCTGCCAGAACCCGCGGCGGAGCCTGCCGACTGTACAGCTC GTCCATGCC-3'). The *Frt*-flanked kanamycin-resistant gene was amplified using the following primers: forward 5'-TTACCCGGGCATATGAATATCCTCT TAG-3'; reverse 5'-TTAGGATCTGTAGGCTGGAGCTGCTTCG-3'. The resulting fragment was digested with *XmaI* and *BamHI*. The *mCherry* fragment and the kanamycin fragment were cloned into pUC18 between *SacI* and *BamHI* sites.

The λ -Red recombination was performed as mentioned in the previous section using the primers: forward 5'-TATCAAAGAAGATTTTCAAATTTAATCAG AACATTGTCTATCGTAACTGTAGGCTGGAGCTGCTTCG-3'; reverse 5'-AC CTGTTGTAGCGGTTTTAATAAATGCTCAGCTTCTACGGTAAATTTATC GCGCTGCCAGAACCCGCGG-3'. The presence of the gene fusion was verified using oligonucleotides 5'-CGTTGGCACCTACCAGAAAG-3' and 5'-ATGCCTGCCGT AAGATCGAG-3'. The sequence of the fusion gene in this strain was confirmed by DNA sequencing.

The *dnaX(t)-YPet:mCherry-dnaN* strain was constructed using P1 transduction by transducing the *dnaX-YPet* fusion²⁶ together with the adjacent *kanR* gene into a strain that contains the *mCherry-dnaN* gene fusion. The presence of the *dnaX-YPet* gene fusion after transduction was verified using the oligonucleotides: 5'-GAGCCTGCCAATGAGTTATC-3' and 5'-GGCTTGCTTCATCA GGTAC-3' and similarly the *mCherry-dnaN* fusion using 5'-CGTTGGCACCT ACCAGAAAG-3' and 5'-ATGCCTGCCGTAAGATCGAG-3'. The sequences of the fusions in this strain were confirmed by DNA sequencing.

Supplementary Tables 2 and 3 provide an overview of the plasmids used, as well as a summary of the different strains. The cell morphology and the doubling times of the fusion strains in LB and M9-glycerol growth medium were compared with AB1157 wild type. No significant differences were observed (Supplementary Table 1 and Supplementary Fig. 1a). The doubling times of the cells in the microfluidic device were similar (slightly faster) compared with cells grown in a shake flask (Supplementary Fig. 2). We also confirmed that in the absence of IPTG (the experimental condition used during long time-lapse microscopy), no DNA-bound foci were detected for the *YPet-dnaN:tetR-mCerulean* strain (Supplementary Fig. 1b).

M9 growth medium used in experiments. The M9 growth medium used in experiments is as follows. One litre of M9 growth medium contains 10.5 g l⁻¹ of autoclaved M9 broth (Sigma-Aldrich); 0.1 mM of autoclaved CaCl₂ (Sigma-Aldrich); 0.1 mM of autoclaved MgSO₄ (J.T.Baker); 0.3% of filter-sterilized glycerol (Sigma-Aldrich) as carbon source; 0.1 g l⁻¹ of filter-sterilized five amino acids, namely L-threonine, L-leucine, L-proline, L-histidine and L-arginine (all from Sigma-Aldrich) and 10 μ l of 0.5% filter-sterilized thiamine (Sigma-Aldrich).

Microfluidics for extended time-lapse microscopy. We use our own design⁴³ of the previously published microfluidic device known as the mother machine⁴⁴ for cell immobilization during long time-lapse experiments. The reader is referred to

Moolman *et al.*⁴³ for a detailed description of the complete fabrication process. Here, we only briefly outline the main steps involved. First, we use electron-beam lithography in combination with dry etching techniques to create the structure in silicon. Next, we make a negative mould of this structure in polydimethylsiloxane (PDMS). The PDMS mould is then used to fabricate the positive structure in PDMS, which is subsequently used for experiments.

Preparation of cells for microscopy. Cells were streaked on Luria-Bertani (LB) plates containing the appropriate antibiotic. Single colonies from these plates were inoculated overnight at 37 °C with shaking in M9 medium supplemented with 0.3% glycerol (Gly), essential nutrients together with the appropriate antibiotics. The subsequent day, the overnight culture was subcultured into the same medium and grown at 37 °C with shaking until an OD₆₀₀ ~ 0.2 was reached. Cells were concentrated by centrifugation for 2 min at 16,100 g. The subsequent steps are dependent on the type of microscopy experiment performed as outlined next.

For agarose pad experiments, the supernatant was decanted and the pellet was resuspended in 100 μ l M9-Gly supplemented with essential nutrients. The resuspended cells were subsequently vortexed for 2 s and immobilized on an M9-Gly 1.5% agarose pad between two coverslips. (The coverslips were ultrasonically cleaned in acetone and isopropyl alcohol and burned by a flame to minimize the fluorescent background before use).

For microfluidic device experiments, the supernatant was decanted and the pellet was resuspended in 50 μ l M9-Gly with essential nutrients and injected into the microfluidic device. After injection into the device, the device was centrifuged for 10 min at 2,500 g (Eppendorf 5810R) so as to load the cells into the growth channels. Following centrifugation, the device was mounted on the microscope with tubing attached and incubated for ~45 min at 37 °C. After incubation, fresh M9-Gly with essential nutrients and the appropriate antibiotics are flushed through the device. The syringe containing the medium is then attached to an automated syringe pump to continuously infuse fresh M9-Gly, essential nutrients and 0.2 mg ml⁻¹ bovine serum albumin (BSA) through the device at a rate of 0.5 ml h⁻¹.

Microscope setup. All the images were acquired on a commercial Nikon Ti microscope equipped with a Nikon CFI Apo TIRF \times 100, 1.49NA oil immersion objective and an Andor iXon 897 Electron Multiplying Charge Coupled Device (EMCCD) camera operated by a personal computer (PC) running Nikon NIS-elements software. Cell outlines were imaged using the standard Nikon brightfield halogen lamp and condenser components. The fluorescence excitation was performed using custom-built laser illumination. A Cobolt Fandango 515 nm continuous wave diode-pumped solid-state laser was used to excite YPet; Cobolt Jive 561 nm continuous wave diode-pumped solid-state laser was used to excite mCherry and PAmCherry, respectively. PAmCherry was activated by a Votran Stradus 405 nm. All the three laser beams were combined using dichroic mirrors (Chroma ZT405sp-xxr, 575dcspxr) and subsequently coupled into a single-mode optical fibre (KineFLEX). The output of the fibre was expanded and focused onto the back focal plane of the objective mounted on the microscope. Notch filters (Semrock NF03-405E, NF03-514E, NF03-561E) were used to eliminate any laser light leaking onto the camera. The emission of the different fluorescent proteins was projected onto the central part of the EMCCD camera using custom filter sets: Chroma z561, ET605/52m, zt561rdc (mCherry), Chroma z514, ET540/30m, zt514rd (YPet), Chroma zet405, ET480/40m, zt405rdc (CFP). A custom design commercial temperature control housing (Okolabs) enclosing the microscope body maintained the temperature at 37 °C. Sample position was controlled with a Nikon stage (TI-S-ER Motorized Stage Encoded, MEC56100) together with the Nikon Perfect Focus System to eliminate Z-drift during image acquisition.

Cell lysate preparation for intensity calibration. The cell lysate used for single-molecule intensity calibration was prepared as follows. Cells were grown overnight at 37 °C with shaking in M9 medium supplemented with 0.3% glycerol (Gly), essential nutrients together with the appropriate antibiotics. The subsequent day, the overnight culture was subcultured into the same medium and grown at 37 °C with shaking until an OD₆₀₀ ~ 0.5 was reached. The cells were collected by centrifugation at 6,000 g (Beckman JLA 9.1000 rotor) for 15 min. Cells were subsequently resuspended in 5 ml M9-Gly and essential nutrients. The cell suspension was French pressed (Constant Systems) twice at 20,000 p.s.i. The cell lysate was then spun down at 30,000 g (Beckman JA-17 rotor) for 35 min. The supernatant was shock-frozen using liquid nitrogen and kept at 37 °C until needed.

Data acquisition. All data acquisition was performed on the same microscope setup. Image acquisition was performed with Nikon NIS-elements software. The acquisition protocol was dependent on the type of experiment performed as outlined next.

Long time-lapse experiments were conducted as follows. The cell outlines were imaged using standard brightfield illumination. Subsequently, the sample was excited by laser excitation (515 nm) with an intensity of ~ 5 W cm⁻² as calculated according to Grünwald *et al.*⁶⁴ The exposure time was set to 80 ms. The camera gain was set to 100. Brightfield and fluorescence images were acquired every 2.5 min. Data spanning ~ 10 h of measurement were acquired overnight.

We conducted two types of PALM experiments. First, we determined the bleaching characteristic of PAmCherry under our experimental conditions, and second, we measured the unloading time of a single β_2 -clamp. PALM images were acquired as follows. First, the cell outlines were imaged by taking a single phase-contrast (PH) image using a commercial Nikon external phase ring configuration. The sample was then excited for a single frame (400 ms exposure time) by a 561 nm laser with an intensity of $\sim 5 \text{ W cm}^{-2}$, calculated according to Grünwald *et al.*⁶⁴ This image was used to determine the auto-fluorescence level due to the sample before activation. Photoactivation of PAmCherry was done with a single pulse (5 s) of 405 nm with an intensity of $\sim 2.5 \text{ W cm}^{-2}$, calculated according to Grünwald *et al.*⁶⁴ Subsequently, a post-activation time-lapse of images were acquired using the 561 nm laser at the same intensity at a frame rate of either $\sim 700 \text{ ms}$ (bleaching experiments) or 5 s (unloading experiments) with an exposure time of 400 ms per frame. Camera gain was set to 100.

Image analysis of long time-lapse experiments. Images were analysed with custom-written MATLAB software (MathWorks). Before any analysis, we subtract the uneven background using a rolling-ball filter⁶⁵ and subsequently corrected for illumination heterogeneity by using the previously measured laser beam profile⁶⁶. We also align the brightfield and fluorescence signals with respect to each other with 1-pixel accuracy. X–Y drift is corrected in both the fluorescence and brightfield images by tracking a fiducial marker in the PDMS to within 1 pixel.

Each drift-corrected region of interest, consisting out of a single growth channel, is analysed individually. The brightfield images are used to determine the cell poles of all the cells in a given frame. For the fluorescence signal, a kymograph of the fluorescence signal is constructed by summation of the pixel intensities per image perpendicular to the channel direction for each frame. This results in summed intensity information as a function of time per growth channel (Fig. 2c). We make use of the generated kymographs to determine individual replication and division cycles per cell (Fig. 2d). A post-processing step is subsequently performed to eliminate cells that did not match the following selection criteria: correct cell length, sufficient growth characteristics, observation of a complete cell cycle, clear fluorescence signal that both starts and ends in a diffuse state (Fig. 2d).

The fluorescence images of the detected individual cells that pass the above selection criteria are analysed further. We base our fluorescence analysis on an image of an individual bacterium with its long axis aligned with the horizontal direction of the image. The width of the image is equal to the length of the bacterium. We fix the height of the image such that a sufficient area above and below the bacterium is included that is indicative of the auto-fluorescence of the sample. We analyse the fluorescent intensity counts of a single bacterium using the individual fluorescence kymographs of each cell (summed line-profiles) by calculating three types of image content for a specific bacterium, namely ‘background’, ‘foci’ and ‘cytoplasm’ (Supplementary Fig. 3a). In brief, we first estimate the background fluorescence from the sample using the signal outside the bacterium. We did not have to take into account auto-fluorescence from the bacterium itself, as we conducted our experiments using minimal medium, which results in negligible levels of cellular auto-fluorescence (Supplementary Fig. 3b). The intensity outside the bacterium is used for a threshold with the remaining pixels intensities being representative of the total bacterium fluorescent counts. We subsequently separate ‘cytoplasm’ and ‘foci’ signals by determining the median of the summed line profiles. The signal significantly above this value is attributed to foci, whereas the remainder (lower values) are treated as the fluorescence signal from the cytoplasm (Supplementary Fig. 3a). This results in an integrated intensity value for the foci and also for the cytoplasm.

Image analysis of PALM experiments. PALM data was analysed using custom-written MATLAB software (MathWorks) in combination with the freely available MicrobeTracker software⁶⁷. Before any spot analysis, the fluorescence images are subjected to illumination correction and to alignment with respect to the phase-contrast (PH) images. The resulting corrected and aligned fluorescence images are then used during further analysis.

Using the PH image, the different cells are detected in the field of view and their respective outlines are determined using MicrobeTracker. Subsequently, using the spot detection algorithm as described in Olivo-Marin⁶⁸, the spots in each individual image of the fluorescence time-lapse series are detected, and the integrated intensity is determined by summing the pixel values of each spot⁶⁹. The integrated intensities of the spots are followed as function of time. This results in individual time-lapse integrated intensity traces of single molecules (Fig. 3c). The cell outlines as determined previously are overlaid with the fluorescence images. Any foci that are not situated in a bacterial cell (false positives) are rejected from further analysis. Only cells that had a clear fluorescence intensity focus were analysed. This focus is indicative of DNA-bound clamps and thus DNA replication. Foci that exhibit multiple steps in fluorescence intensity are also rejected. For the remainder of the foci, the time it takes from the start of the data acquisition until spot disappearance is recorded (Fig. 3c). These calculated time differences are indicative of molecule unloading (or bleaching, depending on the time of acquisition) and analysed further as described in the following section.

Monte Carlo simulation of β_2 -loading and unloading dynamics. For illustrative purposes, we perform Monte Carlo simulations (Fig. 5) starting with no clamps loaded and no primers formed ($N_{\beta_2} = 0$), and assuming the loading rate to be much faster than the rate of primer formation ($N_p \approx 0$). In each small time-step dt , we let $N_{\beta_2}\beta_2 + 1$ with probability $dt t_p^{-1}$ and $N_{\beta_2}\beta_2 - 1$ with probability $N_{\beta_2} dt t_{\text{unload}}^{-1}$. This is repeated until the replication time is reached, upon which the primer formation rate t_p^{-1} is set to zero.

References

1. Indiani, C., McInerney, P., Georgescu, R., Goodman, M. F. & O'Donnell, M. A sliding-clamp toolbelt binds high- and low-fidelity DNA polymerases simultaneously. *Mol. Cell* **19**, 805–815 (2005).
2. López de Saro, F. J. Regulation of interactions with sliding clamps during DNA replication and repair. *Curr. Genomics* **10**, 206–215 (2009).
3. Vivona, J. B. & Kelman, Z. The diverse spectrum of sliding clamp interacting proteins. *FEBS Lett.* **546**, 167–172 (2003).
4. Napolitano, R., Janel-Bintz, R., Wagner, J. & Fuchs, R. P. All three SOS-inducible DNA polymerases (Pol II, Pol IV and Pol V) are involved in induced mutagenesis. *EMBO J.* **19**, 6259–6265 (2000).
5. Wagner, J., Fujii, S., Gruz, P., Nohmi, T. & Fuchs, R. P. The beta clamp targets DNA polymerase IV to DNA and strongly increases its processivity. *EMBO Rep.* **1**, 484–488 (2000).
6. López de Saro, F. J. & O'Donnell, M. Interaction of the beta sliding clamp with MutS, ligase, and DNA polymerase I. *Proc. Natl Acad. Sci. USA* **98**, 8376–8380 (2001).
7. Kobayashi, S., Valentine, M. R., Pham, P., O'Donnell, M. & Goodman, M. F. Fidelity of Escherichia coli DNA polymerase IV. Preferential generation of small deletion mutations by dNTP-stabilized misalignment. *J. Biol. Chem.* **277**, 34198–34207 (2002).
8. Tang, M. *et al.* Roles of E. coli DNA polymerases IV and V in lesion-targeted and untargeted SOS mutagenesis. *Nature* **404**, 1014–1018 (2000).
9. Jergic, S. *et al.* A direct proofreader-clamp interaction stabilizes the pol III replicase in the polymerization mode. *EMBO J.* **32**, 1322–1333 (2013).
10. Dalrymple, B. P., Kongsuwan, K., Wijffels, G., Dixon, N. E. & Jennings, P. A. A universal protein-protein interaction motif in the eubacterial DNA replication and repair systems. *Proc. Natl Acad. Sci. USA* **98**, 11627–11632 (2001).
11. Katayama, T., Kubota, T., Kurokawa, K., Crooke, E. & Sekimizu, K. The initiator function of DnaA protein is negatively regulated by the sliding clamp of the E. coli chromosomal replicase. *Cell* **94**, 61–71 (1998).
12. Kato, J. & Katayama, T. Hda, a novel DnaA-related protein, regulates the replication cycle in Escherichia coli. *EMBO J.* **20**, 4253–4262 (2001).
13. Kurz, M., Dalrymple, B., Wijffels, G. & Kongsuwan, K. Interaction of the sliding clamp beta-subunit and Hda, a DnaA-related protein. *J. Bacteriol.* **186**, 3508–3515 (2004).
14. Ozaki, S. *et al.* A replicase clamp-binding dynamin-like protein promotes colocalization of nascent DNA strands and equipartitioning of chromosomes in E. coli. *Cell Rep.* **4**, 985–995 (2013).
15. Simmons, L. A., Davies, B. W., Grossman, A. D. & Walker, G. C. Beta clamp directs localization of mismatch repair in Bacillus subtilis. *Mol. Cell* **29**, 291–301 (2008).
16. Kong, X., Onrust, R., O'Donnell, M. & Kuriyan, J. Three-dimensional structure of the β subunit of E. coli DNA polymerase III holoenzyme: a sliding DNA clamp. *Cell* **69**, 425–437 (1992).
17. Johnson, A. & O'Donnell, M. Cellular DNA replicases: components and dynamics at the replication fork. *Annu. Rev. Biochem.* **74**, 283–315 (2005).
18. McHenry, C. S. DNA replicases from a bacterial perspective. *Annu. Rev. Biochem.* **80**, 403–436 (2011).
19. Turner, J., Hingorani, M. M., Kelman, Z. & O'Donnell, M. The internal workings of a DNA polymerase clamp-loading machine. *EMBO J.* **18**, 771–783 (1999).
20. Hedglin, M., Kumar, R. & Benkovic, S. J. Replication clamps and clamp loaders. *Cold Spring Harb. Perspect. Biol.* **5**, a010165 (2013).
21. Simonetta, K. R. *et al.* The mechanism of ATP-dependent primer-template recognition by a clamp loader complex. *Cell* **137**, 659–671 (2009).
22. Hayner, J. N. & Bloom, L. B. The β sliding clamp closes around dna prior to release by the Escherichia coli clamp loader γ complex. *J. Biol. Chem.* **288**, 1162–1170 (2013).
23. Downey, C. D. & McHenry, C. S. Chaperoning of a replicative polymerase onto a newly assembled DNA-bound sliding clamp by the clamp loader. *Mol. Cell* **37**, 481–491 (2010).
24. Leu, F. P., Hingorani, M. M., Turner, J. & O'Donnell, M. The delta subunit of DNA polymerase III holoenzyme serves as a sliding clamp unloader in Escherichia coli. *J. Biol. Chem.* **275**, 34609–34618 (2000).
25. McInerney, P., Johnson, A., Katz, F. & O'Donnell, M. Characterization of a triple DNA polymerase replisome. *Mol. Cell* **27**, 527–538 (2007).
26. Reyes-Lamothe, R., Sherratt, D. J. & Leake, M. C. Stoichiometry and architecture of active DNA replication machinery in Escherichia coli. *Science* **328**, 498–501 (2010).

27. Lia, G., Michel, B. & Allemand, J. Polymerase exchange during Okazaki fragment synthesis observed in living cells. *Science* **335**, 328–331 (2012).
28. Ogawa, T. & Okazaki, T. Discontinuous DNA replication. *Annu. Rev. Biochem.* **49**, 421–457 (1980).
29. Tougu, K. & Marians, K. J. The interaction between helicase and primase sets the replication fork clock. *J. Biol. Chem.* **271**, 21398–21405 (1996).
30. Burgers, P. M., Kornberg, A. & Sakakibara, Y. The *dnaN* gene codes for the beta subunit of DNA polymerase III holoenzyme of *Escherichia coli*. *Proc. Natl Acad. Sci. USA* **78**, 5391–5395 (1981).
31. Yao, N. *et al.* Clamp loading, unloading and intrinsic stability of the PCNA, β and gp45 sliding clamps of human, *E. coli* and T4 replicases. *Genes Cells* **1**, 101–113 (1996).
32. Yuzhakov, A., Turner, J. & O'Donnell, M. Replisome assembly reveals the basis for asymmetric function in leading and lagging strand replication. *Cell* **86**, 877–886 (1996).
33. Tanner, N. A. *et al.* *E. coli* DNA replication in the absence of free β clamps. *EMBO J.* **30**, 1830–1840 (2011).
34. Su'etsugu, M. & Errington, J. The replicase sliding clamp dynamically accumulates behind progressing replication forks in *Bacillus subtilis* cells. *Mol. Cell* **41**, 720–732 (2011).
35. Hess, S. T., Girirajan, T. P. K. & Mason, M. D. Ultra-high resolution imaging by fluorescence photoactivation localization microscopy. *Biophys. J.* **91**, 4258–4272 (2006).
36. Betzig, E. *et al.* Imaging intracellular fluorescent proteins at nanometer resolution. *Science* **313**, 1642–1645 (2006).
37. Dulin, D., Lipfert, J., Moolman, M. C. & Dekker, N. H. Studying genomic processes at the single-molecule level: introducing the tools and applications. *Nat. Rev. Genet.* **14**, 19–22 (2013).
38. Duderstadt, K. E., Reyes-Lamothe, R., van Oijen, A. M. & Sherratt, D. J. Replication-fork dynamics. *Cold Spring Harb. Perspect. Biol.* **6**, a010157 (2013).
39. Xia, T., Li, N. & Fang, X. Single-molecule fluorescence imaging in living cells. *Annu. Rev. Phys. Chem.* **64**, 459–480 (2013).
40. Persson, F., Barkefors, I. & Elf, J. Single molecule methods with applications in living cells. *Curr. Opin. Biotechnol.* **24**, 737–744 (2013).
41. Gahlmann, A. & Moerner, W. E. Exploring bacterial cell biology with single-molecule tracking and super-resolution imaging. *Nat. Rev. Microbiol.* **12**, 9–22 (2014).
42. Wang, X., Reyes-Lamothe, R. & Sherratt, D. J. Visualizing genetic loci and molecular machines in living bacteria. *Biochem. Soc. Trans.* **36**, 749–753 (2008).
43. Moolman, M. C., Huang, Z., Krishnan, S. T., Kerssemakers, J. W. J. & Dekker, N. H. Electron beam fabrication of a microfluidic device for studying submicron-scale bacteria. *J. Nanobiotechnol.* **11**, 12 (2013).
44. Wang, P. *et al.* Robust growth of *Escherichia coli*. *Curr. Biol.* **20**, 1099–1103 (2010).
45. Nguyen, A. W. & Daugherty, P. S. Evolutionary optimization of fluorescent proteins for intracellular fret. *Nat. Biotechnol.* **23**, 355–360 (2005).
46. Reyes-Lamothe, R., Possoz, C., Danilova, O. & Sherratt, D. J. Independent positioning and action of *Escherichia coli* replisomes in live cells. *Cell* **133**, 90–102 (2008).
47. Michelsen, O., Teixeira de Mattos, M. J., Jensen, P. R. & Hansen, F. G. Precise determinations of C and D periods by flow cytometry in *Escherichia coli* K-12 and B/r. *Microbiology* **149**, 1001–1010 (2003).
48. Coffman, V. C. & Wu, J. Counting protein molecules using quantitative fluorescence microscopy. *Trends Biochem. Sci.* **37**, 499–506 (2012).
49. Swulius, M. T. & Jensen, G. J. The helical MreB cytoskeleton in *Escherichia coli* MC1000/ple7 is an artifact of the N-Terminal yellow fluorescent protein tag. *J. Bacteriol.* **194**, 6382–6386 (2012).
50. Landgraf, D., Okumus, B., Chien, P., Baker, T. A. & Paulsson, J. Segregation of molecules at cell division reveals native protein localization. *Nat. Methods* **9**, 480–482 (2012).
51. Subach, F. V. *et al.* Photoactivatable mcherry for high-resolution two-color fluorescence microscopy. *Nat. Methods* **6**, 153–159 (2009).
52. Yuan, Q. & McHenry, C. S. Cycling of the *E. coli* lagging strand polymerase is triggered exclusively by the availability of a new primer at the replication fork. *Nucleic Acids Res.* **42**, 1747–1756 (2014).
53. Marians, K. J. Prokaryotic DNA replication. *Annu. Rev. Biochem.* **61**, 673–719 (1992).
54. Stukenberg, P. T., Turner, J. & O'Donnell, M. An explanation for lagging strand replication: polymerase hopping among DNA sliding clamps. *Cell* **78**, 877–887 (1994).
55. Reyes-Lamothe, R. *et al.* High-copy bacterial plasmids diffuse in the nucleoid-free space, replicate stochastically and are randomly partitioned at cell division. *Nucleic Acids Res.* **42**, 1042–1051 (2014).
56. Wijffels, G., Dalrymple, B., Kongsuwan, K. & Dixon, N. E. Conservation of eubacterial replicases. *IUBMB Life* **57**, 413–419 (2005).
57. Su'etsugu, M., Shimuta, T.-R., Ishida, T., Kawakami, H. & Katayama, T. Protein associations in DnaA-ATP hydrolysis mediated by the Hda-replicase clamp complex. *J. Biol. Chem.* **280**, 6528–6536 (2005).
58. Mott, M. L. & Berger, J. M. DNA replication initiation: mechanisms and regulation in bacteria. *Nat. Rev. Microbiol.* **5**, 343–354 (2007).
59. Georgescu, R. E., Yao, Y.N. & O'Donnell, M. Single-molecule analysis of the *Escherichia coli* replisome and use of clamps to bypass replication barriers. *FEBS Lett.* **584**, 2596–2605 (2010).
60. Lenhart, J. S., Sharma, A., Hingorani, M. M. & Simmons, L. A. DnaN clamp zones provide a platform for spatiotemporal coupling of mismatch detection to DNA replication. *Mol. Microbiol.* **87**, 553–568 (2013).
61. Uphoff, S., Reyes-Lamothe, R., Garza de Leon, F., Sherratt, D. J. & Kapanidis, A. N. Single-molecule DNA repair in live bacteria. *Proc. Natl Acad. Sci. USA* **110**, 8063–8068 (2013).
62. Thomason, L. C., Costantino, N. & Court, D. L. *E. coli* genome manipulation by P1 transduction. *Curr. Protoc. Mol. Biol.* Chapter 1, Unit 1.17 (2007).
63. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl Acad. Sci. USA* **97**, 6640–6645 (2000).
64. Grünwald, D., Shenoy, S. M., Burke, S. & Singer, R. H. Calibrating excitation light fluxes for quantitative light microscopy in cell biology. *Nat. Protoc.* **3**, 1809–1814 (2008).
65. Sternberg, Biomedical image processing. *IEEE Comput.* **16**, 22–34 (1983).
66. Taniguchi, Y. *et al.* Quantifying *E. coli* proteome and transcriptome with single-molecule sensitivity in single cells. *Science* **329**, 533–538 (2010).
67. Sliusarenko, O., Heinritz, J. & Emonet, T. High-throughput, subpixel precision analysis of bacterial morphogenesis and intracellular spatio-temporal dynamics. *Mol. Microbiol.* **80**, 612–627 (2011).
68. Olivo-Marin, J. C. Extraction of spots in biological images using multiscale products. *Pattern. Recognit.* **35**, 1989–1996 (2002).
69. Ulbrich, M. H. & Isacoff, E. Y. Subunit counting in membrane-bound proteins. *Nat. Methods* **4**, 319–321 (2007).
70. Georgescu, R. E. *et al.* Structure of a sliding clamp on DNA. *Cell* **132**, 43–54 (2008).

Acknowledgements

We thank Ilja Westerlaken for her assistance with the French Press for cell lysate preparation, Hugo Snippet for initial help with the Southern blotting, and Bronwen Cross for initial help with the western blot, as well as Jasper van Veen, Kasper de Leeuw and Liliane Jimenez van Hoom for initial help with constructing the *PAmCherry-DnaN* strain. We thank Anjana Badrinarayanan for providing the *mCherry-DnaN* strain. We thank Jan Lipfert, Anne Meyer, Tim Blosser, Yaron Caspi, Antoine van Oijen and Theo van Laar for stimulating discussions. We thank Francesco Perdaci, David Dulin, David Gruenwald, Chirlmin Joo and Stanley Dinesh Chandradoss for initial help with microscopy. We thank Bernd Rieger for initial help with image analysis. We acknowledge contributions from Jelle van der Does, Dimitri de Roos and Jaap Beekman for instrumentation and infrastructure. This work was supported by the European Community's Seventh Framework Programme FP7/2007–2013 under grant agreement no. 241548 (MitoSys) and a Vici grant by the Netherlands Organization for Scientific Research (NWO), both to N.H.D., a Wellcome Trust Program Grant WT083469 to D.J.S., a TU Delft startup grant to M.D. and the Natural Sciences & Engineering Research Council of Canada (#435521-13) for R.R.-L.

Author contributions

M.C.M. and N.H.D. designed the research and the experiments. M.C.M. and S.T.K. undertook the experiments. S.T.K. constructed the strains. P.T. performed the western blot. S.T.K. performed the Southern blot. M.C.M. and J.W.J.K. wrote the software to analyse all the microscopy data. M.C.M., A.v.d.B. and M.D. analysed the data. A.v.d.B. performed the Monte Carlo simulation. R.R.-L. and D.J.S. provided strains and contributed to the planning and discussion of the work. M.C.M. and N.H.D. wrote the paper.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Moolman, M. C. *et al.* Slow unloading leads to DNA-bound β_2 -sliding clamp accumulation in live *Escherichia coli* cells. *Nat. Commun.* **5**:5820 doi: 10.1038/ncomms6820 (2014).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>