

## ERRATUM: Inhibitors of histone deacetylases induce tumor-selective apoptosis through activation of the death receptor pathway

A Insinga, S Monestiroli, S Ronzoni, V Gelmetti, F Marchesi, A Viale, L Altucci, C Nervi, S Minucci & P G Pelicci  
*Nat. Med.* 11, 71–76 (2005)

In **Fig. 2a**, the second and fourth graphs in the lower row should be marked *Tnfrsf10b* and *Tnfrsf6*, respectively. In **Fig. 2b**, the gene symbol *Tp53* should read *Tp53<sup>-/-</sup>*.

In **Methods**, the administration of VPA and TSA for *in vivo* antitumor activity studies should be “intraperitoneally for 21 d at a dose of 400 mg/kg twice daily and 1 mg/kg/d, respectively.”

## ERRATUM: Inhibition of respiratory syncytial virus infection with intranasal siRNA nanoparticles targeting the viral NS1 gene

W Zhang, H Yang, X Kong, S Mohapatra, H San Juan-Vergara, G Hellermann, S Behera, R Singam, R F Lockey & S S Mohapatra  
*Nat. Med.* 11, 56–62 (2005)


The open square in **Fig. 1b** refers to results from A549 cells, not from uninfected cells.

## ERRATUM: The clock gene *Per2* influences the glutamatergic system and modulates alcohol consumption


R Spanagel, G Pendyala, C Abarca, T Zghoul, C Sanchis-Segura, M C Magnone, J Lascorz, M Depner, D Holzberg, M Soyka, S Schreiber, F Matsuda, M Lathrop, G Schumann & U Albrecht  
*Nat. Med.* 11, 35–42 (2005)

In the print version of this article, **Figures 4d, 4f** and **5a** were inadvertently truncated. The correct versions appear below.

4d



4f



5a

