

Impact of Ovarian Hormones on the Modulation of the Serotonin Transporter by Fluvoxamine

Saloua Benmansour^{*1}, Jonathan P Piotrowski¹, Alfonso V Altamirano¹ and Alan Frazer^{1,2}

¹Department of Pharmacology, University of Texas Health Science Center at San Antonio, San Antonio, TX, USA; ²South Texas Veterans Health Care System, San Antonio, TX, USA

Most preclinical studies examining the mechanism(s) of action of antidepressants are carried out using male animals. Blockade of serotonin transporter (SERT) function by selective serotonin reuptake inhibitors (SSRIs) is the initial event that triggers a not completely understood process that results in clinical improvement in depression. To investigate whether there are differences in the ability of SSRIs to inhibit the SERT between male and female rats at different phases of the estrous cycle, clearance of locally applied serotonin (5-HT) was measured by *in vivo* chronoamperometry. Local application of the SSRI, fluvoxamine, directly into the CA3 area of hippocampus increased significantly 5-HT clearance time parameters in male rats and female rats in estrus or diestrus, but not in proestrus. The contribution of ovarian steroids to this result was investigated in ovariectomized (OVX) rats treated with estradiol benzoate (EB) and/or progesterone (P). In OVX-control rats, fluvoxamine increased clearance time parameters, whereas EB and/or P treatment blocked this effect, consistent with what was seen in female rats in proestrus. This effect was gender-specific, since treatment of castrated rats with EB/P had no effect on the ability of fluvoxamine to slow 5-HT clearance. The time course of hormonal effects showed that 1–60 min after local application of 17- β -estradiol (E₂) into the CA3 region of OVX rats, fluvoxamine had no effect on clearance time of 5-HT. E₂-BSA mimicked E₂'s effects at 10 min but not at 60 min. Pretreatment with estrogen receptor antagonists blocked the effects of E₂. The finding that acutely both estradiol and progesterone can inhibit the ability of an SSRI to slow the clearance of 5-HT, may have important implications for the use of SSRIs in women.

Neuropsychopharmacology (2009) **34**, 555–564; doi:10.1038/npp.2008.23; published online 5 March 2008

Keywords: serotonin transporter; estradiol; progesterone; fluvoxamine; chronoamperometry; hippocampus

INTRODUCTION

Depression is more common in females than males (Kessler, 2003) and in women, may be more common at times of low estradiol levels (Fink *et al*, 1996; Halbreich *et al*, 1992; Pearlstein, 1995). Depressive episodes in women may be more recurrent, longer in length, and associated with more functional impairment than those in men (Burt and Stein, 2002; Yonkers, 2003). In spite of this, preclinical data examining the effects of antidepressants (ADs) on either behavioral, physiological, or neurochemical parameters, have mainly used males subjects until recently. Pharmacological interventions involving estrogens (and/or progesterone) in depressed women have been proposed to correct presumed steroid hormone deficiencies or fluctuations. Hormonal (or estrogen alone) therapy has also been reported to have AD effects in perimenopausal and

postmenopausal women (Zweifel and O'Brien, 1997; Soares *et al*, 2001; Huttner and Shepherd, 2003), although this has not been a universal finding (Saletu *et al*, 1995). Pre-menopausal depressed women may have better response to selective serotonin reuptake inhibitors (SSRIs) than to tricyclic ADs (Kornstein *et al*, 2000). However, other studies did not confirm these findings (Quitkin *et al*, 2002; Scheibe *et al*, 2003). Adjunctive treatment with estrogen in addition to an SSRI may be beneficial in depressed menopausal patients (Yonkers, 2003), although, again, this has not been found in all studies (Huttner and Shepherd, 2003). For multiple reasons, SSRIs are the treatment of choice for depression during perimenopause and menopause.

Blockade of serotonin transporter (SERT) function by SSRIs is the initial event that triggers a still not completely understood process that results in clinical improvement after repeated treatment (Lenox and Frazer, 2002). Reports analyzing the action of estrogen *in vivo* on SERT uptake sites and mRNA levels are inconclusive. Increases, decreases, and no change have been reported (Attali *et al*, 1997; Bethea *et al*, 1998; Krajnak *et al*, 2003; McQueen *et al*, 1997; Mendelson *et al*, 1993; Pecins-Thompson *et al*, 1998; Zhou *et al*, 2002). Discrepancies in these studies may be explained by variability in the treatment paradigm used,

*Correspondence: Dr S Benmansour, Department of Pharmacology (MC 7764), University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78229-3900, USA, Tel: +1 210 567 4203; Fax: +1 210 567 4303; E-mail: benmansour@uthscsa.edu
Received 5 September 2007; revised 10 January 2008; accepted 28 January 2008

such as the time elapsed between ovariectomy and estrogen treatment, the duration of estrogen treatment, and the brain region evaluated.

Estrogen receptors (ER) as well as progesterone receptors are members of the superfamily of hormone-regulated nuclear receptors. Depending on the onset and duration of the effect, ovarian hormones can act either by a genomic mechanism, using classical nuclear transcription factor receptors (Katzenellenbogen, 2000), or by a non-genomic, less well-characterized, membrane signaling mechanism (Kelly *et al*, 1999; Nadal *et al*, 2000). The genomic pathway requires at least 30–60 min for its effects to manifest and is associated with changes in protein synthesis. However, the 'nuclear hormone receptors' are not restricted to the nucleus; they can be found within the nucleus, cytoplasm, and in the cell membrane. It has more recently become clear that estrogen can elicit rapid, within seconds to minutes, signaling events that are not mediated by classical genomic pathways. Steroid-bovine serum albumin (BSA) conjugates have become useful tools to study the membrane-mediated action of steroids, since they are biologically active and do not penetrate the plasma membrane of the cell readily (Blackmore and Lattanzio, 1991; Ramirez and Zheng, 1996).

To begin providing a comprehensive understanding of how hormones such as estrogen or progesterone influence the effect of SSRIs, it is important to determine whether they alter their inhibitory effect on the SERT. This study describes how SSRIs acutely affect SERT function in female rats in comparison with the effects observed in male rats, and how ovarian hormones influence these effects. SERT function was measured by *in vivo* chronoamperometry in female rats at different stages of the estrous cycle, as well as in ovariectomized (OVX) rats treated with estradiol and/or progesterone. The time course of the effects caused by local administration of 17- β -estradiol (E_2), E_2 -BSA, and ER antagonists was also investigated.

MATERIALS AND METHODS

Animals

Intact females, OVX females, intact males, or castrated males (all Sprague-Dawley rats; 250–350 g, Harlan, Indianapolis, IN) were housed on a 12:12 h light/dark cycle with lights on at 0700 h and with food and water provided *ad libitum*. All animal procedures were in strict accordance with the National Institutes of Health Guide for Care and Use of Laboratory Animals, and were approved by the local Institutional Care and Use Committee. All efforts were made to minimize both the number of animals used and stress or discomfort to the animal during the experimental procedure.

Female rats are sexually mature at about 8–9 weeks and weigh about 200 g at that age. Vaginal smears in intact female rats were monitored daily for the presence of predominantly nucleated cells (proestrus), cornified cells (estrus), or leukocytes (diestrus). Only females showing four consecutive regular estrous cycles (4–5 days) were used in these studies. Classification of the estrous cycle was based on the vaginal smear on the day of the experiment and the prior vaginal smear history. For comparison purposes, male rats were also studied. OVX and castrated rats were allowed

a 2- to 3-week recovery after surgery, before starting treatment, in order to provide time for endogenous hormones to decline and the animals to adapt to their loss.

Hormone Treatments

Because circulating amounts of E_2 do not remain elevated for very long after systemic injection of E_2 , a slower-release esterified form of E_2 , namely estradiol benzoate (EB), was used for this part of the study. The treatment paradigm was chosen to achieve estrogen and progesterone plasma levels that were similar to those characterized in proestrus (Gibbs, 1996). For combination treatment, EB (25 μ g in 100 μ l peanut oil; Sigma-Aldrich, St Louis, MO) was administered subcutaneously to OVX or castrated rats, followed 48 h later by subcutaneous injection of progesterone (P, 500 μ g in 100 μ l peanut oil; Sigma-Aldrich). The experiments were carried out 24 h after administration of P. When given separately, the hormones were administered at these doses with the experiments carried out 72 h after administration of EB or 24 h after P. Control OVX rats received peanut oil injections.

In Vivo Chronoamperometry

Chronoamperometric recordings were carried out *in vivo* in the CA3 region of the hippocampus as described previously (Benmansour *et al*, 1999, 2002). Multi-barrel micropipettes were filled with either 5-HT (200 μ M; Sigma Chemicals Co., St Louis, MO), fluvoxamine (400 μ M; Pharmacia and Upjohn, Kalamazoo, MI), citalopram (400 μ M; Forest Laboratories, Jersey City, NJ), E_2 (dissolved in PBS containing ethanol at a final concentration less than 0.0001%; Sigma-Aldrich), ICI 182,780, or tamoxifen (Tocris, Ellisville, MO), or β -estradiol 6-(*O*-carboxymethyl)oxime:BSA (E_2 -BSA; Sigma-Aldrich). The latter was filtered as described by Stevis *et al* (1999), using a Microcon cartridge with a 3-kDa cut-off (Amicon, Beverly, MA) to eliminate any free E_2 from the E_2 -BSA solution. 5-HT was delivered by pressure ejection. After obtaining reproducible electrochemical signals to 5-HT, the ability of fluvoxamine or citalopram (four times the amount of 5-HT applied) to block 5-HT clearance was measured by its local application 60–90 s before pressure ejection of 5-HT. High-speed chronoamperometric recordings were made using the Fast-12 system (Censet, Lexington, KY).

Several parameters are obtained from the electrochemical signal produced by exogenous applications of 5-HT. Shown in this study is clearance time parameter, T_{80} , which is the time required for the peak amplitude to be reduced by 80%. Also analyzed in this study is clearance rate, T_C , which is based on the slope of the initial pseudolinear (from T_{20} to T_{60}) portion of the decaying signal (Zahniser *et al*, 1999).

Autoradiographic Procedures

Serotonin uptake sites were detected using [3 H]cyanoimipramine ([3 H]-CN-IMI) (Benmansour *et al*, 1999, 2002). Brain sections (20 μ m) were incubated with 1 nM [3 H]-CN-IMI (80–85 Ci/mmol; American Radiolabeled Chemicals Inc., St Louis, MO) for 24 h at 4°C in a buffer containing 50 mM Tris (pH 7.4) and 150 mM NaCl. Non-specific

binding was defined using 5 μ M sertraline, and was approximately 5% of total binding. Following incubation, sections were washed in buffer for 60 min at 4°C and were then dipped in distilled water. Dried, slide-mounted sections and calibrated [³H] standards (American radiolabelled Chemicals Inc.) were placed into spring-loaded cassettes and apposed to tritium-sensitive film, Kodak Biomax MR (Kodak, Rochester, NY) at room temperature for 6 weeks. Optical densities of brain images were converted to femtomoles per milligram protein using the NIH-IMAGE software. Measurements were taken at the level of plate 30 of the atlas of Paxinos and Watson (1986).

Homogenate-Binding Procedures

The binding of [³H]citalopram to hippocampal membranes was determined according to the method developed by D'Amato *et al* (1987), with a few modifications. In brief, rats were decapitated and the hippocampus was dissected out and homogenized in 25 ml of ice-cold buffer (50 mM Tris-HCl, 120 mM NaCl, and 5 mM KCl, pH 7.4). The homogenate was centrifuged at 30 000g for 10 min at 4°C. The resulting pellet was resuspended in buffer and centrifuged again twice. The homogenate was incubated with a saturating concentration of [³H]citalopram (5 nM, 72 Ci/mmol; PerkinElmer, Boston, MA) for 75 min in the absence or presence of fluvoxamine (3×10^{-10} – 1×10^{-6} M) at 24°C in a final volume of 500 μ l. Incubation was terminated by adding ice-cold buffer and filtration through Whatman GF/B filters (pretreated with 0.1% polyethylenamine) using a 24-channel cell harvester. Radioactivity trapped in the filters was measured using a Packard 1900 TR liquid scintillation counter. Protein content was determined by the method of Bradford (1976). A non-linear regression fit of the data using the Graphpad Prism Software (Intuitive Software for Science, San Diego, CA) allowed to determine the IC₅₀ values and calculate the inhibition constants K_i according to the Cheng–Prusoff equation. The K_d value used in this study to calculate K_i was determined from saturation binding data for [³H]citalopram (0.1–10 nM) in hippocampal membranes prepared from male or female rats in proestrus. K_d values (average of three independent experiments for each group, each performed in triplicate) for [³H]citalopram binding to the SERT were similar in male rats (0.87 ± 0.10 nM) or females in proestrus (0.93 ± 0.09 nM). Therefore, an average value for K_d of 0.9 nM was used to calculate K_i for all the groups.

Hormones Levels

OVX rats were treated with EB/P or vehicle as described above in the hormone treatment section. Trunk blood was collected (72 h after administration of EB or 24 h after P) for measurement of hormone levels in serum by radioimmunoassay (RIA). Estradiol concentrations in rat serum were determined by RIA of duplicate samples using the DSL ultrasensitive estradiol RIA kit (DSL-4800), according to the manufacturer's instructions; progesterone concentrations in the rat serum were determined by RIA of duplicate samples using the DSL progesterone-coated tube RIA kit (DSL-3900) according to the manufacturer's protocol (Adams *et al*, 2001; Gore *et al*, 2000).

Statistical Analysis

Data were analyzed by *t*-test or one-way ANOVA followed by Newman–Keuls *post hoc* multiple comparisons, with significance determined at $p < 0.05$.

RESULTS

Effect of Fluvoxamine on SERT Function During the Estrous Cycle

The first experiment was designed to examine the ability of fluvoxamine to inhibit SERT function in female rats in diestrus, proestrus, and estrus in comparison with the effects observed in male rats. A representative 5-HT signal generated by local application of exogenous serotonin in the CA3 region is shown in Figure 1. As expected, fluvoxamine prolonged the clearance time of 5-HT in male rats (Benmansour *et al*, 1999, 2002). By contrast, in female rats in proestrus, local application of fluvoxamine did not prolong the clearance time of 5-HT. The effect of fluvoxamine on the 5-HT clearance time parameter, T_{80} , in males, or in females in proestrus, estrus, or diestrus, is shown in Figure 2a. In male rats, as well as in female rats in estrus or diestrus,




Figure 1 Representative 5-HT electrochemical signals illustrating the effect of locally applied fluvoxamine in the CA3 region of hippocampus of a male rat or a female rat in proestrus. The signal was generated by local application of 5-HT (2.6–10 pmol). Fluvoxamine ($4 \times$ the amount of 5-HT) was pressure-ejected 60–90 s before the next application of 5-HT. For clarity, only oxidation signals are shown.




Figure 2 Effects of fluvoxamine on the clearance time, T_{80} , and the clearance rate, T_c , in male and female rats at different stages of the estrous cycle. Electrochemical recordings were carried out in the CA3 region of dorsal hippocampus of male rats or female rats in proestrus, estrus or diestrus. Fluvoxamine was pressure-ejected 60–90s before the second application of 5-HT. (a) Bars and brackets represent mean T_{80} value (in sec) before and after fluvoxamine application; \pm SEM ($n = 8-10$). (b) Bars and brackets represent mean T_c value (in nM/s) before and after fluvoxamine application; \pm SEM ($n = 8-10$). * $p < 0.05$, paired Student's t -test comparing pre-fluvoxamine value with post-fluvoxamine value for each group. The amount of 5-HT ejected was adjusted so as to obtain comparable signal amplitudes (0.25–0.4 μ M) in the various groups of animals. This approach enables clearance rates to be compared easily among the treatment groups, as clearance rate is dependent on signal amplitude. Consequently, the volume (and therefore the amount) of 5-HT was varied. The mean amount of 5-HT in pmoles \pm SEM pressure ejected in each group was not significantly different between groups and was as follows: male, 6.3 ± 3 ; female proestrus, 14.7 ± 4 ; female estrus, 10.0 ± 3 ; and female diestrus, 7.2 ± 2 . Fluvoxamine was always given at four times the amount of 5-HT applied. There were no significant correlations between the volume of 5-HT ejected and fluvoxamine-induced effects on the T_{80} parameter (data not shown). Thus, whether fluvoxamine had an effect or not, was independent of the volume (or amount) of 5-HT ejected.

fluvoxamine slowed clearance of 5-HT, as demonstrated by a significant increase in T_{80} value. By contrast, in female rats in proestrus, the inhibitory effect of fluvoxamine on clearance time was abolished. The clearance rate, T_c did not change significantly upon local application of fluvoxamine in males or females at any stage of the cycle (Figure 2b).

SERT Density and Affinity for Fluvoxamine During the Estrous Cycle

The density of SERT binding sites was measured in the same rats using quantitative autoradiography of [3 H]-CN-

Table 1 [3 H]-CN-IMI Binding in Males and Females Rats at Different Phases of the Estrous Cycle

Groups	Region		
	Hippocampus (CA3)	Cortex (parietal)	Raphe nu (dorsal)
Male ($n = 6$)	510 ± 14^a	331 ± 16	2343 ± 170
F—estrus ($n = 8$)	494 ± 14	322 ± 13	2377 ± 126
F—diestrus ($n = 12$)	473 ± 13	313 ± 15	2370 ± 154
F—proestrus ($n = 8$)	475 ± 17	319 ± 5	2084 ± 126

Abbreviations: F, female; [3 H]-CN-IMI, [3 H]-cyanoimipramine.

Data are expressed as femtomoles/mg protein.

^aEach value is mean \pm SEM.




Figure 3 Representative data illustrating the potency of fluvoxamine to displace [3 H]citalopram (5 nM) binding to hippocampal membranes prepared from female rats in proestrus or estrus or from male rats. In each experiment, each point is calculated in triplicate. The number of rats (n) was 9 for male, 5 for estrus, and 5 for proestrus. Displacement curves show that the ability of fluvoxamine to displace [3 H]citalopram binding was similar in males and females in proestrus or estrus.

IMI binding (1 nM). This ligand concentration is about $8 \times$ the K_D value for [3 H]-CN-IMI (Kovachich *et al*, 1988) so that SERT densities obtained approximate B_{max} values. Specific binding in several brain areas was quantified in males or in females in proestrus, estrus, or diestrus, and the data are presented in Table 1. SERT binding sites were similar in males and females in all the areas analyzed. There was no fluctuation in binding sites in females during the estrous cycle.

The ability of fluvoxamine to block 5-HT clearance was reduced in female rats in proestrus as compared with that in female rats in estrus or male rats (Figures 1 and 2). This effect could result from a lower affinity (ie, higher K_i value) of fluvoxamine for SERT binding sites in rats in proestrus as compared with that in the two other groups. To test this hypothesis, the potency of fluvoxamine to displace [3 H]citalopram (5 nM) binding to hippocampal membranes prepared from female rats in proestrus or estrus or in male rats was measured. The ability of fluvoxamine to displace [3 H]citalopram binding was similar in males and in females in proestrus or in estrus (Figure 3). Comparison of

fluvoxamine's K_i values (3.84 ± 0.56 ; 3.43 ± 0.34 ; 3.33 ± 0.40 nM for males ($n=9$); females in estrus ($n=5$), or in proestrus ($n=5$), respectively) showed no differences between these groups (see details for calculation under Materials and Methods). Thus, the potency of fluvoxamine to compete for SERT-binding sites was similar in female rats in proestrus or estrus or in male rats.

Effects of Acute Ovarian Hormone Treatments on SERT Function and on the Effect of Fluvoxamine

Many parameters change during the estrus cycle in addition to circulating levels of estrogen or progesterone. To evaluate whether results obtained in female rats in proestrus were due to the influence of estrogen and/or progesterone on SERT function, OVX rats were treated with either EB and/or P. Concentrations of hormones in the serum were measured in order to verify that the hormone treatment regimen achieved estrogen and progesterone plasma levels characterized in proestrus. Mean (\pm SEM) serum levels of estradiol ($E_2 = 52.7 \pm 14.21$ pg/ml) and progesterone ($P = 11.2 \pm 1.91$ ng/ml) were measured by RIA after administration of the combination treatment EB/P ($n=6$). Control rats ($n=4$) received similar injections of peanut oil ($E_2 = 8.5 \pm 1.68$ pg/ml; $P = 2.5 \pm 1.24$ ng/ml). During the estrus cycle, E_2 peak levels of about 50 pg/ml occur in *pm* proestrus (Gibbs, 1996); this is in good agreement with the E_2 level found. Values for P are, if anything, somewhat lower than those found in proestrus.

The ability of fluvoxamine to modulate 5-HT clearance in these groups of rats is shown in Figure 4a. In control OVX rats, the clearance time parameter, T_{80} , was significantly increased by local application of fluvoxamine even though the clearance rate parameter, T_C , was not (Figure 4a and b). However, the ability of fluvoxamine to prolong the clearance time of serotonin was abolished by either EB or P treatment alone or by the combination of the two hormones. Similarly, local application into the CA3 region of another SSRI, citalopram, increased T_{80} value ($169 \pm 15\%$ of the pre-citalopram value) in OVX-control rats ($n=4$). This effect was blocked in OVX rats treated with EB/P ($n=4$, T_{80} value post-citalopram was $112 \pm 8\%$ of the pre-citalopram value).

In addition, EB treatment increased baseline (ie, pre-fluvoxamine) T_{80} value and decreased clearance rate, T_C , significantly from that measured in OVX rats (Figure 4b). The other treatment paradigms did not produce this effect.

To determine whether the effects of hormones are gender specific, castrated male rats were treated with EB/P, as described under Materials and Methods, and the effect of locally applied fluvoxamine was analyzed. T_{80} values in castrated rats treated with vehicle: 73 ± 12 s pre- and 108 ± 16 s post-fluvoxamine were similar to those measured in castrated male treated with EB/P (81 ± 11 s pre- and 123 ± 13 s post-fluvoxamine) The effect of fluvoxamine on T_{80} was significant ($p < 0.01$, paired *t*-test) and of similar magnitude in the two groups. Thus, the inhibitory effect of fluvoxamine on SERT function in castrated male rats was not affected by treatment with EB/P as it was in ovariectomized female rats.




Figure 4 Effects of fluvoxamine on the clearance time, T_{80} , and clearance rate, T_C , in OVX rats treated with ovarian hormones. Electrochemical recordings were performed in the CA3 region of dorsal hippocampus of OVX rats treated with EB, P, the combination EB/P, or peanut oil, as described under Materials and Methods. Fluvoxamine was pressure-ejected 60–90 s before the second application of 5-HT. (a) Bars and brackets represent mean T_{80} values (in seconds) before and after fluvoxamine application; \pm SEM ($n=7-9$). (b) Bars and brackets represent mean T_C values (in nM/s) before and after fluvoxamine application; \pm SEM ($n=7-9$). * $p < 0.05$, paired Student's *t*-test comparing each group's pre-fluvoxamine value with its post-fluvoxamine value. ** $p < 0.05$, Newman-Keuls comparing pre-fluvoxamine values for OVX hormones-treated with OVX-CTR. In this experiment too, the amount of 5-HT ejected was adjusted so as to obtain comparable signal amplitudes (0.25–0.4 μ M) in the various groups of animals, as described in the legend to Figure 2. The mean amount of 5-HT in pmoles \pm SEM pressure ejected in each group was not significantly different between groups and was as follows: OVX-CTR, 4.1 ± 1 ; OVX-EB + P, 9.8 ± 3 ; OVX-EB, 6.4 ± 2 ; and OVX-P, 4.2 ± 2 .

Time Course of the Effect of Local Application of 17- β -Estradiol on the Effect of Fluvoxamine

In this experiment, the time course of the effect of direct local application of E_2 into the CA3 region of the hippocampus on fluvoxamine's blockade of 5-HT clearance was investigated in OVX rats. Fluvoxamine's ability to block 5-HT clearance was examined from 1 to 120 min after local application of E_2 (10 or 57 pmoles) or PBS as the control vehicle. Changes in the T_{80} value obtained after application of fluvoxamine are presented in Figure 5a. Vehicle-treated rats or those administered various amounts of E_2 had the effect of fluvoxamine measured both prior to hormone administration (the time 0 value in Figure 5) as well as afterwards. The effect of fluvoxamine was consistent




Figure 5 Time course of the effect of local application of E_2 on fluvoxamine's ability to increase T_{80} and its modulation by ER antagonists. (a) The time course was examined from 1 to 120 min after local application of E_2 (57 or 10 pmoles) or PBS as the control vehicle. Time 0 represent the effect of fluvoxamine measured prior to giving the hormones or PBS. The vehicle for E_2 was PBS with <0.0001% ethanol. For these experiments, the same amount of 5-HT was pressure-ejected into a rat at 1 or 10 min time points after zero. It was not possible to measure all time points in each rat, since the signal sometimes took longer than 10 min to be stable after administration of fluvoxamine and 5-HT. The peak amplitudes for 5-HT obtained in the control rats at time 0 vs time point 90–120 min (0.34 ± 0.03 and $0.27 \pm 0.02 \mu\text{M}$, respectively) were not significantly different, nor were the T_{80} values (77 ± 10 and 101 ± 12 s, respectively). (b) E_2 (20 pmoles) was locally applied with or without ER antagonists and the effect of fluvoxamine was measured at 0 (before E_2 administration) or 10 or 40–60 min after local application of E_2 . Pretreatment with the ER antagonists ICI 182,780 or tamoxifen (60 pmoles) was performed 10 min before local application of E_2 . Bars and brackets (a, b) represent the increase in T_{80} values (in seconds) after the application of fluvoxamine; \pm SEM ($n = 4-12$). * $p < 0.05$, paired *t*-test comparing each post- E_2 (or post-PBS) treatment with the corresponding pretreatment value (ie, time 0 value). ** $p < 0.05$, Newman-Keuls comparing fluvoxamine-induced increase in T_{80} values obtained with antagonist + E_2 with values obtained with E_2 alone at a similar time point.

over the time of the experiment in rats receiving the vehicle, that is, in these rats, fluvoxamine consistently increased T_{80} value by about 45–65 s over the 2 h that this experiment was carried out (Figure 5). However, 1 min after local application of E_2 at 57 pmoles, fluvoxamine's ability to prolong the clearance time of 5-HT (as shown by an increase in T_{80} value) was blocked completely. This effect of E_2 persisted for 60 min and then dissipated such that the effect of fluvoxamine was restored 90–120 min after administration




Figure 6 Effects of the E_2 -BSA conjugate on fluvoxamine's ability to increase 5-HT clearance time. The ability of fluvoxamine to increase T_{80} was examined at time 0 (before application of E_2 -BSA or BSA alone) or 10 or 40–60 min after local application of BSA or E_2 -BSA (calculated so as to contain 20 pmoles E_2). Bars and brackets represent the increase in T_{80} values (in seconds) after the application of fluvoxamine; \pm SEM ($n = 6-8$). * $p < 0.05$, paired *t*-test comparing the increase in T_{80} after fluvoxamine application for each post-drug treatment with the corresponding value before drug treatment.

of E_2 . The lower amount of E_2 (10 pmoles) had no effect on fluvoxamine's ability to inhibit 5-HT clearance at either early or later time points (Figure 5a). Preliminary experiments carried out with 20 pmoles of E_2 generated data similar to that seen with 57 pmoles (data not shown).

To determine whether this inhibitory effect of local application of E_2 is mediated by membrane or intracellular ERs, two ER antagonists (tamoxifen and ICI 182,780; Xiao *et al*, 2003) were tested. In addition, the effect of a membrane impermeable estradiol-BSA conjugate (E_2 -BSA) (Blackmore and Lattanzio, 1991; Ramirez and Zheng, 1996) was measured as well. The drugs were given 10 min prior to addition of E_2 and their effects were measured either shortly (10 min) or at later times (40–60 min) after administration of E_2 or E_2 -BSA. E_2 was used at 20 pmol in these experiments, as it was the minimum effective dose for both rapid (1–10 min) and later developing effects (30–60 min). As expected, E_2 alone blocked the ability of fluvoxamine to induce an increase in the clearance time for 5-HT at 10 min as well as at 40–60 min post- E_2 (Figure 5b). Both ICI 182,780 and tamoxifen blocked this inhibitory effect of E_2 at 10 min. The effect of ICI 182,780 persisted at 40–60 min, whereas the inhibitory effect of tamoxifen was no longer observed at this time (Figure 5b). Local application of E_2 -BSA abolished the fluvoxamine-induced increase in the T_{80} value only at the early time point (10 min after E_2 -BSA application) and not at the later time, 40–60 min (Figure 6). BSA alone had no effect on the ability of fluvoxamine to increase T_{80} value (Figure 6).

DISCUSSION

These results show that the ability of fluvoxamine to inhibit clearance time of 5-HT *in vivo* is markedly diminished in female rats during proestrus. This is not due to proestrus

lowering the affinity of fluvoxamine for the SERT. This effect of proestrus seems to be due either to estrogen and/or progesterone, as administration of either hormone to OVX rats, but not to castrated male rats, produced effects similar to those seen in proestrus. In addition, estradiol alone altered 5-HT clearance rate. Results obtained from local application of E_2 , E_2 -BSA and ER antagonists directly into the CA3 region of the hippocampus indicates that both intracellular as well as membrane ERs may be involved in the ability of E_2 to inhibit the effect of fluvoxamine.

The hippocampus is an appropriate area to study the interaction between ovarian steroids and SSRIs on SERT function. There are considerable data demonstrating a role for the hippocampus in the pathophysiology of depression (Campbell and Macqueen, 2004), particularly with respect to changes in hippocampal volume (Campbell *et al*, 2004). The hippocampus receives a dense serotonergic innervation (Jacobs and Azmitia, 1992) and it is the initial increase in either 5-HT and/or NE caused by ADs that is thought to trigger changes responsible for their beneficial clinical effects (Delgado *et al*, 1999; Miller *et al*, 1996). The increase in 5-HT is due to the inhibition of the SERT, the initial cellular target for SSRIs (Lenox and Frazer, 2002). Finally, the hippocampus is an estrogen responsive area in the brain, with humans containing mRNAs for both $ER\alpha$ and $ER\beta$ (Osterlund *et al*, 2000a, b).

The detection of neurotransmitter injected into brain by a sensor located some distance from the site of injection is a complex phenomenon. Factors that influence the shape of the signals detected include not only uptake of the transmitter, but also its diffusion in obstructed extracellular space (Nicholson and Phillips, 1981), as well as, perhaps, metabolism. The clearance time parameter (T_{80}) used to quantitatively analyze the serotonin electrochemical signal was selected because it is known to reflect primarily the uptake process rather than metabolism or diffusion and also it clearly and reproducibly detects effects of locally applied competitive uptake inhibitors (Benmansour *et al*, 1999, 2002; Cass *et al*, 1993; Hoffman and Gerhardt, 1998; Zahniser *et al*, 1999). Whereas local application of uptake inhibitors have consistent and significant effects on clearance time parameters, they do not consistently alter clearance rates as measured using *in vivo* chronoamperometry, even though it might be expected that they would. In a very careful study, the clearance of dopamine by the dopamine transporter (DAT) was shown to be altered by local application of inhibitors of the DAT, with the effect being dependent on the initial basal amplitude of the DA signal (Zahniser *et al*, 1999). Perhaps not surprisingly, the most pronounced increase in T_{80} caused by uptake inhibitors was found at the lowest basal amplitude. The effects of DAT inhibitors on clearance rate were less consistent, but there were trends for clearance rate to increase in the low basal amplitude group, to decrease in the high basal amplitude group, and not to change in the medium basal amplitude group. None of these effects were statistically significant. Thus, although one might expect clearance rate to change, it does not after local application of uptake inhibitors. Why this is so is not clear. With local application of drugs, it is difficult to predict the concentration achieved and consequently the occupancy of the SERT; this is especially so, as local application of drugs does not

allow steady state to develop. Further, we are not using concentrations of 5-HT that saturate the SERT (Daws *et al*, 2005). It is well established that local application of SSRIs raises extracellular 5-HT (eg, Kreiss and Lucki, 1995). Consequently, when sub-saturating concentrations of both 5-HT and SSRIs are present, the rate of uptake of 5-HT by SERT not occupied by SSRIs would be expected to increase due to elevated 5-HT. Thus an expected decrease in clearance rate by SERT that is occupied by SSRIs may be compensated by an increase in clearance rate by those not occupied.

During proestrus or following treatment of OVX rats with estradiol or progesterone either alone or in combination, fluvoxamine no longer increased clearance time for exogenously administered 5-HT (Figures 2 and 4). These hormonal effects could possibly reflect a conformational change in the transporter that reduces the accessibility of inhibitors to the binding site. The clearance time parameter, T_{80} , was increased after treatment with EB (Figure 4). This effect was similar to that obtained after local application of fluvoxamine in OVX-control rats. Such an effect of estradiol might account, at least in part, for the observation that depression may be more common when estradiol levels are low (Fink *et al*, 1996; Halbreich *et al*, 1992; Pearlstein, 1995) and that estrogen can have AD effects (Zweifel and O'Brien, 1997; Soares *et al*, 2001; Huttner and Shepherd, 2003). However, it is difficult to reconcile this effect of estrogen with the higher incidence of depression in women (Kessler, 2003). Further adding estrogen therapy to SSRI treatment may be beneficial in depressed menopausal women (Yonkers, 2003). Yet our data shows that estrogen, at least when given acutely, inhibits the inhibitory effect of SSRIs on SERT function, which is not consistent with such a clinical effect. Further research examining the effect of more chronic treatment with hormones on SERT function and the effect of SSRIs on such function should help to clarify such issues.

Treatment with EB not only increased clearance time parameters, but also reduced the clearance rate for 5-HT (Figure 4). Estrogen can act at several levels of the serotonergic system. For example, it can inhibit reuptake by monoaminergic transporters *in vitro* (Michel *et al*, 1987). Also, estrogen has been shown to interact with either 5-HT_{1A} or 5HT_{2A} receptors in several brain areas (Fink *et al*, 1996; Osterlund *et al*, 1999; Raap *et al*, 2000). The present study shows that estrogen not only blocked the ability of an SSRI to induce prolongation of the 5-HT signal, but also decreased SERT function (Figure 4). These changes could be interpreted as a result of an EB-induced change in the affinity of the SERT for 5-HT. EB could have induced a change in the phosphorylation state of the SERT and/or changes in protein association (ie, syntaxin 1A, PP2Ac) that could support changes in SERT trafficking and catalytic modulation (Blakely *et al*, 2005). On the other hand, EB treatment could have induced changes in the characteristics of the extracellular space (ie, tortuosity) such that pressure-ejected 5-HT is 'moving slower' both to reach the electrode tip and to be removed from it. This is, however, unlikely, since the rise time of the 5-HT signal was not increased in EB-treated OVX rats (data not shown). Furthermore, the experiment carried out with treated with EB castrated rats showing blockade of 5-HT clearance after fluvoxamine

application is yet another argument in favor of the specificity of the effect of hormones.

Gender differences exist in many of the actions of estrogen in the brain, and the process of sexual differentiation appears to affect many brain regions outside of the traditional brain areas involved in the reproductive function (McEwen, 2001). Our data show that treatment of castrated rats with the combination EB/P did not alter the ability of fluvoxamine to increase the T_{80} value of 5-HT clearance, whereas treatment of OVX rats with EB/P blocked fluvoxamine induced increase in T_{80} . Thus, the modulations by ovarian hormones on the SSRI-SERT interaction are gender specific. It was shown that estradiol was able to affect SERT expression in female as well as male rat brain (McQueen *et al*, 1999). It is possible that the differences are due to the treatment protocol (hormonal treatment 2 weeks after gonadectomy as opposed to treatment immediately after gonadectomy, as well as time of hormonal acute exposure (72 h as opposed to 36 h)).

Our data indicate that progesterone, similar to estrogen, blocked fluvoxamine's ability to prolong the 5-HT signal (Figure 4). This finding is consistent with other studies showing that not only estradiol but also progesterone can alter serotonergic neurotransmission. A stimulatory role for progesterone in the regulation of 5-HT synthesis has been demonstrated (Jahn and Deis, 1987), even though progesterone appears to have no effect on the translation of the serotonin-synthesizing enzyme, tryptophan hydroxylase (Betha *et al*, 2000). Our data show that progesterone, similar to estrogen, interferes with the ability of an SSRI to block SERT function. Progesterone however had no effect on basal SERT function.

The analysis of SERT binding site densities indicates no change during the rat estrous cycle (Table 1). [3 H]-CN-IMI is a lipophilic ligand and could label not only membrane SERT but intracellular SERT as well. The number of SERTs located on the plasma membrane is controlled through kinase- and phosphatase-linked pathways, as shown in transfected cells (Qian *et al*, 1997; Ramamoorthy and Blakely, 1999; Ramamoorthy *et al*, 1998). Recently, in an elegant study using surface biotinylation of purified synaptosomes from rat midbrain, a reduction of the fraction of cell-surface expression of SERT protein in the plasma membrane of synaptosomes following either protein kinase C activation or p38 mitogen-activated protein kinase was demonstrated (Samuel *et al*, 2005). Therefore, a redistribution of SERT from the intracellular pool to the membrane surface could result from treatment with EB. This hypothesis remains to be tested.

Using *in vivo* voltammetry, it was shown that local application of E_2 into the nucleus accumbens of OVX rats induced rapid modulation of dopamine activity (Thompson and Moss, 1994). Consistent with this, the time course of the effect of locally applied E_2 into the CA3 region of the hippocampus of OVX rats showed that fluvoxamine's ability to block the clearance of 5-HT was abolished as early as 1–10 min after E_2 application as well as at later time points (20–60 min) (Figure 5). Data obtained after local application of the membrane-impermeable E_2 -BSA conjugate provide further support for the involvement of membrane ERs, since it mimicked only the early effects obtained after E_2 application (Figure 6). Modulation of fluvoxamine's ability

to block 5-HT clearance by E_2 was blocked by ER antagonists. ICI 182,780 blocked the effects of E_2 at early as well as at later time points, whereas tamoxifen blocked only the early effect of E_2 . It is possible that tamoxifen was removed from the hippocampus faster than ICI 182,780, or that the concentration used did not antagonize intracellular ER. These data indicate that the E_2 -induced blockade of fluvoxamine's effects on SERT is likely mediated via genomic as well as non-genomic mechanisms.

In conclusion, the main finding of the present study demonstrates that acute treatment with either EB or P impacts the ability of SSRIs to alter the function of what is widely considered their initial cellular target—the SERT. There is at least one interesting potential clinical implication of these data. Although pharmacotherapy of depression with SSRIs or other drugs is more effective than treatment with placebo, 50–70% of patients have less than optimal to no response to an initial trial with an AD (Fava *et al*, 2006; Rush *et al*, 2006; Trivedi *et al*, 2006). If hormones such as estrogen or progesterone reduce the ability of SSRIs to block the SERT, then some patients may be, for genetic reasons, unusually sensitive to such hormonal effects, which could contribute to non-response.

ACKNOWLEDGEMENTS

This work was supported by funds from NARSAD and the Department of Veterans Affairs. We thank Dr Andrea Gore for help with analysis of hormones levels and Kelly Sharp and Dr Georgianna Gould for technical assistance.

DISCLOSURE/CONFLICT OF INTEREST

Dr Benmansour, Mr Piotrowski, and Mr Altamirano report no biomedical financial interests or potential conflicts of interest. Dr Frazer reports that he has been on advisory boards for Cyberonics Inc. and H Lundbeck A/S and that he has consulted and/or received research support for preclinical studies from Forest Research Institute, Eli Lilly and Company, and Wyeth Pharmaceuticals, but no support for this study was received by any pharmaceutical company.

REFERENCES

- Adams MM, Morrison JH, Gore AC (2001). *N*-methyl-D-aspartate receptor mRNA levels change during reproductive senescence in the hippocampus of female rats. *Exp Neurol* 170: 171–179.
- Attali G, Weizman A, Gil-Ad I, Rehavi M (1997). Opposite modulatory effects of ovarian hormones on rat brain dopamine and serotonin transporters. *Brain Res* 756: 153–159.
- Benmansour S, Cecchi M, Morilak DA, Gerhardt GA, Javors MA, Gould GG *et al* (1999). Effects of chronic antidepressant treatments on serotonin transporter function, density, and mRNA level. *J Neurosci* 19: 10494–10501.
- Benmansour S, Owens WA, Cecchi M, Morilak DA, Frazer A (2002). Serotonin clearance *in vivo* is altered to a greater extent by antidepressant-induced downregulation of the serotonin transporter than by acute blockade of this transporter. *J Neurosci* 22: 6766–6772.
- Betha CL, Gundlach C, Mirkes SJ (2000). Ovarian steroid action in the serotonin neural system of macaques. *Novartis Found Symp* 230: 112–130, discussion 130–133.

- Bethea CL, Pecins-Thompson M, Schutzer WE, Gundlach C, Lu ZN (1998). Ovarian steroids and serotonin neural function. *Mol Neurobiol* **182**: 87–123.
- Blackmore PF, Lattanzio FA (1991). Cell surface localization of a novel non-genomic progesterone receptor on the head of human sperm. *Biochem Biophys Res Commun* **1811**: 331–336.
- Blakely RD, Defelice LJ, Galli A (2005). Biogenic amine neurotransmitter transporters: just when you thought you knew them. *Physiology (Bethesda)* **20**: 225–231.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Anal Biochem* **72**: 248–254.
- Burt VK, Stein K (2002). Epidemiology of depression throughout the female life cycle. *J Clin Psychiatry* **63**(Suppl 7): 9–15.
- Campbell S, MacQueen G (2004). The role of the hippocampus in the pathophysiology of major depression. *J Psychiatry Neurosci* **296**: 417–426.
- Campbell S, Marriott M, Nahmias C, MacQueen GM (2004). Lower hippocampal volume in patients suffering from depression: a meta-analysis. *Am J Psychiatry* **1614**: 598–607.
- Cass WA, Zahniser NR, Flach KA, Gerhardt GA (1993). Clearance of exogenous dopamine in rat dorsal striatum and nucleus accumbens: role of metabolism and effects of locally applied uptake inhibitors. *J Neurochem* **616**: 2269–2278.
- D'Amato RJ, Largent BL, Snowman AM, Snyder SH (1987). Selective labeling of serotonin uptake sites in rat brain by [³H]citalopram contrasted to labeling of multiple sites by [³H]imipramine. *J Pharmacol Exp Ther* **2421**: 364–371.
- Daws LC, Montanez S, Owens WA, Gould GG, Frazer A, Toney GM et al (2005). Transport mechanisms governing serotonin clearance *in vivo* revealed by high-speed chronoamperometry. *J Neurosci Methods* **143**: 49–62.
- Delgado PL, Miller HL, Salomon RM, Licinio J, Krystal JH, Moreno FA et al (1999). Tryptophan-depletion challenge in depressed patients treated with desipramine or fluoxetine: implications for the role of serotonin in the mechanism of antidepressant action. *Biol Psychiatry* **462**: 212–220.
- Fava M, Rush AJ, Wisniewski SR, Nierenberg AA, Alpert JE, McGrath PJ et al (2006). A comparison of mirtazapine and nortriptyline following two consecutive failed medication treatments for depressed outpatients: a STAR*D report. *Am J Psychiatry* **1637**: 1161–1172.
- Fink G, Sumner BE, Rosie R, Grace O, Quinn JP (1996). Estrogen control of central neurotransmission: effect on mood, mental state, and memory. *Cell Mol Neurobiol* **163**: 325–344.
- Gibbs RB (1996). Fluctuations in relative levels of choline acetyltransferase mRNA in different regions of the rat basal forebrain across the estrous cycle: effects of estrogen and progesterone. *J Neurosci* **163**: 1049–1055.
- Gore AC, Yeung G, Morrison JH, Oung T (2000). Neuroendocrine aging in the female rat: the changing relationship of hypothalamic gonadotropin-releasing hormone neurons and N-methyl-D-aspartate receptors. *Endocrinology* **14112**: 4757–4767.
- Halbreich U, Piletz J, Halaris A (1992). Influence of gonadal hormones on neurotransmitters, receptor, cognition and mood. *Clin Neuropharmacol* **15**(Suppl 1 Part A): 590A–591A.
- Hoffman AF, Gerhardt GA (1998). *In vivo* electrochemical studies of dopamine clearance in the rat substantia nigra: effects of locally applied uptake inhibitors and unilateral 6-hydroxydopamine lesions. *J Neurochem* **701**: 179–189.
- Huttner RP, Shepherd JE (2003). Gonadal steroids, selective serotonin reuptake inhibitors, and mood disorders in women. *Med Clin North Am* **875**: 1065–1076.
- Jacobs BL, Azmitia EC (1992). Structure and function of the brain serotonin system. *Physiol Rev* **721**: 165–229.
- Jahn GA, Deis RP (1987). A possible dual regulation of prolactin release by the serotonergic system in rats at pro-oestrus and during late pregnancy: role of ovarian hormones. *J Endocrinol* **1123**: 367–374.
- Katzenellenbogen BS (2000). Mechanisms of action and cross-talk between estrogen receptor and progesterone receptor pathways. *J Soc Gynecol Investig* **71**(Suppl): S33–S37.
- Kelly MJ, Lagrange AH, Wagner EJ, Ronnekleiv OK (1999). Rapid effects of estrogen to modulate G protein-coupled receptors via activation of protein kinase A and protein kinase C pathways. *Steroids* **641–2**: 64–75.
- Kessler RC (2003). Epidemiology of women and depression. *J Affect Disord* **741**: 5–13.
- Kornstein SG, Schatzberg AF, Thase ME, Yonkers KA, McCullough JP, Keitner GI et al (2000). Gender differences in treatment response to sertraline vs imipramine in chronic depression. *Am J Psychiatry* **1579**: 1445–1452.
- Kovachich GB, Aronson CE, Brunswick DJ, Frazer A (1988). Quantitative autoradiography of serotonin uptake sites in rat brain using [³H]cyanoimipramine. *Brain Res* **4541–2**: 78–88.
- Krajnak K, Rosewell KL, Duncan MJ, Wise PM (2003). Aging, estradiol and time of day differentially affect serotonin transporter binding in the central nervous system of female rats. *Brain Res* **9901–2**: 87–94.
- Kreiss DS, Lucki I (1995). Effects of acute and repeated administration of antidepressant drugs on extracellular levels of 5-hydroxytryptamine measured *in vivo*. *J Pharmacol Exp Ther* **274**: 866–876.
- Lenox RH, Frazer A (2002). *Mechanism of Action of Antidepressants and Mood-Stabilizers*. Lippincott Williams & Wilkins: Philadelphia.
- McEwen BS (2001). Invited review: estrogens effect on the brain: multiple sites and molecular mechanisms. *J Appl Physiol* **91**: 2785–2801.
- McQueen JK, Wilson H, Fink G (1997). Estradiol-17 beta increases serotonin transporter (SERT) mRNA levels and the density of SERT-binding sites in female rat brain. *Brain Res Mol Brain Res* **451**: 13–23.
- McQueen JK, Wilson H, Sumner BE, Fink G (1999). Serotonin transporter (SERT) mRNA and binding site densities in male rat brain affected by sex steroids. *Brain Res Mol Brain Res* **63**: 241–247.
- Mendelson SD, McKittrick CR, McEwen BS (1993). Autoradiographic analyses of the effects of estradiol benzoate on [³H]paroxetine binding in the cerebral cortex and dorsal hippocampus of gonadectomized male and female rats. *Brain Res* **6011–2**: 299–302.
- Michel MC, Rother A, Hiemke C, Ghraf R (1987). Inhibition of synaptosomal high-affinity uptake of dopamine and serotonin by estrogen agonists and antagonists. *Biochem Pharmacol* **3619**: 3175–3180.
- Miller HL, Delgado PL, Salomon RM, Berman R, Krystal JH, Heninger GR et al (1996). Clinical and biochemical effects of catecholamine depletion on antidepressant-induced remission of depression. *Arch Gen Psychiatry* **532**: 117–128.
- Nadal A, Ropero AB, Laribi O, Maillet M, Fuentes E, Soria B (2000). Nongenomic actions of estrogens and xenoestrogens by binding at a plasma membrane receptor unrelated to estrogen receptor alpha and estrogen receptor beta. *Proc Natl Acad Sci USA* **9721**: 11603–11608.
- Nicholson C, Phillips JM (1981). Ion diffusion modified by tortuosity and volume fraction in the extracellular microenvironment of the rat cerebellum. *J Physiol* **321**: 225–257.
- Osterlund MK, Grandien K, Keller E, Hurd YL (2000a). The human brain has distinct regional expression patterns of estrogen receptor alpha mRNA isoforms derived from alternative promoters. *J Neurochem* **754**: 1390–1397.
- Osterlund MK, Gustafsson JA, Keller E, Hurd YL (2000b). Estrogen receptor beta (ERbeta) messenger ribonucleic acid (mRNA) expression within the human forebrain: distinct distribution pattern to ERalpha mRNA. *J Clin Endocrinol Metab* **8510**: 3840–3846.

- Osterlund MK, Overstreet DH, Hurd YL (1999). The flinders sensitive line rats, a genetic model of depression, show abnormal serotonin receptor mRNA expression in the brain that is reversed by 17beta-estradiol. *Brain Res Mol Brain Res* 741-2: 158-166.
- Paxinos G, Watson C (1986). *The Rat Brain in Stereotaxic Coordinates*. Academic: New York, NY.
- Pearlstein TB (1995). Hormones and depression: what are the facts about premenstrual syndrome, menopause, and hormone replacement therapy? *Am J Obstet Gynecol* 1732: 646-653.
- Pecins-Thompson M, Brown NA, Bethea CL (1998). Regulation of serotonin re-uptake transporter mRNA expression by ovarian steroids in rhesus macaques. *Brain Res Mol Brain Res* 531-2: 120-129.
- Qian Y, Galli A, Ramamoorthy S, Risso S, DeFelice LJ, Blakely RD (1997). Protein kinase C activation regulates human serotonin transporters in HEK-293 cells via altered cell surface expression. *J Neurosci* 171: 45-57.
- Quitkin FM, Stewart JW, McGrath PJ, Taylor BP, Tisminetzky MS, Petkova E et al (2002). Are there differences between women's and men's antidepressant responses? *Am J Psychiatry* 15911: 1848-1854.
- Raap DK, DonCarlos L, Garcia F, Muma NA, Wolf WA, Battaglia G et al (2000). Estrogen desensitizes 5-HT(1A) receptors and reduces levels of G(z), G(i1) and G(i3) proteins in the hypothalamus. *Neuropharmacology* 3910: 1823-1832.
- Ramamoorthy S, Blakely RD (1999). Phosphorylation and sequestration of serotonin transporters differentially modulated by psychostimulants. *Science* 285: 763-766.
- Ramamoorthy S, Melikian HE, Qian Y, Blakely RD (1998). Biosynthesis, N-glycosylation, and surface trafficking of biogenic amine transporter proteins. *Methods Enzymol* 296: 347-370.
- Ramirez VD, Zheng J (1996). Membrane sex-steroid receptors in the brain. *Front Neuroendocrinol* 174: 402-439.
- Rush AJ, Trivedi MH, Wisniewski SR, Nierenberg AA, Stewart JW, Warden D et al (2006). Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry* 163: 1905-1917.
- Saletu B, Brandstatter N, Metka M, Stamenkovic M, Anderer P, Semlitsch HV et al (1995). Double-blind, placebo-controlled, hormonal, syndromal and EEG mapping studies with transdermal oestradiol therapy in menopausal depression. *Psychopharmacology (Berl)* 122: 321-329.
- Samuvel DJ, Jayanthi LD, Bhat NR, Ramamoorthy S (2005). A role for p38 mitogen-activated protein kinase in the regulation of the serotonin transporter: evidence for distinct cellular mechanisms involved in transporter surface expression. *J Neurosci* 251: 29-41.
- Scheibe S, Preuschhof C, Cristi C, Bagby RM (2003). Are there gender differences in major depression and its response to antidepressants? *J Affect Disord* 753: 223-235.
- Soares CN, Almeida OP, Joffe H, Cohen LS (2001). Efficacy of estradiol for the treatment of depressive disorders in perimenopausal women: a double-blind, randomized, placebo-controlled trial. *Arch Gen Psychiatry* 58: 529-534.
- Stevis PE, Deecher DC, Suhadolnik L, Mallis LM, Frail DE (1999). Differential effects of estradiol and estradiol-BSA conjugates. *Endocrinology* 140: 5455-5458.
- Thompson TL, Moss RL (1994). Estrogen regulation of dopamine release in the nucleus accumbens: genomic- and nongenomic-mediated effects. *J Neurochem* 625: 1750-1756.
- Trivedi MH, Fava M, Wisniewski SR, Thase ME, Quitkin F, Warden D et al (2006). Medication augmentation after the failure of SSRIs for depression. *N Engl J Med* 354: 1243-1252.
- Xiao L, Jackson LR, Becker JB (2003). The effect of estradiol in the striatum is blocked by ICI 182,780 but not tamoxifen: pharmacological and behavioral evidence. *Neuroendocrinology* 774: 239-245.
- Yonkers KA (2003). Special issues related to the treatment of depression in women. *J Clin Psychiatry* 64(Suppl 18): 8-13.
- Zahniser NR, Larson GA, Gerhardt GA (1999). *In vivo* dopamine clearance rate in rat striatum: regulation by extracellular dopamine concentration and dopamine transporter inhibitors. *J Pharmacol Exp Ther* 2891: 266-277.
- Zhou W, Koldzic-Zivanovic N, Clarke CH, de Beun R, Wassermann K, Bury PS et al (2002). Selective estrogen receptor modulator effects in the rat brain. *Neuroendocrinology* 751: 24-33.
- Zweifel JE, O'Brien WH (1997). A meta-analysis of the effect of hormone replacement therapy upon depressed mood. *Psychoneuroendocrinology* 22: 189-212.