

Sex-Dependent Antipsychotic Capacity of 17 β -Estradiol in the Latent Inhibition Model: A Typical Antipsychotic Drug in Both Sexes, Atypical Antipsychotic Drug in Males

Michal Arad*¹ and Ina Weiner¹

¹Department of Psychology, Tel-Aviv University, Tel-Aviv, Israel

The estrogen hypothesis of schizophrenia suggests that estrogen is a natural neuroprotector in women and that exogenous estrogen may have antipsychotic potential, but results of clinical studies have been inconsistent. We have recently shown using the latent inhibition (LI) model of schizophrenia that 17 β -estradiol exerts antipsychotic activity in ovariectomized (OVX) rats. The present study sought to extend the characterization of the antipsychotic action of 17 β -estradiol (10, 50 and 150 μ g/kg) by testing its capacity to reverse amphetamine- and MK-801-induced LI aberrations in gonadally intact female and male rats. No-drug controls of both sexes showed LI, ie, reduced efficacy of a previously non-reinforced stimulus to gain behavioral control when paired with reinforcement, if conditioned with two but not five tone-shock pairings. In both sexes, amphetamine (1 mg/kg) and MK-801 (50 μ g/kg) produced disruption (under weak conditioning) and persistence (under strong conditioning) of LI, modeling positive and negative/cognitive symptoms, respectively. 17 β -estradiol at 50 and 150 μ g/kg potentiated LI under strong conditioning and reversed amphetamine-induced LI disruption in both males and females, mimicking the action of typical and atypical antipsychotic drugs (APDs) in the LI model. 17 β -estradiol also reversed MK-induced persistent LI, an effect mimicking atypical APDs and NMDA receptor enhancers, but this effect was observed in males and OVX females but not in intact females. These findings indicate that in the LI model, 17 β -estradiol exerts a clear-cut antipsychotic activity in both sexes and, remarkably, is more efficacious in males and OVX females where it also exerts activity considered predictive of anti-negative/cognitive symptoms.

Neuropsychopharmacology (2010) **35**, 2179–2192; doi:10.1038/npp.2010.89; published online 7 July 2010

Keywords: amphetamine; antipsychotic drugs; MK-801; estrogen; latent inhibition; schizophrenia

INTRODUCTION

Epidemiological and life cycle studies have indicated that women with schizophrenia have a more favorable illness course than men during reproductive years, characterized by later onset of symptoms, lower symptom severity and better response to antipsychotic drugs (APDs); however, the menopausal period is associated with increased symptom severity and reduced sensitivity to treatment (Hafner, 2003; Hafner *et al*, 1989; Kulkarni, 2009; Kulkarni *et al*, 1996, 2008b; Riecher-Rossler and de Geyter, 2007; Seeman and Lang, 1990). This has led to the suggestion that exogenous estrogen may have therapeutic potential in women with schizophrenia, but results of clinical studies have been inconclusive (Akhondzadeh *et al*, 2003; Korhonen *et al*, 1995; Kulkarni *et al*, 2008a; Kulkarni *et al*, 1996; Kulkarni

et al, 2008b; Kulkarni *et al*, 2001; Mortimer, 2007; Riecher-Rossler, 2002; Riecher-Rossler and de Geyter, 2007; Riecher-Rossler and Hafner, 1993; Riecher-Rossler *et al*, 1994). Interestingly, estradiol improved psychotic symptoms also in schizophrenic men (Kulkarni, 2009).

Support for the antipsychotic action of estrogen can be derived from animal studies showing that 17 β -estradiol blocks/reduces the behavioral response to the pro-psychotic dopaminergic drugs, amphetamine and cocaine, (Becker and Beer, 1986; Becker and Rudick, 1999; Bedard *et al*, 1983; Bedard *et al*, 1978; Gordon and Diamond, 1981; Hafner *et al*, 1991; Naik *et al*, 1978; Segarra *et al*, 2009) and potentiates APD-induced catalepsy (Bedard *et al*, 1982; Chiodo *et al*, 1979; De Ryck *et al*, 1982; Di Paolo *et al*, 1979; Nicoletti *et al*, 1983; Palermo-Neto and Dorce, 1990). 17 β -estradiol has been also shown to improve cognitive performance in humans and rodents of both sexes (Barnes *et al*, 2006; Daniel and Bohacek, 2010; Frick, 2009; Gogos *et al*, 2006; Kitamura *et al*, 2009; Packard *et al*, 1996; Sherwin *et al*, 2009; Soderstrom *et al*, 2009). Such action would be beneficial in schizophrenia which is characterized by profound cognitive deficits (Barch and Carter, 2008).

*Correspondence: M Arad, Department of Psychiatry, University of Maryland School of Medicine, Room 916 MSTF, 685 West Baltimore Street, Baltimore, MD 21201, USA, Tel: +410 706 6799, Fax: +410 706 4002, E-mail: marad@psych.umaryland.edu
Received 10 March 2010; revised 25 May 2010; accepted 27 May 2010

To date, the study of the antipsychotic capacity of estrogen in formal animal models of schizophrenia has been limited (Chavez *et al*, 2009; Gogos *et al*, 2010; Hafner *et al*, 1991; Sutcliffe *et al*, 2008; Van den Buuse and Eikelis, 2001).

We have recently provided evidence for antipsychotic properties of estrogen using the latent inhibition (LI) model of schizophrenia (Arad and Weiner 2008, 2009, 2010). LI is one of the best-documented cross-species manifestations of attentional selectivity in associative learning (Lubow, 1989; Mackintosh, 1975), whereby repeated non-reinforced pre-exposure to the to-be-conditioned stimulus interferes with its subsequent efficacy to generate conditioned response. Amphetamine-induced disruption of LI is considered to model the inability to ignore irrelevant stimuli associated with positive symptoms of schizophrenia (Weiner, 2003). Conversely, rodents treated with NMDA receptor antagonists such as MK-801 or PCP that produce and exacerbate negative symptoms and cognitive deficits (Javitt and Zukin, 1991; Krystal *et al*, 2003), persist in expressing LI under conditions that prevent/reduce LI expression in untreated rats (Gaisler-Salomon *et al*, 2008; Gaisler-Salomon and Weiner, 2003; Lipina *et al*, 2005; Palsson *et al*, 2005). Thus, persistent LI induced by MK-801 has been suggested to model attentional perseveration associated with negative/cognitive symptoms of schizophrenia (Gaisler-Salomon *et al*, 2008; Weiner and Arad 2009). Both typical and atypical APDs restore LI in amphetamine-treated rats, and this is paralleled by their capacity to restore LI in naive animals under conditions that do not yield robust LI in no-drug controls (Weiner and Arad 2009). MK-801-induced persistent LI is reversed by atypical but not typical APDs, as well as by glycinergic NMDA enhancers, consistent with the differential efficacy of these treatments in improving negative/cognitive symptoms (Harvey *et al*, 2005; Heresco-Levy *et al*, 2005).

In support of the antipsychotic action of 17 β -estradiol, we (Arad and Weiner 2008, 2009, 2010) showed that (a) disruption and restoration of LI are associated with low and high levels of endogenous estrogen; (b) behaviorally inactive dose of 17 β -estradiol restores the capacity of ineffective APD doses to block amphetamine-induced LI disruption; and (c) 17 β -estradiol given on its own reverses amphetamine-induced LI disruption in ovariectomized (OVX) as well as in sham-operated female rats. Our aim here was to extend the characterization of the antipsychotic action of 17 β -estradiol by testing its capacity to reverse amphetamine-induced LI disruption and MK-801-induced LI persistence in gonadally intact female and male rats. In addition, we tested whether 17 β -estradiol on its own potentiates LI.

MATERIALS AND METHODS

Animals

Female and male Wistar rats, 3-month old, bred in our laboratory were housed by sex, four per cage under reversed cycle lighting (lights on: 07:00–19:00 h) with ad lib access to food and water except for the duration of the LI experiments. All experimental protocols conformed to the guidelines of the Institutional Animal Care and Use Committee of Tel-Aviv University, Israel, and to the

guidelines of the NIH (animal welfare assurance number A5010–01, expires on 30 September 2011). All efforts were made to minimize the number of animals used and their suffering.

Ovariectomy (OVX)

Female rats were bilaterally ovariectomized under isoflurane (Nichols Piramal, UK) anesthesia. After shaving the abdominal area, a midline incision was made through the skin and muscle layer. Fallopian tubes were ligated by a nylon thread, after which the ovaries were carefully removed. Sutures of muscle layer and skin were removed 10 days later. Rats were allowed additional 3 weeks of recovery after removal of the sutures before the beginning of water restriction (see below). Within the 3-week recovery period, about a week after removal of sutures, vaginal smears were collected daily in the morning for 8 days in sham and OVX rats, to confirm presence or discontinuation of estrous cycle, respectively. Phases of the estrous cycle were determined by the morphology of cells in the vaginal smear under a light microscope (Marcondes *et al*, 2002). Sham-operated female and male controls (in experiment 4) underwent an identical surgical procedure without ovaries' removal. Sham females with two regular 4-day cycles in succession and OVX rats without estrous cycle were used for behavioral testing.

Apparatus and Procedure

LI was measured in a thirst-motivated conditioned emotional response procedure in Campden Instruments rodent test chambers with a retractable bottle, each enclosed in a ventilated sound-attenuating chest. When the bottle was not present, the hole was covered with a metal lid. The pre-exposed (PE) to-be-conditioned stimulus was a 10 s, 80 dB, 2.8 kHz tone produced by a Sonalert module. Shock was supplied through the floor by a Campden Instruments shock generator and shock scrambler set at 0.5 mA intensity and 1 s duration. Licks were detected by a Campden Instruments drinkometer. Equipment programming and data recording were controlled by the computer.

At 10 days before the beginning of the LI procedure, rats were put on a 23 h water restriction schedule and handled for about 2 min daily for 5 days. On the next 5 days, rats were trained to drink in the experimental chamber for 15–20 min/day. Water in the test apparatus was given in addition to the daily ration of 1 h given in the home cages. The LI procedure was conducted on days 11–14 and consisted of four stages given 24 h apart:

Pre-exposure. With the bottle removed, the PE rats received 40 tone presentations with an inter-stimulus interval of 40 s. The non-pre-exposed (NPE) rats were confined to the chamber for an identical period of time without receiving the tone.

Conditioning. With the bottle removed, rats received two (weak conditioning) or five (strong conditioning) tone-shock pairings given 5 min apart. Shock immediately followed tone termination. Weak conditioning produces LI in non-treated controls and thus allows the demonstration of treatment-induced LI disruption. This level of conditioning was therefore used with amphetamine (experiments 5 and 6).

Conversely, strong conditioning prevents LI in non-treated controls and thus allows the demonstration of treatment-induced abnormally persistent LI. This level of conditioning was used with MK-801 (experiments 7, 8, and 9). Both levels were used with 17 β -estradiol administration on its own (experiments 1, 2, 3, and 4) to determine if 17 β -estradiol disrupts and/or potentiates LI.

Rebaseline. Rats were given a 15 min drinking session as in initial training.

Test. Each rat was placed in the chamber and allowed to drink from the bottle. When the rat completed 75 licks the tone was presented for 5 min. The following times were recorded: Time to first lick, time to complete 1–50 licks, time to complete 51–75 licks (before the tone onset), and time to complete 76–100 licks (after the tone onset). Times to complete 76–100 licks were submitted to logarithmic transformation to allow parametric ANOVA. Longer log times indicate stronger suppression of drinking. LI is defined as significantly shorter log times to complete 76–100 licks of the PE compared with NPE rats.

Drug and Hormone Administration

Amphetamine, MK-801, and 17 β -estradiol were administered in a volume of 1 ml/kg. Amphetamine (Sigma, Switzerland) was dissolved in saline and administered i.p. at a dose of 1 mg/kg. MK-801 (dizocilpine; Merck Research Laboratories, USA) was dissolved in saline and administered i.p. at a dose of 50 μ g/kg. These doses are routinely used in our LI studies (Arad and Weiner 2010; Barak *et al*, 2009; Black *et al*, 2008). 17 β -estradiol (Sigma, Israel) was dissolved in corn oil and administered s.c. at doses of 10, 50, and 150 μ g/kg. These doses were used in our previous studies; they mimic low hormonal levels in female rats as seen during metestrus–diestrus, high levels as seen during proestrus–estrus, and very high levels as seen during the last trimester of pregnancy (Arad and Weiner, 2009, 2010; Galea *et al*, 2001; Gibbs *et al*, 1998; Nofrey *et al*, 2008; Van den Buuse and Eikelis, 2001; Walf and Frye, 2009). All compounds were administered before pre-exposure and conditioning sessions at 30 min interval for amphetamine and MK-801, and 120 min for 17 β -estradiol. No-drug controls received the corresponding vehicles. In all experiments, rebaseline and test sessions were conducted in a drug-free state.

Experimental Design

Experiments 1 and 2. Tested the effects of 17 β -estradiol on LI under weak conditioning (forty pre-exposures and two conditioning trials) in gonadally intact female and male rats. The experiments included 40 females (exp 1; n/group = 5) and 40 males (exp 2; n/group = 5) divided into eight experimental groups in a 2 \times 4 design with main factors of pre-exposure (0,40) and hormonal treatment (0, 10, 50, and 150 μ g/kg 17 β -estradiol).

Experiments 3 and 4. Tested the effects of 17 β -estradiol on LI under strong conditioning (forty pre-exposures and five conditioning trials) in gonadally intact female and male

rats. The experiments included 49 females (exp 3; n/group = 6–7) and 62 males (exp 4; n/group = 7–8) divided into eight experimental groups in a 2 \times 4 design with main factors of pre-exposure (0,40) and hormonal treatment (0, 10, 50, and 150 μ g/kg 17 β -estradiol).

Experiments 5 and 6. Tested the capacity of 17 β -estradiol to reverse amphetamine-induced LI disruption (under weak conditioning) in gonadally intact female and male rats. The experiments included 92 females (exp 5; n/group = 7–8) and 94 males (exp 6; n/group = 7–8) divided into twelve experimental groups in a 2 \times 2 \times 3 design with main factors of pre-exposure (0,40), pro-psychotic treatment (0 and 1 mg/kg amphetamine), and hormonal treatment (0, 50, and 150 μ g/kg 17 β -estradiol).

Experiments 7, 8, and 9. Tested the capacity of 17 β -estradiol to reverse MK-801-induced LI persistence (under strong conditioning) in gonadally intact female and male rats as well as in OVX female rats. The experiments included 73 intact females (exp 7; n/group = 6–7), 85 intact males (exp 8; n/group = 7–8), and 73 OVX females (exp 9; n/group = 6–7) divided into twelve experimental groups in a 2 \times 2 \times 3 design with main factors of pre-exposure (0,40), pro-schizophrenia treatment (0 and 50 μ g/kg MK-801), and hormonal treatment (0, 50, and 150 μ g/kg 17 β -estradiol).

Statistical Analysis

Time to complete 51–75 licks (before tone onset) and mean log times to complete 76–100 licks (after tone onset) were analyzed using two-way ANOVAs with main factors of pre-exposure and hormonal treatment (experiments 1, 2, 3, and 4) and three-way ANOVAs with main factors of pre-exposure, treatment, and hormonal treatment (experiments 5, 6, 7, 8, and 9). In cases of significant interactions involving the factor of pre-exposure, LSD *post-hoc* comparisons were used to assess the difference between the PE and NPE groups within each treatment condition.

It is important to mention that in all experiments, females and males were run in different systems to avoid potential interference. Thus, we could not analyze their data in the same ANOVA. However, given that we had four main aims in this study, we introduced the results as well as the figures under each aim as different parts of the same experiment to allow a visual comparison between the females' and males' data.

RESULTS

There were no differences between the experimental groups in the time required to complete 51–75 licks (A period; all *p*'s > 0.05) in any of the experiments (overall mean A periods were 6.65, 7.54, 10.25, 9.57, 7.13, 6.92, 7.02, 6.31, and 9.44 for experiments 1, 2, 3, 4, 5, 6, 7, 8, and 9, respectively).

Experiments 1 and 2: Effects of 17 β -Estradiol on LI Under Weak Conditioning (40 Pre-Exposures and Two Conditioning Trials) in Female and Male Rats

The two parts of Figure 1, a (exp 1) and b (exp 2), present the mean log times to complete 76–100 licks (after tone

onset) of the PE and NPE female (Figure 1a) and male (Figure 1b) rats treated with oil, 10, 50, or 150 μ g/kg 17 β -estradiol. As can be seen in both parts of Figure 1, oil-treated female and male rats showed LI, whereas rats treated with 10 μ g/kg of 17 β -estradiol did not show LI. The higher doses of 17 β -estradiol (50 and 150 μ g/kg) spared LI. ANOVAs yielded significant main effects of pre-exposure (females: $F_{(1,32)}=55.9$, $p<0.0001$; males: $F_{(1,32)}=45.4$, $p<0.0001$) and hormonal treatment (females: $F_{(3,32)}=3.0$, $p<0.05$; males: $F_{(3,32)}=3.5$, $p<0.05$), as well as a significant pre-exposure \times hormonal treatment interaction (females: $F_{(3,32)}=8.5$, $p<0.0005$; males: $F_{(3,32)}=8.6$, $p<0.0005$). *Post-hoc* comparisons for each of the analyses confirmed the presence of LI in female and male rats given oil, 50, or 150 μ g/kg 17 β -estradiol (p 's <0.01), but not after 10 μ g/kg 17 β -estradiol.

Experiments 3 and 4: Effects of 17 β -Estradiol on LI under strong Conditioning (40 Pre-Exposures and five Conditioning Trials) in Female and Male Rats

The two parts of Figure 2, a (exp 3) and b (exp 4), present the mean log times to complete 76–100 licks (after tone onset) of the PE and NPE female (Figure 2a) and male (Figure 2b) rats treated with oil, 10, 50, and 150 μ g/kg 17 β -estradiol. As can be seen, oil-treated female and male rats did not show LI as expected under strong conditioning. 10 μ g/kg of 17 β -estradiol had no effect on LI, whereas female and male rats treated with 50 and 150 μ g/kg of 17 β -estradiol persisted in showing LI. ANOVAs yielded a significant main effect of pre-exposure (females: $F_{(1,41)}=8.1$, $p<0.01$; males: $F_{(1,54)}=18.7$, $p<0.0001$), as well as a significant pre-exposure \times hormonal treatment interaction (females: $F_{(3,41)}=5.7$, $p<0.005$; males: $F_{(3,54)}=4.2$, $p<0.01$). *Post-hoc* comparisons for each of the analyses confirmed the presence of LI in female and male rats given 50 or 150 μ g/kg 17 β -estradiol (p 's <0.01), but not in other conditions.

Experiments 5 and 6: Effects of 17 β -Estradiol on Amphetamine-Induced LI Disruption Under Weak Conditioning (40 Pre-Exposures and two Conditioning Trials) in Female and Male Rats

The two parts of Figure 3, a (exp 5) and b (exp 6), present the mean log times to complete 76–100 licks (after tone onset) of the PE and NPE female (Figure 3a) and male (Figure 3b) rats treated with saline or amphetamine (1 mg/kg), and pre-treated with oil, 50, or 150 μ g/kg 17 β -estradiol. As can be seen, under weak conditioning, vehicle-injected female and male rats exhibited LI, whereas amphetamine-injected female and male rats did not exhibit LI. Both doses of 17 β -estradiol reversed amphetamine-induced LI disruption in females and males. On their own, both doses of 17 β -estradiol spared LI in females and males. ANOVA for females yielded significant main effects of pre-exposure ($F_{(1,80)}=98.9$, $p<0.0001$) and treatment ($F_{(1,80)}=7.2$, $p<0.01$), as well as a significant pre-exposure \times treatment \times hormonal treatment interaction ($F_{(2,80)}=3.2$, $p<0.05$). ANOVA for males yielded a significant main effect of pre-exposure ($F_{(1,82)}=55.9$, $p<0.0001$), as well as a significant pre-exposure \times treatment \times hormonal treatment interaction ($F_{(2,82)}=7.4$, $p<0.005$). *Post-hoc* comparisons for each of the analyses confirmed the presence of LI in saline-injected female and male rats pre-treated with 0, 50, or 150 μ g/kg 17 β -estradiol and in amphetamine-injected female and male rats pre-treated with 50 or 150 μ g/kg 17 β -estradiol (p 's <0.01), but not in the other conditions.

Experiments 7, 8, and 9: Effects of 17 β -Estradiol on MK-801-induced LI Persistence Under Strong Conditioning (40 Pre-Exposures and five Conditioning Trials) in Female, Male, and OVX Female Rats

The three parts of Figure 4, a (exp 7), b (exp 8), and c (exp 9), present the mean log times to complete 76–100 licks




Figure 1 Effects of 17 β -estradiol on LI under weak conditioning (forty pre-exposures and two conditioning trials) in female and male rats. Mean (\pm SEM) log time to complete 76–100 licks (after the tone onset) of the PE and the NPE female (a) and male (b) rats administered with 0, 10, 50, or 150 μ g/kg of 17 β -estradiol (oil, 10, 50, or 150 μ g/kg, respectively). Asterisks indicate a significant difference between the PE and NPE groups, namely, presence of LI (** $p<0.01$).




Figure 2 Effects of 17 β -estradiol on LI under strong conditioning (forty pre-exposures and five conditioning trials) in female and male rats. Mean (\pm SEM) log times to complete 76–100 licks (after the tone onset) of the PE and the NPE female (a) and male (b) rats administered with 0, 10, 50, or 150 μ g/kg of 17 β -estradiol (oil, 10, 50, or 150 μ g/kg, respectively). Asterisks indicate a significant difference between the PE and NPE groups, namely, presence of LI (** $p < 0.01$).




Figure 3 Effects of 17 β -estradiol on amphetamine (amph)-induced LI disruption under weak conditioning (forty pre-exposures and two conditioning trials) in female and male rats. Mean (\pm SEM) log times to complete 76–100 licks (after the tone onset) of the PE and the NPE saline- or amph-injected female (a) and male (b) rats, administered with 0, 50, or 150 μ g/kg of 17 β -estradiol (oil, E 50 μ g/kg, or E 150 μ g/kg, respectively). Asterisks indicate a significant difference between the PE and NPE groups, namely, presence of LI (** $p < 0.01$).

(after tone onset) of the PE and NPE female (Figure 4a), male (Figure 4b), and OVX female (Figure 4c) rats treated with saline or MK-801 (50 μ g/kg), and pre-treated with oil, 50, or 150 μ g/kg 17 β -estradiol. As expected with strong conditioning, LI was absent in vehicle-injected rats, whereas MK-801-injected rats persisted in showing LI. In MK-801-treated female rats, both doses of 17 β -estradiol were without any effect, whereas in males and OVX females, both doses reversed MK-801-induced LI persistence. On its own, 17 β -estradiol led to LI in gonadally intact female and male rats, but not in OVX female rats. ANOVA for female rats yielded significant main effects of pre-exposure ($F_{(1,61)} = 58.0$, $p < 0.0001$), treatment ($F_{(1,61)} = 17.7$, $p < 0.0001$), and hormonal treatment ($F_{(2,61)} = 5.6$, $p < 0.01$),

as well as a significant pre-exposure \times treatment \times hormonal treatment interaction ($F_{(2,61)} = 3.4$, $p < 0.05$). ANOVA for male rats yielded a significant main effect of pre-exposure ($F_{(1,73)} = 33.2$, $p < 0.0001$), as well as a significant pre-exposure \times treatment \times hormonal treatment interaction ($F_{(2,73)} = 11.6$, $p < 0.0001$). ANOVA for OVX female rats yielded significant main effects of pre-exposure ($F_{(1,61)} = 6.0$, $p < 0.05$) and treatment ($F_{(1,61)} = 27.7$, $p < 0.0001$), as well as a near significant pre-exposure \times treatment \times hormonal treatment interaction ($F_{(2,61)} = 2.97$, $p = 0.059$). *Post-hoc* comparisons for each of the analyses confirmed the presence of LI in saline-injected female and male rats given 50 or 150 μ g/kg 17 β -estradiol, in MK-801-injected female rats given 0, 50, or 150 μ g/kg 17 β -estradiol, and in MK-801-injected rats




Figure 4 Effects of 17 β -estradiol on MK-801 (MK)-induced LI persistence under strong conditioning (forty pre-exposures and five conditioning trials) in female, male, and OVX female rats. Mean (\pm SEM) log times to complete 76–100 licks (after the tone onset) of the PE and the NPE saline (sal)- or MK-injected female (a), male (b), and OVX female (c) rats, administered with 0, 50, or 150 μ g/kg of 17 β -estradiol (oil, E 50, or E 150, respectively). Asterisks indicate a significant difference between the PE and NPE groups, namely, presence of LI (* p < 0.05; ** p < 0.01).

given 0 μ g/kg 17 β -estradiol (p 's < 0.05), but not in the other conditions.

DISCUSSION

The present study sought to assess whether estrogen exerts an antipsychotic action, as indexed by the LI model, in gonadally intact female and male rats. Using a non-pharmacological and two acute pharmacological LI models, we showed that: (1) under conditions yielding LI in non-treated controls, 10 μ g/kg 17 β -estradiol disrupted LI, whereas higher doses (50 and 150 μ g/kg) were without any effect in both sexes; (2) 17 β -estradiol at higher doses restored LI under conditions preventing LI in non-treated controls, an effect considered predictive for activity against positive symptoms, in both sexes; (3) 17 β -estradiol at higher doses reversed amphetamine-induced disruption of LI, an effect considered predictive for activity against positive symptoms, in both sexes; and (4) 17 β -estradiol at higher doses reversed MK-801-induced persistent LI, an effect considered predictive for activity against negative/cognitive symptoms, in gonadally intact male and OVX rats, but not in gonadally intact female rats (see Table 1). These findings indicate that in the LI model, 17 β -estradiol exerts a clear-cut antipsychotic activity in both sexes and, remarkably, is more efficacious in males and OVX females, in which it also exerts activity considered predictive of anti-negative symptoms/pro-cognitive action.

Bimodal Effect of 17 β -Estradiol on LI

17 β -estradiol, administered on its own, produced a dose-dependent bimodal effect on LI in both female and male rats, with the low dose disrupting LI and the two higher

doses restoring LI under conditions (strong conditioning) that disrupted LI in controls.

Potentialiation or facilitation of LI under conditions of weak or absent LI in controls by APDs is a widely used index of antipsychotic activity. It is produced by a wide range of typical and atypical APDs and is also seen in normal humans (Barrett *et al*, 2004; Dunn *et al*, 1993; McCartan *et al*, 2001; Shadach *et al*, 1999; Weiner, 2003; Weiner and Arad 2009; Weiner and Feldon 1987; Williams *et al*, 1996; Williams *et al*, 1997). The fact that higher doses of 17 β -estradiol produced a pattern typically found for APDs suggests that at these doses, 17 β -estradiol may have antipsychotic properties.

The LI-facilitating effect of 17 β -estradiol was restricted to high doses as the low dose of 10 μ g/kg disrupted LI in both sexes. We have previously obtained this same bimodal effect of 17 β -estradiol on LI in OVX rats (Arad and Weiner, 2010) and Nofrey *et al* (2008) have also reported LI-disrupting effect of 10 μ g/kg 17 β -estradiol in OVX rats. The fact that 17 β -estradiol produces contrasting effects on LI at low and high doses implies that estradiol acts dose-dependently on different neural substrates. One likely substrate is the dopaminergic system. APD-induced LI potentiation is mediated by blockade of DA transmission within the nucleus accumbens (NAC) at the time of conditioning (Gray *et al*, 1995a; Gray *et al*, 1997; Joseph *et al*, 2000; Warburton *et al*, 1996; Weiner 2003). Consequently, our results imply that the higher doses of 17 β -estradiol, which potentiated LI, reduced mesolimbic DA function. Conversely, low dose, which disrupted LI, may have acted by increasing DA release within the NAC as does the amphetamine (Warburton *et al*, 1996). Both reduction and increase of striatal dopaminergic function by 17 β -estradiol have been reported for all indices of dopaminergic activity, including receptor levels/binding, membrane

Table 1 Summary of Experimental Design and Results

	Tone-shock pairings	Sex	Pro-psychotic treatment	Doss of 17 β -estradiol			
				Vehicle	10 μ g/kg	50 μ g/kg	150 μ g/kg
Experiments 1 and 2	2	F	—	LI	No LI	LI	LI
		M	—	LI	No LI	LI	LI
Experiments 3 and 4	5	F	—	No LI	No LI	LI	LI
		M	—	No LI	No LI	LI	LI
Experiments 5 and 6	2	F	Sal	LI	—	LI	LI
			Amph	No LI	—	LI	LI
		M	Sal	LI	—	LI	LI
			Amph	No LI	—	LI	LI
Experiments 7–9	5	F	Sal	No LI	—	LI	LI
			MK	LI	—	LI	LI
			Sal	No LI	—	LI	LI
		M	MK	LI	—	No LI	No LI
			Sal	No LI	—	No LI	No LI
			MK	LI	—	No LI	No LI
OVX (F)	Sal	No LI	—	No LI	No LI		
	MK	LI	—	No LI	No LI		

Abbreviations: Amph, amphetamine; F, female; LI, latent inhibition; M, male; MK, MK-801; OVX, ovariectomy; Sal, saline; —, has not been administered.

dopamine transporter levels, and release, depending on dose and treatment paradigm (Arvin *et al*, 2000; Becker, 1999; Chavez *et al*, 2010; Di Paolo, 1994; Disshon *et al*, 1998; Dluzen and Horstink, 2003; McDermott, 1993; Morissette *et al*, 2008; Morissette and Di Paolo, 1993; Peris *et al*, 1991; Shieh and Yang, 2008; Thompson and Moss, 1994; Yu *et al*, 2009; Zhou *et al*, 2002). It has been suggested that anti-dopaminergic effects are primarily exerted by high doses of estrogen or chronic administration, whereas pro-dopaminergic actions are more associated with lower levels of estrogen (Barber *et al*, 1976; Becker, 1999; Bedard *et al*, 1977; Chavez *et al*, 2010; Cyr *et al*, 2002; Di Paolo, 1994; Di Paolo *et al*, 1981; Hruska and Silbergeld, 1980; McEwen and Alves, 1999; Riddoch *et al*, 1971; Yu *et al*, 2009). Although the specific mechanisms by which 17 β -estradiol exerts its dose-dependent effects on LI observed here remain to be elucidated, if both effects are indeed DA-mediated, this would imply that low 17 β -estradiol doses exert a pro-psychotic action.

An alternative mechanism for the dose-dependent effect of 17 β -estradiol can be derived from our finding that a similar dose-dependent bimodal effect is produced by atypical APDs, such as risperidone, which disrupts LI at low doses and potentiates LI at higher doses (Weiner *et al*, 2003). The LI disruptive effect of atypical APDs is distinct from that of amphetamine because it occurs in the pre-exposure stage and is mediated by 5HT_{2A} antagonism, whereas the disruptive effect of amphetamine occurs in the conditioning stage and is mediated by enhanced DA transmission. As estrogen modulates brain serotonergic activity and specifically the 5HT_{2A} receptors (Fink *et al*, 1998; Sumner and Fink, 1997), this could be the mechanism underlying the LI disruptive action of low 17 β -estradiol, and thus reflect an atypical antipsychotic action of this hormone. Clearly it is of interest to determine whether low estrogen is pro-psychotic or antipsychotic. Testing the effects of APDs on 17 β -estradiol-induced LI disruption

would be one straightforward way to answer this question; In addition, these alternatives can be teased out by assessing at which stage 17 β -estradiol acts to disrupt LI.

Reversal of Disrupted LI: Putative Efficacy for Positive Symptoms

Amphetamine-induced LI disruption and its reversal by both typical and atypical APDs in the male rodent is a long standing model of positive symptoms (Weiner, 2003; Weiner and Arad, 2009). Although gender differences in response to amphetamine and other psychostimulants have been widely reported (for review see Fattore *et al*, 2008), such differences are not evident in the pro-psychotic action of amphetamine in the LI model (Arad and Weiner, 2010). Disruption of LI reflects a selective attention deficit, whereby animals lose the capacity to ignore the irrelevant stimulus, and is also observed in amphetamine-treated humans and high-schizotypal humans, (Braunstein-Bercovitz *et al*, 2002; Gray *et al*, 1992b; Salgado *et al*, 2000; Swerdlow *et al*, 2003; Thornton *et al*, 1996) as well as in acutely psychotic schizophrenia patients (Baruch *et al*, 1988; Gray *et al*, 1992a; Gray *et al*, 1995b; Rasclé *et al*, 2001; Swerdlow *et al*, 2005). A failure to inhibit attention to irrelevant stimuli is likely to give rise to aberrantly increased salience perception and distractibility that are associated with psychotic symptoms (Kapur *et al*, 2005; Weiner and Arad, 2009). 17 β -estradiol at 50 and 150 μ g/kg doses, which potentiated LI under conditions that disrupted LI in untreated gonadally intact rats (strong conditioning), also reversed amphetamine-induced LI disruption, as typically found with APDs. The present results replicate our recent finding (Arad and Weiner, 2010) that 17 β -estradiol prevents amphetamine from disrupting LI in gonadally intact female rats and extends this action of 17 β -estradiol to gonadally intact male rats.

Reversal of amphetamine-induced LI disruption by APDs, like APD-induced potentiation of LI, is mediated by blockade of DA transmission within the NAC at the time of conditioning (Gray *et al*, 1995a; Gray *et al*, 1997; Joseph *et al*, 2000; Warburton *et al*, 1996; Weiner, 2003). Consequently, our findings that acute high doses of 17 β -estradiol blocked the effects of amphetamine as well as potentiated LI on its own support the notion that 17 β -estradiol reduces mesolimbic DA function. Taken together, the efficacy of 17 β -estradiol to alleviate non-pharmacologically and pharmacologically induced LI disruption is indicative of its therapeutic capacity for positive symptoms in schizophrenia.

Previous studies showed that in OVX rats, 17 β -estradiol can reverse psychosis-mimicking abnormalities induced by dopamine agonists, including hyperactivity, stereotypy and circling (Becker and Beer, 1986; Becker and Rudick, 1999; Bedard *et al*, 1983; Bedard *et al*, 1978; Earley and Leonard, 1978; Euvrard *et al*, 1979; Euvrard *et al*, 1980; Gordon, 1980; Gordon and Diamond, 1981; Naik *et al*, 1978), disrupted PPI (Gogos *et al*, 2010), and disrupted LI (Arad and Weiner, 2009, 2010). To the best of our knowledge, this is the first demonstration that 17 β -estradiol exerts an antipsychotic action in gonadally intact rats of both sexes. These outcomes indicate that 17 β -estradiol possesses antipsychotic properties in both sexes and, by extension, that such action is independent of endogenous estrogen levels.

Reversal of Abnormally Persistent LI: Putative Efficacy for Negative and Cognitive Symptoms

As NMDA antagonists induce in addition to positive symptoms, also negative symptoms and cognitive impairments characteristic of endogenous schizophrenia (Javitt and Zukin, 1991; Krystal *et al*, 2003; Lahti *et al*, 1995; Malhotra *et al*, 1997; Tamminga, 1998), their behavioral effects in animals, with the exception of locomotor hyperactivity, are usually used as pharmacological models of negative and/or cognitive symptoms (Bakshi *et al*, 1994; Javitt and Zukin, 1991; Moghaddam and Jackson, 2003; Nilsson *et al*, 2001; Sams-Dodd, 1996; Swerdlow *et al*, 1996). Cognitive and behavioral inflexibility that is often observed following NMDA blockade in rats and humans has been argued to be particularly relevant to the modeling of negative/cognitive symptoms (Carlsson *et al*, 1999; Carlsson and Carlsson, 1990b; Krystal *et al*, 2000; Moghaddam *et al*, 1997).

As shown previously (Gaisler-Salomon *et al*, 2008; Gaisler-Salomon and Weiner, 2003; Lipina *et al*, 2005), under strong conditioning, MK-801 induced persistent LI in male rats. In other words, whereas vehicle-injected PE male rats switched in the conditioning stage to respond according to the stimulus-reinforcement contingency, MK-801-injected PE rats persisted in responding according to the stimulus-no-event contingency acquired in pre-exposure, in line with many findings showing that NMDA receptor blockade induces behavioral and cognitive inflexibility (Carlsson and Carlsson, 1990a; Jentsch and Taylor, 2001; Moghaddam *et al*, 1997; Svensson, 2000; van der Meulen *et al*, 2003). The same effect of MK-801 on LI was shown here, for the first time, in gonadally intact and OVX females. Thus, like amphetamine, MK-801 produced its schizophrenia-

relevant effect in a sex-independent manner. However, unlike amphetamine-induced disrupted LI, in gonadally intact rats, MK-801-induced persistent LI was ameliorated by 17 β -estradiol in male but not female rats. The latter is rather remarkable as it suggests that 17 β -estradiol is more efficient in ameliorating MK-801 effects on LI in the absence of estrogen. This suggestion was supported by our finding that 17 β -estradiol was also effective in reversing MK-801-induced persistent LI in OVX rats. The basis for the sex-dependent sensitivity of MK-801-induced persistent LI to 17 β -estradiol in gonadally intact rats is unclear. Greater sensitivity of gonadally intact females than males to NMDA antagonists, such as MK-801, requiring higher 17 β -estradiol doses to block its effects, has been reported but for much higher, neurodegeneration-producing MK-801 doses (Andine *et al*, 1999; Auer, 1996; de Olmos *et al*, 2008; Dribben *et al*, 2003; Fix *et al*, 1995; Honack and Loscher, 1993; Wozniak *et al*, 1998). The sex-dependent effects might also be dependent on the behavioral phenomenon tested, as 17 β -estradiol was found to reverse the disruptive effects of the NMDA antagonist PCP in the novel object-recognition test in gonadally intact female rats (Sutcliffe *et al*, 2007).

The mechanism/s by which 17 β -estradiol reverses MK-801-induced persistence of LI remain to be investigated, but two options can be suggested based on known pharmacological reversals of this abnormality. Reversal of MK-801-induced persistent LI by atypical APDs is mediated by 5HT_{2A} receptor antagonism (Gaisler-Salomon and Weiner, 2003; Weiner and Arad, 2009). In addition, MK-801-induced persistent LI is reversed by a wide variety of compounds enhancing NMDA receptor function (Black *et al*, 2009; Gaisler-Salomon *et al*, 2008).

The relationship between estrogen and serotonin is well documented. Studies in intact rats of both sexes have shown that estrogen modulates brain serotonergic activity (Fink *et al*, 1998) and that during high estrogen levels, 5HT_{2A} density is increased (Sumner and Fink 1997). 17 β -estradiol treatment in gonadectomized female and male rats increased 5HT_{2A} mRNA levels and receptors' density in the frontal cortex and NAC (Cyr *et al*, 1998; Fink *et al*, 1998; Sumner and Fink, 1995; Sumner and Fink, 1998). Importantly, the same effect is produced in intact male rats by clozapine (Buckland *et al*, 1997). Clinical studies reported that raloxifene, a second generation selective estrogen receptor modulator, mimics 17 β -estradiol effect on 5HT_{2A} receptor binding by increasing 5HT_{2A} receptor binding levels and its mRNA levels in the frontal cortex, amygdala, NAC, and striatum (Cyr *et al*, 2002).

There is also a well-established relationship between estrogen and NMDA receptor function. In OVX rats, 17 β -estradiol increases hippocampal NMDA receptors and glutamate transmission, whereas in the striatum, frontal cortex, and NAC, 17 β -estradiol decreases NMDA receptor binding. Furthermore, 17 β -estradiol increases agonist binding but decreases antagonist binding of NMDA receptors in the hippocampus (Cyr *et al*, 2000; Cyr *et al*, 2001; Daniel and Dohanich, 2001; El-Bakri *et al*, 2004; Gazzaley *et al*, 1996; Weiland, 1992; Woolley and Schwartzkroin, 1998; Woolley *et al*, 1997). Although these effects could explain the reversal of MK-801-induced persistent LI in OVX rats seen here, it is not clear whether similar effects are exerted in intact male brains. However,

17 β -estradiol does enhance glutamatergic transmission in hippocampal slices taken from male rats (Teyler *et al*, 1980).

17 β -Estradiol in LI—An Antipsychotic with a Sex-Dependent Action

Although the mechanisms underlying the effects of 17 β -estradiol seen here remain to be investigated, our results show that at certain doses this agent possesses an antipsychotic profile in the LI model. Moreover, this profile is sex dependent. Specifically, 17 β -estradiol possesses a profile of a typical APD (consisting of LI potentiation on its own and reversal of amphetamine-induced LI disruption) in both sexes and a profile of an atypical APD (consisting of LI potentiation on its own, reversal of amphetamine-induced disrupted LI and reversal of MK801-induced persistent LI), in male rats.

In psychological terms, in all of the models, 17 β -estradiol targeted selectively the processes responsible for attentional selectivity (in the PE groups) without affecting associative capacity (in the NPE groups). Furthermore, reversal of disrupted and persistent LI can be seen as normalization of two poles of dysfunctional attentional control. On the one hand, 17 β -estradiol strengthens/restores the capacity to ignore irrelevant stimuli in amphetamine-treated rats and on the other hand, 17 β -estradiol strengthens/restores the capacity to dis-ignore irrelevant stimuli when they become relevant, enabling flexible re-deployment of attentional resources according to current situational demands, in MK-801-treated rats. The former would be beneficial in normalizing aberrantly increased salience perception and distractibility that are associated with psychotic symptoms (Gray *et al*, 1991; Kapur, 2003; Smith *et al*, 2006; Swerdlow and Koob, 1987; Weiner and Joel, 2002), whereas the latter would be beneficial in reducing cognitive inflexibility and inattention that are associated with negative/cognitive symptoms (Carlsson and Carlsson, 1990b; Krystal *et al*, 2003; Moghaddam *et al*, 1997; Weiner, 2003). Given that 17 β -estradiol reverses amphetamine-induced LI disruption in both sexes, but MK-801-induced LI persistence only in male and OVX rats, the implications of our results for the clinic are that 17 β -estradiol would be beneficial against positive symptoms in both sexes but against negative/cognitive symptoms in male patients and in women with very low estrogen levels, eg, during menopause.

This conclusion seems in line with positive reports from the clinic. Efficacy of 17 β -estradiol against positive symptoms in schizophrenic women has been shown in several studies (Kulkarni *et al*, 2008a; Kulkarni *et al*, 1996; Kulkarni *et al*, 2008b; Kulkarni *et al*, 2001; Riecher-Rossler and de Geyer, 2007), as it was reported to be less effective or ineffective against negative symptoms (Akhondzadeh *et al*, 2003; Kulkarni *et al*, 2008a). In an attempt to reconcile inconsistent reports in the literature regarding efficacy of estradiol treatment in schizophrenic women, Mortimer (2007) suggested that such treatment would be effective in schizophrenic women who suffer from severe estrogen deficiency. Interestingly, reduced levels of plasma estrogen were found in both male (Huber *et al*, 2005) and female (Bergemann *et al*, 2005; Huber *et al*, 2004; Huber *et al*, 2001; Riecher-Rossler *et al*, 1994) schizophrenia patients, and recently a small study of men with schizophrenia who

received oral estradiol valerate also showed an abatement in psychotic symptoms (Kulkarni, 2009). The role of estrogen in schizophrenia has been supported by the finding that variation in the estrogen receptor alpha gene and cortical estrogen receptor alpha mRNA is associated with schizophrenia (Perlman *et al*, 2005; Perlman *et al*, 2004; Weickert *et al*, 2008; Wong and Weickert, 2009) and it has been suggested that the brain response to circulating estrogen may be altered in schizophrenia (Weickert *et al*, 2008). Taken together with our animal data, estrogen seems to have the potential to be useful in the treatment of schizophrenia.

In summary, our previous (Arad and Weiner, 2010) and present data are clear in showing that estrogen can exert antipsychotic activity, reversing hyperdopaminergia-induced behavioral abnormality in gonadally intact rats of both sexes and in OVX rats, and reversing hypoglutamatergia-induced abnormality in male and OVX rats. By extension, these results suggest that estrogen can be viewed as an effective treatment not only for positive symptoms in women with schizophrenia, but also for a wide spectrum of symptoms in women and men with schizophrenia, including negative/cognitive symptoms. Unfortunately, the risk of inducing cancers has limited the applicability of estrogen in humans for use in modulating the central nervous system neurotransmission (Rossouw *et al*, 2002), although recent studies found no increased risk (Anderson *et al*, 2006; Stefanick *et al*, 2006; Stevenson, 2009). Importantly in this context, our previous data showed that co-administration of a physiological dose of 17 β -estradiol and APD augments APD efficacy, and in fact may be more effective than raising the dose of APD. Further research is required to determine the correct dose and duration of the use of 17 β -estradiol as monotherapy, as well as an adjunctive therapy. Our present results raise the possibility that variable outcomes in the clinic may be because of the differences in the dosage of estrogen and, in general, alert to the importance of dose-response studies. Finally, new estrogenic compounds acting selectively in the brain may provide a safer, non-feminizing approach for the treatment of schizophrenia (Cyr *et al*, 2002; Kulkarni, 2009).

ACKNOWLEDGEMENTS

We wish to thank the Joseph Sagol Fellowship Program in Humanities and Social Sciences at Tel-Aviv University for their funding (MA).

DISCLOSURE

The authors declare no conflict of interest.

REFERENCES

- Akhondzadeh S, Nejatiasafa AA, Amini H, Mohammadi MR, Larijani B, Kashani L *et al* (2003). Adjunctive estrogen treatment in women with chronic schizophrenia: a double-blind, randomized, and placebo-controlled trial. *Prog Neuropsychopharmacol Biol Psychiatry* 27: 1007–1012.
- Anderson GL, Chlebowski RT, Rossouw JE, Rodabough RJ, McTiernan A, Margolis KL *et al* (2006). Prior hormone therapy and breast cancer risk in the Women's Health Initiative

- randomized trial of estrogen plus progestin. *Maturitas* 55: 103–115.
- Andine P, Widermark N, Axelsson R, Nyberg G, Olofsson U, Martensson E *et al* (1999). Characterization of MK-801-induced behavior as a putative rat model of psychosis. *J Pharmacol Exp Ther* 290: 1393–1408.
- Arad M, Weiner I (2008). Fluctuation of latent inhibition along the estrous cycle in the rat: modeling the cyclicity of symptoms in schizophrenic women? *Psychoneuroendocrinology* 33: 1401–1410.
- Arad M, Weiner I (2009). Disruption of latent inhibition induced by ovariectomy can be reversed by estradiol and clozapine as well as by co-administration of haloperidol with estradiol but not by haloperidol alone. *Psychopharmacology (Berl)* 206: 731–740.
- Arad M, Weiner I (2010). Contrasting effects of increased and decreased dopamine transmission on latent inhibition in ovariectomized rats and their modulation by 17 β -estradiol: an animal model of menopausal psychosis? *Neuropsychopharmacology* 35: 1570–1582.
- Arvin M, Fedorkova L, Disshon KA, Dluzen DE, Leipheimer RE (2000). Estrogen modulates responses of striatal dopamine neurons to MPP(+): evaluations using *in vitro* and *in vivo* techniques. *Brain Res* 872: 160–171.
- Auer RN (1996). Effect of age and sex on N-methyl-D-aspartate antagonist-induced neuronal necrosis in rats. *Stroke* 27: 743–746.
- Bakshi VP, Swerdlow NR, Geyer MA (1994). Clozapine antagonizes phencyclidine-induced deficits in sensorimotor gating of the startle response. *J Pharmacol Exp Ther* 271: 787–794.
- Barak S, Arad M, De Levie A, Black MD, Griebel G, Weiner I (2009). Pro-cognitive and antipsychotic efficacy of the α 7 nicotinic partial agonist SSR180711 in pharmacological and neurodevelopmental latent inhibition models of schizophrenia. *Neuropsychopharmacology* 34: 1753–1763.
- Barber PV, Arnold AG, Evans G (1976). Recurrent hormone dependent chorea: effects of oestrogens and progestogens. *Clin Endocrinol (Oxf)* 5: 291–293.
- Barch DM, Carter CS (2008). Measurement issues in the use of cognitive neuroscience tasks in drug development for impaired cognition in schizophrenia: a report of the second consensus building conference of the CNTRICS initiative. *Schizophr Bull* 34: 613–618.
- Barnes P, Staal V, Muir J, Good MA (2006). 17- β -estradiol administration attenuates deficits in sustained and divided attention in young ovariectomized rats and aged acyclic female rats. *Behav Neurosci* 120: 1225–1234.
- Barrett SL, Bell R, Watson D, King DJ (2004). Effects of amisulpride, risperidone and chlorpromazine on auditory and visual latent inhibition, prepulse inhibition, executive function and eye movements in healthy volunteers. *J Psychopharmacol* 18: 156–172.
- Baruch I, Hemsley DR, Gray JA (1988). Differential performance of acute and chronic schizophrenics in a latent inhibition task. *J Nerv Ment Dis* 176: 598–606.
- Becker JB (1999). Gender differences in dopaminergic function in striatum and nucleus accumbens. *Pharmacol Biochem Behav* 64: 803–812.
- Becker JB, Beer ME (1986). The influence of estrogen on nigrostriatal dopamine activity: behavioral and neurochemical evidence for both pre- and postsynaptic components. *Behav Brain Res* 19: 27–33.
- Becker JB, Rudick CN (1999). Rapid effects of estrogen or progesterone on the amphetamine-induced increase in striatal dopamine are enhanced by estrogen priming: a microdialysis study. *Pharmacol Biochem Behav* 64: 53–57.
- Bedard P, Boucher R, Di Paolo T, Labrie F (1983). Biphasic effect of estradiol and domperidone on lingual dyskinesia in monkeys. *Exp Neurol* 82: 172–182.
- Bedard P, Dankova J, Boucher R, Langelier P (1978). Effect of estrogens on apomorphine-induced circling behavior in the rat. *Can J Physiol Pharmacol* 56: 538–541.
- Bedard P, Langelier P, Villeneuve A (1977). Oestrogens and extrapyramidal system. *Lancet* 2: 1367–1368.
- Bedard PJ, Malouin F, Dipaolo T, Labrie F (1982). Estradiol, TRH and striatal dopaminergic mechanisms. *Prog Neuropsychopharmacol Biol Psychiatry* 6: 555–561.
- Bergemann N, Mundt C, Parzer P, Pakrasi M, Eckstein-Mannspenger U, Haisch S *et al* (2005). Estrogen as an adjuvant therapy to antipsychotics does not prevent relapse in women suffering from schizophrenia: results of a placebo-controlled double-blind study. *Schizophr Res* 74: 125–134.
- Black MD, Varty GB, Arad M, Barak S, De Levie A, Boulay D *et al* (2008). Procognitive and antipsychotic efficacy of glycine transport 1 inhibitors (GlyT1) in acute and neurodevelopmental models of schizophrenia: latent inhibition studies in the rat. *Psychopharmacology (Berl)* 202: 385–396.
- Black MD, Varty GB, Arad M, Barak S, De Levie A, Boulay D *et al* (2009). Procognitive and antipsychotic efficacy of glycine transport 1 inhibitors (GlyT1) in acute and neurodevelopmental models of schizophrenia: latent inhibition studies in the rat. *Psychopharmacology (Berl)* 202: 385–396.
- Braunstein-Bercovitz H, Rammesayer T, Gibbons H, Lubow RE (2002). Latent inhibition deficits in high-schizotypal normals: symptom-specific or anxiety-related? *Schizophr Res* 53: 109–121.
- Buckland PR, D'Souza U, Maher NA, McGuffin P (1997). The effects of antipsychotic drugs on the mRNA levels of serotonin 5HT_{2A} and 5HT_{2C} receptors. *Brain Res Mol Brain Res* 48: 45–52.
- Carlsson A, Waters N, Carlsson ML (1999). Neurotransmitter interactions in schizophrenia—therapeutic implications. *Biol Psychiatry* 46: 1388–1395.
- Carlsson M, Carlsson A (1990a). Interactions between glutamatergic and monoaminergic systems within the basal ganglia—implications for schizophrenia and Parkinson's disease. *Trends Neurosci* 13: 272–276.
- Carlsson M, Carlsson A (1990b). Schizophrenia: a subcortical neurotransmitter imbalance syndrome? *Schizophr Bull* 16: 425–432.
- Chavez C, Gogos A, Jones ME, van den Buuse M (2009). Psychotropic drug-induced locomotor hyperactivity and prepulse inhibition regulation in male and female aromatase knockout (ArKO) mice: role of dopamine D1 and D2 receptors and dopamine transporters. *Psychopharmacology (Berl)* 206: 267–279.
- Chavez C, Hollaus M, Scarr E, Pavey G, Gogos A, van den Buuse M (2010). The effect of estrogen on dopamine and serotonin receptor and transporter levels in the brain: an autoradiography study. *Brain Res* 1321: 51–59.
- Chiodo LA, Caggiula AR, Saller CF (1979). Estrogen increases both spiperone-induced catalepsy and brain levels of [³H]spiperone in the rat. *Brain Res* 172: 360–366.
- Cyr M, Bosse R, Di Paolo T (1998). Gonadal hormones modulate 5-hydroxytryptamine_{2A} receptors: emphasis on the rat frontal cortex. *Neuroscience* 83: 829–836.
- Cyr M, Calon F, Morissette M, Di Paolo T (2002). Estrogenic modulation of brain activity: implications for schizophrenia and Parkinson's disease. *J Psychiatry Neurosci* 27: 12–27.
- Cyr M, Ghribi O, Di Paolo T (2000). Regional and selective effects of oestradiol and progesterone on NMDA and AMPA receptors in the rat brain. *J Neuroendocrinol* 12: 445–452.
- Cyr M, Ghribi O, Thibault C, Morissette M, Landry M, Di Paolo T (2001). Ovarian steroids and selective estrogen receptor modulators activity on rat brain NMDA and AMPA receptors. *Brain Res Brain Res Rev* 37: 153–161.

- Daniel JM, Bohacek J (2010). The critical period hypothesis of estrogen effects on cognition: insights from basic research. *Biochim Biophys Acta* (e pub ahead of print).
- Daniel JM, Dohanich GP (2001). Acetylcholine mediates the estrogen-induced increase in NMDA receptor binding in CA1 of the hippocampus and the associated improvement in working memory. *J Neurosci* 21: 6949–6956.
- de Olmos S, Bueno A, Bender C, Lorenzo A, de Olmos J (2008). Sex differences and influence of gonadal hormones on MK801-induced neuronal degeneration in the granular retrosplenial cortex of the rat. *Brain Struct Funct* 213: 229–238.
- De Ryck M, Hruska RE, Silbergeld EK (1982). Estrogen and haloperidol-induced versus handling-related catalepsy in male rats. *Pharmacol Biochem Behav* 17: 1027–1035.
- Di Paolo T (1994). Modulation of brain dopamine transmission by sex steroids. *Rev Neurosci* 5: 27–41.
- Di Paolo T, Carmichael R, Labrie F, Raynaud JP (1979). Effects of estrogens on the characteristics of [3H]spiroperidol and [3H]RU24213 binding in rat anterior pituitary gland and brain. *Mol Cell Endocrinol* 16: 99–112.
- Di Paolo T, Poyet P, Labrie F (1981). Effect of chronic estradiol and haloperidol treatment on striatal dopamine receptors. *Eur J Pharmacol* 73: 105–106.
- Disshon KA, Boja JW, Dluzen DE (1998). Inhibition of striatal dopamine transporter activity by 17 β -estradiol. *Eur J Pharmacol* 345: 207–211.
- Dluzen D, Horstink M (2003). Estrogen as neuroprotectant of nigrostriatal dopaminergic system: laboratory and clinical studies. *Endocrine* 21: 67–75.
- Dribben W, Nemmers B, Nardi A, Taylor G, Olney J, Farber N (2003). Chronic but not acute estradiol treatment protects against the neurodegenerative effects of N-methyl-D-aspartate receptor antagonists. *Endocrine* 21: 53–58.
- Dunn LA, Atwater GE, Kilts CD (1993). Effects of antipsychotic drugs on latent inhibition: sensitivity and specificity of an animal behavioral model of clinical drug action. *Psychopharmacology (Berl)* 112: 315–323.
- Earley CJ, Leonard BE (1978). Behavioural studies on the effects of d-amphetamine and estradiol benzoate alone and in combination. *Psychopharmacology (Berl)* 56: 179–183.
- El-Bakri NK, Islam A, Zhu S, Elhassan A, Mohammed A, Winblad B *et al* (2004). Effects of estrogen and progesterone treatment on rat hippocampal NMDA receptors: relationship to Morris water maze performance. *J Cell Mol Med* 8: 537–544.
- Euvrard C, Labrie F, Boissier JR (1979). Effect of moxestrol on haloperidol-induced changes in striatal acetylcholine levels and dopamine turn-over. *Commun Psychopharmacol* 3: 329–334.
- Euvrard C, Oberlander C, Boissier JR (1980). Antidopaminergic effect of estrogens at the striatal level. *J Pharmacol Exp Ther* 214: 179–185.
- Fattore L, Altea S, Fratta W (2008). Sex differences in drug addiction: a review of animal and human studies. *Womens Health (Lond Engl)* 4: 51–65.
- Fink G, Sumner BE, McQueen JK, Wilson H, Rosie R (1998). Sex steroid control of mood, mental state and memory. *Clin Exp Pharmacol Physiol* 25: 764–775.
- Fix AS, Wozniak DF, Truex LL, McEwen M, Miller JP, Olney JW (1995). Quantitative analysis of factors influencing neuronal necrosis induced by MK-801 in the rat posterior cingulate/retrosplenial cortex. *Brain Res* 696: 194–204.
- Frick KM (2009). Estrogens and age-related memory decline in rodents: what have we learned and where do we go from here? *Horm Behav* 55: 2–23.
- Gaisler-Salomon I, Diamant L, Rubin C, Weiner I (2008). Abnormally persistent latent inhibition induced by MK801 is reversed by risperidone and by positive modulators of NMDA receptor function: differential efficacy depending on the stage of the task at which they are administered. *Psychopharmacology (Berl)* 196: 255–267.
- Gaisler-Salomon I, Weiner I (2003). Systemic administration of MK-801 produces an abnormally persistent latent inhibition which is reversed by clozapine but not haloperidol. *Psychopharmacology (Berl)* 166: 333–342.
- Galea LA, Wide JK, Barr AM (2001). Estradiol alleviates depressive-like symptoms in a novel animal model of postpartum depression. *Behav Brain Res* 122: 1–9.
- Gazzaley AH, Weiland NG, McEwen BS, Morrison JH (1996). Differential regulation of NMDAR1 mRNA and protein by estradiol in the rat hippocampus. *J Neurosci* 16: 6830–6838.
- Gibbs RB, Burke AM, Johnson DA (1998). Estrogen replacement attenuates effects of scopolamine and lorazepam on memory acquisition and retention. *Horm Behav* 34: 112–125.
- Gogos A, Kwek P, Chavez C, van den Buuse M (2010). Estrogen treatment blocks 8-hydroxy-2-dipropylaminotetralin- and apomorphine-induced disruptions of prepulse inhibition: involvement of dopamine D1 or D2 or serotonin 5-HT1A, 5-HT2A, or 5-HT7 receptors. *J Pharmacol Exp Ther* 333: 218–227.
- Gogos A, Nathan PJ, Guille V, Croft RJ, van den Buuse M (2006). Estrogen prevents 5-HT1A receptor-induced disruptions of prepulse inhibition in healthy women. *Neuropsychopharmacology* 31: 885–889.
- Gordon JH (1980). Modulation of apomorphine-induced stereotypy by estrogen: time course and dose response. *Brain Res Bull* 5: 679–682.
- Gordon JH, Diamond BI (1981). Antagonism of dopamine supersensitivity by estrogen: neurochemical studies in an animal model of tardive dyskinesia. *Biol Psychiatry* 16: 365–371.
- Gray JA, Feldon J, Rawlins JNP, Hemsley DR, Smith AD (1991). The neuropsychology of schizophrenia. *Behav Brain Sci* 14: 1–84.
- Gray JA, Joseph MH, Hemsley DR, Young AM, Warburton EC, Boulguez P *et al* (1995a). The role of mesolimbic dopaminergic and retrohippocampal afferents to the nucleus accumbens in latent inhibition: implications for schizophrenia. *Behav Brain Res* 71: 19–31.
- Gray JA, Moran PM, Grigoryan G, Peters SL, Young AM, Joseph MH (1997). Latent inhibition: the nucleus accumbens connection revisited. *Behav Brain Res* 88: 27–34.
- Gray NS, Hemsley DR, Gray JA (1992a). Abolition of latent inhibition in acute, but not chronic, schizophrenics. *Neurol Psychiatr Brain Res* 1: 83–89.
- Gray NS, Pickering AD, Hemsley DR, Dawling S, Gray JA (1992b). Abolition of latent inhibition by a single 5mg dose of d-amphetamine in man. *Psychopharmacology (Berl)* 107: 425–430.
- Gray NS, Pilowsky LS, Gray JA, Kerwin RW (1995b). Latent inhibition in drug naive schizophrenics: relationship to duration of illness and dopamine D2 binding using SPET. *Schizophr Res* 17: 95–107.
- Hafner H (2003). Gender differences in schizophrenia. *Psychoneuroendocrinology* 28(Suppl 2): 17–54.
- Hafner H, Behrens S, De Vry J, Gattaz WF (1991). An animal model for the effects of estradiol on dopamine-mediated behavior: implications for sex differences in schizophrenia. *Psychiatry Res* 38: 125–134.
- Hafner H, Riecher A, Maurer K, Loffler W, Munk-Jorgensen P, Stromgren E (1989). How does gender influence age at first hospitalization for schizophrenia? A transnational case register study. *Psychol Med* 19: 903–918.
- Harvey PD, Rabinowitz J, Eerdeken M, Davidson M (2005). Treatment of cognitive impairment in early psychosis: a comparison of risperidone and haloperidol in a large long-term trial. *Am J Psychiatry* 162: 1888–1895.
- Heresco-Levy U, Javitt DC, Ebstein R, Vass A, Lichtenberg P, Bar G *et al* (2005). D-serine efficacy as add-on pharmacotherapy to

- risperidone and olanzapine for treatment-refractory schizophrenia. *Biol Psychiatry* 57: 577–585.
- Honack D, Loscher W (1993). Sex differences in NMDA receptor mediated responses in rats. *Brain Res* 620: 167–170.
- Hruska RE, Silbergeld EK (1980). Increased dopamine receptor sensitivity after estrogen treatment using the rat rotation model. *Science* 208: 1466–1468.
- Huber TJ, Borsutzky M, Schneider U, Emrich HM (2004). Psychotic disorders and gonadal function: evidence supporting the oestrogen hypothesis. *Acta Psychiatr Scand* 109: 269–274.
- Huber TJ, Rollnik J, Wilhelms J, von zur Muhlen A, Emrich HM, Schneider U (2001). Estradiol levels in psychotic disorders. *Psychoneuroendocrinology* 26: 27–35.
- Huber TJ, Tettenborn C, Leifke E, Emrich HM (2005). Sex hormones in psychotic men. *Psychoneuroendocrinology* 30: 111–114.
- Javitt DC, Zukin SR (1991). Recent advances in the phencyclidine model of schizophrenia. *Am J Psychiatry* 148: 1301–1308.
- Jentsch JD, Taylor JR (2001). Impaired inhibition of conditioned responses produced by subchronic administration of phencyclidine to rats. *Neuropsychopharmacology* 24: 66–74.
- Joseph MH, Peters SL, Moran PM, Grigoryan GA, Young AM, Gray JA (2000). Modulation of latent inhibition in the rat by altered dopamine transmission in the nucleus accumbens at the time of conditioning. *Neuroscience* 101: 921–930.
- Kapur S (2003). Psychosis as a state of aberrant salience: a framework linking biology, phenomenology, and pharmacology in schizophrenia. *Am J Psychiatry* 160: 13–23.
- Kapur S, Mizrahi R, Li M (2005). From dopamine to salience to psychosis—linking biology, pharmacology and phenomenology of psychosis. *Schizophr Res* 79: 59–68.
- Kitamura N, Araya R, Kudoh M, Kishida H, Kimura T, Murayama M *et al* (2009). Beneficial effects of estrogen in a mouse model of cerebrovascular insufficiency. *PLoS One* 4: e5159.
- Korhonen S, Saarijarvi S, Aito M (1995). Successful estradiol treatment of psychotic symptoms in the premenstrual phase: a case report. *Acta Psychiatr Scand* 92: 237–238.
- Krystal JH, Bennett A, Abi-Saab D, Belger A, Karper LP, D'Souza DC *et al* (2000). Dissociation of ketamine effects on rule acquisition and rule implementation: possible relevance to NMDA receptor contributions to executive cognitive functions. *Biol Psychiatry* 47: 137–143.
- Krystal JH, D'Souza DC, Mathalon D, Perry E, Belger A, Hoffman R (2003). NMDA receptor antagonist effects, cortical glutamatergic function, and schizophrenia: toward a paradigm shift in medication development. *Psychopharmacology (Berl)* 169: 215–233.
- Kulkarni J (2009). Oestrogen—a new treatment approach for schizophrenia? *Med J Aust* 190: S37–S38.
- Kulkarni J, de Castella A, Fitzgerald PB, Gurvich CT, Bailey M, Bartholomeusz C *et al* (2008a). Estrogen in severe mental illness: a potential new treatment approach. *Arch Gen Psychiatry* 65: 955–960.
- Kulkarni J, de Castella A, Smith D, Taffe J, Keks N, Copolov D (1996). A clinical trial of the effects of estrogen in acutely psychotic women. *Schizophr Res* 20: 247–252.
- Kulkarni J, Gurvich C, Gilbert H, Mehmedbegovic F, Mu L, Marston N *et al* (2008b). Hormone modulation: a novel therapeutic approach for women with severe mental illness. *Aust N Z J Psychiatry* 42: 83–88.
- Kulkarni J, Riedel A, de Castella AR, Fitzgerald PB, Rolfe TJ, Taffe J *et al* (2001). Estrogen—a potential treatment for schizophrenia. *Schizophr Res* 48: 137–144.
- Lahti AC, Koffel B, LaPorte D, Tamminga CA (1995). Subanesthetic doses of ketamine stimulate psychosis in schizophrenia. *Neuropsychopharmacology* 13: 9–19.
- Lipina T, Labrie V, Weiner I, Roder J (2005). Modulators of the glycine site on NMDA receptors, D: -serine and ALX 5407, display similar beneficial effects to clozapine in mouse models of schizophrenia. *Psychopharmacology (Berl)* 179: 54–67.
- Lubow RE. Latent Inhibition and Conditioned Attention Theory. Cambridge University Press: Cambridge University Press, 1989.
- Mackintosh NJ (1975). A theory of attention: Variations in the associability of stimuli with reinforcement. *Psychol Rev* 82: 276–298.
- Malhotra AK, Pinals DA, Adler CM, Elman I, Clifton A, Pickar D *et al* (1997). Ketamine-induced exacerbation of psychotic symptoms and cognitive impairment in neuroleptic-free schizophrenics. *Neuropsychopharmacology* 17: 141–150.
- Marcondes FK, Bianchi FJ, Tanno AP (2002). Determination of the estrous cycle phases of rats: some helpful considerations. *Braz J Biol* 62: 609–614.
- McCartan D, Bell R, Green JF, Campbell C, Trimble K, Pickering A *et al* (2001). The differential effects of chlorpromazine and haloperidol on latent inhibition in healthy volunteers. *J Psychopharmacol* 15: 96–104.
- McDermott JL (1993). Effects of estrogen upon dopamine release from the corpus striatum of young and aged female rats. *Brain Res* 606: 118–125.
- McEwen BS, Alves SE (1999). Estrogen actions in the central nervous system. *Endocr Rev* 20: 279–307.
- Moghaddam B, Adams B, Verma A, Daly D (1997). Activation of glutamatergic neurotransmission by ketamine: a novel step in the pathway from NMDA receptor blockade to dopaminergic and cognitive disruptions associated with the prefrontal cortex. *J Neurosci* 17: 2921–2927.
- Moghaddam B, Jackson ME (2003). Glutamatergic animal models of schizophrenia. *Ann N Y Acad Sci* 1003: 131–137.
- Morissette M, Al Sweidi S, Callier S, Di Paolo T (2008). Estrogen and SERM neuroprotection in animal models of Parkinson's disease. *Mol Cell Endocrinol* 290: 60–69.
- Morissette M, Di Paolo T (1993). Effect of chronic estradiol and progesterone treatments of ovariectomized rats on brain dopamine uptake sites. *J Neurochem* 60: 1876–1883.
- Mortimer AM (2007). Relationship between estrogen and schizophrenia. *Expert Rev Neurother* 7: 45–55.
- Naik SR, Kelkar MR, Sheth UK (1978). Attenuation of stereotyped behaviour by sex steroids. *Psychopharmacology (Berl)* 57: 211–214.
- Nicoletti F, Ferrara N, Patti F, Viglianesi M, Rampello L, Bianchi A *et al* (1983). Influence of sex steroids and prolactin on haloperidol-induced catalepsy. *Brain Res* 279: 352–358.
- Nilsson M, Waters S, Waters N, Carlsson A, Carlsson ML (2001). A behavioural pattern analysis of hypoglutamatergic mice—effects of four different antipsychotic agents. *J Neural Transm* 108: 1181–1196.
- Nofrey BS, Ben-Shahar OM, Brake WG (2008). Estrogen abolishes latent inhibition in ovariectomized female rats. *Brain Cogn* 66: 156–160.
- Packard MG, Kohlmaier JR, Alexander GM (1996). Posttraining intrahippocampal estradiol injections enhance spatial memory in male rats: interaction with cholinergic systems. *Behav Neurosci* 110: 626–632.
- Palermo-Neto J, Dorce VA (1990). Influences of estrogen and/or progesterone on some dopamine related behavior in rats. *Gen Pharmacol* 21: 83–87.
- Palsson E, Klamer D, Wass C, Archer T, Engel JA, Svensson L (2005). The effects of phencyclidine on latent inhibition in taste aversion conditioning: differential effects of preexposure and conditioning. *Behav Brain Res* 157: 139–146.
- Peris J, Decambre N, Coleman-Hardee ML, Simpkins JW (1991). Estradiol enhances behavioral sensitization to cocaine and amphetamine-stimulated striatal [3H]dopamine release. *Brain Res* 566: 255–264.

- Perlman WR, Tomaskovic-Crook E, Montague DM, Webster MJ, Rubinow DR, Kleinman JE *et al* (2005). Alteration in estrogen receptor alpha mRNA levels in frontal cortex and hippocampus of patients with major mental illness. *Biol Psychiatry* **58**: 812–824.
- Perlman WR, Webster MJ, Kleinman JE, Weickert CS (2004). Reduced glucocorticoid and estrogen receptor alpha messenger ribonucleic acid levels in the amygdala of patients with major mental illness. *Biol Psychiatry* **56**: 844–852.
- Rasclé C, Mazas O, Vaiva G, Tournant M, Raybois O, Goudemand M *et al* (2001). Clinical features of latent inhibition in schizophrenia. *Schizophr Res* **51**: 149–161.
- Riddoch D, Jefferson M, Bickerstaff ER (1971). Chorea and the oral contraceptives. *Br Med J* **4**: 217–218.
- Riecher-Rossler A (2002). Oestrogen effects in schizophrenia and their potential therapeutic implications—review. *Arch Womens Ment Health* **5**: 111–118.
- Riecher-Rossler A, de Geyter C (2007). The forthcoming role of treatment with oestrogens in mental health. *Swiss Med Wkly* **137**: 565–572.
- Riecher-Rossler A, Hafner H (1993). Schizophrenia and oestrogens—is there an association? *Eur Arch Psychiatry Clin Neurosci* **242**: 323–328.
- Riecher-Rossler A, Hafner H, Stumbaum M, Maurer K, Schmidt R (1994). Can estradiol modulate schizophrenic symptomatology? *Schizophr Bull* **20**: 203–214.
- Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML *et al* (2002). Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA* **288**: 321–333.
- Salgado JV, Hetem LA, Vidal M, Graeff FG, Danion JM, Sandner G (2000). Reduction of latent inhibition by D-amphetamine in a conditioned suppression paradigm in humans. *Behav Brain Res* **117**: 61–67.
- Sams-Dodd F (1996). Phencyclidine-induced stereotyped behaviour and social isolation in rats: a possible animal model of schizophrenia. *Behav Pharmacol* **7**: 3–23.
- Seeman MV, Lang M (1990). The role of estrogens in schizophrenia gender differences. *Schizophr Bull* **16**: 185–194.
- Segarra AC, Agosto-Rivera JL, Febo M, Lugo-Escobar N, Menéndez-Delmestre R, Puig-Ramos A *et al* (2009). Estradiol: a key biological substrate mediating the response to cocaine in female rats. *Horm Behav* **58**: 33–43.
- Shadach E, Feldon J, Weiner I (1999). Clozapine-induced potentiation of latent inhibition is due to its action in the conditioning stage: implications for the mechanism of action of antipsychotic drugs. *Int J Neuropsychopharmacol* **2**: 283–291.
- Sherwin BB, Chertkow H, Schipper H, Nasreddine Z (2009). A randomized controlled trial of estrogen treatment in men with mild cognitive impairment. *Neurobiol Aging* (e pub ahead of print).
- Shieh KR, Yang SC (2008). Effects of estradiol on the stimulation of dopamine turnover in mesolimbic and nigrostriatal systems by cocaine- and amphetamine-regulated transcript peptide in female rats. *Neuroscience* **154**: 1589–1597.
- Smith A, Li M, Becker S, Kapur S (2006). Dopamine, prediction error and associative learning: a model-based account. *Network* **17**: 61–84.
- Soderstrom I, Strand M, Ingridsson AC, Nasic S, Olsson T (2009). 17beta-estradiol and enriched environment accelerate cognitive recovery after focal brain ischemia. *Eur J Neurosci* **29**: 1215–1224.
- Stefanick ML, Anderson GL, Margolis KL, Hendrix SL, Rodabough RJ, Paskett ED *et al* (2006). Effects of conjugated equine estrogens on breast cancer and mammography screening in postmenopausal women with hysterectomy. *JAMA* **295**: 1647–1657.
- Stevenson JC (2009). Hormone replacement therapy and cardiovascular disease revisited. *Menopause Int* **15**: 55–57.
- Summer BE, Fink G (1995). Estrogen increases the density of 5-hydroxytryptamine(2A) receptors in cerebral cortex and nucleus accumbens in the female rat. *J Steroid Biochem Mol Biol* **54**: 15–20.
- Sumner BE, Fink G (1997). The density of 5-hydroxytryptamine2A receptors in forebrain is increased at pro-oestrus in intact female rats. *Neurosci Lett* **234**: 7–10.
- Sumner BE, Fink G (1998). Testosterone as well as estrogen increases serotonin2A receptor mRNA and binding site densities in the male rat brain. *Brain Res Mol Brain Res* **59**: 205–214.
- Sutcliffe JS, Marshall KM, Neill JC (2007). Influence of gender on working and spatial memory in the novel object recognition task in the rat. *Behav Brain Res* **177**: 117–125.
- Sutcliffe JS, Rhaman F, Marshall KM, Neill JC (2008). Oestradiol attenuates the cognitive deficit induced by acute phencyclidine treatment in mature female hooded-Lister rats. *J Psychopharmacol* **22**: 918–922.
- Svensson TH (2000). Dysfunctional brain dopamine systems induced by psychotomimetic NMDA-receptor antagonists and the effects of antipsychotic drugs. *Brain Res Brain Res Rev* **31**: 320–329.
- Swerdlow NR, Bakshi V, Geyer MA (1996). Seroquel restores sensorimotor gating in phencyclidine-treated rats. *J Pharmacol Exp Ther* **279**: 1290–1299.
- Swerdlow NR, Koob GF (1987). Dopamine, schizophrenia, mania and depression: toward a unified hypothesis of cortico-striato-pallido-thalamic function. *Behav Brain Sci* **10**: 197–245.
- Swerdlow NR, Stephany N, Wasserman LC, Talledo J, Sharp R, Auerbach PP (2003). Dopamine agonists disrupt visual latent inhibition in normal males using a within-subject paradigm. *Psychopharmacology (Berl)* **169**: 314–320.
- Swerdlow NR, Stephany N, Wasserman LC, Talledo J, Sharp R, Minassian A *et al* (2005). Intact visual latent inhibition in schizophrenia patients in a within-subject paradigm. *Schizophr Res* **72**: 169–183.
- Tamminga CA (1998). Schizophrenia and glutamatergic transmission. *Crit Rev Neurobiol* **12**: 21–36.
- Teyler TJ, Vardaris RM, Lewis D, Rawitch AB (1980). Gonadal steroids: effects on excitability of hippocampal pyramidal cells. *Science* **209**: 1017–1018.
- Thompson TL, Moss RL (1994). Estrogen regulation of dopamine release in the nucleus accumbens: genomic- and nongenomic-mediated effects. *J Neurochem* **62**: 1750–1756.
- Thornton JC, Dawe S, Lee C, Capstick C, Corr PJ, Cotter P *et al* (1996). Effects of nicotine and amphetamine on latent inhibition in human subjects. *Psychopharmacology (Berl)* **127**: 164–173.
- Van den Buuse M, Eikelis N (2001). Estrogen increases prepulse inhibition of acoustic startle in rats. *Eur J Pharmacol* **425**: 33–41.
- van der Meulen JA, Bilbija L, Joosten RN, de Bruin JP, Feenstra MG (2003). The NMDA-receptor antagonist MK-801 selectively disrupts reversal learning in rats. *Neuroreport* **14**: 2225–2228.
- Walf AA, Frye CA (2009). Estradiol reduces anxiety- and depression-like behavior of aged female mice. *Physiol Behav* **99**: 169–174.
- Warburton EC, Mitchell SN, Joseph MH (1996). Calcium dependence of sensitised dopamine release in rat nucleus accumbens following amphetamine challenge: implications for the disruption of latent inhibition. *Behav Pharmacol* **7**: 119–129.
- Weickert CS, Miranda-Angulo AL, Wong J, Perlman WR, Ward SE, Radhakrishna V *et al* (2008). Variants in the estrogen receptor alpha gene and its mRNA contribute to risk for schizophrenia. *Hum Mol Genet* **17**: 2293–2309.

- Weiland NG (1992). Estradiol selectively regulates agonist binding sites on the N-methyl-D-aspartate receptor complex in the CA1 region of the hippocampus. *Endocrinology* **131**: 662–668.
- Weiner I (2003). The ‘two-headed’ latent inhibition model of schizophrenia: modeling positive and negative symptoms and their treatment. *Psychopharmacology (Berl)* **169**: 257–297.
- Weiner I, Arad M (2009). Using the pharmacology of latent inhibition to model domains of pathology in schizophrenia and their treatment. *Behav Brain Res* **204**: 369–386.
- Weiner I, Feldon J (1987). Facilitation of latent inhibition by haloperidol in rats. *Psychopharmacology (Berl)* **91**: 248–253.
- Weiner I, Joel D (2002). Dopamine in schizophrenia: dysfunctional information processing in basal ganglia-thalamocortical split circuits. In: Di Chiara G (ed). *Handbook of Experimental Pharmacology*. Springer: Berlin. pp 418–472.
- Weiner I, Schiller D, Gaisler-Salomon I (2003). Disruption and potentiation of latent inhibition by risperidone: the latent inhibition model of atypical antipsychotic action. *Neuropsychopharmacology* **28**: 499–509.
- Williams JH, Wellman NA, Geaney DP, Cowen PJ, Feldon J, Rawlins JN (1996). Antipsychotic drug effects in a model of schizophrenic attentional disorder: a randomized controlled trial of the effects of haloperidol on latent inhibition in healthy people. *Biol Psychiatry* **40**: 1135–1143.
- Williams JH, Wellman NA, Geaney DP, Feldon J, Cowen PJ, Rawlins JN (1997). Haloperidol enhances latent inhibition in visual tasks in healthy people. *Psychopharmacology (Berl)* **133**: 262–268.
- Wong J, Weickert CS (2009). Transcriptional interaction of an estrogen receptor splice variant and ErbB4 suggests convergence in gene susceptibility pathways in schizophrenia. *J Biol Chem* **284**: 18824–18832.
- Woolley CS, Schwartzkroin PA (1998). Hormonal effects on the brain. *Epilepsia* **39**(Suppl 8): S2–S8.
- Woolley CS, Weiland NG, McEwen BS, Schwartzkroin PA (1997). Estradiol increases the sensitivity of hippocampal CA1 pyramidal cells to NMDA receptor-mediated synaptic input: correlation with dendritic spine density. *J Neurosci* **17**: 1848–1859.
- Wozniak DF, Dikranian K, Ishimaru MJ, Nardi A, Corso TD, Tenkova T *et al* (1998). Disseminated corticolimbic neuronal degeneration induced in rat brain by MK-801: potential relevance to Alzheimer’s disease. *Neurobiol Dis* **5**: 305–322.
- Yu PL, Wu CI, Lee TS, Pan WH, Wang PS, Wang SW (2009). Attenuation of estradiol on the reduction of striatal dopamine by amphetamine in ovariectomized rats. *J Cell Biochem* **108**: 1318–1324.
- Zhou W, Cunningham KA, Thomas ML (2002). Estrogen regulation of gene expression in the brain: a possible mechanism altering the response to psychostimulants in female rats. *Brain Res Mol Brain Res* **100**: 75–83.