

Epigenetic Mechanisms in Stroke and Epilepsy

Jee-Yeon Hwang¹, Kelly A Aromolaran¹ and R Suzanne Zukin^{*,1}

¹Dominick P. Purpura Department of Neuroscience, Albert Einstein College of Medicine, New York, NY, USA

Epigenetic remodeling and modifications of chromatin structure by DNA methylation and histone modifications represent central mechanisms for the regulation of neuronal gene expression during brain development, higher-order processing, and memory formation. Emerging evidence implicates epigenetic modifications not only in normal brain function, but also in neuropsychiatric disorders. This review focuses on recent findings that disruption of chromatin modifications have a major role in the neurodegeneration associated with ischemic stroke and epilepsy. Although these disorders differ in their underlying causes and pathophysiology, they share a common feature, in that each disorder activates the gene silencing transcription factor REST (repressor element 1 silencing transcription factor), which orchestrates epigenetic remodeling of a subset of 'transcriptionally responsive targets' implicated in neuronal death. Although ischemic insults activate REST in selectively vulnerable neurons in the hippocampal CA1, seizures activate REST in CA3 neurons destined to die. Profiling the array of genes that are epigenetically dysregulated in response to neuronal insults is likely to advance our understanding of the mechanisms underlying the pathophysiology of these disorders and may lead to the identification of novel therapeutic strategies for the amelioration of these serious human conditions.

Neuropsychopharmacology Reviews (2013) **38**, 167–182; doi:10.1038/npp.2012.134; published online 15 August 2012

Keywords: epigenetics; global ischemia; status epilepticus; gene silencing factor REST; microRNAs; neurodegeneration

INTRODUCTION

Epigenetics is influenced by environmental cues and has a role in all aspects of neuronal function, from embryogenesis and early brain development to tissue-specific gene expression and global gene silencing; therefore, it is reasonable to believe that dysregulation of epigenetics would have a significant role in neuropsychiatric disorders such as ischemic stroke and epilepsy (Levenson and Sweatt, 2005; Graff *et al*, 2011). Gene transcription is regulated by methylation of DNA at CpG dinucleotides and covalent modifications of highly conserved amino-acid residues in core histone proteins, which in turn modifies chromatin structure and the accessibility of chromatin to the core transcriptional machinery (Wood *et al*, 2006; Berger, 2007; Robison and Nestler, 2011). Chromatin-modifying proteins catalyze post-translational modifications, such as acetylation, methylation, ubiquitination, phosphorylation, SUMOylation, and adenosine diphosphate (ADP) ribosylation, which serve as epigenetic tags that promote or repress gene transcription (for review, see (Jenuwein and Allis, 2001; Kouzarides, 2007; Berger, 2007; Brandl

et al, 2009)). Moreover, emerging evidence points to a widespread role for microRNAs as modulators of translation of target genes at early stages of brain development, and are critical to dendritogenesis, synapse formation and synapse maturation (Schratt, 2009a; Siegel *et al*, 2011b). The epigenome is large and complex and the modifications can work together in various combinations to create different biological responses. This diverse and dynamic repertoire provides a mechanism by which genetic predisposition, experience, and changes in neural function can operate together to create cellular and behavioral responses. Dysregulation of the vast epigenome could contribute to the complex cause for neurological disease and disorders.

This article reviews the role of epigenetic modifications of genes involved in synaptic function and plasticity and how dysregulation of these genes contribute to pathological conditions with cognitive impairments. In addition, the article highlights the potential power of drugs that target the epigenetic machinery in developing new therapeutic approaches for two neurological disorders, stroke and epilepsy. These two disorders are of particular interest because unlike late-onset, progressive neurodegenerative disorders such as Alzheimer's, Parkinson's, and Huntington's disease, stroke and epilepsy can arise from a single traumatic event. Moreover, there is considerable comorbidity in these two disorders; 11% of adult onset epilepsy occurs as a result of a

*Correspondence: Dr RS Zukin, Dominick P. Purpura Department of Neuroscience, Albert Einstein College of Medicine, 1300 Morris Park Ave, New York, NY 10461, USA. Tel: +1 718 430 2160, Fax: +1 718 430 8930, E-mail: Suzanne.zukin@einstein.yu.edu

Received 3 April 2012; revised 26 May 2012; accepted 30 May 2012

stroke (Rakhade and Jensen, 2009), indicating that there may be common triggers and mechanisms underlying impaired memory for these two disorders.

EPIGENETIC MODIFICATIONS

Epigenetic modifications such as DNA methylation and histone modifications reflect environmental influences that are not 'hard-wired' into the DNA sequence and are critical to genomic reprogramming during normal brain development, tissue-specific gene expression and global gene silencing (Levenson and Sweatt, 2005; Graff *et al*, 2011). During normal brain development, the connections between neurons and neuronal circuitry undergo remodeling in response to environmental cues. An emerging view is that environmental factors can also contribute to disruptions in synaptic signaling and neuronal survival (Day and Sweatt, 2011). Thus, epigenetic modifications such as methylation of DNA and histones are associated not only with learning and memory, but also with deficits in memory arising as a consequence of ischemic stroke or seizures.

DNA Methylation

DNA methylation is a key epigenetic signaling tool used by the cell to lock genes in the 'off' position. Properly established and maintained patterns of DNA methylation are essential for normal brain development, lineage commitment, and normal functioning of the adult organism (Robertson, 2005; Guibert *et al*, 2009; Senner, 2011). DNA methylation provides stable indexing marks that extend over long chromosomal domains, giving rise to 'memorized' states of gene expression that may be inherited from one cell generation to the next (Borrelli *et al*, 2008; Day and Sweatt, 2011). Methylation of mammalian DNA is catalyzed by DNA methyltransferases (DNMTs), chromatin-modifying proteins that establish and maintain unique DNA methylation patterns in different cells in different contexts. Three active members of the DNMT family (DNMT1, DNMT3a, and DNMT3b) are expressed in mammals. Although DNMT1 ensures faithful maintenance of DNA methylation, DNMT3a and DNMT3b are essential for *de novo* DNA methylation (Okano *et al*, 1999; Song *et al*, 2012). While DNA methylation at a gene promoter is typically associated with transcriptional repression, DNA methylation in gene-bodies is typically associated with active gene transcription (Ball *et al*, 2009; Laurent *et al*, 2010).

Aberrant DNA methylation is implicated in a wide range of human diseases, including cancer, imprinting disease, repeat-instability neurobehavioral diseases such as Fragile X syndrome (Robertson, 2005), neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, and disorders such as ischemic stroke and epilepsy (Petrij *et al*, 1995; Bartsch *et al*, 2005; Roelfsema *et al*, 2005; Urdinguio *et al*, 2009; Qureshi and Mehler, 2010a).

Histone Modifications and Transcriptional Regulation

Nucleosomes are the basic unit of chromatin and are composed of 147bp DNA wrapped around a histone octamer (two each of H2A, H2B, H3, H4), linker histone (H1) and histone variants (H2A.X, H2A.Z, H3.3) (Ruthenburg *et al*, 2007). Each histone has a central domain and unstructured N-terminal tails that contain multiple potential sites for post-translational modifications including acetylation, methylation, phosphorylation, ubiquitination, SUMOylation, and ADP ribosylation. These marks are added or removed by a specific set of chromatin-modifying enzymes (Ruthenburg *et al*, 2007). Distinct, site-specific histone modifications act sequentially or in combination to form a 'histone code' read by other proteins to bring about distinct downstream events, which in turn mediate active gene transcription or repression (Ruthenburg *et al*, 2007).

Histone Acetyltransferases

Histone acetyltransferases (HATs) confer acetyl moieties to lysine and arginine residues in the N-terminal tails of core histone proteins. Acetylation relaxes chromatin structure, enabling the transcriptional machinery to access the transcriptional start site. HATs not only modify histones, but also other substrates such as transcription factors (Kouzarides, 2000), and are therefore also referred to as lysine acetyltransferases to reflect their array of targets (Aka *et al*, 2011). Although HATs can affect global gene expression by acetylating large regions of chromatin, including coding and non-promoter regions, they can also act in a gene-specific manner by acetylating core histones at specific gene promoters (Dekker and Haisma, 2009; Selvi *et al*, 2010).

Histone acetylation has a critical role in memory and synaptic function and its dysregulation is linked to neurological disease. For example, mutations in the genes encoding either of the two well-known HATs, CBP and p300, leads to Rubinstein-Taybi syndrome, an autism spectrum disorder (ASD) characterized by intellectual disabilities and physical abnormalities (Petrij *et al*, 1995; Bartsch *et al*, 2005; Roelfsema *et al*, 2005; Urdinguio *et al*, 2009). In addition, CBP has a role in the pathophysiology of Huntington disease (Jiang *et al*, 2006).

Histone Deacetylases

The control of histone acetylation depends on the balance between two sets of chromatin-modifying enzymes, HATs and histone deacetylases (HDACs). The HDAC superfamily is made up of four classes (class I, II, III, IV), with class I and II being the most studied with prominent expression in the brain (for review see: (Gray and Ekstrom, 2001; de Ruijter *et al*, 2003; Morrison *et al*, 2007; Haggarty and Tsai, 2011)). HDACs remove acetyl groups from core histone proteins, thereby condensing the chromatin and repressing

transcription. HDACs also remove acetyl moieties from other proteins such as transcription factors and thereby alter cellular function (Morrison *et al*, 2007; Morris *et al*, 2010). HDACs do not bind DNA directly, but rather function as part of a multi-protein complex to recognize other factors such as transcriptional activators and repressors, which direct them to their target genes (de Ruijter *et al*, 2003; Carey and La Thangue, 2006).

Histone Methylation

Methylation occurs not only on cytosine bases in DNA, but also on lysine and arginine residues in the N-terminal tails of histone proteins (Scharf and Imhof, 2011). The impact of marks of methylation is complex in that its impact is determined in a site-specific manner and by the number of methylation marks that are conferred at a specific residue. The impact of methylation is site-specific, in that it can promote either gene activation or gene repression, depending on the exact lysine or arginine that is marked. Activation is signaled by methylation of lysines 4, 36, and 79 on histone 3 (H3K4, H3K36, H3K79) and repression is signaled by methylation of lysines 9 and 27 on histone 3 and lysine 20 on histone 4 (H3K9, H3K27, H4K20) (Kouzarides, 2007; Borrelli *et al*, 2008). Methylation marks are reversible in that they are removed by histone demethylases such as lysine-specific demethylase 1 (LSD1) (Lee *et al*, 2005; Shi, 2005). Rather than changing chromatin structure directly, the main function of methylation is to attract proteins that can modify chromatin (Cloos, 2011). Methylation itself is also important in cognitive functions and intellectual disabilities (Tahiliani *et al*, 2007). Moreover, histone demethylases are implicated in diseases such as cancer, autism, X-linked mental retardation, and schizophrenia (Cloos, 2011).

MicroRNAs in Epigenetic Remodeling

microRNAs are short, single-stranded RNAs that regulate gene expression post-transcriptionally, typically by binding recognition motifs within the 3'-untranslated regulatory regions (UTRs) of target mRNAs (Siegel *et al*, 2011b). Although best known for their role in regulation of translation (Kasinski and Slack, 2011; Salta and De, 2012), microRNAs can also regulate RNA transcription (Kim *et al*, 2008; Benhamed *et al*, 2012). Importantly, microRNAs regulate expression of key epigenetic regulators including DNMTs, HDACs, polycomb group proteins, and MeCP2 (Sato *et al*, 2011). Moreover, a subset of microRNAs such as let-7a, miR-9, miR-34a, miR-124, miR-137, miR-148, and miR-203 are regulated by epigenetic mechanisms including DNA methylation and histone modification (Sato *et al*, 2011). Interestingly, although miR-9 represses REST (repressor element 1 silencing transcription factor) and miR-9* represses CoREST, REST itself represses miR-9 and miR-9* (Packer *et al*, 2008). Thus, microRNAs can participate in 'epigenetic-miRNA regulatory circuits' (Sato *et al*, 2011).

Emerging evidence points to a widespread role for microRNAs as key modulators of target gene expression in neurons. microRNAs have a pivotal role in the late stages of neural differentiation, maintenance, and plasticity (Schratt, 2009b) and early stages of brain development including dendritogenesis, synaptogenesis, and synapse maturation. A key feature of neuronal microRNAs is their ability to regulate networks of neuronal or nonneuronal genes in an activity- or experience-dependent manner (Schratt, 2009b). Moreover, microRNAs achieve spatial specificity of gene expression in that they can regulate mRNA translation locally in the synaptodendritic compartment in an activity-dependent manner (Schratt, 2009b). As an example, miR-132 is critical for hippocampal spine formation and the inflammatory response (Shaked *et al*, 2009; Magill *et al*, 2010). miR-132 is thought to function together with FMRP to regulate translation in dendrites. Dysregulation of miR-132 and its target genes is implicated in the pathophysiology of Fragile X syndrome (Edbauer *et al*, 2010) and schizophrenia (Miller *et al*, 2012).

STROKE

Stroke is the leading cause of disabilities in adults in the United States, and is the second leading cause of death worldwide (Moskowitz *et al*, 2010; Ofengeim *et al*, 2011). Of the 600 000 new victims each year, 30% die and another nearly 30% become severely and permanently disabled. Most phase III clinical trials conducted to date have failed. Currently, there is no known treatment or cure for the neurodegeneration associated with this debilitating and devastating disease. It is thus critical that new molecular mechanistic information be revealed and be brought into the translational arena with the goal of identifying new therapeutic strategies for neuroprotection, rescue, and repair.

Transient forebrain or global ischemia arises as a consequence of cardiac arrest, cardiac surgery, profuse bleeding, near-drowning, and carbon monoxide poisoning and affects ~200 000 Americans each year (Liou *et al*, 2003; Moskowitz *et al*, 2010; Ofengeim *et al*, 2011). Neurological sequelae include deficits in learning and memory. Although all forebrain areas experience oxygen and glucose deprivation, a brief ischemic insult causes cell death primarily of CA1 pyramidal neurons, critical to memory formation and retrieval. Neuronal death, exhibiting characteristics of apoptosis and necrosis, is not detected until 2–3 days after ischemia (Liou *et al*, 2003; Moskowitz *et al*, 2010; Ofengeim *et al*, 2011). By 1 week after ischemia, the CA1 pyramidal cell layer can be virtually ablated. During the ischemic episode, cells exhibit a transient, early rise in intracellular Ca^{2+} , depolarize and become inexcitable; ambient glutamate rises by ~4-fold to ~2 μ M. After reperfusion, cells appear morphologically normal, exhibit normal intracellular Ca^{2+} and regain the ability to generate action potentials for 24–72 h after the insult. A myriad of death cascades are activated, including caspase-dependent and independent apoptosis, mitochondrial release of cytochrome *c*,

Smac/DIABLO and AIF, excitotoxicity, lipid peroxidation, and swelling (Liou *et al*, 2003; Lo *et al*, 2003; Zukin *et al*, 2004). Ultimately, there is a late rise in intracellular Ca^{2+} and Zn^{2+} , and death of CA1 neurons ensues, exhibiting hallmarks of apoptosis and necrosis. Although the molecular mechanisms underlying ischemia-induced death are not definitively understood, the substantial delay between insult and neuronal death is consistent with a role for transcriptional changes in cell death (Liou *et al*, 2003; Moskowitz *et al*, 2010; Ofengeim *et al*, 2011).

Transcriptional Regulation by REST During Stroke

REST, also known as NRSF (neuron-specific silencing factor), is a gene-silencing transcription factor that is widely expressed during embryogenesis and has a strategic role in terminal neuronal differentiation (Roopra *et al*, 2001; Ballas and Mandel, 2005; Ooi and Wood, 2007). In pluripotent stem cells and neural progenitors, REST actively represses a large array of coding and noncoding neuron-specific genes important to synaptic plasticity and structural remodeling, including synaptic vesicle proteins, channels, neurotransmitter receptors, and microRNAs that regulate networks of neuronal or non-neuronal genes. Bioinformatics analysis predicts nearly 2000 putative REST targets in the mammalian genome, including both coding and noncoding genes (Bruce, 2004; Conaco *et al*, 2006; Qureshi and Mehler, 2009). The interplay between REST and microRNAs is a new, emerging topic (Conaco *et al*, 2006; Singh *et al*, 2008; Johnson *et al*, 2008; Packer *et al*, 2008). During the late stages of neuronal differentiation, loss of REST is critical to acquisition of the neuronal phenotype (Su *et al*, 2006).

In mature neurons REST is quiescent, but can be activated in selectively vulnerable hippocampal neurons by insults such as global ischemia (Calderone *et al*, 2003; Formisano *et al*, 2007; Noh *et al*, 2012) and epileptic seizures (Palm *et al*, 1998; Garriga-Canut *et al*, 2006; Gillies *et al*, 2009) and aberrantly accumulates in the nucleus of selectively vulnerable striatal neurons in Huntington's disease (Zuccato *et al*, 2003, 2007).

A fundamental mechanism by which REST silences target genes is that of epigenetic remodeling (Borrelli *et al*, 2008). REST binds the RE1 element of target genes and recruits CoREST (Andres *et al*, 1999; Ballas *et al*, 2001) and mSin3A (Naruse *et al*, 1999; Huang *et al*, 1999; Grimes *et al*, 2000; Roopra *et al*, 2001), corepressor platforms which in turn recruit HDACs-1 and 2. HDACs deacetylate core histone proteins and affect dynamic and reversible gene silencing (Roopra *et al*, 2001; Ballas and Mandel, 2005; Ooi and Wood, 2007). REST mediates long-term gene silencing by associating with the site-specific histone methyltransferase G9a, which promotes dimethylation of histone 3 at lysine 9 (H3K9me2) *via* CoREST-dependent (Lunyak *et al*, 2002; Ballas *et al*, 2005) and -independent (Roopra *et al*, 2004) mechanisms and with LSD1, which removes mono- and dimethyl groups from lysine 4 of H3 (Lee *et al*, 2005; Shi,

2005). In addition, REST recruits methyl-CpG-binding protein 2 (MeCP2) (Lunyak *et al*, 2002; Ballas *et al*, 2005), a transcriptional repressor, which binds to hotspots of methylated CpG dinucleotides in gene promoters where it complexes with other repressors (Feng and Nestler, 2010). Importantly, MeCP2 is a reader of epigenetic marks on histones and DNA (Borrelli *et al*, 2008).

A prevailing view has been that REST is a master transcriptional regulator of neuronal genes in pluripotent stem cells and neural progenitors and that loss of REST during the late stages of neural differentiation by ubiquitin-based proteasomal degradation (Westbrook *et al*, 2008; Guardavaccaro *et al*, 2008) is required for acquisition of the neural phenotype (Ballas *et al*, 2005). An earlier paper by our group broadened this view (Calderone *et al*, 2003). Experiments involving molecular and genetic approaches showed that ischemic insults trigger activation of REST in mature hippocampal neurons destined to die and that the increase in REST correlates with a decrease in histone acetylation and gene silencing of GluA2. This is significant in that the GluA2 subunit prevents Ca^{2+} influx *via* AMPA receptors (AMPA), is essential to synaptogenesis, long-lasting forms of synaptic plasticity and neuronal death (Kwak and Weiss, 2006; Liu and Zukin, 2007). Acute knockdown of REST in hippocampal slices subjected to oxygen glucose deprivation (OGD), an *in vitro* model of ischemia, prevented GluA2 downregulation and neuronal death (Calderone *et al*, 2003). While compelling, these findings raised new questions. Are other synaptic proteins regulated by REST in insulted neurons? Are REST and corepressors recruited to the promoters of target genes and, if so, does the corepressor complex orchestrate epigenetic remodeling and gene silencing? Is REST causally related to neuronal death in a clinically relevant model of ischemic stroke?

A more recent study by our group employed a targeted ChIP-chip analysis with a microarray containing all known functional targets of REST to identify genes with altered REST occupancy in CA1 tissue from animals subjected to global ischemia (Noh *et al*, 2012). Of note, REST was enriched at promoters of a subset of target genes vital to synaptic function, of which *gria2*, the gene that encodes the AMPAR subunit GluA2, was the most affected. These findings represent an advance over previous studies, in which they show that, in addition to *gria2*, other genes important to synaptic plasticity and structural remodeling are functional targets of REST in fully differentiated neurons destined to die. Moreover, these results reinforce the notion that, although REST can repress thousands of putative target genes, the subset that is transcriptionally responsive is cell-type- and context-specific.

To address the mechanism by which REST silences target genes, a more in-depth analysis of *gria2* was performed. The *gria2* gene was the leading candidate in the ChIP-chip analysis (Noh *et al*, 2012). Moreover, the role of GluA2 silencing and expression of GluA2-lacking, Ca^{2+} -permeable AMPARs in neuronal death are well-established (Kwak and Weiss, 2006; Liu and Zukin, 2007; Isaac *et al*, 2007).

A critical factor in ischemic cell death is GluA2-lacking AMPARs, which mediate influx of toxic Ca^{2+} and Zn^{2+} in vulnerable CA1 neurons (Pellegrini-Giampietro *et al*, 1992; Gorter *et al*, 1997; Opitz *et al*, 2000; Calderone *et al*, 2003). The GluA2 subunit renders AMPARs impermeable to Ca^{2+} / Zn^{2+} , modifies channel kinetics and conductance and receptor targeting to synapses (Liu and Zukin, 2007). Thus, an alteration in *gria2* gene expression would have profound implications for synaptic plasticity and neuronal survival (Liu and Cull-Candy, 2000; Liu *et al*, 2004; Liu and Zukin, 2007). GluA2 is a known gene target of REST (Myers *et al*, 1998; Huang *et al*, 1999).

Ischemia induced enrichment of REST, mSin3A, CoREST, G9a, and MeCP2, at the *gria2* promoter in postischemic CA1 (but not CA3). CoREST and mSin3A serve as corepressor platforms that recruit HDACs 1 and 2, key components of the REST complex, which promote removal of acetyl groups from core histone proteins H3 and H4. Ischemia induced site-specific deacetylation of H3, indicative of gene repression. REST also recruits G9a (Ooi and Wood, 2007; Qureshi and Mehler, 2009), a protein that promotes site-specific methylation of H3. Ischemia induced methylation of H3 at lysine 9, but not lysine 4, an epigenetic signature of gene repression. These findings indicate that the REST complex is active in insulted CA1 and document epigenetic remodeling of the REST target GluA2 in neurons destined to die (Noh *et al*, 2012) (Figure 1a).

The affinity of REST on *gria2* could be due to the fact that it is well known in the chromatin world that the epigenetic landscape at promoter sites determines the binding affinity of DNA-binding proteins/regulatory proteins, which act to repress or promote active gene transcription. The enrichment or depletion of DNA-binding proteins at a given promoter

site depends not only on the cellular or nuclear abundance, but also on binding affinity of the regulatory element for the binding protein (Bruce *et al*, 2009; Yu *et al*, 2011). An additional and nontrivial point to account for the increased REST affinity is that of multivalency, the situation in which two or more DNA-binding proteins or epigenetic marks bound at the same promoter act synergistically to promote repression (or activation) and alter the affinity/capacity for a given DNA-binding protein, relative to the affinity observed in the absence of the marks (Ruthenburg *et al*, 2007).

To address a causal relation between REST-dependent epigenetic remodeling and neuronal death, the impact of the HDAC inhibitor trichostatin A (TSA) was examined (Wang *et al*, 2012; Noh *et al*, 2012). A single, acute injection of the HDAC inhibitor TSA, administered to animals before or after an ischemic episode, ameliorated neuronal injury. This finding documents a role for epigenetic remodeling in the neuronal death associated with ischemic stroke. Moreover, it has important clinical implications and suggests that HDAC inhibitors may be a promising avenue for intervention in the neurodegeneration associated with ischemic stroke. To address a role for REST in ischemia-induced neuronal death, two additional strategies were undertaken (Noh *et al*, 2012). We showed that RNAi-mediated knockdown or inhibition of REST by dominant-negative REST prevented epigenetic remodeling, GluA2 silencing and neuronal death. These findings indicate that REST is causally related to epigenetic remodeling and neuronal death and adds ischemic stroke to the growing list of diseases involving REST dysregulation (Figure 1a). Moreover, the study has broad implications for our understanding of molecular mechanisms underlying neurodegenerative disorders.




Figure 1. Model showing REST-dependent epigenetic remodeling of the *gria2* promoter in response to ischemic stroke or seizures. Global ischemia (left) or seizures (right) activate REST. REST binds to the RE1 element within the promoter of its target gene *gria2* and recruits mSin3A and CoREST, HDACs-1/2, G9a and MeCP2. The REST-corepressor complex promotes epigenetic remodeling of core histone proteins at the *gria2* promoter. This, in turn, represses GluA2 expression, leading to formation of GluA2-lacking, Ca^{2+} -permeable AMPARs. Modified with permission from (Noh *et al*, 2012).

DNA Methylation in Animal Models of Stroke

Several studies have explored global alterations in DNA methylation associated with animal models of ischemic stroke. In one study, induction of focal ischemia by means of transient MCAO markedly increased DNA methylation and neuronal death in the cerebral cortex of wild-type mice. Administration of 5-aza-2'-deoxycytidine, a demethylase and broad-spectrum inhibitor of DNMT, conferred neuroprotection in wild-type mice (Endres *et al.*, 2000). Moreover, mice heterozygous for the gene encoding DNMT1 (*Dnmt1^{S/+}*) exhibited reduced methylation and increased resistance to ischemic damage, consistent with the concept that enhanced DNA methylation, and presumably gene silencing, are associated with enhanced susceptibility to brain injury (Endres *et al.*, 2000). Although transgenic mice with reduced levels of DNMT1 were resistant to cerebral ischemia-induced brain injury, mice with complete deletion of DNMT1 did not show protection (Endres *et al.*, 2000). These findings suggest that the levels of DNMT expression and DNA methylation must be tightly regulated to protect neurons from ischemic insults.

In recent years, new technologies have enabled identification and quantitative assessment of alterations in DNA methylation at specific loci in the genome in response to ischemic insults. A study that links ischemia to alterations in DNA methylation at promoters of specific genes focused on the Na⁺-K⁺-2Cl⁻ (NKCC1) co-transporter type-1 gene (Lee *et al.*, 2010). It is well established that NKCC1 is important in GABA(A) receptor-mediated responses that switch from depolarization to hyperpolarization during postnatal development. NKCC1 is downregulated during postnatal development (Rivera *et al.*, 1999), but can be reactivated in adult neurons in response to focal ischemic injury (Pond *et al.*, 2006). Consistent with this, DNA methylation of the NKCC1 promoter increased during postnatal development, but decreased in response to ischemic injury (Lee *et al.*, 2010). These findings are consistent with the notion that DNA methylation of the NKCC1 gene promoter is responsible for the increase in NKCC1 expression in postischemic neurons.

Histone Modifications and Transcriptional Regulation in Stroke

In the past decade HDAC inhibitors have emerged as potential therapeutics for stroke. A number of studies have shown the ability of HDAC inhibitors to afford neuroprotection in the MCAO model of ischemic stroke (Gibson and Murphy, 2010). In one study, Chiarugi and colleagues reported that the potent broad-spectrum HDAC inhibitor suberoylanilide hydroxamic acid (SAHA) administered intraperitoneally to mice at times after induction of MCAO prevented deacetylation of core histone H3, promoted expression of prosurvival proteins Bcl-2 and Hsp70 and reduced infarct volume, indicating a neuroprotective action for SAHA (Faraco *et al.*, 2006).

Several additional studies indicate that the broad spectrum HDAC inhibitor TSA confers protection against

ischemia-induced neuronal death. One study showed that TSA promotes neuronal viability following OGD *in vitro* and reduces neuronal injury in wild-type mice subjected to MCAO *in vivo* (Wang *et al.*, 2012). Neuroprotection by TSA in these models occurred through activation of the transcription factor, nuclear factor erythroid 2 (Nrf2) and its downstream targets (Wang *et al.*, 2012). As an example, Nrf2 binds the promoter of the antioxidant sulfiredoxin and increases its expression, which promotes neuronal survival in the face of oxidative stress (Soriano *et al.*, 2009). The second study showed that a single, acute injection of TSA administered to animals after global ischemia *in vivo*, a clinically relevant model of ischemic stroke, ameliorates neuronal injury (Noh *et al.*, 2012), further emphasizing the role of histone acetylation in ischemia-induced neuronal death. A more recent study demonstrated that not only SAHA, but also the class 1 HDAC inhibitor MS-275, afford protection against OGD-induced neuronal death, promote functional recovery of axons, and preserve white matter cellular architecture (Baltan *et al.*, 2011b). Moreover, HDAC inhibitors conserved ATP levels and reduced excitotoxicity, thereby preserving white matter structure and function (Baltan *et al.*, 2011b).

Histone methylation and demethylation also affect the pathology of global ischemia. As an example, transient, forebrain global ischemia upregulates the site-specific demethylase LSD1, which removes methyl groups from lysine 4 on core histone protein H3 (H3K4) (Zhang *et al.*, 2010). The upregulation occurs in hippocampal CA1 and dentate gyrus (DG), as well as in cerebral cortex. Of these regions, highest levels of LSD1 were observed in the DG, a region relatively resistant to neuronal death. Although the increase in LSD1 expression was rapid in the hippocampal CA1 and DG (<1 h), it was induced later (by 6 h) in cortex. These differences could contribute to differences in neuronal viability observed in the hippocampus *vs* cortex after ischemia.

Another study that revealed a role for histone methylation in the neuronal death associated with global ischemia, focused on the gene encoding μ -opioid receptor (MOR-1). MOR-1 is a member of the G protein-coupled receptor superfamily and is activated by endogenous opioid peptides (Kieffer and Gaveriaux-Ruff, 2002). In rats subjected to global ischemia, MOR-1 is downregulated in the hippocampal CA1 (Formisano *et al.*, 2007). Ischemia promoted REST binding to the MOR-1 promoter by 12 h after ischemia, assessed by ChIP, consistent with the possibility that REST mediates MOR-1 downregulation. Moreover, ischemia promoted dimethylation of lysine 9 (H3K9), but not lysine 4 (H3K4) on histone 3 at the MOR-1 promoter, an epigenetic signature of gene repression. MOR-1 is expressed on GABAergic inhibitory interneurons of the CA1, where it inhibits the release of GABA onto pyramidal cells and thereby increases neuronal excitability. Thus, downregulation of MOR-1 in selectively vulnerable CA1 neurons after ischemia is neuroprotective, and may represent a failed attempt to promote neuronal survival (Formisano *et al.*, 2007).

The Role of microRNAs in Stroke

Recent studies indicate striking alterations in microRNA expression in response to ischemic stroke and ischemic preconditioning in animal models and in young patients who suffer stroke (for review see (Rink and Khanna, 2011)). Neuronal death resulting from focal ischemia induced experimentally in animals occurs *via* caspase-dependent and -independent mechanisms (Du *et al*, 2004; Yuan *et al*, 2009; Liu *et al*, 2009; Siegel *et al*, 2010). The X-linked inhibitor of apoptosis (XIAP) is an endogenous inhibitor of caspases that has a pivotal role in the response to ischemic insults. Under physiological conditions, XIAP expression is higher in the brains of female *vs* male mice and is significantly decreased in response to stroke in females, but not males (Siegel *et al*, 2011a). A recent study provides evidence that miR-23a is a key regulator of XIAP expression. Ischemia promotes an increase in miR-23a expression and its enrichment at the 3'-UTR of XIAP in the cortex of female, but not male, mice. These findings implicate a possible role for miR-23a in the severity and functional outcome of stroke in a gender-specific manner (Siegel *et al*, 2011a).

Heat-shock proteins (HSP) are a class of functionally related proteins involved in protein folding and unfolding and cell survival. Dramatic upregulation of HSPs in response to stress is known as 'the heat-shock response'. The HSPs of the 70 kDa family, HSP72 (cytosol), GRP75 (mitochondria), and GRP78/BIP (endoplasmic reticulum), are evolutionarily conserved proteins that promote neuronal survival/protection in animal models of stroke (Rajdev *et al*, 2000; Hoehn *et al*, 2001; Kudo *et al*, 2008; Oida *et al*, 2008). A recent study identified miR-181 as a possible regulator of several HSP70 family members and validated HSP GRP78/BIP to be a target of miR-181 in response to ischemic brain injury. In this study, miR-181 increased in the core, where cells die, but decreased in the penumbra, where cells survive. The increase in miR-181a was associated with a decrease in GRP78, implicating translational repression. Although overexpression of miR-181a exacerbated neuronal death, reduced miR-181a activity increased GRP78 protein expression and reduced neuronal death (Ouyang *et al*, 2012). These data implicate miR-181 expression in repression of GRP78 and induction of neuronal death.

EPILEPSY

Epilepsy, defined as recurrent unprovoked seizures, affects 1–2% of the population world-wide (Hauser *et al*, 1993) and ranks second worldwide in prevalence among all neurological disorders (Rakhade and Jensen, 2009). Cognitive impairment, particularly memory disruption, is a major complicating feature of epilepsy. Sensory, motor, or autonomic function can also be affected (Fisher *et al*, 2005; Berg *et al*, 2010). Current therapies are largely unsatisfactory, as they suppress seizures, but do not affect

the underlying cause, are effective in only a subset of affected individuals, and are often toxic (Duncan *et al*, 2006; Galanopoulou *et al*, 2012). Epilepsy refers to a constellation of disorders and can be classified according to characteristic symptoms and signs, seizure type, cause, age of onset, and electroencephalographic patterns (Berg *et al*, 2010). For many years it was thought that epilepsy occurred exclusively as a result of cortical damage (McNamara, 1999). It is now known that sporadic epilepsy can also arise as a consequence of traumatic brain injury, stroke, abnormalities in brain wiring, toxic-metabolic etiologies, inflammation, autoimmunity, or an imbalance in the ratio of inhibitory to excitatory synaptic transmission (Berg *et al*, 2010). Epilepsy can also arise as a consequence of spontaneous or inherited gene mutations. Interestingly, the latter includes a vast number of channelopathies, in which recurrent seizures arise from single-point mutations, deletions, duplications, or expansions in genes encoding a component of a voltage- or ligand-gated ion channels, but also many other genes that directly or indirectly control brain development and neuronal function and activity (Catterall *et al*, 2008, 2010; Galanopoulou *et al*, 2012).

Epileptogenesis is the process whereby a neural network acquires the ability to manifest recurrent epileptic seizures *de novo* and in response to neuronal insults, as well as the development and progression of epilepsy (Rakhade and Jensen, 2009; Galanopoulou *et al*, 2012). Epileptogenesis has been conceptualized as a 'cascade' of molecular and cellular events (Rakhade and Jensen, 2009). In the first phase of epilepsies owing to structural/metabolic etiologies, seizures may develop on the order of minutes to days in response to a neuronal insult such as traumatic brain injury, and are accompanied by acute changes to neuronal networks and activation of immediate early genes (Rakhade and Jensen, 2009). Next, a silent or latent-stage ensues, involving alterations in transcription, neuronal death, and re-wiring, or inflammation. In addition, changes in neurogenesis, sprouting, and reorganization of neuronal networks contribute to the development of spontaneous seizures that increase in frequency over time and can last for weeks to months or years (Rakhade and Jensen, 2009). Emerging evidence indicates that alterations not only in transcription, but also epigenetic mechanisms, including DNA methylation, histone code modifications, chromatin remodeling, and modulation of the epigenetic machinery by noncoding RNAs are involved in the pathogenesis of human epilepsy and in the process of epileptogenesis (Qureshi and Mehler, 2010b).

Comorbidity with Autism

Children with epilepsy often have intellectual and developmental disabilities and co-morbidity of epilepsy with disorders such as anxiety, attention-deficit hyperactivity disorder and/or autism (Russ *et al*, 2012). A prominent comorbidity is that of children diagnosed with both autism and epilepsy (Berg *et al*, 2010; Tuchman *et al*, 2010; Bolton

et al, 2011). Approximately 20–30% of children with autism have seizures, with the highest rate in autistic children with severe intellectual disabilities (Amiet *et al*, 2008; Berg *et al*, 2010; Tuchman *et al*, 2010). Moreover, six percent of children who suffer a seizure very early in life (aged 1–2 years) develop ASDs with intellectual disabilities (Clarke *et al*, 2005). Although the mechanisms responsible for this high co-morbidity are as yet unknown, it is postulated that dysregulation of epigenetic programming may be involved.

As an example, Rett syndrome is an ASD, a neurobehavioral disorder characterized by mental retardation, stereotyped behaviors and recurrent seizures (Jian *et al*, 2006). Approximately 80% of children diagnosed with Rett syndrome exhibit seizures. Rett syndrome typically arises as a consequence of loss-of-function mutations in the X-linked gene MeCP2 (Amir *et al*, 1999; Abrahams and Geschwind, 2008). This finding links seizures not only to ASDs, but also to deficits in the epigenetic machinery. However, the correlation between onset and severity of epilepsy and MeCP2 genotype in Rett syndrome patients is controversial. Although some studies report such a correlation (Jian *et al*, 2006; Nectoux *et al*, 2008), others do not (Nissenkorn *et al*, 2010).

Other examples of ASDs that present with seizures are Prader–Willi syndrome and Angelman syndrome, ASDs that present with intellectual disabilities and developmental delays (Abrahams and Geschwind, 2008). Given that DNA methylation is critical for genomic imprinting, and that the defective gene in both syndromes lies within imprinted domains at 15q11–q13, it is likely that dysregulation of DNA methylation has a role. Although Prader–Willi syndrome arises from loss of paternally expressed genes in this region, Angelman syndrome arises from the loss of a single gene, *ube3a*, the gene encoding maternally expressed ubiquitin protein ligase E3A, which is imprinted only in the brain (Abrahams and Geschwind, 2008). Ube3a is required for maintaining plasticity during experience-dependent neocortical development, consistent with the notion that loss of neocortical plasticity contributes to the cognitive deficits associated with Angelman syndrome (Yashiro *et al*, 2009).

Transcriptional Regulation by REST in Epilepsy

The original paper that linked seizures to epigenetic remodeling of synaptic proteins was pioneered by Timmusk and colleagues who showed that kainate-induced seizures in rats, an animal model of status epilepticus, induces REST mRNA expression rapidly (<3 h after onset of seizures) in the hippocampal CA3 (Palm *et al*, 1998). A follow-up study by Dingledine and colleagues using a related animal model of status epilepticus, showed that seizures promote deacetylation of core histone protein H4 (a mark of gene repression) at the RE1 site of the *gria2* promoter (gene encoding the AMPAR subunit GluA2), while promoting an increase in acetylation of H4 (a mark of open chromatin and active gene transcription) at the promoter of brain-derived neurotrophic factor BDNF (Huang *et al*, 2002; Tsankova

et al, 2004). Although GluA2 expression was decreased, leading to an increase in GluA2-lacking, Ca²⁺-permeable AMPARs at CA3 synapses and neuronal death in CA3 (Grooms *et al*, 2000; Sanchez *et al*, 2001; Huang *et al*, 2002), BDNF expression was increased (Kokaia *et al*, 1995; Binder *et al*, 1999; Huang *et al*, 2002; Tsankova *et al*, 2004). Alterations in expression of these proteins contribute to the pathophysiology of recurrent seizures. Administration of the HDAC inhibitor TSA to rats before induction of seizures prevented and rapidly reversed the increase in histone deacetylation at the *gria2* promoter and blunted the down-regulation of GluA2 (Huang *et al*, 2002). These findings link epigenetic remodeling to seizure-induced silencing of GluA2 and, indirectly, seizure-induced neuronal death. Because *gria2* is a known target of REST, these findings also implicate REST-dependent epigenetic remodeling of GluA2 in response to seizures (Huang *et al*, 1999) (Figure 1b).

Another REST target implicated in epilepsy is the hyperpolarization-activated cyclic adenosine monophosphate-gated channel type 1 (HCN1). In normal brain, the HCN1 channel mediates the I(h) current, and keeps in check the intrinsic excitability of pyramidal cell dendrites in the entorhinal cortex and hippocampus (Lewis and Chetkovich, 2011). Accordingly, HCN1 null mice exhibit increased dendritic excitability and susceptibility to seizures in an experimental model of temporal lobe epilepsy (Huang *et al*, 2009). A recent study showed that seizures promote an increase in REST expression, enrichment of REST at the *hcn1* promoter, dimethylation of lysine 9 on core histone protein H3 (H3K9), an epigenetic mark of gene repression, and silencing of the *hcn1* gene (McClelland *et al*, 2011). Thus, REST-dependent epigenetic remodeling represses HCN1 expression and I(h) currents, leading to increased dendritic excitability and epileptiform activity. Disruption of REST binding to the *hcn1* promoter by means of decoy oligodeoxynucleotides prevented *hcn1* repression and restored its ability to maintain normal levels of dendritic excitability. Collectively, these studies indicate that REST is causally related to HCN1 repression, decreased dendritic excitability, and enhanced epileptiform activity in entorhinal cortical layer III pyramidal neurons (Shah *et al*, 2004; Jung *et al*, 2007, 2011) (Figure 1b).

DNA Methylation in Epilepsy

Recent studies provide evidence that aberrant DNA methylation may contribute to the pathophysiology of epilepsy. In the adult brain, the extracellular matrix protein Reelin is important for normal synaptic plasticity, dendritic morphology, and cognitive function (Herz and Chen, 2006). In the developing brain, Reelin helps maintain the correct laminar structure of granule cells in the DG. Brain specimens from humans diagnosed with temporal lobe epilepsy commonly show dispersion of granule cells, an anatomical hallmark of epilepsy, and a deficiency in expression of Reelin (Haas *et al*, 2002; Heinrich *et al*, 2006). A recent study revealed hypermethylation at the *reelin* promoter (Kobow *et al*, 2009), suggesting that

aberrant DNA methylation is causally related to the low levels of Reelin in the DG and link aberrant methylation to the pathophysiology of human epilepsy.

Another recent study performed an unbiased, genome-wide study of altered DNA methylation in tissue samples from the CA3 of control, epileptic tolerant and status epilepticus mice (Miller-Delaney *et al*, 2012). Epileptic tolerance is an evolutionarily conserved form of cerebral plasticity in which a brief period of seizures ('epileptic preconditioning') confers transient tolerance to a subsequent bout of seizures in the brain. A total of 321 genes showed altered DNA methylation in animals subjected to status epilepticus or to epileptic preconditioning (Miller-Delaney *et al*, 2012). Of these, 293 genes (> 90%) showed a decrease in DNA methylation (hypomethylation). Only 15 genes, many of which were associated with nuclear function such as DNA binding and transcriptional regulation, were hypermethylated. These results were surprising, given that epileptic tolerance elicits transcriptional silencing it was thought that DNA would exhibit hypermethylation. Importantly, the genome-wide methylation profiles revealed upregulation of novel genes, not previously associated with epilepsy, such as the polycomb group proteins, in the vulnerable CA3 of epileptic tolerant animals. Polycomb proteins act *via* epigenetic mechanisms to silence potential mediators of neuronal death and promote cellular arrest, enabling adult neurons to survive neuronal insults (Zukin, 2010).

Histone Modification and Transcriptional Regulation in Epilepsy

Emerging evidence indicates that aberrant histone modifications and altered gene expression are hallmark features of animal models of status epilepticus and humans with temporal lobe epilepsy. As an example, a recent study showed that status epilepticus in rats promotes hyperacetylation of H4 and phosphorylation of H3 (marks of active transcription) at the *c-fos*, *c-jun*, and *cbp* promoters and increases *c-fos*, *c-jun*, and CBP expression (Sng *et al*, 2006). *c-fos* and *c-jun* are immediate early genes that are activated rapidly and transiently in response to cellular stimuli and promote cell growth, differentiation, and neuronal survival. CBP is a HAT and essential co-activator of the prosurvival transcription factor CREB. Pretreatment of rats with curcumin, an active component of turmeric with antioxidant, anti-inflammatory, and neuroprotective properties, and a HAT inhibitor, attenuated histone modifications, expression of immediate early genes *c-fos* and *c-jun*, and severity of status epilepticus (Sng *et al*, 2006). Together, these data suggest that histone modifications such as acetylation and phosphorylation have a pivotal role in regulation of genes involved in neuronal survival.

Another recent study showed that HDAC2, an HDAC implicated in brain development (Guan *et al*, 2009; Morris *et al*, 2010), is upregulated in humans with temporal lobe epilepsy and in animal models of status epilepticus (Huang *et al*, 2012). This is significant in that HDAC2 is a negative

TABLE 1 microRNAs with Altered Expression in Epilepsy

Upregulated microRNAs in epilepsy		
Jimenez-Mateos <i>et al</i> (2011)	Hu <i>et al</i> (2011)	Song <i>et al</i> (2011)
miR-10b	miR-22	miR-9*
miR-21	miR-24	miR-23a
miR-27a	miR-26a	miR-23b
miR-29a	miR-29b	miR-24
miR-30e	miR-30c	miR-27b
miR-132	miR-34a	miR-29a
miR-134	miR-99a	miR-99a
miR-139	miR-101-1	miR-126
miR-146b	miR-124a	miR-127
miR-153	miR-125a	miR-132
miR-199	miR-125b	miR-134
miR-200a	miR-132	miR-137
miR-219	miR-145	miR-139-5p
miR-323	miR-150	miR-140*
miR-326	miR-151	miR-146a
miR-375	miR-196b	miR-324-5p
miR-425	miR-199a	miR-379
miR-451	miR-213	miR-383
miR-487b	miR-375	
miR-507		
miR-509		
miR-518d		
Downregulated microRNAs in epilepsy		
Jimenez-Mateos <i>et al</i> (2011)	Hu <i>et al</i> (2011)	Song <i>et al</i> (2011)
miR-133b	miR-10b	miR-98
miR-330	miR-21	miR-185
miR-374	miR-25	miR-352
miR-381	miR-29a	let-7E
miR-422b	miR-181b	let-7D
miR-497	miR-181c	
miR-657	miR-215	

regulator of cognitive function (Graff *et al*, 2012). Accordingly, HDAC2 represses genes associated with synaptic plasticity and memory formation (*erg1*, *creb1*, *bdnf*, NMDA receptor subunits, *grin1*, *grin2a*, *grin2b*, and *c-fos*). These findings implicate upregulation of HDAC2 in the cognitive impairments associated with temporal lobe epilepsy.

Methylation of lysine residues in the N-terminal histone tail is dynamically regulated by the opposing actions of histone methyltransferases such as G9a and histone demethylases such as LSD1 and KDM5C (SMCX). KDM5C removes di- and tri-methyl marks from lysine 4 of H3 and represses gene transcription. Mutations in the histone demethylase KDM5C are linked to X-linked mental retardation and epilepsy (Tahiliani *et al*, 2007). KDM5C effects epigenetic remodeling and gene silencing by assembling with the transcriptional repressor REST and its corepressors

(CoREST, HDAC1 and -2, and G9a). Notably, KDM5C promotes REST-dependent repression of a subset of target genes such as BDNF and sodium channel type 2A (SCN2A) (Tahiliani *et al*, 2007). These genes are of interest because increases in their activity are implicated in the pathophysiology of epilepsy (Binder *et al*, 2001; Shi *et al*, 2011). Inactivating mutations in KDM5C impairs REST-dependent silencing of neuronal genes such as BDNF and SCN2A, thereby intensifying seizures and contributing to X-linked mental retardation.

In addition, heat-shock 27-kDa-associated protein 1 (HSPBAP1), a member of the histone demethylase family JmjC, is expressed in the anterior temporal neocortex of patients with intractable epilepsy, but not in normal controls (Xi *et al*, 2007; Cloos, 2011). Although it is not yet known by which mechanism upregulation of HSPBAP1 contributes to epilepsy or neuronal death, an attractive scenario is that HSPBAP1 represses expression of the protective protein HSP27, thereby exacerbating neuronal death.

The Role of microRNAs in Epilepsy

Recently, the ability of status epilepticus to alter the microRNA expression profile in the selectively vulnerable hippocampal CA3 was evaluated by means of microRNA array analysis, an unbiased, epigenome-wide approach to detect alterations in microRNA expression (Jimenez-Mateos *et al*, 2011; Hu *et al*, 2011; Song *et al*, 2011). In Table 1, we summarize alterations in microRNA expression associated with status epilepticus from three independent studies. In all three studies, miR-132 was strikingly upregulated in the hippocampal CA3 after status epilepticus (Table 1, underlined). miR-24, miR-29a, miR-99a, miR-134, and miR-357 were increased in at least two of the three studies (Table 1, boldface), identifying these microRNAs as regulated in response to seizures. In contrast, the three studies showed no consensus (or overlap) in downregulated microRNAs (Table 1, Lower panel). GO analysis classified upregulated microRNAs as components of ERK-MAPK signaling, TGF- β signaling, and/or tight junctions (Hu *et al*, 2011). A subset of the downregulated microRNAs are implicated in synaptic plasticity, and T-cell receptor signaling.

Alterations in expression of many of the microRNAs identified in the array analysis (miR-132, miR-let-7, miR-23a/b, miR-34a, miR-22, miR-125a, miR-21) were individually validated. For example, miR-132 was shown to be increased in the CA3 in an animal model of status epilepticus. *In vivo* microinjection of antagomirs against miR-132 depleted miR-132 in CA3. A functional role of miR-132 in anti-inflammation has been identified (Shaked *et al*, 2009). Since inflammation and blood-brain barrier opening are implicated in epileptogenesis, it is thought that increased miR-132 may contribute to epileptogenesis (Vezzani *et al*, 2011). In addition, miR-146, another inflammation-associated microRNA, is increased in reactive astrocytes (Aronica *et al*, 2010). These studies document involvement of microRNAs in regulation not only in neurons, but also in astroglia.

Preliminary studies in animal models of stroke and epilepsy indicate that specific microRNAs may serve as potential biomarkers for these neurological disorders (Liu *et al*, 2010). A recent study investigated microRNA expression profiles in the hippocampus and blood of animals 24h after they were subjected to global ischemia or kainic acid-induced status epilepticus. Analysis of their results identified specific microRNAs that changed at least two-fold in response to these neuronal insults. Following ischemia, miR-182 was upregulated and miR-223 and miR-210 were downregulated in both blood and brain; in addition, four microRNAs (miR-29c, miR-34b-3p, miR-98, miR-122) were downregulated in both blood and brain after epilepsy (Liu *et al*, 2010). These data support the possibility that a subset of microRNAs may prove useful as biomarkers in that blood levels of these microRNAs correlate with their levels in the injured hippocampus.

NOVEL THERAPEUTIC STRATEGIES

In the past decade, an explosion of studies has linked epigenetic modifications to human diseases such as ischemic stroke and epilepsy. Spurred with this information, the pharmaceutical industry has promoted a number of drugs that target the epigenetic machinery as potential therapeutic strategies to ameliorate the symptoms associated with stroke and epilepsy. An important focus has been on drugs that can penetrate the brain-blood barrier and have minimal toxicity. Several such drugs have been developed and are approved by the US Food and Drug Administration (FDA). The most common drugs in clinical trials to date are DNMT inhibitors and HDAC inhibitors (Table 2).

DNMT Inhibitors

The use of DNMT inhibitors as a pharmacological strategy is a quickly growing field. As an example, mice subjected to MCAO exhibit aberrant DNA methylation; administration of the DNMT inhibitor 5-aza-2'-deoxycytidine affords neuroprotection (Endres *et al*, 2000). Although DNMT inhibitors have not, as yet, been tested in animal models of status epilepticus or in humans with epilepsy, the finding of aberrant DNA methylation in status epilepticus (Kobow *et al*, 2009; Miller-Delaney *et al*, 2012) suggests that pharmacological agents that reverse DNA methylation are a high priority. The first generation of DNMT inhibitors developed for humans (Table 2) block DNMT function by preventing unbinding of DNMT from DNA. This, in turn, prevents methylation at neighboring sites on the DNA, leading to global demethylation, which would account, at least in part, for off-target effects (Kelly *et al*, 2010). Use of DNMT inhibitors for amelioration of neurological disorders may be difficult for a number of reasons. First, stable and bioavailable compounds would need to penetrate the blood-brain barrier to reach the affected brain region. These drugs would need to be of high potency and specificity to minimize off-target effects. Second, as these drugs are known to work well

TABLE 2 Drugs that Target the Epigenetic Machinery

	DNMT inhibitors/HDAC inhibitors	Developmental stage	Specificity	Basic research in epilepsy or stroke
DNMT inhibitors	5-Azacytidin (azacytidine, Vidaza)	Approved 2004	DNMTs	
	5-Aza-2-deoxycytidin (Dacogen, decitabine)	Approved 2006	DNMTs	Endres <i>et al</i> (2000, 2001)
	MG98	Phase I/II	DNMT I	
	Zebularine	Preclinical	DNMTs	
HDAC inhibitors	Valproic acid (VPA)	Approved 1986	Class I and II	Wang <i>et al</i> (2011); Xuan <i>et al</i> (2012)
	SAHA (Zolinza, vorinostat)	Approved 2006	Class I and II	Faraco <i>et al</i> (2006); Baltan <i>et al</i> (2011a, b)
	Romidepsin (Depsipeptid)	Approved 2009	Class I	
	LBH589 (Panobinostat)	Phase III	Class I and II	
	MS-275 (Entinostat)	Phase II	Class I	
	MGCD0103 (Mocetinostat)	Phase II	Class I	
	CI-994	Phase II	Class I and II	
	PXD101 (Belinostat)	Phase II	Class I and II	
	Baceca	Phase II	Class I	
	Savicol	Phase II	NA	
	PCI-24781 (Abexinostat)	Phase II	Class I and II	
	ITF2357 (Givinostat)	Phase II	NA	
	Kevertin	Phase I	Class I	
	CUDI-101	Phase I	Class I and II	
	AR-42	Phase I	Class I and II	
	4SC-202	Phase I	Class I	
ACY-1215	Phase I	Class II		
Sulforaphane	Phase I	Class I and II		

in rapidly dividing cells, they may be less efficacious in treating postmitotic cells such as neurons. Third, DNMT inhibitors would need to be continually administered over days or weeks (Kelly *et al*, 2010). Recent studies have therefore focused on development of second-generation DNMT inhibitors, which have increased specificity and reduced toxicity. An example is MG98, a second-generation antisense oligonucleotide, which inhibits translation of the mRNA encoding human DNMT1 (Plummer *et al*, 2009). This and related second-generation DNMT inhibitors hold promise for the amelioration of the neuronal death and cognitive deficits associated with stroke and epilepsy.

HDAC Inhibitors

HDAC inhibitors are actively and intensively being evaluated in pre-clinical studies and clinical trials for their ability to intervene in the neurodegeneration and cognitive deficits associated with stroke and epilepsy. In pre-clinical studies, four main classes of HDAC are under investigation: the short-chain fatty acids, such as sodium butyrate, phenylbutyrate, and valproic acid, the hydroxamic acids, such as TSA and SAHA, the epoxyketones, such as trapoxin, and the benzamides (Abel and Zukin, 2008). SAHA and TSA promote neuronal survival in animal models of ischemia (Gibson and Murphy, 2010; Baltan *et al*, 2011b; Wang *et al*, 2012; Noh *et al*, 2012) and epilepsy (Huang *et al*, 2002; Kobow and Blumcke, 2011). Intriguingly, valproic acid, which is commonly prescribed as an anticonvulsant drug,

was subsequently discovered to be an HDAC inhibitor (Gottlicher *et al*, 2001). In spite of their neuroprotective actions in epilepsy and stroke, valproic acid and other broad-spectrum HDAC inhibitors can cause global DNA demethylation (Dong *et al*, 2010) and histone acetylation (Eleuteri *et al*, 2009), resulting in cell death. These findings suggest that HDAC inhibitors are relevant and feasible targets for developing clinical drugs, but at the same time, point to the need to develop new HDAC inhibitors with greatly enhanced specificity and reduced toxicity.

FUTURE RESEARCH DIRECTIONS

The epigenetics of brain disorders is a new, emerging topic with great potential for development of novel therapeutic strategies for intervention in the neurodegeneration and cognitive deficits associated with stroke and epilepsy. In the past decade, remarkable progress has been made in documenting a role for epigenetics in brain disorders such as seizures and ischemia. A case in point is the discovery that the gene-silencing transcription factor REST is activated in adult neurons in response to ischemia and orchestrates epigenetic remodeling and silencing of subsets of transcriptionally responsive target genes. Also of note, microRNAs are emerging as key modulators of post-transcriptional gene regulation critical to normal brain development. In the past several years, neural-specific microRNAs have been discovered; many of these act to

suppress networks of neuronal or non-neuronal genes. In addition, microRNAs can act in synaptodendritic compartments to regulate local translation of proteins important to synaptic plasticity and neuronal survival (Wang *et al*, 2010). Dysregulation of microRNAs in neurodegenerative diseases and in response to neuronal insults can contribute to neuronal death (Johnson *et al*, 2008; Hebert *et al*, 2009).

The past decade has witnessed an explosion of new information focused on site-specific epigenetic modifications of histone proteins, the concept that modifications at multiple sites can act synergistically to more dramatically promote transcriptional activation or repression ('multivalency'), the discovery of new transcription factors and other chromatin-modifying proteins and the mechanisms by which they activate or repress target genes. Moreover, extraordinary progress has been made in developing new technologies to examine epigenome-wide alterations in DNA methylation, histone modifications, and transcription factor occupancy in neurologic diseases and disorders.

Important commonalities between the mechanisms underlying ischemia and seizures have been discovered. In both disorders, the gene silencing transcription factor REST is activated in selectively vulnerable mature hippocampal neurons and orchestrates epigenetic remodeling and silencing of a subset of neuron-specific target genes. This observation raises the possibility that new therapeutic strategies used to treat one disorder such as epilepsy may also hold promise for treatment of other brain disorders such as stroke and possibly even neurodegenerative disorders such as Huntington's disease.

microRNAs represent an exciting alternative mode of post-transcriptional regulation in brain in response to neuronal insults such as ischemia or seizures. An important future direction would be to identify not only subsets of microRNAs that are up- or downregulated in these pathological conditions, but also the downstream targets of the microRNAs and their potential impact on neuronal survival or death. Armed with this knowledge, one could potentially develop novel therapeutic strategies to target/modulate specific microRNAs. microRNAs are particularly interesting targets for pharmaceutical intervention in that a single microRNA can influence networks of neuronal and/or nonneuronal genes (Fasanaro *et al*, 2010). One interesting application is the use of microRNAs as biomarkers for the diagnosis of neurological diseases. As an example, recent studies have identified specific microRNAs that are upregulated in the blood within hours after a heart attack and are currently investigating the use of these microRNAs as potential biomarkers for myocardial infarction (Salic and De Windt, 2012). microRNAs are widely used in several types of cancer, with one study identifying microRNAs that correlate with the clinical outcome of breast cancer patients (Wu *et al*, 2012). Currently, microRNAs are undergoing evaluation for possible clinical use as biomarkers for neurological diseases (de Planell-Saguer and Rodicio, 2011).

The brain expresses 70% of known microRNAs, each of which exhibits distinctive spatio-temporal expression (Cao *et al*, 2006; de Planell-Saguer and Rodicio, 2011). Dysregu-

lation of microRNA expression have been found in animal models of Alzheimer's disease (Hebert *et al*, 2009). If indeed altered levels of microRNAs are found to correlate with the severity of the disease, they could become valuable tools to diagnose this disease early and be able to administer treatment. There are several microRNAs that are altered in response to brain injury after stroke or seizures (Liu *et al*, 2010). In one study of young stroke patients, these profiles correlated with the type of stroke they suffered and the progress of recovery (Tan *et al*, 2009). These studies demonstrate that microRNAs, if specific and readily detectable at the appropriate times, could be powerful tools not only for the diagnosis, but also the treatment of neurological disorders.

The past decade has witnessed the application of drugs that target the epigenetic machinery in a variety of disorders. Although HDAC inhibitors and other drugs that target the epigenetic machinery are FDA-approved and in use clinically to treat patients with late-stage cancer and autoimmune disorders, use of these drugs to treat brain disorders such as ischemia and epilepsy is a new and burgeoning field. The finding that a single, acute injection of the HDAC inhibitor TSA administered to animals after an ischemic episode ameliorates neuronal injury has important clinical implications and suggests that HDAC inhibitors (Noh *et al*, 2012) may be a promising avenue for intervention in the neurodegeneration associated with ischemic stroke. Unsolved problems include the development of drugs that have the ability to penetrate the blood-brain barrier and ameliorate neuronal death and cognitive deficits, with minimal toxicity. This is important because although it might not be possible to prevent the initial stroke or seizure, if treated properly, drugs that target the epigenetic machinery may serve to limit or prevent further neuronal death, thereby limiting the severity of the insult.

Although compelling, these findings raise new questions. Does epigenetic remodeling promote susceptibility to neuronal insults such as seizures and ischemia, and/or do epigenetic modifications occur in response to neuronal insults? Findings from studies of neurological diseases with long-term, progressive neurodegeneration, such as Huntington's disease and Alzheimer's disease, indicate that epigenetic dysregulation may do both. Studies performed to date show that ischemic stroke promotes epigenetic dysregulation, which in turn promotes neuronal death. Recent findings add ischemic stroke and epilepsy to the growing list of disorders that involve dysregulation of chromatin-modifying proteins.

ACKNOWLEDGEMENTS

We would like to acknowledge all of the authors whose valuable work we could not include due to the limited number of citations allowed. This work was supported by NIH grant NS46742 to RSZ, AHA grant 12POST9420019 to KA and a generous grant from the FM Kirby Foundation. RSZ is the FM Kirby Professor in Neural Repair and Protection.

DISCLOSURE

The authors declare no conflict of interest.

REFERENCES

- Abel T, Zukin RS (2008). Epigenetic targets of HDAC inhibition in neurodegenerative and psychiatric disorders. *Curr Opin Pharmacol* **8**: 57–64.
- Abrahams BS, Geschwind DH (2008). Advances in autism genetics: on the threshold of a new neurobiology. *Nat Rev Genet* **9**: 341–355.
- Aka JA, Kim GW, Yang XJ (2011). K-acetylation and its enzymes: overview and new developments. *Handb Exp Pharmacol* **206**: 1–12.
- Amiet C, Gourfinkel-An I, Bouzamondo A, Tordjman S, Baulac M, Lechat P et al (2008). Epilepsy in autism is associated with intellectual disability and gender: evidence from a meta-analysis. *Biol Psychiatry* **64**: 577–582.
- Amir RE, Van dv I, Wan M, Tran CQ, Francke U, Zoghbi HY (1999). Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. *Nat Genet* **23**: 185–188.
- Andres ME, Burger C, Peral-Rubio MJ, Battaglioli E, Anderson ME, Grimes J et al (1999). CoREST: a functional corepressor required for regulation of neural-specific gene expression. *Proc Natl Acad Sci USA* **96**: 9873–9878.
- Aronica E, Fluiter K, Iyer A, Zurolo E, Vreijling J, van Vliet EA et al (2010). Expression pattern of miR-146a, an inflammation-associated microRNA, in experimental and human temporal lobe epilepsy. *Eur J Neurosci* **31**: 1100–1107. **Evidence showing the possible involvement of microRNA in the modulation of astroglial inflammatory response.**
- Ball MP, Li JB, Gao Y, Lee JH, LeProust EM, Park IH et al (2009). Targeted and genome-scale strategies reveal gene-body methylation signatures in human cells. *Nat Biotechnol* **27**: 361–368.
- Ballas N, Battaglioli E, Atouf F, Andres ME, Chenoweth J, Anderson ME et al (2001). Regulation of neuronal traits by a novel transcriptional complex. *Neuron* **31**: 353–365.
- Ballas N, Grunseich C, Lu DD, Spohr JC, Mandel G (2005). REST and its corepressors mediate plasticity of neuronal gene chromatin throughout neurogenesis. *Cell* **121**: 645–657.
- Ballas N, Mandel G (2005). The many faces of REST oversee epigenetic programming of neuronal genes. *Curr Opin Neurobiol* **15**: 500–506.
- Baltan S, Bachleda A, Morrison RS, Murphy SP (2011a). Expression of histone deacetylases in cellular compartments of the mouse brain and the effects of ischemia. *Transl Stroke Res* **2**: 411–423.
- Baltan S, Murphy SP, Danilov CA, Bachleda A, Morrison RS (2011b). Histone deacetylase inhibitors preserve white matter structure and function during ischemia by conserving ATP and reducing excitotoxicity. *J Neurosci* **31**: 3990–3999.
- Bartsch O, Schmidt S, Richter M, Morlot S, Seemanova E, Wiebe G et al (2005). DNA sequencing of CREBBP demonstrates mutations in 56% of patients with Rubinstein-Taybi syndrome (RSTS) and in another patient with incomplete RSTS. *Hum Genet* **117**: 485–493.
- Benhamed M, Herbig U, Ye T, Dejean A, Bischof O (2012). Senescence is an endogenous trigger for microRNA-directed transcriptional gene silencing in human cells. *Nat Cell Biol* **14**: 266–275.
- Berg AT, Berkovic SF, Brodie MJ, Buchhalter J, Cross JH, van Emde BW et al (2010). Revised terminology and concepts for organization of seizures and epilepsies: report of the ILAE commission on classification and terminology, 2005–2009. *Epilepsia* **51**: 676–685.
- Berger SL (2007). The complex language of chromatin regulation during transcription. *Nature* **447**: 407–412.
- Binder DK, Croll SD, Gall CM, Scharfman HE (2001). BDNF and epilepsy: too much of a good thing? *Trends Neurosci* **24**: 47–53.
- Binder DK, Routbort MJ, McNamara JO (1999). Immunohistochemical evidence of seizure-induced activation of trk receptors in the mossy fiber pathway of adult rat hippocampus. *J Neurosci* **19**: 4616–4626.
- Bolton PF, Carcani-Rathwell I, Hutton J, Goode S, Howlin P, Rutter M (2011). Epilepsy in autism: features and correlates. *Br J Psychiatry* **198**: 289–294.
- Borrrelli E, Nestler EJ, Allis CD, Sassone-Corsi P (2008). Decoding the epigenetic language of neuronal plasticity. *Neuron* **60**: 961–974.
- Brandl A, Heinzel T, Kramer OH (2009). Histone deacetylases: salesmen and customers in the post-translational modification market. *Biol Cell* **101**: 193–205.
- Bruce AW (2004). Genome-wide analysis of repressor element 1 silencing transcription factor/neuron-restrictive silencing factor (REST/NRSF) target genes. *Proc Natl Acad Sci USA* **101**: 10458–10463.
- Bruce AW, Lopez-Contreras AJ, Flicek P, Down TA, Dhani P, Dillon SC et al (2009). Functional diversity for REST (NRSF) is defined by *in vivo* binding affinity hierarchies at the DNA sequence level. *Genome Res* **19**: 994–1005.
- Calderone A, Jover T, Noh K.-M., Tanaka H, Yokota H, Lin Y et al (2003). Ischemic insults de-repress the gene silencer rest in neurons destined to die. *J Neurosci* **23**: 2112–2121.
- Cao X, Yeo G, Muotri AR, Kuwabara T, Gage FH (2006). Noncoding RNAs in the mammalian central nervous system. *Annu Rev Neurosci* **29**: 77–103.
- Carey N, La Thangue NB (2006). Histone deacetylase inhibitors: gathering pace. *Curr Opin Pharmacol* **6**: 369–375.
- Catterall WA, Dib-Hajj S, Meisler MH, Pietrobon D (2008). Inherited neuronal ion channelopathies: new windows on complex neurological diseases. *J Neurosci* **28**: 11768–11777.
- Catterall WA, Kalume F, Oakley JC (2010). NaV1.1 channels and epilepsy. *J Physiol* **588**: 1849–1859.
- Clarke DF, Roberts W, Daraksan M, Dupuis A, McCabe J, Wood H et al (2005). The prevalence of autistic spectrum disorder in children surveyed in a tertiary care epilepsy clinic. *Epilepsia* **46**: 1970–1977.
- Cloos P (2011). The role of histone demethylases in disease. In: Roach HI, Bronner F, Oreffo ROC (eds). *Epigenetic aspects of chronic diseases*. Springer: London, pp 75–93.
- Conaco C, Otto S, Han JJ, Mandel G (2006). Reciprocal actions of REST and a microRNA promote neuronal identity. *Proc Natl Acad Sci USA* **103**: 2422–2427.
- Day JJ, Sweatt JD (2011). Epigenetic mechanisms in cognition. *Neuron* **70**: 813–829.
- Dekker FJ, Haisma HJ (2009). Histone acetyl transferases as emerging drug targets. *Drug Discov Today* **14**: 942–948.
- de Planell-Saguer M, Rodicio MC (2011). Analytical aspects of microRNA in diagnostics: a review. *Anal Chim Acta* **699**: 134–152.
- de Ruijter AJ, van Gennip AH, Caron HN, Kemp S, van Kuilenburg AB (2003). Histone deacetylases (HDACs): characterization of the classical HDAC family. *Biochem J* **370**: 737–749.
- Dong E, Chen Y, Gavin DP, Grayson DR, Guidotti A (2010). Valproate induces DNA demethylation in nuclear extracts from adult mouse brain. *Epigenetics* **5**: 730–735. **Shows toxicity of valproic acid caused by global DNA demethylation.**
- Du L, Bayir H, Lai Y, Zhang X, Kochanek PM, Watkins SC et al (2004). Innate gender-based proclivity in response to cytotoxicity and programmed cell death pathway. *J Biol Chem* **279**: 38563–38570.
- Duncan JS, Sander JW, Sisodiya SM, Walker MC (2006). Adult epilepsy. **367**: 1087–1100.
- Edbauer D, Neilson JR, Foster KA, Wang CF, Seeburg DP, Battersby MN et al (2010). Regulation of synaptic structure and function by FMRP-associated microRNAs miR-125b and miR-132. *Neuron* **65**: 373–384.
- Eleuteri S, Monti B, Brignani S, Contestabile A (2009). Chronic dietary administration of valproic acid protects neurons of the rat nucleus basalis magnocellularis from ibotenic acid neurotoxicity. *Neurotox Res* **15**: 127–132.
- Endres M, Fan G, Meisel A, Dimagl U, Jaenisch R (2001). Effects of cerebral ischemia in mice lacking DNA methyltransferase 1 in post-mitotic neurons. *Neuroreport* **12**: 3763–3766.
- Endres M, Meisel A, Biniszkiwicz D, Namura S, Prass K, Ruscher K et al (2000). DNA methyltransferase contributes to delayed ischemic brain injury. *J Neurosci* **20**: 3175–3181. **Evidence that increased DNA methylation is associated with cerebral ischemia; DNMT deletion mice and DNMT inhibitor shows neuronal protection against stroke.**
- Faraco G, Pancani T, Formentini L, Mascagni P, Fossati G, Leoni F et al (2006). Pharmacological inhibition of histone deacetylases by suberoylanilide hydroxamic acid specifically alters gene expression and reduces ischemic injury in the mouse brain. *Mol Pharmacol* **70**: 1876–1884. **Shows rescue of neuron against ischemia-induced cell death by HDAC inhibitor, SAHA.**
- Fasanaro P, Greco S, Ivan M, Capogrossi MC, Martelli F (2010). microRNA: emerging therapeutic targets in acute ischemic diseases. *Pharmacol Ther* **125**: 92–104.
- Feng J, Nestler EJ (2010). MeCP2 and drug addiction. *Nat Neurosci* **13**: 1039–1041.
- Fisher RS, van Emde Boas W, Blume W, Elger C, Genton P, Lee P et al (2005). Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* **46**: 470–472.
- Formisano L, Noh KM, Miyawaki T, Mashiko T, Bennett MV, Zukin RS (2007). Ischemic insults promote epigenetic reprogramming of mu opioid receptor expression in hippocampal neurons. *Proc Natl Acad Sci USA* **104**: 4170–4175.
- Galanopoulou AS, Buckmaster PS, Staley KJ, Moshe SL, Perucca E, Engel Jr J et al (2012). Identification of new epilepsy treatments: issues in preclinical methodology. *Epilepsia* **53**: 571–582.
- Garriga-Canut M, Schoenike B, Qazi R, Bergendahl K, Daley TJ, Pfender RM et al (2006). 2-Deoxy-D-glucose reduces epilepsy progression by NRSF-CtBP-dependent metabolic regulation of chromatin structure. *Nat Neurosci* **9**: 1382–1387.

- Gibson CL, Murphy SP (2010). Benefits of histone deacetylase inhibitors for acute brain injury: a systematic review of animal studies. *J Neurochem* **115**: 806–813.
- Gillies S, Haddley K, Vasiliou S, Bubbs VJ, Quinn JP (2009). The human neurokinin B gene, TAC3, and its promoter are regulated by Neuron Restrictive Silencing Factor (NRSF) transcription factor family. *Neuropeptides* **43**: 333–340.
- Gorter JA, Petrozino JJ, Aronica EM, Rosenbaum DM, Opitz T, Bennett MV et al (1997). Global ischemia induces downregulation of GluR2 mRNA and increases AMPA receptor-mediated Ca²⁺ influx in hippocampal CA1 neurons of gerbil. *J Neurosci* **17**: 6179–6188.
- Gottlicher M, Minucci S, Zhu P, Kramer OH, Schimpf A, Giavera S et al (2001). Valproic acid defines a novel class of HDAC inhibitors inducing differentiation of transformed cells. *EMBO J* **20**: 6969–6978. **Report discovered valproic acid as a HDAC inhibitor.**
- Graff J, Kim D, Dobbin MM, Tsai LH (2011). Epigenetic regulation of gene expression in physiological and pathological brain processes. *Physiol Rev* **91**: 603–649.
- Graff J, Rei D, Guan JS, Wang WY, Seo J, Hennig KM et al (2012). An epigenetic blockade of cognitive functions in the neurodegenerating brain. *Nature* **483**: 222–226.
- Gray SG, Ekstrom TJ (2001). The human histone deacetylase family. *Exp Cell Res* **262**: 75–83.
- Grimes JA, Nielsen SJ, Battaglioli E, Miska EA, Speh JC, Berry DL et al (2000). The co-repressor mSin3A is a functional component of the REST-CoREST repressor complex. *J Biol Chem* **275**: 9461–9467.
- Grooms SY, Opitz T, Bennett MV, Zukin RS (2000). Status epilepticus decreases glutamate receptor 2 mRNA and protein expression in hippocampal pyramidal cells before neuronal death. *Proc Natl Acad Sci USA* **97**: 3631–3636.
- Guan JS, Haggarty SJ, Giacometti E, Dannenberg JH, Joseph N, Gao J et al (2009). HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature* **459**: 55–60.
- Guardavaccaro D, Frescas D, Dorrello NV, Peschiaroli A, Multani AS, Cardozo T et al (2008). Control of chromosome stability by the beta-TrCP-REST-Mad2 axis. *Nature* **452**: 365–369.
- Guibert S, Fome T, Weber M (2009). Dynamic regulation of DNA methylation during mammalian development. *Epigenomics* **1**: 81–98.
- Haas CA, Dudeck O, Kirsch M, Huszka C, Kann G, Pollak S et al (2002). Role for reelin in the development of granule cell dispersion in temporal lobe epilepsy. *J Neurosci* **22**: 5797–5802.
- Haggarty SJ, Tsai LH (2011). Probing the role of HDACs and mechanisms of chromatin-mediated neuroplasticity. *Neurobiol Learn Mem* **96**: 41–52.
- Hauser WA, Annegers JF, Kurland LT (1993). Incidence of epilepsy and unprovoked seizures in Rochester, Minnesota: 1935–1984. *Epilepsia* **34**: 453–468.
- Hebert SS, Horre K, Nicolai L, Bergmans B, Papadopoulou AS, Delacourte A et al (2009). MicroRNA regulation of Alzheimer's amyloid precursor protein expression. *Neurobiol Dis* **33**: 422–428.
- Heinrich C, Nitta N, Flubacher A, Muller M, Fahrner A, Kirsch M et al (2006). Reelin deficiency and displacement of mature neurons, but not neurogenesis, underlie the formation of granule cell dispersion in the epileptic hippocampus. *J Neurosci* **26**: 4701–4713.
- Herz J, Chen Y (2006). Reelin, lipoprotein receptors and synaptic plasticity. *Nat Rev Neurosci* **7**: 850–859.
- Hoehn B, Ringer TM, Xu L, Giffard RG, Sapolsky RM, Steinberg GK et al (2001). Overexpression of HSP72 after induction of experimental stroke protects neurons from ischemic damage. *J Cereb Blood Flow Metab* **21**: 1303–1309.
- Hu K, Zhang C, Long L, Long X, Feng L, Li Y et al (2011). Expression profile of microRNAs in rat hippocampus following lithium-pilocarpine-induced status epilepticus. *Neurosci Lett* **488**: 252–257. **One of three studies of microRNA array after status epilepticus.**
- Huang Y, Doherty JJ, Dingleline R (2002). Altered histone acetylation at glutamate receptor 2 and brain-derived neurotrophic factor genes is an early event triggered by status epilepticus. *J Neurosci* **22**: 8422–8428. **Report showing GluA2 is downregulated by induction of histone deacetylation in the GluA2 promoter and rescued by HDAC inhibitor after status epilepticus.**
- Huang Y, Myers SJ, Dingleline R (1999). Transcriptional repression by REST: recruitment of Sin3A and histone deacetylase to neuronal genes. *Nat Neurosci* **2**: 867–872. **Recent study showing HDAC2 is associated with pathology of epilepsy.**
- Huang Y, Zhao F, Wang L, Yin H, Zhou C, Wang X (2012). Increased expression of histone deacetylases 2 in temporal lobe epilepsy: a study of epileptic patients and rat models. *Synapse* **66**: 151–159.
- Huang Z, Walker MC, Shah MM (2009). Loss of dendritic HCN1 subunits enhances cortical excitability and epileptogenesis. *J Neurosci* **29**: 10979–10988.
- Isaac JT, Ashby M, McBain CJ (2007). The role of the GluR2 subunit in AMPA receptor function and synaptic plasticity. *Neuron* **54**: 859–871.
- Jenuwein T, Allis CD (2001). Translating the histone code. *Science* **293**: 1074–1080.
- Jian L, Nagarajan L, de KN, Ravine D, Bower C, Anderson A et al (2006). Predictors of seizure onset in Rett syndrome. *J Pediatr* **149**: 542–547.
- Jiang H, Poirier MA, Liang Y, Pei Z, Weiskittel CE, Smith WW et al (2006). Depletion of CBP is directly linked with cellular toxicity caused by mutant huntingtin. *Neurobiol Dis* **23**: 543–551.
- Jimenez-Mateos EM, Bray I, Sanz-Rodriguez A, Engel T, McKiernan RC, Mouri G et al (2011). miRNA expression profile after status epilepticus and hippocampal neuroprotection by targeting miR-132. *Am J Pathol* **179**: 2519–2532.
- Johnson R, Zuccato C, Belyaev ND, Guest DJ, Cattaneo E, Buckley NJ (2008). A microRNA-based gene dysregulation pathway in Huntington's disease. *Neurobiol Dis* **29**: 438–445.
- Jung S, Jones TD, Lugo Jr JN, Sheerin AH, Miller JW, D'Ambrosio R et al (2007). Progressive dendritic HCN channelopathy during epileptogenesis in the rat pilocarpine model of epilepsy. *J Neurosci* **27**: 13012–13021.
- Jung S, Warner LN, Pitsch J, Becker AJ, Poulos NP (2011). Rapid loss of dendritic HCN channel expression in hippocampal pyramidal neurons following status epilepticus. *J Neurosci* **31**: 14291–14295.
- Kasinski AL, Slack FJ (2011). Epigenetics and genetics. MicroRNAs en route to the clinic: progress in validating and targeting microRNAs for cancer therapy. *Nat Rev Cancer* **11**: 849–864.
- Kelly TK, De Carvalho DD, Jones PA (2010). Epigenetic modifications as therapeutic targets. *Nat Biotechnol* **28**: 1069–1078.
- Kieffer BL, Gaveriaux-Ruff C (2002). Exploring the opioid system by gene knockout. *Prog Neurobiol* **66**: 285–306.
- Kim DH, Saetrom P, Snove Jr O, Rossi JJ (2008). MicroRNA-directed transcriptional gene silencing in mammalian cells. *Proc Natl Acad Sci USA* **105**: 16230–16235.
- Kobow K, Blumcke I (2011). The methylation hypothesis: do epigenetic chromatin modifications play a role in epileptogenesis? *Epilepsia* **52**(Suppl 4): 15–19.
- Kobow K, Jeske I, Hildebrandt M, Hauke J, Hahnen E, Buslei R et al (2009). Increased reelin promoter methylation is associated with granule cell dispersion in human temporal lobe epilepsy. *J Neuropathol Exp Neurol* **68**: 356–364. **First direct evidence showing aberrant DNA methylation in pathogenesis of epilepsy.**
- Kokaia M, Ernfors P, Kokaia Z, Elmer E, Jaenisch R, Lindvall O (1995). Suppressed epileptogenesis in BDNF mutant mice. *Exp Neurol* **133**: 215–224.
- Kouzarides T (2000). Acetylation: a regulatory modification to rival phosphorylation? *EMBO J* **19**: 1176–1179.
- Kouzarides T (2007). Chromatin modifications and their function. *Cell* **128**: 693–705.
- Kudo T, Kanemoto S, Hara H, Morimoto N, Morihara T, Kimura R et al (2008). A molecular chaperone inducer protects neurons from ER stress. *Cell Death Differ* **15**: 364–375.
- Kwak S, Weiss JH (2006). Calcium-permeable AMPA channels in neurodegenerative disease and ischemia. *Curr Opin Neurobiol* **16**: 281–287.
- Laurent L, Wong E, Li G, Huynh T, Tsigiris A, Ong CT et al (2010). Dynamic changes in the human methylome during differentiation. *Genome Res* **20**: 320–331.
- Lee HA, Hong SH, Kim JW, Jang IS (2010). Possible involvement of DNA methylation in NKCC1 gene expression during postnatal development and in response to ischemia. *J Neurochem* **114**: 520–529. **Shows DNA methylation changes in the NKCC gene promoter during development and in ischemic injury which is responsible for NKCC1 gene expression.**
- Lee MG, Wynder C, Cooch N, Shiekhattar R (2005). An essential role for CoREST in nucleosomal histone 3 lysine 4 demethylation. *Nature* **437**: 432–435.
- Levenson JM, Sweatt JD (2005). Epigenetic mechanisms in memory formation. *Nat Rev Neurosci* **6**: 108–118.
- Lewis AS, Chetkovich DM (2011). HCN channels in behavior and neurological disease: too hyper or not active enough? *Mol Cell Neurosci* **46**: 357–367.
- Liou AK, Clark RS, Henshall DC, Yin XM, Chen J (2003). To die or not to die for neurons in ischemia, traumatic brain injury and epilepsy: a review on the stress-activated signaling pathways and apoptotic pathways. *Prog Neurobiol* **69**: 103–142.
- Liu DZ, Tian Y, Ander BP, Xu H, Stamova BS, Zhan X et al (2010). Brain and blood microRNA expression profiling of ischemic stroke, intracerebral hemorrhage, and kainate seizures. *J Cereb Blood Flow Metab* **30**: 92–101. **Evidence of microRNAs as biomarkers for ischemic stroke, intracerebral hemorrhage and seizure.**
- Liu F, Li Z, Li J, Siegel C, Yuan R, McCullough LD (2009). Sex differences in caspase activation after stroke. *Stroke* **40**: 1842–1848.
- Liu S, Lau L, Wei J, Zhu D, Zou S, Sun HS et al (2004). Expression of Ca²⁺-permeable AMPA receptor channels primes cell death in transient forebrain ischemia. *Neuron* **43**: 43–55.
- Liu SJ, Zukin RS (2007). Ca²⁺-permeable AMPA receptors in synaptic plasticity and neuronal death. *Trends Neurosci* **30**: 126–134.
- Liu SQ, Cull-Candy SG (2000). Synaptic activity at calcium-permeable AMPA receptors induces a switch in receptor subtype. *Nature* **405**: 454–458.

- Lo EH, Dalkara T, Moskowitz MA (2003). Mechanisms, challenges and opportunities in stroke. *Nat Rev Neurosci* **4**: 399–415.
- Lunyak VV, Burgess R, Prefontaine GG, Nelson C, Sze SH, Chenoweth J *et al* (2002). Corepressor-dependent silencing of chromosomal regions encoding neuronal genes. *Science* **298**: 1747–1752.
- Magill ST, Cambronne XA, Luikart BW, Lioy DT, Leighton BH, Westbrook GL *et al* (2010). microRNA-132 regulates dendritic growth and arborization of newborn neurons in the adult hippocampus. *Proc Natl Acad Sci USA* **107**: 20382–20387.
- McClelland S, Flynn C, Dube C, Richichi C, Zha Q, Ghestem A *et al* (2011). Neuron-restrictive silencer factor-mediated hyperpolarization-activated cyclic nucleotide gated channelopathy in experimental temporal lobe epilepsy. *Ann Neurol* **70**: 454–464. **Important follow-up study showing REST represses specific target gene (Hcn1) expression epigenetically after status epilepticus.**
- McNamara JO (1999). Emerging insights into the genesis of epilepsy. *Nature* **399**: A15–A22.
- Miller BH, Zeier Z, Xi L, Lanz TA, Deng S, Strathmann J *et al* (2012). MicroRNA-132 dysregulation in schizophrenia has implications for both neurodevelopment and adult brain function. *Proc Natl Acad Sci USA* **109**: 3125–3130.
- Miller-Delaney SF, Das S, Sano T, Jimenez-Mateos EM, Bryan K, Buckley PG *et al* (2012). Differential DNA methylation patterns define status epilepticus and epileptic tolerance. *J Neurosci* **32**: 1577–1588. **First genome-wide DNA methylation study showing altered DNA methylation in promoters of genes after status epilepticus and in epileptic tolerance.**
- Morris MJ, Karra AS, Monteggia LM (2010). Histone deacetylases govern cellular mechanisms underlying behavioral and synaptic plasticity in the developing and adult brain. *Behav Pharmacol* **21**: 409–419.
- Morrison BE, Majdzadeh N, D'Mello SR (2007). Histone deacetylases: focus on the nervous system. *Cell Mol Life Sci* **64**: 2258–2269.
- Moskowitz MA, Lo EH, Iadecola C (2010). The science of stroke: mechanisms in search of treatments. *Neuron* **67**: 181–198.
- Myers SJ, Peters J, Huang Y, Comer MB, Barthel F, Dinglede R (1998). Transcriptional regulation of the GluR2 gene: neural-specific expression, multiple promoters, and regulatory elements. *J Neurosci* **18**: 6723–6739.
- Naruse Y, Aoki T, Kojima T, Mori N (1999). Neural restrictive silencer factor recruits mSin3 and histone deacetylase complex to repress neuron-specific target genes. *Proc Natl Acad Sci USA* **96**: 13691–13696.
- Nectoux J, Bahi-Buisson N, Guellec I, Coste J, De RN, Rosas H *et al* (2008). The p.Val66Met polymorphism in the BDNF gene protects against early seizures in Rett syndrome. *Neurology* **70**: 2145–2151.
- Nissenkorn A, Gak E, Vecsler M, Reznik H, Menascu S, Ben ZB (2010). Epilepsy in Rett syndrome—the experience of a National Rett Center. *Epilepsia* **51**: 1252–1258.
- Noh KM, Hwang JY, Follenzi A, Athanasiadou R, Miyawaki T, Grealley JM *et al* (2012). Repressor element-1 silencing transcription factor (REST)-dependent epigenetic remodeling is critical to ischemia-induced neuronal death. *Proc Natl Acad Sci USA* **109**: E962–E971. **Shows REST binds a subset of transcriptionally responsive genes and recruits corepressors which together orchestrate epigenetic remodeling and gene silencing.**
- Ofengeim D, Miyawaki T, Zukin RS (2011). Molecular and cellular mechanisms of ischemia-induced neuronal death. In: Mohr JP, Wolf P, Grotta JC, Moskowitz MA, Mayberg M, von Kummer R (eds) *Stroke: Pathophysiology, Diagnosis and Management*. Churchill Livingstone Elsevier: Philadelphia. pp 75–109.
- Oida Y, Shimazawa M, Imaizumi K, Hara H (2008). Involvement of endoplasmic reticulum stress in the neuronal death induced by transient forebrain ischemia in gerbil. *Neuroscience* **151**: 111–119.
- Okano M, Bell DW, Haber DA, Li E (1999). DNA methyltransferases Dnmt3a and Dnmt3b are essential for *de novo* methylation and mammalian development. *Cell* **99**: 247–257.
- Ooi L, Wood IC (2007). Chromatin crosstalk in development and disease: lessons from REST. *Nat Rev Genet* **8**: 544–554.
- Opitz T, Grooms SY, Bennett MV, Zukin RS (2000). Remodeling of alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid receptor subunit composition in hippocampal neurons after global ischemia. *Proc Natl Acad Sci USA* **97**: 13360–13365.
- Ouyang YB, Lu Y, Yue S, Xu LJ, Xiong XX, White RE *et al* (2012). miR-181 regulates GRP78 and influences outcome from cerebral ischemia *in vitro* and *in vivo*. *Neurobiol Dis* **45**: 555–563. **Evidence showing miR-181 as a possible repressor of HSP70 family which contributes to neuronal death and rescue this phenomenon by antagomir or inhibitor of miR-181a.**
- Packer AN, Xing Y, Harper SQ, Jones L, Davidson BL (2008). The bifunctional microRNA miR-9/miR-9* regulates REST and CoREST and is downregulated in Huntington's disease. *J Neurosci* **28**: 14341–14346.
- Palm K, Belluardo N, Metsis M, Timmus T (1998). Neuronal expression of zinc finger transcription factor REST/NRSF/XBR gene. *J Neurosci* **18**: 1280–1296.
- Pellegrini-Giampietro DE, Zukin RS, Bennett MV, Cho S, Pulsinelli WA (1992). Switch in glutamate receptor subunit gene expression in CA1 subfield of hippocampus following global ischemia in rats. *Proc Natl Acad Sci USA* **89**: 10499–10503.
- Petrij F, Giles RH, Dauwerse HG, Saris JJ, Hennekam RC, Masuno M *et al* (1995). Rubinstein-Taybi syndrome caused by mutations in the transcriptional co-activator CBP. *Nature* **376**: 348–351.
- Plummer R, Vidal L, Griffin M, Lesley M, de BJ, Coulthard S *et al* (2009). Phase I study of MG98, an oligonucleotide antisense inhibitor of human DNA methyltransferase 1, given as a 7-day infusion in patients with advanced solid tumors. *Clin Cancer Res* **15**: 3177–3183.
- Pond BB, Berglund K, Kuner T, Feng G, Augustine GJ, Schwartz-Bloom RD (2006). The chloride transporter Na(+)-K(+)-Cl- cotransporter isoform-1 contributes to intracellular chloride increases after *in vitro* ischemia. *J Neurosci* **26**: 1396–1406.
- Qureshi IA, Mehler MF (2009). Regulation of non-coding RNA networks in the nervous system—what's the REST of the story? *Neurosci Lett* **466**: 73–80.
- Qureshi IA, Mehler MF (2010a). Emerging role of epigenetics in stroke: part 1: DNA methylation and chromatin modifications. *Arch Neurol* **67**: 1316–1322.
- Qureshi IA, Mehler MF (2010b). Epigenetic mechanisms underlying human epileptic disorders and the process of epileptogenesis. *Neurobiol Dis* **39**: 53–60.
- Rajdev S, Hara K, Kokubo Y, Mestril R, Dillmann W, Weinstein PR *et al* (2000). Mice overexpressing rat heat shock protein 70 are protected against cerebral infarction. *Ann Neurol* **47**: 782–791.
- Rakhade SN, Jensen FE (2009). Epileptogenesis in the immature brain: emerging mechanisms. *Nat Rev Neurol* **5**: 380–391.
- Rink C, Khanna S (2011). MicroRNA in ischemic stroke etiology and pathology. *Physiol Genomics* **43**: 521–528.
- Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtinen H, Lamsa K *et al* (1999). The K+/Cl- co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* **397**: 251–255.
- Robertson KD (2005). DNA methylation and human disease. *Nat Rev Genet* **6**: 597–610.
- Robison AJ, Nestler EJ (2011). Transcriptional and epigenetic mechanisms of addiction. *Nat Rev Neurosci* **12**: 623–637.
- Roelfsema JH, White SJ, Ariyurek Y, Bartholdi D, Niedrist D, Papadia F *et al* (2005). Genetic heterogeneity in Rubinstein-Taybi syndrome: mutations in both the CBP and EP300 genes cause disease. *Am J Hum Genet* **76**: 572–580.
- Roopra A, Huang Y, Dinglede R (2001). Neurological disease: listening to gene silencers. *Mol Interv* **1**: 219–228.
- Roopra A, Qazi R, Schoenike B, Daley TJ, Morrison JF (2004). Localized domains of G9a-mediated histone methylation are required for silencing of neuronal genes. *Mol Cell* **14**: 727–738.
- Russ SA, Larson K, Halfon N (2012). A national profile of childhood epilepsy and seizure disorder. *Pediatrics* **129**: 256–264.
- Ruthenburg AJ, Li H, Patel DJ, Allis CD (2007). Multivalent engagement of chromatin modifications by linked binding modules. *Nat Rev Mol Cell Biol* **8**: 983–994.
- Salic K, De Windt LJ (2012). MicroRNAs as biomarkers for myocardial infarction. *Curr Atheroscler Rep* **14**: 193–200.
- Salta E, De SB (2012). Non-coding RNAs with essential roles in neurodegenerative disorders. *Lancet Neurol* **11**: 189–200.
- Sanchez RM, Koh S, Rio C, Wang C, Lamperti ED, Sharma D *et al* (2001). Decreased glutamate receptor 2 expression and enhanced epileptogenesis in immature rat hippocampus after perinatal hypoxia-induced seizures. *J Neurosci* **21**: 8154–8163.
- Sato F, Tsuchiya S, Meltzer SJ, Shimizu K (2011). MicroRNAs and epigenetics. *FEBS J* **278**: 1598–1609.
- Scharf AN, Imhof A (2011). Every methyl counts—epigenetic calculus. *FEBS Lett* **585**: 2001–2007.
- Schratt G (2009a). Fine-tuning neural gene expression with microRNAs. *Curr Opin Neurobiol* **19**: 213–219.
- Schratt G (2009b). microRNAs at the synapse. *Nat Rev Neurosci* **10**: 842–849.
- Selvi BR, Cassel JC, Kundu TK, Boutillier AL (2010). Tuning acetylation levels with HAT activators: therapeutic strategy in neurodegenerative diseases. *Biochim Biophys Acta* **1799**: 840–853.
- Senner CE (2011). The role of DNA methylation in mammalian development. *Reprod Biomed Online* **22**: 529–535.
- Shah MM, Anderson AE, Leung V, Lin X, Johnston D (2004). Seizure-induced plasticity of h channels in entorhinal cortical layer III pyramidal neurons. *Neuron* **44**: 495–508.

- Shaked I, Meerson A, Wolf Y, Avni R, Greenberg D, Gilboa-Geffen A *et al* (2009). MicroRNA-132 potentiates cholinergic anti-inflammatory signaling by targeting acetylcholinesterase. *Immunity* **31**: 965–973.
- Shi X, Yasumoto S, Kurahashi H, Nakagawa E, Fukasawa T, Uchiya S *et al* (2011). Clinical spectrum of SCN2A mutations. *Brain Dev* **34**: 541–545.
- Shi YJ (2005). Regulation of LSD1 histone demethylase activity by its associated factors. *Mol Cell* **19**: 1–8.
- Siegel C, Li J, Liu F, Benashski SE, McCullough LD (2011a). miR-23a regulation of X-linked inhibitor of apoptosis (XIAP) contributes to sex differences in the response to cerebral ischemia. *Proc Natl Acad Sci USA* **108**: 11662–11667. **Suggests miR-23a as a regulator of XIAP sex differences in response to stroke.**
- Siegel G, Saba R, Schratt G (2011b). microRNAs in neurons: manifold regulatory roles at the synapse. *Curr Opin Genet Dev* **21**: 491–497.
- Siegel C, Turtzo C, McCullough LD (2010). Sex differences in cerebral ischemia: possible molecular mechanisms. *J Neurosci Res* **88**: 2765–2774.
- Singh SK, Kagalwala MN, Parker-Thornburg J, Adams H, Majumder S (2008). REST maintains self-renewal and pluripotency of embryonic stem cells. *Nature* **453**: 223–227.
- Sng JC, Taniura H, Yoneda Y (2006). Histone modifications in kainate-induced status epilepticus. *Eur J Neurosci* **23**: 1269–1282. **Shows histone H4 hyperacetylation and histone H3 phosphorylation in c-fos gene promoter and histone H3 phosphorylation in c-jun gene promoter after status epilepticus.**
- Song J, Teplova M, Ishibe-Murakami S, Patel DJ (2012). Structure-based mechanistic insights into DNMT1-mediated maintenance DNA methylation. *Science* **335**: 709–712. **One of three studies of microRNA array after status epilepticus.**
- Song YJ, Tian XB, Zhang S, Zhang YX, Li X, Li D *et al* (2011). Temporal lobe epilepsy induces differential expression of hippocampal miRNAs including let-7e and miR-23a/b. *Brain Res* **1387**: 134–140.
- Soriano FX, Baxter P, Murray LM, Sporn MB, Gillingwater TH, Hardingham GE (2009). Transcriptional regulation of the AP-1 and Nrf2 target gene sulfiredoxin. *Mol Cells* **27**: 279–282.
- Su X, Gopalakrishnan V, Stearns D, Aldape K, Lang FF, Fuller G *et al* (2006). Abnormal expression of REST/NRSF and Myc in neural stem/progenitor cells causes cerebellar tumors by blocking neuronal differentiation. *Mol Cell Biol* **26**: 1666–1678.
- Tahiliani M, Mei P, Fang R, Leonor T, Rutenberg M, Shimizu F *et al* (2007). The histone H3K4 demethylase SMCX links REST target genes to X-linked mental retardation. *Nature* **447**: 601–605.
- Tan KS, Armugam A, Sepramaniam S, Lim KY, Setyowati KD, Wang CW *et al* (2009). Expression profile of microRNAs in young stroke patients. *PLoS One* **4**: e7689.
- Tsankova NM, Kumar A, Nestler EJ (2004). Histone modifications at gene promoter regions in rat hippocampus after acute and chronic electroconvulsive seizures. *J Neurosci* **24**: 5603–5610.
- Tuchman R, Cuccaro M, Alessandri M (2010). Autism and epilepsy: historical perspective. *Brain Dev* **32**: 709–718.
- Urdinguio RG, Sanchez-Mut JV, Esteller M (2009). Epigenetic mechanisms in neurological diseases: genes, syndromes, and therapies. *Lancet Neurol* **8**: 1056–1072.
- Vezzani A, French J, Bartfai T, Baram TZ (2011). The role of inflammation in epilepsy. *Nat Rev Neurol* **7**: 31–40.
- Wang B, Zhu X, Kim Y, Li J, Huang S, Saleem S *et al* (2012). Histone deacetylase inhibition activates transcription factor Nrf2 and protects against cerebral ischemic damage. *Free Radic Biol Med* **52**: 928–936. **Recent report showing HDAC inhibitor, TSA contribute to enhancement of neuronal cell viability in ischemic insult.**
- Wang DO, Martin KC, Zukin RS (2010). Spatially restricting gene expression by local translation at synapses. *Trends Neurosci* **33**: 173–182.
- Wang Z, Leng Y, Tsai LK, Leeds P, Chuang DM (2011). Valproic acid attenuates blood-brain barrier disruption in a rat model of transient focal cerebral ischemia: the roles of HDAC and MMP-9 inhibition. *J Cereb Blood Flow Metab* **31**: 52–57.
- Westbrook TF, Hu G, Ang XL, Mulligan P, Pavlova NN, Liang A *et al* (2008). SCFbeta-TRCP controls oncogenic transformation and neural differentiation through REST degradation. *Nature* **452**: 370–374.
- Wood MA, Hawk JD, Abel T (2006). Combinatorial chromatin modifications and memory storage: a code for memory? *Learn Mem* **13**: 241–244.
- Wu X, Somlo G, Yu Y, Palomares MR, Xuejun LA, Zhou W *et al* (2012). *De novo* sequencing of circulating miRNAs identifies novel markers predicting clinical outcome of locally advanced breast cancer. *J Transl Med* **10**: 42.
- Xi ZQ, Sun JJ, Wang XF, Li MW, Liu XZ, Wang LY *et al* (2007). HSPBAP1 is found extensively in the anterior temporal neocortex of patients with intractable epilepsy. *Synapse* **61**: 741–747.
- Xuan A, Long D, Li J, Ji W, Hong L, Zhang M *et al* (2012). Neuroprotective effects of valproic acid following transient global ischemia in rats. *Life Sci* **90**: 463–468.
- Yashiro K, Riday TT, Condon KH, Roberts AC, Bernardo DR, Prakash R *et al* (2009). Ube3a is required for experience-dependent maturation of the neocortex. *Nat Neurosci* **12**: 777–783.
- Yu HB, Johnson R, Kunarso G, Stanton LW (2011). Coassembly of REST and its cofactors at sites of gene repression in embryonic stem cells. *Genome Res* **21**: 1284–1293.
- Yuan M, Siegel C, Zeng Z, Li J, Liu F, McCullough LD (2009). Sex differences in the response to activation of the poly (ADP-ribose) polymerase pathway after experimental stroke. *Exp Neurol* **217**: 210–218.
- Zhang YZ, Zhang QH, Ye H, Zhang Y, Luo YM, Ji XM *et al* (2010). Distribution of lysine-specific demethylase 1 in the brain of rat and its response in transient global cerebral ischemia. *Neurosci Res* **68**: 66–72.
- Zuccato C, Belyaev N, Conforti P, Ooi L, Tartari M, Papadimou E *et al* (2007). Widespread disruption of repressor element-1 silencing transcription factor/neuron-restrictive silencer factor occupancy at its target genes in Huntington's disease. *J Neurosci* **27**: 6972–6983.
- Zuccato C, Tartari M, Crotti A, Goffredo D, Valenza M, Conti L *et al* (2003). Huntingtin interacts with REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes. *Nat Genet* **35**: 76–83.
- Zukin RS (2010). Eradicating the mediators of neuronal death with a fine-tooth comb. *Sci Signal* **3**: e20.
- Zukin RS, Jover T, Yokota H, Calderone A, Simionescu M, Lau C.G. (2004). Molecular and cellular mechanisms of ischemia-induced neuronal death. In: Mohr JP, Choi DW, Grotta JC, Weir B, Wolf PA (eds) *Stroke: Pathophysiology, Diagnosis, and Management*. Churchill Livingstone: Philadelphia. pp 829–854.