

# 1The Modified Pharaoh Approach: Stingless bees mummify beetle

## 2parasites alive

3

4Mark K Greco <sup>1,6</sup>, Dorothee Hoffmann <sup>4</sup>, Anne Dollin <sup>5</sup>, Michael Duncan <sup>6</sup>, Robert Spooner-

5Hart <sup>6</sup>, Peter Neumann <sup>1,2,3</sup>

6

7<sup>1</sup> Swiss Bee Research Centre, Agroscope Liebefeld-Posieux Research Station ALP, CH-

83033 Bern, Switzerland

9<sup>2</sup> Eastern Bee Research Institute of Yunnan Agricultural University, Heilongtan, Kunming,

10Yunnan Province, China

11<sup>3</sup> Department of Zoology and Entomology, Rhodes University, 61440 Grahamstown,

12Republic of South Africa

13<sup>4</sup> Department of Zoology, Martin-Luther-Universität Halle-Wittenberg, Hoher Weg 4, 06099

14Halle (Saale), Germany

15<sup>5</sup> Australian Native Bee Research Centre, North Richmond, NSW, Australia

16<sup>6</sup> Centre for Plant and Food Science, School of Natural Sciences, University of Western

17Sydney, Richmond, NSW 1797, Australia

18

19**Social insect colonies usually live in nests, which are often invaded by parasitic species<sup>1</sup>.**

20**Workers from these colonies use different defence strategies to combat invaders<sup>1</sup>.**

21 Nevertheless, some parasitic species are able to bypass primary colony defences due to their  
22 morphology and behaviour<sup>1-3</sup>. In particular, some beetle nest invaders cannot be killed or  
23 removed by workers of social bees<sup>2-5</sup>, thus creating the need for alternative social defence  
24 strategies to ensure colony survival. Here we show, using Diagnostic Radioentomology<sup>6</sup>, that  
25 stingless bee workers *Trigona carbonaria*, immediately mummify invading destructive nest  
26 parasites *Aethina tumida* alive, with a mixture of resin, wax and mud, thereby preventing  
27 severe damage to the colony. In sharp contrast to the responses of honeybee<sup>7</sup> and bumblebee  
28 colonies<sup>8</sup>, the rapid live mummification strategy of *T. carbonaria* effectively prevents beetle  
29 parasite advancements and removes their ability to reproduce. The convergent evolution of  
30 live mummification by stingless bees and social encapsulation by honeybees<sup>3</sup> suggests that  
31 colonies of social bees generally rely on, secondary defence mechanisms when harmful nest  
32 intruders cannot be killed or ejected easily. This process is analogous to immune responses  
33 within organisms.

34 Social insects live in colonies and usually construct nests which are often attractive to  
35 parasites. Some parasites feed on stored food or brood and can destroy colonies<sup>3</sup> thus generating the  
36 need for efficient defence mechanisms. While some Coleopteran nest intruders are harmless<sup>8-12</sup>,  
37 others can be damaging parasites<sup>4</sup>. Parasitising beetle species pose particular difficulties for their  
38 social insect hosts because their hard exoskeletons protect them from direct primary defence  
39 strategies such as biting or stinging. The small hive beetle, *Aethina tumida* (Coleoptera:  
40 Nitidulidae), is a parasite and scavenger of honeybee (*Apis mellifera*) colonies endemic to sub-

41 Saharan Africa<sup>2,5,7,13</sup>. It has become an invasive species<sup>14</sup> with well established populations in North  
42 America and Australia<sup>13,15</sup>. It lives within *A. mellifera* nests and feeds on brood, stored food and  
43 dead bees<sup>5,7,16,17</sup>. Frequently, the feeding small hive beetle larvae cause the complete destruction of  
44 the nest<sup>5,7</sup> however, the presence of adult small hive beetles alone can be detrimental to colonies of  
45 European honeybees<sup>18</sup>. This obviously creates demand for efficient defence mechanisms against  
46 intrusion and reproduction by adult small hive beetles.

47        Unlike other parasites, small hive beetles are easily detected and can be vigorously attacked  
48 by honeybee workers<sup>19</sup>. Nevertheless, adult small hive beetles can bypass primary defences of the  
49 bees and easily intrude weak or strong host colonies<sup>5,7</sup> because it is difficult for honeybees to kill or  
50 eject them<sup>3,5</sup> due to the beetles' hard exoskeletons and defensive behaviours, such as the turtle  
51 defence posture or by dropping from combs<sup>3,7</sup>. Cape honeybees, *A. m. capensis*, display secondary  
52 defence mechanisms by encapsulating small hive beetles in tombs made from tree resin (propolis),  
53 which the bees collect for use as a nest cavity sealant<sup>3</sup>. Despite the lack of co-evolution between  
54 host and parasite, European honeybees also encapsulate small hive beetles in propolis tombs<sup>20</sup>  
55 suggesting that encapsulation appears to be part of the general secondary defence of honeybee  
56 colonies.

57        Recent evidence suggests that small hive beetles also parasitise colonies of other social bees.  
58 In fact, small hive beetles have been found naturally infesting commercial bumblebee colonies,  
59 *Bombus impatiens*, in the field<sup>21</sup> and in greenhouses<sup>8</sup> in North America. Natural small hive beetle  
60 infestations were reported in colonies of stingless bees, *Dactylurina staudingerii*, in West Africa<sup>22</sup>  
61 and small hive beetle larvae were also observed in a *T. carbonaria* colony that had recently died

62(Anne Dollin, personal observations) in Australia. Odour cues from stored nest products could  
63attract host-searching adult small hive beetles. We therefore expect colonies of stingless bees to be  
64attractive to small hive beetles and, possibly, suitable for their reproduction. Analogous to  
65honeybees, stingless bees use batumen (a mixture of wax, plant resins and mud) to seal nest  
66cavities<sup>23</sup>, thus similar to honeybees, stingless bees may also show alternative secondary defence  
67mechanisms against harmful nest intruders. Here, we evaluated the defence behaviour of an  
68Australian species of stingless bee, *T. carbonaria*, against hive-intruding small hive beetles.

69 Laboratory reared<sup>24</sup> adult small hive beetles, with BaSO<sub>4</sub>-marked elytra, were introduced to  
70the entrances of five *T. carbonaria* hives (N=10 each hive) via a transparent plastic tube<sup>3,8</sup>. All hives  
71were CT scanned in a human body scanner (General Electric HiSpeed 64 Slice, General Electric  
72Company) at 5 min intervals for 90 min<sup>25</sup>. To assess small hive beetle distribution within the hives,  
73we used BeeView 3D rendering software (Disect Systems Ltd; Suffolk, UK). Two dimensional  
74images were performed to enable precise measurement of small hive beetle positions and 3D  
75images were performed to provide spatial representation of small hive beetles with respect to hive  
76structures. One hive was randomly selected after scanning and snap frozen with LN<sub>2</sub> for visual  
77screening to compare positions of small hive beetles with respect to scanned images.

78 Upon introduction of small hive beetles, bees from all *T. carbonaria* hives immediately  
79coated beetles with batumen. The vigorous attacks by workers (Fig. 1) caused the beetles to remain  
80motionless, with their heads tucked underneath the pronotum and legs and antennae pressed tightly  
81to the body (= turtle defence posture<sup>3</sup>). When not attacked, beetles progressed further into the hive.  
82However, most *T. carbonaria* bees continuously attacked the small hive beetles, thereby keeping

83them in the turtle defence posture. While six small hive beetles did not manage to progress into the  
84hives and were mummified on the spot, others were able to progress further. In one hive, two small  
85hive beetles reached a distance of 170 mm from the hive entrance, just beneath the brood (Fig. 2A).  
86All forward advancements by beetles ceased within 10 min of their introduction into the hive (Fig.  
872B). The dissection of one hive confirmed the positions of small hive beetles (N = 10) in relation to  
88its scanned images.

89       When colonies of social bees are invaded by nest parasites which are difficult to kill or eject,  
90the host colony faces a dilemma. Successful parasite reproduction must be prevented but direct  
91physical attacks alone are not always sufficient to kill defensive opponents like adult small hive  
92beetles<sup>3</sup>. The encapsulation process of adult small hive beetles in honeybee colonies combines  
93prison construction and guarding which usually lasts 1-4 days<sup>3</sup>. Beetles mimic worker bee begging  
94behaviour and are fed by worker bees<sup>27</sup>, thus allowing enough time for beetle mating to occur<sup>27</sup>. Our  
95data clearly show that the stingless bees, *T. carbonaria*, use live mummification of parasitic small  
96hive beetles, the “Alternative Pharaoh Approach”, as an effective and fast secondary defence  
97mechanism to prevent successful parasite reproduction. While social encapsulation of small  
98intruders in wax or propolis confinements has been described from *Bombus* and *Apis*<sup>28</sup>, to our  
99knowledge, this is the first report of live mummification of nest intruders in colonies of social bees.  
100Our experiment shows that live beetle mummification by *T. carbonaria* takes as little as 10 min Fig.  
1012B, suggesting that this behaviour can be more effective than that of honeybees. When small hive  
102beetles adopt the turtle defence posture most of the honeybee guards leave the beetles, which then  
103scurry into hiding<sup>3,19</sup>. In contrast, most *T. carbonaria* bees continuously attack the small hive

104beetles, thereby keeping them in the turtle defence posture. This enables other workers to mummify  
105the beetles alive with batumen whilst they remain motionless Fig.3. Therefore, it appears that the  
106combination of continuous attacks and quick recruitment of mummifying bees underlies this  
107efficient secondary colony defence mechanism of *T. carbonaria*. There have however, been reports  
108of heat-stressed *T. carbonaria* colonies being destroyed by small hive beetles in Australia (Mark  
109Greco, personal observations), suggesting that this invasive species may still pose some threat to  
110native pollinators Fig. 4.

111 In conclusion, single bees, are not able to kill or eject beetle parasites alone. Only a team  
112with individuals performing specific tasks (e.g. wrestling or gluing in the case of live  
113mummification) can overcome parasite advancements. Live mummification of small hive beetles by  
114stingless bees has probably evolved as a secondary defence mechanism to prevent successful  
115reproduction of nest parasites. This process is a social analogue to immune responses within  
116organisms. It is clearly effective, because small hive beetles are quickly immobilised and prevented  
117from successful reproduction. This seems especially important in light of the high reproductive  
118potential of small hive beetles<sup>24</sup>. The convergent evolution of live mummification of nest parasites  
119in stingless bees and social encapsulation in honeybees is another striking example of evolution  
120between insect societies and their parasites.

121

1221. Breed MD, Guzman-Novoa E, Hunt GJ (2004) Defensive behaviour of honey bees:

123 Organization, Genetics, and Comparisons with Other Bees. Annual Review of Entomology

124 49:271–98

1252. El-Niweiri MAA, El-Sarrag MS, Neumann P (2008) Filling the Sudan gap: the northernmost  
 126 natural distribution limit of small hive beetles. *Journal of Apicultural Research*: 47(3):184–185
1273. Neumann P, Pirk CWW, Hepburn HR, Solbrig AJ, Ratnieks FLW, Elzen PJ, Baxter JR (2001)  
 128 Social encapsulation of beetle parasites by Cape honeybee colonies (*Apis mellifera capensis*  
 129 Esch.). *Naturwissenschaften* 88: 214-216
1304. Schmid-Hempel, P. 1998 Parasites in social insects. Princeton University Press.
1315. Lundie AE (1940) The small hive beetle *Aethina tumida*, Science Bulletin 220, Department of.  
 132 Agriculture and Forestry, Government Printer, Pretoria.
1336. Greco MK, Spooner-Hart R, Holford P (2005) A new technique for monitoring *Trigona*  
 134 *carbonaria* nest contents, brood and activity using X-ray computerised tomography. *Journal of*  
 135 *Apicultural Research* 44:97-100
1367. Neumann P, Elzen PJ (2004) The biology of the small hive beetle (*Aethina tumida* Murray,  
 137 Coleoptera: Nitidulidae): Gaps in our knowledge of an invasive species. *Apidologie* 35: 229-  
 138 247
1398. Hoffmann D, Pettis JS, Neumann P (2008) Potential host shift of the small hive beetle (*Aethina*  
 140 *tumida*) to bumblebee colonies (*Bombus impatiens*). *Insectes Soc* 55:153–162
1419. Lea AM (1910) Australian and Tasmanian Coleoptera inhabiting or resorting to the nests of  
 142 ants, bees and termites. *Proc R Soc Victoria [NS]* 23: 116-230.
14310. Lea AM (1912) Australian and Tasmanian Coleoptera inhabiting or resorting to the nests of  
 144 ants, bees and termites. *Proc R Soc Victoria [NS]* 25 Suppl: 31-78.

14511. Neumann P, Ritter W (2004) A scientific note on the association of *Cychramus luteus*  
 146 (Coleoptera: Nitidulidae) with honeybee (*Apis mellifera*) colonies. *Apidologie* 35: 665-666
14712. Haddad N, Esser J, Neumann P (2008) Association of *Cryptophagus hexagonalis* (Coleoptera:  
 148 Cryptophagidae) with honey bee colonies (*Apis mellifera*). *Journal of Apicultural Research*  
 149 47:189-190
15013. Neumann P, Ellis JD (2008) The small hive beetle (*Aethina tumida* Murray, Coleoptera:  
 151 Nitidulidae): distribution, biology and control of an invasive species. *Journal of Apicultural*  
 152 *Research* 47:181–183
15314. Elzen PJ, Baxter JR, Westervelt D, Randall C, Delaplane KS, Cutts L, Wilson WT (1999) Field  
 154 control and biology studies of a new pest species, *Aethina tumida* Murray (Coleoptera,  
 155 Nitidulidae) attacking European honey bees in the Western hemisphere, *Apidologie* 30:361–366
15615. Spiewok S, Pettis JS, Duncan M, Spooner-Hart R, Westervelt D, Neumann P (2007) Small hive  
 157 beetle, *Aethina tumida*, populations I: Infestation levels of honeybee colonies, apiaries and  
 158 regions. *Apidologie* 38:595–605
15916. Schmolke MD (1974) A study of *Aethina tumida*: the small hive beetle, Project Report,  
 160 University of Rhodesia
16117. Spiewok S, Neumann P (2006a) Cryptic low-level reproduction of small hive beetles in  
 162 honeybee colonies. *Journal of Apicultural Research* 45:47-48
16318. Ellis J.D., Hepburn H.R., Delaplane K., Neumann P., Elzen P.J. (2003a) The effects of adult  
 164 small hive beetles, *Aethina tumida* (Coleoptera: Nitidulidae), on nests and flight activity of Cape  
 165 and European honey bees (*Apis mellifera*). *Apidologie* 34, 399-408.



16619. Elzen PJ, Baxter JR, Neumann P, Solbrig A, Pirk CWW, Hepburn HR, Westervelt D, Randall C  
167 (2001) Behaviour of African and European subspecies of *Apis mellifera* toward the small hive  
168 beetle, *Aethina tumida*. Journal of Apicultural Research 40:40-41
16920. Ellis JD, Hepburn HR, Ellis AM, Elzen PJ (2003b) Social encapsulation of the small hive beetle  
170 (*Aethina tumida* Murray) by European honeybees (*Apis mellifera* L.). Insectes Sociaux 50:286–  
171 291
17221. Spiewok S, Neumann P (2006b) Infestation of commercial bumblebee (*Bombus impatiens*) field  
173 colonies by small hive beetles (*Aethina tumida*). Ecological Entomology 31:623-628
17422. Mutsaers M (2006) Beekeepers' observations on the small hive beetle (*Aethina tumida*) and  
175 other pests in bee colonies in West and East Africa. In: Proceedings of the 2<sup>nd</sup> EurBee  
176 conference, Prague, Czech Republic. p 44
17723. Michener CD (1961) Observations on the nests and behaviour of *Trigona* in Australia and New  
178 Guinea (Hymenoptera: Apidae). American Museum of Novitates 2026 (08-05-1961): 1-45
17924. Muerrle TM, Neumann P (2004) Mass production of small hive beetles (*Aethina tumida*  
180 Murray, Coleoptera: Nitidulidae). Journal of Apicultural Research 43:144-145
18125. Greco MK, Bell M, Spooner-Hart R, Holford P (2006) X-ray computerized tomography as a  
182 new method for monitoring *Amegilla holmesi* nest structure, nesting behaviour and adult female  
183 activity. Entomologia Experimentalis et Applicata 120:71-76
18426. Breed, M. D. 2003. Nestmate Recognition Assays As a Tool for Population and Ecological  
185 Studies in Eusocial Insects: A Review. J. Kansas Entomol. Soc. 76: 539-550

18627. Ellis JD, Pirk C, Hepburn HR, Kastberger G, Elzen PJ (2002) Small hive beetles survive in  
187 honeybee prisons by behavioural mimicry. *Naturwissenschaften* 89: 326-328

18828. Michener, C.D. (1974) *The social behaviour of the bees; A comparative study*. Harvard  
189 University Press, pp. 404.

190

191**Acknowledgements** We thank Macarthur Diagnostic Imaging for donating time on the CT scanner  
192and for use of their Campbelltown facility.

193

194**Author contributions** M.G., M.D., D.H. and P.N. performed the experiment. P.N. and M.G. wrote  
195the paper. All authors designed the experiment, discussed the results, analysed the data and  
196commented on the manuscripts.

197

198**Author information** Correspondence and requests for materials should be addressed to M.G.  
199(mark.greco@alp.admin.ch).

200

201**Figure 1:** A *T. carbonaria* worker mummifies a live small hive beetle by gluing bits of batumen on  
202its elytra and legs.

203



204

205

206

207

208

209

210

211



212

213 **Figure 2:** Live mummification of adult small hive beetles in *T. carbonaria* hives visualised by CT

214 scans: (a) 3D CT image of *T. carbonaria* brood (single arrow) and two small hive beetles below

215 brood (double arrows); (b) 2D CT image of small hive beetles (short arrows) in entrance of

216 *T. carbonaria* hive demonstrating no change in position after 10 min.



217

218

219


220

221

222

223

224 **Figure 3:** A 3D pseudocolour CT scan image of a *T. carbonaria* hive, detailing brood (b) and live  
225 mummified small hive beetles (four white oval bodies) in entrance (e).



226

227

228

229

230

231

232

233**Figure 4:** Photograph of a *T. carbonaria* hive invaded by reproducing small hive beetles, detailing  
234brood (b) and small hive beetle larvae (L). The hive became vulnerable to invasion after being  
235weakened as a result of extreme ambient temperature (48°C).



236

237

238

239