



Elevated CO₂ and O₃ Effects on Fine-Root Life Span in Ponderosa Pine

Donald L. Phillips¹, Mark G. Johnson¹, David T. Tingey¹, Marjorie J. Storm²
¹ US Environmental Protection Agency, National Health & Environmental Effects Laboratory, Corvallis, OR;
² Dynamac International, Inc., c/o US EPA, Corvallis, OR

Abstract

Atmospheric carbon dioxide (CO₂) and ozone (O₃) concentrations are rising, which may have opposing effects on tree C balance and allocation to fine roots. More information is needed on interactive CO₂ and O₃ effects on roots, particularly fine-root life span, a critical demographic parameter and determinant of soil C and N pools and cycling rates. We conducted a study in which ponderosa pine (*Pinus ponderosa*) seedlings were exposed to two levels of CO₂ and O₃ in sun-lit controlled-environment terracoms for three years. Minirhizotrons were used to monitor individual fine roots in three soil horizons every 28 days. Proportional hazards regression was used to analyze effects of CO₂, O₃, diameter, depth, and season of root initiation on fine-root survivorship. More fine roots were produced in the elevated CO₂ treatment than in ambient CO₂. Median life spans varied from 140-448 days depending on the season of root initiation. Elevated CO₂, increasing root diameter, and increasing root depth all significantly increased fine-root survivorship and median life span. Life span was slightly, but not significantly, lower in elevated O₃, and increased O₃ did not reduce the effect of elevated CO₂. These results indicate the potential for elevated CO₂ to increase the number of fine roots and their residence time in the soil, which is also affected by root diameter, root depth, and phenology.

Introduction

- CO₂ and O₃ are generally increasing worldwide
- Leaves are sites of uptake and direct action, but CO₂ and O₃ can affect C assimilation and allocation with consequences for whole plant
- Fine roots are especially important because of:
 - role in water and nutrient uptake
 - significant contribution to NPP and its responses to CO₂ and O₃
- CO₂ and O₃ have potentially offsetting effects of on fine roots (allocation, growth, mortality)
- Fine root life span is an important determinant of soil C pools and cycling rates
- Multi-year studies are needed on CO₂ & O₃ effects on fine root longevity

Hypotheses


1. Elevated CO₂ alone will increase life span
2. Elevated O₃ will alone decrease life span
3. Elevated CO₂ and O₃ together will show additive compensatory effects on life span
4. Life span will increase with root diameter, soil depth, and vary with season of root formation

Methods


- 3 year experiment on ponderosa pine seedlings
- Sun-lit controlled environment mesocosms at EPA lab in Corvallis, OR
- 2-way factorial (2 CO₂ x 2 O₃ with 3 replicates)
 - CO₂: ambient (AC), elevated (EC; AC+270 ppm)
 - O₃: low (LO; 0 ppm-h seasonal SUM06) high (HO; 10-26 ppm-h seasonal SUM06)
- Minirhizotron tubes at 4 depths
- Each fine root (<= 2 mm diameter) was measured in images recorded every 28 days
- Proportional hazards regression was used to analyze the effects on fine-root life span of:
 - CO₂ and O₃ treatments
 - season of root formation, horizon, diameter



Mesocosm array



Mesocosm cross-section



Minirhizotron imaging system



Minirhizotron image with roots

Results and Discussion


- 46% more individual roots were produced in elevated CO₂
- Elevated CO₂ significantly increased median life span
- Effect was not due to differences in diameter or depth

- Median life span was 10% longer at low ozone but the difference was not significant
- High ozone treatment may have been too mild for significant effect

- Median life span increased 6-fold with depth
- Possible factors: more stable temperature & moisture, reduced herbivory, lower maintenance respiration in cooler temperatures

- Median life span increased 3-fold over diameter categories
- Possible factors: functional differences, vulnerability to stresses (e.g., parasitism, drought), construction costs, respiration rates, root branch order

- Median life span varied >3-fold over seasonal cohorts
- Late summer/fall cohorts lived longest; winter/early spring cohort had shortest life spans
- Over all cohorts, monthly survival rates were greatest in Jan-Apr, periods of low soil temp. and peak soil moisture



Conclusions

- Elevated CO₂ led to longer-lived fine roots in growing ponderosa pine seedlings, as did increasing depth & diameter
- No compensatory effects of elevated CO₂ and elevated O₃ on root life span were observed
- Fine root C turnover at end of study was lower for EC (290 g/m²/yr) than for AC (374 g/m²/yr)
- Thus, elevated CO₂ may lead to increased root C residence time in the soil (at least for tree seedlings)

This poster does not reflect EPA policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.