

of monoclonal antibodies specific for type II collagen together with lipopolysaccharide and is characterized by an early innate phase followed by a later adaptive phase. Compared with wild-type mice, the onset of CAIA in T-bet-deficient mice was delayed and was less severe in both the initial and the late phases. Mice that are deficient in RAG2 (recombination-activating gene 2), which completely lack B and T cells, developed early joint inflammation, but the later phase was attenuated. And mice that are deficient in both T-bet and RAG2 were resistant to the induction of arthritis.

To further assess the cellular components of the innate phase, the authors carried out adoptive-transfer experiments. Transfer of wild-type DCs to T-bet-deficient mice on day 1 of CAIA induction restored inflammatory arthritis, and wild-type DCs restored the

early phase response when transferred to mice that are deficient in both T-bet and RAG2. Consistent with this role for DCs in the early inflammatory phase, the authors showed that T-bet-deficient DCs produced less of the pro-inflammatory mediators interleukin-1 α and CC-chemokine ligand 3 (CCL3) and more CCL17, a ligand for the T_H2-cell-specific chemokine receptor CCR4, than wild-type DCs. Moreover, the ability of DCs to prime naive T cells *in vivo* was impaired in the absence of T-bet.

In light of these results, T-bet could be a target to control inflammation at multiple points in both innate and adaptive immune responses.

Lucy Bird

ORIGINAL RESEARCH PAPER Wang, J. *et al.* Transcription factor T-bet regulates inflammatory arthritis through its function in dendritic cells. *J. Clin. Invest.* 12 Jan 2006 (doi:10.1172/JCI26631)



are also structurally similar to purine bases and have recently been reported to be a danger signal) and calcium pyrophosphate dihydrate (both of which elicit inflammatory responses; known as gout and pseudogout, respectively) induce caspase-1-dependent IL-1 β secretion by primed macrophages, and this relies on components of the inflammasome, including NALP3. Dixit and colleagues found that NALP3 is also required for pro-caspase-1 activation by another danger signal, ATP (which, *in vivo*, is released into the extracellular environment by dying cells), and after exposure to certain Gram-positive bacteria but not certain Gram-negative bacteria.

It is thought that, similar to recognition by TLRs, pathogen-associated molecular patterns interact with the leucine-rich-repeat domains of NALPs, leading to activation of the inflammasome.

The study by Núñez and colleagues provides evidence (supported in part by the other two studies) that TLRs and NALP3 induce immune responses through recognition of the same or similar microbial structures and that IL-1 β secretion is independently regulated at two stages: the first is TLR dependent and results in nuclear-factor- κ B-mediated transcription of the gene that encodes pro-IL-1 β , and the second is NALP3 dependent and results in cleavage of pro-IL-1 β into IL-1 β .

Taken together, these results show that NALP3 has a crucial role in host defence against certain bacteria and that it might be a proximal sensor of cellular stress and danger signals. In addition, these studies provide insight into the mechanisms of autoinflammatory disorders that involve NALP3.

Davina Dadley-Moore

ORIGINAL RESEARCH PAPERS
Kanneganti, T.-D. *et al.* Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. *Nature* 11 Jan 2006 (doi:10.1038/nature04517) | Martinon, F., Pétrilli, V., Mayor, A., Tardivel, A. & Tschopp, J. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 11 Jan 2006 (doi:10.1038/nature04516) | Mariathasan, S. *et al.* Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 11 Jan 2006 (doi:10.1038/nature04515)

IN BRIEF

T-CELL DEVELOPMENT

Premature expression of chemokine receptor CCR9 impairs T cell development.

Uehara, S. *et al.* *J. Immunol.* **176**, 75–84 (2006)

Only late CD4⁺CD8⁻ double-negative (DN) thymocytes and CD4⁺CD8⁺ double-positive thymocytes express CC-chemokine receptor 9 (CCR9). To investigate whether expression of CCR9 controls the migration of thymocytes and their retention in the thymus, Uehara *et al.* generated transgenic mice in which CCR9 was expressed throughout T-cell development. CD4⁺ and CD8⁺ single-positive (SP) thymocytes could exit the thymus, indicating that CCR9 downregulation is not essential for SP thymocyte emigration. By contrast, ectopic expression of CCR9 by early DN thymocytes partially blocked thymocyte development at the DN3 stage, and CD25⁺ thymocytes (DN2 and DN3 thymocytes) were found throughout the cortex rather than at the subcapsular region. These results indicate that the timing of CCR9 expression is important for normal thymocyte development and that this might be because CCR9 controls CD25⁺ thymocyte localization.

HAEMATOPOIESIS

Glycogen synthase kinase-3 is an *in vivo* regulator of hematopoietic stem cell repopulation.

Trowbridge, J. J. *et al.* *Nature Med.* **12**, 89–98 (2006)

Glycogen synthase kinase 3 (GSK3) regulates several signalling pathways (such as the Notch and WNT signalling pathways) that affect the *in vitro* function of haematopoietic stem cells (HSCs). In this study, non-obese diabetic-severe combined immunodeficient mice were transplanted with HSCs from normal mice in the presence or absence of an ATP-competitive inhibitor of GSK3. Treatment with the GSK3 inhibitor increased recipient haematopoietic-cell reconstitution in both the short and the long term and thereby increased recipient survival. Further analysis showed that the GSK3 inhibitor increased the production of progenitor cells while maintaining the number of HSCs, leading the authors to suggest that treatment with GSK3 inhibitors might be a useful therapeutic approach to increase recipient haematopoietic-cell reconstitution after HSC transplantation.

HAEMATOPOIESIS

Antagonistic effect of CCAAT enhancer-binding protein- α and Pax5 in myeloid or lymphoid lineage choice in common lymphoid progenitors.

Hsu, C.-L. *et al.* *Proc. Natl Acad. Sci. USA* **103**, 672–677 (2006)

Although lymphoid-lineage-committed progenitors, such as common lymphoid progenitors (CLPs) and pro-T cells, are mainly destined to become lymphocytes, they maintain a latent myeloid differentiation potential, which can be initiated by stimulation through ectopic interleukin-2 receptor (IL-2R). This study shows that IL-2R triggering of CLPs and pro-T cells results in rapid upregulation of the myeloid-related transcription factor CCAAT enhancer-binding protein- α (C/EBP α). Moreover, ectopic expression of C/EBP α in these cells leads to myeloid-cell development. C/EBP α expression completely suppressed expression of the B-cell-lineage transcription factor PAX5 in CLPs and pro-T cells. By contrast, IL-2R stimulation of pro-B cells did not lead to lineage conversion, owing to insufficient downregulation of PAX5 expression by C/EBP α , indicating that complete loss of PAX5 is crucial for lymphoid-to-myeloid-lineage conversion.