

## ORIGINAL ARTICLE

**Bid can mediate a pro-apoptotic response to etoposide and ionizing radiation without cleavage in its unstructured loop and in the absence of p53**C Maas, E de Vries, SWG Tait<sup>1</sup> and J Borst

Division of Immunology, The Netherlands Cancer Institute, Amsterdam, The Netherlands

BH3-only protein Bid is a key player in death receptor-induced apoptosis, because it provides the link with the mitochondrial route for caspase activation. In this pathway, Bid is activated upon cleavage by caspase-8. Its BH3 domain-containing carboxy-terminal fragment subsequently provokes mitochondrial outer membrane permeabilization by Bak/Bax activation. Bid has also been implicated in the apoptotic response to ionizing radiation (IR) and the topoisomerase inhibitor etoposide, anti-cancer regimens that cause double-strand (ds)DNA breaks. We confirm the existence of this pathway and show that it is p53-independent. However, the degree of Bid participation in the apoptotic response to dsDNA breaks depends on the nature of cell transformation. We used Bid-deficient mouse embryonic fibroblast (MEF) lines that were reconstituted with Bid to control the cellular background and demonstrated that the Bid-dependent apoptotic pathway induced by IR and etoposide operates in MEFs that are transformed by SV40, but is not evident in E1A/Ras-transformed MEFs. The Bid-dependent apoptotic response in p53-deficient SV40-transformed MEFs contributed to clonogenic execution of the cells, implying relevance for treatment outcome. In these cells, Bid acted in a conventional manner in that it required its BH3 domain to mediate apoptosis in response to IR and etoposide, and triggered apoptotic execution by indirect activation of Bak/Bax, mitochondrial permeabilization and caspase-9 activation. However, the mechanism of Bid activation was unconventional, because elimination of all known or suspected cleavage sites for caspases or other proteolytic enzymes and even complete elimination of its unstructured cleavage loop left Bid's pro-apoptotic role in the response to IR and etoposide unaffected.

*Oncogene* (2011) 30, 3636–3647; doi:10.1038/onc.2011.75; published online 21 March 2011

**Keywords:** Bid; apoptosis; DNA damage; p53

Correspondence: Professor J Borst, Division of Immunology, The Netherlands Cancer Institute, Plesmanlaan 121, Amsterdam 1066 CX, The Netherlands.

E-mail: j.borst@nki.nl

<sup>1</sup>Current address: Department of Immunology, St Jude Children's Research Hospital, Memphis, TN 38105, USA.

Received 11 May 2010; revised and accepted 8 February 2011; published online 21 March 2011

**Introduction**

DNA-damaging regimens can limit the clonogenicity of tumor cells via irreversible cell cycle arrest, mitotic catastrophe or apoptosis (Brown and Wouters, 1999; Schmitt *et al.*, 2000). The apoptotic response to DNA damage relies on Bcl-2 family members, with BH3-domain only proteins and Bak/Bax acting in a pro-apoptotic manner and inhibitory Bcl-2 family counteracting their activity. Upon apoptotic stimulation, BH3-only proteins translocate to the mitochondria, where they activate Bax and/or Bak that enable mitochondrial outer membrane permeabilization (MOMP). This releases cytochrome *c*, Smac/DIABLO and HtrA2/Omi into the cytosol, resulting in the activation of caspase-9 and effector caspases, and apoptotic execution (Wang, 2001).

The p53 tumor suppressor protein has an important role in the apoptotic response to DNA damage. It may directly activate Bax/Bak at the mitochondria and acts as a transcription factor to direct the expression of apoptotic mediators (Chipuk and Green, 2006). These include the BH3-only proteins Puma, Noxa and Bid, as well as Bax and Apaf-1, all of which contribute to the mitochondrial apoptosis pathway. Analysis of Puma- or Noxa-deficient mice corroborated the role of these BH3-only proteins in p53-dependent apoptosis (Villunger *et al.*, 2003). Side-by-side analysis of the responsiveness of *p53*<sup>-/-</sup>, *puma*<sup>-/-</sup>, *noxa*<sup>-/-</sup> and *puma;noxa*<sup>-/-</sup> cells to ionizing radiation (IR) and etoposide furthermore indicated that Puma and Noxa are the main effectors of p53-dependent DNA-damage-induced apoptosis in lymphoid cells and transformed mouse embryonic fibroblasts (MEFs), with Puma having the dominant role (Michalak *et al.*, 2008). However, the combined deletion of Puma and Noxa did not always offer the same level of protection as p53, implicating other p53-regulated BH3-only proteins. Moreover, p53-mutant cells can also undergo DNA-damage-induced apoptosis (Strasser *et al.*, 1994) and the question is which BH3-only proteins activate the mitochondria in that case. Expression of Puma by genotoxic stimuli is strictly p53-dependent, but that of Noxa is not (Yu *et al.*, 2001; Ploner *et al.*, 2008), while Bid is constitutively expressed in many normal and tumor cell types (Krajewska *et al.*, 2002). Therefore, both Noxa and Bid, but possibly also other BH3-only proteins, may contribute to DNA-damage-induced apoptosis in p53 wild-type (WT) or mutant cells. Indeed, in cerebellar neurons, only the

combined deletion of Puma, Bid and Bim prevented IR-induced apoptosis (Ren *et al.*, 2010).

Bid is the requisite BH3-only protein that connects death receptors to the mitochondrial route for caspase activation. In this pathway, Bid is activated upon proteolytic cleavage in its unstructured loop by caspase-8 (Luo *et al.*, 1998). This cleavage facilitates mitochondrial association of the cleaved Bid complex by exposing a glycine residue that becomes myristoylated (Zha *et al.*, 2000). Moreover, it triggers ubiquitination and degradation of the N-terminal fragment of Bid, thus liberating the BH3 domain in the C-terminal fragment to perform its pro-apoptotic function (Tait *et al.*, 2007). Early studies suggested that Bid can also contribute to DNA damage-induced apoptosis: Primary Bid<sup>-/-</sup> MEFs were less sensitive to adriamycin- and 5-fluorouracil-induced apoptosis than WT MEFs (Sax *et al.*, 2002) and introduction of Bid sensitized primary Bid<sup>-/-</sup> MEFs to ultraviolet-, etoposide- and cisplatin-induced apoptosis (Sarig *et al.*, 2003). Because Bid can be cleaved by effector caspases and can participate in a feed-forward loop for MOMP (Slee *et al.*, 2000), it is important to assess whether Bid has the primary role in Bax/Bak activation in such cases. We have demonstrated that Bid was indeed responsible for the primary induction of MOMP upon treatment with IR or etoposide in p53-mutant Jurkat T leukemic cells (Werner *et al.*, 2004).

A prominent study confirmed a role for Bid in the apoptotic response to DNA damage. In telomerase (hTERT)-immortalized Bid<sup>-/-</sup> MEFs, WT Bid reconstitution restored the defective apoptotic response to IR and etoposide to WT levels. Bid acted as a sensor of double-strand (ds)DNA breaks, as it was phosphorylated by Ataxia Telangiectasia, mutated (ATM) when cells were treated with IR or etoposide. This triggered an unsuspected function of Bid in S-phase arrest, which was hypothesized to favor DNA repair and cell survival (Kamer *et al.*, 2005). Bid was therefore proposed to have a dual role in the response to dsDNA breaks: promotion of repair by a delay in S-phase progression, followed by apoptosis in case repair is insufficient (Gross, 2006). Related work used immortalized myeloid progenitor cells to demonstrate Bid's pro-survival role in S-phase arrest, which proved BH3-domain independent (Zinkel *et al.*, 2005). However, an independent study, in which cells from newly developed Bid<sup>-/-</sup> mice were extensively examined, failed to reveal either a pro-apoptotic or a pro-survival role for Bid in the DNA-damage response (Kaufmann *et al.*, 2007). The ensuing debate centered in part on the cell types that had been used for analysis, their genetic background, cell cycle status and/or transformed state, because these might provide clues for the discrepant outcome of the studies (Zinkel *et al.*, 2007).


We demonstrate that Bid can indeed be the requisite apoptotic mediator in response to dsDNA damage, but its participation depends on the nature of cell transformation. SV40-transformed MEFs required Bid for IR-, but not etoposide-induced apoptosis, while down-regulation of p53 rendered them fully reliant on Bid for induction of apoptosis and clonogenic execution by both

IR and etoposide. E1A/Ras-transformed MEFs, however, did not employ this pathway, either in the presence or in the absence of p53. Bid acted conventionally in the p53-independent apoptotic response to IR and etoposide in that it required its BH3 domain and executed the cells via Bak/Bax-dependent MOMP and caspase-9 activation. The mechanism of Bid activation by DNA-damaging regimens was unconventional though, because extensive mutation analysis demonstrated that it was fully independent of proteolytic cleavage in its unstructured loop and even proceeded normally when the loop was deleted.

## Results

### *Bid can convey the apoptotic response to etoposide and IR, in a p53-independent manner*

We found previously that Bid was required for IR- and etoposide-induced apoptosis in p53 mutant Jurkat T leukemic cells (Werner *et al.*, 2004). Therefore, we hypothesized that the p53 status might dictate whether cells depend on Bid for their apoptotic response to dsDNA damage. To examine this, we used immortalized MEFs as a cellular model, to match earlier studies (Kamer *et al.*, 2005; Zinkel *et al.*, 2005). Bid<sup>-/-</sup> MEFs that had been transformed with the whole genome of SV40 were stably transduced with an empty RNA interference (RNAi) vector (EV1) or with a vector encoding p53 short hairpin (sh)RNA to create p53-proficient (control) and p53-deficient (p53-RNAi) variants of the same cells. These cell lines were stably transduced with a Bid expression vector or with the empty vector (EV2) to have each cell line in a WT Bid or a Bid<sup>-/-</sup> version. The lines were created in parallel in the same time frame. This well-controlled set-up excluded differences between experimental groups that were due to the genetic background of the cells, the process of retroviral transduction or other variables not related to expression of Bid or p53. Immunoblotting showed that the starting population of Bid<sup>-/-</sup> MEFs expressed p53 and lacked Bid. It also validated silencing of p53 and expression of Bid in the relevant transduced cells (Figure 1a). Moreover, immunoblotting for Bim, Puma, Mcl-1, Bcl-2, Bcl-x<sub>L</sub> and Bcl-W showed that their expression was comparable in p53 RNAi Bid<sup>-/-</sup> MEFs with and without exogenous Bid (Supplementary Figure 1). The cell lines were treated with etoposide or IR at different doses and analyzed for their apoptotic response after 24 or 48 h, respectively. Apoptosis was read out by flow cytometry as the percentage of cells with cleaved caspase-3. The control Bid<sup>-/-</sup> MEFs (EV1 + EV2) showed an apoptotic response to etoposide, but not to IR. Reconstitution of Bid enhanced etoposide-induced apoptosis and enabled IR-induced apoptosis (Figure 1b). The p53-deficient Bid<sup>-/-</sup> MEFs (p53 RNAi + EV2) did not show an apoptotic response to either etoposide or IR, while Bid reconstitution enabled this response (Figure 1c). These data clearly indicate a role for Bid in the apoptotic response to these DNA-damaging stimuli in SV40-transformed MEFs. These cells require Bid



**Figure 1** The p53 status can influence the reliance on Bid for etoposide- and IR-induced apoptosis. SV40-transformed Bid<sup>-/-</sup> MEFs were retrovirally transduced (Td) with an empty RNAi vector (EV1) or with p53 shRNA, together with an empty expression vector (EV2) or a vector encoding WT Bid. (a) Validation of p53 downregulation by p53 shRNA and Bid expression from the introduced vector in the relevant MEF cell lines, as demonstrated by immunoblotting with antibodies to p53 and Bid on total cell lysates. Immunoblotting for actin served as a loading control. (b, c) Sensitivity of control (EV1-transduced) MEFs (b) and p53-deficient (p53-RNAi) MEFs (c) to apoptosis induction by DNA-damaging regimens. Cells were exposed to the indicated dosages of etoposide or IR. Apoptosis levels were determined as the percentage of cells with cleaved caspase-3 after 24 h (etoposide) or 48 h (IR). Data presented are expressed as means of three independent experiments  $\pm$  s.d. Statistically significant differences between values of EV and Bid samples are indicated for \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ .

for IR-induced apoptosis regardless of their p53 status and they require Bid for etoposide-induced apoptosis when p53 cannot participate in the response. We conclude that Bid can convey the apoptotic response to etoposide and IR in a p53-independent manner. Moreover, the p53 status of transformed cells can determine the extent of their reliance on Bid for the apoptotic response to these stimuli.


#### *Bid can contribute to clonogenic cell death in response to etoposide and IR*

For tumor therapy, the relevant parameter is the clonogenic potential of the tumor cells after treatment. To investigate the importance of Bid-dependent apoptosis in arresting the clonogenicity of transformed cells with DNA-damaging stimuli, we performed clonogenic survival assays. The p53-RNAi Bid<sup>-/-</sup> MEF cell lines expressing Bid or empty control vector (EV) were exposed to different doses of etoposide or IR and

surviving colonies were quantified. The presence of Bid in p53-deficient MEFs reduced clonogenic survival upon etoposide (Figure 2a) or IR treatment (Figure 2b). This demonstrated that in these p53-deficient, SV40-transformed cells, Bid is important for clonogenic cell death following treatment with DNA-damaging anti-cancer regimens.

#### *The nature of cell transformation determines whether Bid mediates a pro-apoptotic response to etoposide or IR*

The discussion regarding the existence of a Bid-dependent apoptotic response to DNA damage focused in part on the cellular background of the WT and Bid<sup>-/-</sup> cells that were compared. However, in our Bid reconstitution set-up, such variables were excluded. Another variable might be the nature of cell transformation. Kamer *et al.* (2005) primarily studied hTERT-immortalized Bid<sup>-/-</sup> MEFs that were reconstituted with Bid and found Bid to



**Figure 2** Bid can contribute to clonogenic cell death in response to etoposide and IR. (a, b) P53-RNAi Bid<sup>-/-</sup> MEFs carrying an empty vector (EV) or stably expressing WT Bid were exposed to the indicated dosages of etoposide (a) or IR (b). Surviving colonies were counted 14 days post-treatment. The relative surviving fraction is plotted against the dose of etoposide or IR. Data presented are representative of two independent experiments.

have an important role in etoposide- and IR-induced apoptosis. Kaufmann *et al.* (2007), however, compared various E1A/Ras-transformed WT and Bid<sup>-/-</sup> MEF lines and found no evidence for a contribution of Bid to these apoptotic responses.

Therefore, to gain further insight, we tested E1A/Ras-transformed Bid<sup>-/-</sup> MEF lines for their response to etoposide and IR, before and after Bid reconstitution and in the presence or absence of p53. Four different cell lines were created, exactly as outlined for SV40-transformed MEFs. Bid reconstitution and p53 downregulation were validated by immunoblotting (Supplementary Figure 2a). The cell lines were exposed to various doses of etoposide and IR. Downregulation of p53, although not complete (Supplementary Figure 2a), reduced the apoptosis sensitivity of the cells to both stimuli (Supplementary Figure 2b). However, Bid reconstitution did not alter the apoptosis sensitivity of these E1A/Ras-transformed cells to etoposide or IR, either in the p53-proficient or in the p53-deficient situation (Supplementary Figure 2b). Our approach excluded cellular background differences and confirmed that Bid does not contribute to the apoptotic response to etoposide or IR in E1A/Ras-transformed MEFs. Moreover, p53 downregulation did not make the Bid-dependent pathway more prevalent.

#### *Bid requires its BH3 domain to indirectly activate Bak/Bax during DNA-damage-induced apoptosis*


The conventional pro-apoptotic function of BH3-only proteins lies in their  $\alpha$ -helical BH3 domain, which can interact with the BH1/BH2 groove of other family members (Letai *et al.*, 2002). To study whether Bid mediated DNA-damage-induced apoptosis by virtue of its BH3 domain, we generated a Bid variant with glycine 94 in its BH3 domain mutated into a glutamate (Bid G94E). It was previously shown that the tBid-C (C-terminal fragment of Bid) with a G94E mutation could not efficiently interact with Bak/Bax or the anti-apoptotic Bcl-2 proteins and displayed reduced pro-apoptotic activity (Willis *et al.*, 2007). SV40-transformed p53 RNAi Bid<sup>-/-</sup> MEF cells were stably transduced to

express either WT Bid or Bid G94E (Figure 3a). The effect of the G94E mutation on Bid function was tested by its impact on death receptor-induced apoptosis. Cells were treated with different doses of tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) in combination with cycloheximide (to inhibit NF- $\kappa$ B activation) and apoptosis was read out as before. TNF $\alpha$ -induced apoptosis was significantly lower in cells expressing the Bid G94E mutant than in cells expressing WT Bid, thus validating the construct (Figure 3b). The Bid G94E mutant was also significantly less effective than WT Bid in inducing apoptosis after treatment with etoposide or IR (Figure 3b), indicating that Bid required a functional BH3 domain to mediate apoptosis in response to these stimuli.

In the indirect model for Bax/Bak activation, inhibitory Bcl-2 family members bind and sequester Bax/Bak, thus preventing them from multimerizing and causing MOMP. BH3-only proteins displace Bax/Bak from this complex by binding to inhibitory Bcl-2 family members with their BH3 domain (Willis *et al.*, 2007). In the direct activation model, Bid activates Bax and/or Bak by a direct physical interaction using its BH3 domain (Letai *et al.*, 2002). A tBid-C mutant with glycine 94 replaced by an alanine (G94A) cannot bind to Bax and Bak but binds all anti-apoptotic Bcl-2 proteins (Willis *et al.*, 2007). To test whether Bid acted in the DNA damage pathway by direct or indirect Bax/Bak activation, SV40-transformed p53-RNAi Bid<sup>-/-</sup> MEFs were reconstituted with WT Bid or G94A Bid (Figure 3c) and their apoptotic response to DNA-damage-induced apoptosis was examined. The Bid G94A mutant was as effective as WT Bid in mediating etoposide- or IR-induced apoptosis (Figure 3d). We conclude from these findings that in p53-deficient SV40-transformed MEFs, Bid mediates apoptosis in response to etoposide and IR by virtue of its BH3 domain and does not directly activate Bax/Bak, but most likely relieves them from inhibition by anti-apoptotic Bcl-2 proteins.

#### *Bid-dependent DNA-damage-induced apoptosis proceeds via the mitochondrial pathway*

To evaluate whether Bid-dependent DNA-damage-induced apoptosis proceeded via the mitochondrial pathway, we overexpressed Bcl-2 in Bid-expressing SV40-transformed p53-RNAi Bid<sup>-/-</sup> MEFs (Figure 4a) and tested the effect on etoposide- and IR-induced apoptosis. Bcl-2 overexpression significantly inhibited both etoposide- and IR-induced apoptosis (Figure 4b). This further supports that Bid acts in this pathway by virtue of BH3-domain-dependent interactions with its family members. It also strongly suggests that Bid-dependent DNA-damage-induced apoptosis required MOMP. Because Bcl-2 may also regulate the permeability of the endoplasmic reticulum membrane (Lam *et al.*, 1994), the inhibitory effect of Bcl-2 did not unambiguously implicate the mitochondria in DNA-damage-induced apoptosis. Therefore, we overexpressed dominant-negative caspase-9 (dnCaspase-9; Figure 4c), which abrogates apoptosome-mediated caspase-9 activation. dnCaspase-9 significantly blocked




**Figure 3** Bid uses its BH3 domain to indirectly activate Bax and/or Bak during etoposide- and IR-induced apoptosis. p53-RNAi Bid<sup>-/-</sup> MEFs were transduced (Td) to stably express WT Bid, the Bid G94E mutant or the Bid G94A mutant. (a, c) Downregulation of p53 by RNAi and Bid protein expression from the introduced vectors were validated by immunoblotting, where actin served as a loading control. (b) p53-RNAi Bid<sup>-/-</sup> MEFs expressing WT Bid or Bid G94E were exposed to the indicated dosages of etoposide or IR, or to TNF $\alpha$  in combination with cycloheximide (CHX) as a validation of the Bid G94E mutant. Apoptosis levels were determined as the percentage of cells with cleaved caspase-3 after 24 h (etoposide), 48 h (IR) or 5 h (TNF $\alpha$  + CHX). Background apoptosis levels, as those observed in p53-RNAi Bid<sup>-/-</sup> MEF carrying EV2, were subtracted from obtained values for all stimuli. (d) p53-RNAi Bid<sup>-/-</sup> MEFs expressing WT Bid or Bid G94A were exposed to the indicated dosages of etoposide or IR and apoptosis was read out as outlined in (b). Data in (b) and (d) are expressed as means of three independent experiments  $\pm$  s.d. Statistically significant differences between values of WT Bid and Bid G94E are indicated in (b) for \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ .

both etoposide- and IR-induced apoptosis (Figure 4d). Together, these observations demonstrate that Bid-dependent DNA-damage-induced apoptosis proceeds via the mitochondrial pathway for caspase activation.

*Bid activation by DNA-damaging stimuli does not require cleavage at aspartates 55, 60 and/or 75*

Bid is an auto-inhibitory molecule that has—in the full-length inactive state—its pro-apoptotic BH3 domain sequestered by residues in its N-terminal region (see


Supplementary Figure 3 for Bid structure; Chou *et al.*, 1999; McDonnell *et al.*, 1999; Tan *et al.*, 1999). Proteolytic cleavage of Bid in its unstructured loop triggers mitochondrial translocation (Gross *et al.*, 1999) and proteasomal degradation of Bid's N-terminal fragment, allowing tBid-C to perform its pro-apoptotic function (Tait *et al.*, 2007). In the death receptor pathway, Bid is cleaved at aspartate 60 (D60) by caspase-8 (Luo *et al.*, 1998). Caspase-2 and -3 can also cleave Bid at D60 (Gross *et al.*, 1999; Guo *et al.*, 2002), whereas Granzyme B cleaves Bid at aspartate 75 (D75; Luo *et al.*, 1998).



**Figure 4** Bid-dependent apoptosis in response to etoposide and IR proceeds via the mitochondria. p53-RNAi Bid<sup>-/-</sup> MEFs that had been transduced to stably express WT Bid were used as such (control) or additionally transduced (Td) to express Bcl-2 or dnCasp-9. (a, c) Downregulation of p53 by RNAi and expression of Bid, Bcl-2 or dnCasp-9 protein from the introduced vectors were validated by immunoblotting on total cell lysates, where actin served as a loading control. (b) Control or Bcl-2-expressing cells were exposed to the indicated dosages of etoposide or IR. Apoptosis levels were determined as the percentage of cells with cleaved caspase-3 after 24 h (etoposide) or 48 h (IR). (d) Control or dnCasp-9-expressing cells were exposed to the indicated dosages of etoposide or IR and apoptosis was read out as indicated for panel (b). Data presented in (b) and (d) are expressed as means of three independent experiments ± s.d. Statistically significant differences between values of control and Bcl-2 or dnCasp-9 samples are indicated for \**P*<0.05, \*\**P*<0.01 and \*\*\**P*<0.001.

Biochemical analysis after treatment of Bid-reconstituted MEF cells with etoposide or IR did not reveal Bid cleavage as assessed in total cell lysates (Supplementary Figure 4 and data not shown). This was not surprising because in our experience only minute amounts of Bid cleavage products are produced in response to these stimuli, which we could only demonstrate biochemically in purified mitochondrial fractions after biosynthetic labeling (Werner *et al.*, 2004 and results not shown). Moreover, because Bid can be cleaved by effector caspases in a feed-forward loop for MOMP (Slee *et al.*, 2000), analysis for the Bid cleavage that is instrumental in initial MOMP induction should be done in cells in which the feed-forward loop is excluded, such as by expression of dnCaspase-9. Also, under those conditions, we could not conclusively detect Bid cleavage products after treatment with etoposide or IR either in total cell lysates or after immunoprecipitation (results not shown). For these reasons and to assess the functional relevance of a possible Bid cleavage, we decided

to use a genetic approach. We first tested whether D60 or D75 were essential in p53-deficient MEFs for the responses to etoposide and IR. SV40-transformed p53-RNAi Bid<sup>-/-</sup> MEFs cells were stably reconstituted with D60E or D75E single-point mutants, or a D60E/D75E double-point mutant of Bid. Cells reconstituted with WT Bid served as control (Figure 5a). Death receptor stimulation by TNF $\alpha$  was used to validate the Bid mutants. In contrast to WT Bid, the Bid D60E mutant could not mediate TNF $\alpha$ -induced apoptosis. Bid D75E mutation did not significantly impede its capacity to relay the apoptotic signal (Figure 5b). This indicates that D60 is the predominant cleavage site for Bid activation in death-receptor-induced apoptosis. Bid D60E/D75E mutation prevented TNF $\alpha$ -induced apoptosis as effectively as the Bid D60E mutation (Figure 5c). Next, we studied the relevance of Bid cleavage at D60 and/or D75 for DNA-damage-induced apoptosis. Single D60E or D75E mutation did not affect etoposide or IR-induced apoptosis (Figure 5b); combined D60E/D75E mutation




**Figure 5** Bid cleavage at aspartate residues in its unstructured loop is not required for apoptosis induction in response to etoposide and IR. p53-RNAi Bid<sup>-/-</sup> MEFs were transduced (Td) to stably express WT Bid, or the Bid mutants Bid D60E, Bid D75E, Bid D60E/D75E or Bid D55E/D60E/D75E. (a, d) Downregulation of p53 by RNAi and expression of WT or mutant Bid protein from the introduced vectors were validated by immunoblotting on total cell lysates, where actin served as a loading control. (b) p53-RNAi Bid<sup>-/-</sup> MEFs expressing WT Bid, Bid D60E or Bid D75E were exposed to the indicated dosages of TNF $\alpha$  + cycloheximide (CHX), etoposide or IR. Apoptosis levels were determined as the percentage of cells with cleaved caspase-3 after 5 h (TNF $\alpha$  + CHX), 24 h (etoposide) or 48 h (IR). (c) P53-RNAi Bid<sup>-/-</sup> MEFs expressing WT Bid or Bid D60E/D75E were exposed to the indicated dosages of TNF $\alpha$  + CHX, etoposide or IR and apoptosis was read out as outlined for panel (b). (e) P53-RNAi Bid<sup>-/-</sup> MEFs expressing WT Bid or Bid D55E/D60E/D75E were exposed to the indicated dosages of etoposide or IR and apoptosis was read out as outlined for panel (b). Data presented in (b), (c) and (e) are expressed as means of three independent experiments  $\pm$  s.d. Statistically significant differences between values of Bid and Bid D60E or D60E/D75E samples are indicated for \* $P$  < 0.05, \*\* $P$  < 0.01 and \*\*\* $P$  < 0.001.

also had no effect (Figure 5c). Potentially, Bid can also be cleaved at aspartate 55 (D55) by caspase-3, as revealed by analysis with the Merops peptidase database (<http://merops.sanger.ac.uk>). We therefore expressed a Bid D55E/D60E/D75E mutant in SV40-transformed p53-RNAi Bid<sup>-/-</sup> MEFs (Figure 5d). This Bid mutant relayed the apoptotic signal as effectively as WT Bid (Figure 5e). Together, these data indicate that Bid does not require cleavage at any of the available aspartate residues in its unstructured loop to induce apoptosis in

response to the DNA-damaging stimuli. These data also excluded that Bid made its contribution to apoptotic execution in a feed-forward loop for MOMP, after cleavage by caspases downstream of the mitochondria.

*Bid activation by DNA-damaging stimuli does not require cleavage at other known protease recognition sites*  
*In vitro* studies have shown that Bid can also be activated through proteolytic cleavage by calpain or different



**Figure 6** Bid cleavage at alternative defined proteolytic cleavage sites in its unstructured loop is not required for apoptosis induction in response to etoposide and IR. p53-RNAi Bid<sup>-/-</sup> MEFs were transduced (Td) to stably express WT Bid, or the Bid Q58A, Bid S65A or Bid G70A/R71A mutants. (a) Downregulation of p53 by RNAi and expression of WT or mutant Bid protein from the introduced vectors were validated by immunoblotting on total cell lysates, where actin served as a loading control. (b) p53-RNAi Bid<sup>-/-</sup> MEFs expressing WT Bid, Bid Q58A, Bid S65A or Bid G70A/R71A were exposed to the indicated dosages of etoposide or IR. After 24 h (etoposide) or 48 h (IR), apoptosis levels were determined as the percentage of cells with cleaved caspase-3. Data are expressed as means of three independent experiments ± s.d.

cathepsins within its unstructured loop (Mandic *et al.*, 2002; Cirman *et al.*, 2004). Calpain cleavage site glycine 70 (G70) and cathepsin cleavage sites glutamine 58 (Q58), serine 65 (S65) and arginine 71 (R71) are highly conserved between species. To investigate whether Bid cleavage at one of these residues was required for

DNA-damage-induced apoptosis in SV40-transformed p53-deficient MEFs, we expressed WT Bid or Bid Q58A, S65A and G70A/R71A mutants in p53-RNAi Bid<sup>-/-</sup> MEFs (Figure 6a) and tested their effects on DNA-damage-induced apoptosis. All mutants were as capable as WT Bid to relay the apoptotic signal in response to etoposide and IR (Figure 6b). We conclude that Bid does not require proteolysis at defined calpain or cathepsin cleavage sites in its unstructured loop to induce apoptosis in response to the DNA-damaging stimuli.

*Bid activation by DNA-damaging stimuli does not involve cleavage within its unstructured loop*


Together, our results so far indicate that activation of Bid during DNA-damage-induced apoptosis does not require proteolysis at a conventional cleavage site within its unstructured loop. To study whether another cleavage site within its unstructured loop might be required for Bid activation, we generated a Bid mutant (Bid w/o loop) with its 37-amino-acid unstructured loop replaced by a short, random stretch of glycines and serines. All potential cleavage sites present in the unstructured loop were thereby removed. The SV40-transformed p53-RNAi Bid<sup>-/-</sup> MEFs were reconstituted with the Bid w/o loop mutant or WT Bid as a control (Figure 7a). In contrast to WT Bid, the Bid w/o loop mutant could not mediate TNF $\alpha$ -induced apoptosis at all, thus validating the construct (Figure 7b). However, the Bid w/o loop mutant was as effective as WT Bid in mediating etoposide- or IR-induced apoptosis (Figure 7b). These data prove that Bid activation during DNA-damage-induced apoptosis does not involve cleavage within its unstructured loop.

**Discussion**

In accord with certain previous works, we demonstrate that Bid can convey the apoptotic signal in response to IR or etoposide, but its participation depends on the cellular background. Participation of BH3-only proteins in the response to apoptotic stimuli is highly cell type- and stimulus-dependent. In the mouse, Puma and—to a lesser extent—Noxa are required for the apoptotic response to DNA damage in primary thymocytes, mature T cells, intestinal epithelial cells and neuronal cells (Michalak *et al.*, 2008). In mature B cells, BH3-only protein Bim was also important and in pre-B cells, combined deletion of the *puma* and *noxa* genes did not reproduce the effect of p53 deletion, indicating that other BH3-only proteins had a role (Erlacher *et al.*, 2005; Michalak *et al.*, 2008). Testing a potential role for Bid, Kaufmann *et al.* (2007) found normal apoptotic responses to IR and etoposide in Bid<sup>-/-</sup> thymocytes, pre-B cells, resting mature T- and B cells, and proliferating T cells. These findings are not in conflict with those of Kamer *et al.* (2005) and Zinkel *et al.* (2005), and emphasize that in these healthy lymphoid cells, Bid is not important for this response. In mouse brain tissue,



however, Bid complemented Bim and Puma in apoptosis induction after IR and the three proteins were collectively responsible for complete apoptotic execution.



In fact, the phenotype of mice with triple deletion of the *bid*, *bim* and *puma* genes suggested that these three proteins are collectively responsible for Bax and Bak activation in response to diverse stimuli in multiple tissues (Ren *et al.*, 2010).

Kamer *et al.* (2005) demonstrated by Bid reconstitution that in hTERT-immortalized MEFs, Bid mediated the apoptotic response to IR and etoposide. We have received hTERT-immortalized Bid<sup>-/-</sup>MEFs from this research group and independently confirmed this finding (results not shown). We report here that also in SV40-transformed MEFs, Bid conveys the apoptotic response to IR and etoposide. Bid most evidently contributed to the etoposide response when p53 was downregulated, but the IR response strongly relied on Bid, both in the presence and in the absence of p53. In E1A/Ras-transformed MEFs, however, Bid did not contribute to the apoptotic response to IR or etoposide, when p53 was either expressed or downregulated. We have hereby extended the study of Kaufmann *et al.* (2007), who compared various WT and Bid<sup>-/-</sup> E1A/Ras-transformed MEF cell lines, but did not perform reconstitution experiments or test the effect of p53 downregulation. We conclude that it is not the p53 status *per se*, but the oncogenic pathways that are active in the MEF cell lines, in combination with the p53 status, that determine their dependence on Bid for apoptosis induction by IR and etoposide.

Bid was reconstituted in its full-length, inactive form, in which it is expressed constitutively in many normal and tumor cell types (Krajewska *et al.*, 2002). We showed that Bid reconstitution did not alter the expression level of various pro- and anti-apoptotic Bcl-2 family proteins (Supplementary Figure 1). Also, Bid reconstitution did not alter the apoptosis sensitivity of transformed MEFs indirectly, because it did not influence the response of E1A/Ras-transformed MEFs to etoposide or IR at all. This is in agreement with the fact that Bid needs to be activated to perform its pro-apoptotic function. We found that in SV40-transformed MEFs, Bid's role in apoptosis induction in response to etoposide and IR was conventional, in the sense that it needed an intact BH3 domain, indirectly activated Bax and/or Bak, and relied on MOMP and caspase-9 to activate effector caspases. Bid activation did not involve

**Figure 7** Bid cleavage in its unstructured loop is not required for apoptosis induction in response to etoposide and IR. p53-RNAi Bid<sup>-/-</sup> MEFs were transduced (Td) to stably express WT Bid or Bid with the unstructured loop exchanged for a random stretch of Gly and Ser amino acids (Bid w/o loop). (a) Downregulation of p53 by RNAi and expression of WT or mutant Bid protein from the introduced vectors were validated by immunoblotting on total cell lysates, where actin served as a loading control. (b) p53-RNAi Bid<sup>-/-</sup> MEFs expressing WT Bid or the Bid w/o loop mutant were exposed to the indicated dosages of TNF $\alpha$  + cycloheximide (CHX), etoposide or IR. After 5 h (TNF $\alpha$  + CHX), 24 h (etoposide) or 48 h (IR), apoptosis levels were determined as the percentage of cells with cleaved caspase-3. Data are expressed as means of three independent experiments  $\pm$  s.d. Statistically significant differences between values of Bid and Bid w/o loop are indicated for \*\* $P < 0.01$  and \*\*\* $P < 0.001$ .

cleavage at defined caspase cleavage sites. Accordingly, the D59 caspase cleavage site in mouse Bid was found to be irrelevant for its supporting role in apoptosis induction by etoposide in primary MEFs (Sarig *et al.*, 2003). Bid can also participate in the apoptotic response to DNA-damaging anti-cancer regimens in human tumor cells. In p53 mutant Jurkat T-acute lymphoblastic leukemia cells, Bid depletion experiments indicated that Bid conveyed the apoptotic response to IR and etoposide. In this response, Bid can act downstream of the apoptosome in an amplification loop for mitochondrial activation (Shelton *et al.*, 2009). However, by expression of dominant-negative caspases and RNAi, we showed that Bid also acted upstream of the apoptosome in a caspase-independent manner (Werner *et al.*, 2004). In HeLa cervix carcinoma cells, which have a dysfunctional p53 pathway due to human papillomavirus-encoded E6 and E7 proteins, Bid downregulation by RNAi was found to decrease etoposide-, doxorubicin- and oxaliplatin-induced apoptosis. RNAi for Puma showed that Bid acted in concert with Puma in execution of the apoptotic response to oxaliplatin (Koehler *et al.*, 2008). Furthermore, in the response to oxaliplatin, Bid cleavage at D59 was not required for apoptosis induction (Anguissola *et al.*, 2009). These data agree that Bid can be activated in response to DNA damage in a caspase-independent manner. We now show that in SV40-transformed MEFs, defined cleavage sites for calpains or cathepsins also did not have a role in Bid activation. Moreover, even complete removal of the unstructured loop did not affect the pro-apoptotic response of Bid to IR or etoposide.

Although it is difficult to rule out Bid activation by cleavage outside its unstructured loop, the combined data suggest that it might be full-length Bid that induces apoptosis in response to the dsDNA break inducers etoposide and IR in the MEFs examined here. In the case of oxaliplatin stimulation of HeLa cells, a Bid-FRET probe indicated that full-length Bid translocated to the mitochondria in a caspase- and calpain-independent manner. We have also examined Bid translocation in the context of our study, but found the data difficult to quantify, because the different cells within a population responded to IR or etoposide in an asynchronous manner over the protracted period of 8–24 h after the initial stimulation that was analyzed. It was also not possible to determine whether mitochondrial association of Bid was the cause or consequence of apoptosis induction or induction of mitochondrial permeability (in case of cells expressing dnCaspase-9). Full-length Bid has previously been implicated in anoikis and glutamate-induced apoptosis in neurons (Valentijn and Gilmore, 2004; Ward *et al.*, 2006). Also, it has been found that full-length Bid can be transported to the mitochondria, with the help of PACS-2 (Simmen *et al.*, 2005). It has been proposed that Bid is activated in the nucleus to mediate DNA-damage-induced apoptosis, on the basis of several observations. First, full-length Bid was seen to translocate from the cytosol to the nucleus and from the nucleus to the cytosol in response to DNA-damaging anti-cancer drugs in HeLa cells (Oberkovitz *et al.*, 2007; Anguissola *et al.*,

2009). In addition, enforced nuclear retention of Bid through fusion of Bid with a nuclear localization signal inhibited etoposide-induced apoptosis (Oberkovitz *et al.*, 2007). Thus, DNA damage may activate full-length Bid to translocate from the nucleus to the mitochondria and induce MOMP. What might trigger this presumed re-localization is not known. The Bid phosphorylation that occurs in response to dsDNA break inducers is not required for its role in apoptosis induction (Kamer *et al.*, 2005). Because full-length Bid is auto-inhibited, it must be activated to expose its BH3 domain (Tait *et al.*, 2007). How this can occur when Bid is not cleaved in its unstructured loop is unclear. Potentially, post-translational modification can induce a conformational change that enables Bid to expose its BH3 domain and participate in apoptosis. This modification must occur outside the loop, as its removal did not affect Bid activity.

As a potential mediator of DNA-damage-induced apoptosis, Bid might function as a tumor suppressor. Zinkel *et al.* (2005) postulated such a role for Bid, which was supported by their previous finding that Bid<sup>-/-</sup> mice were at the risk of developing a fatal myeloproliferative disorder upon aging (Zinkel *et al.*, 2003). This has also been debated, but further work will have to point out whether loss of Bid can collaborate with oncogenic mutations to promote cellular transformation. We conclude that depending on the nature of cell transformation, Bid can be the decisive factor for apoptotic death of tumor cells in response to DNA-damaging anti-cancer regimens and can contribute to loss of clonogenicity independent of the p53 pathway. Bid hereby represents a common intermediate between death receptor- and DNA-damage-induced apoptosis. It may therefore, in certain tumor types, be a determinant for therapeutic outcome upon combined treatment with death receptor agonists and DNA-damaging anti-cancer regimens.

## Materials and methods

### Cells and stimulation

SV40-immortalized Bid<sup>-/-</sup> MEFs were originally from the laboratory of Dr S Korsmeyer (Harvard Medical School, Boston, MA, USA). The derivative Bid<sup>-/-</sup> MEF lines stably expressing empty vectors, the p53-targeting short hairpin RNA, Bid WT and Bid mutant complementary DNA (cDNA) were generated by retroviral transduction. Cells over-expressing Bcl-2 or the dnCaspase-9 active site mutant C287A were manufactured similarly. Cell lines were cultured in Dulbecco's modified Eagle's medium supplemented with 8% fetal bovine serum, 2 mM L-glutamine and antibiotics. Human recombinant TNF $\alpha$ , cycloheximide and etoposide were purchased from Sigma-Aldrich (St Louis, MO, USA). For apoptosis assays, cells were stimulated with the indicated dosages of TNF $\alpha$  with cycloheximide, etoposide or IR for the indicated time periods at 37 °C with 5% CO<sub>2</sub>. Irradiation of cells was performed with a <sup>137</sup>Cs source (415 Ci; Von Gahlen Nederland BV, Zevenaar, The Netherlands) at an absorbed dose rate of approximately 0.66 Gy/min.

### Constructs

The cDNAs encoding Bid mutants Bid D60E, Bid D75E, Bid D60E/D75E, Bid D55E/D60E/D75E, Bid Q58A, Bid S65A,

Bid G70A/R71A, Bid G94E and Bid G94A were generated with a Quickchange Site-directed Mutagenesis Kit using full-length human Bid cDNA as a template (Stratagene, La Jolla, CA, USA). The Bid w/o loop mutant was generated by inverse mutagenesis PCR from full-length human Bid cDNA with the following primers: forward, 5'-GGTTCAGGTTTCATCAGGTCAAGAAGACATCATCCGGAATATT-3' and reverse, 5'-TGATGAACCTGAACCACCTGGCAGCTCGTGGCCAGTGCGTC-3' (<http://openwetware.org>). The unstructured cleavage loop was hereby replaced by a 12-amino-acid stretch of glycine and serine residues, anticipated to display similar flexibility. WT Bid-, Bid mutant- and Bcl-2 cDNAs were cloned into the retroviral vector LZRS-IRES-Zeocin/pBR and dnCaspase-9 cDNA was cloned into the retroviral vector LZRS-IRES-GFP. Both vectors are derivatives of the LZRS-pBMN-LacZ vector, which was provided by Dr GP Nolan (Stanford University School of Medicine, Stanford, CA, USA). RNAi for p53 was performed using shRNA (complementary sense and antisense oligonucleotides with p53-targeting sequence 5'-GTACATGTGTAATAGCTCC-3') cloned into the retroviral vector pRETRO-SUPER, with a puromycin resistance cassette. All constructs were verified by dideoxynucleotide sequencing.

#### Retroviral gene transduction

To produce retrovirus, LZRS and pRETRO-SUPER constructs were transfected into the 293T human embryonic kidney cell-derived packaging cell line Phoenix-Eco, using FuGENE 6 transfection reagent according to the manufacturer's instructions (Roche Molecular Biochemicals, Mannheim, Germany). After 48 h, virus-containing supernatant was harvested. Cells were incubated twice with fresh viral supernatant, for 8 h and overnight. The next day, viral supernatant was removed and cells were cultured in fresh medium. Cells were selected 3 days after transduction with 300 µg/ml zeocin (Invitrogen, Carlsbad, CA, USA) when LZRS-IRES-Zeocin/pBR constructs were used or with 10 µg/ml puromycin (Sigma-Aldrich) when pRETRO-SUPER-p53 short hairpin RNA construct was used. Cells transduced with the LZRS-dnCaspase-9-IRES-GFP construct were sorted for GFP expression using a MoFlo high-speed cell sorter (Cytomation, Fort Collins, CO, USA). Cell lines that were compared side-by-side in apoptosis experiments were created in parallel in the same time frame, including empty vector and WT Bid-reconstituted control cell lines. The cell lines were validated by immunoblotting for p53 downregulation and Bid expression as shown.

#### Apoptosis assay

Assessment of the percentage of cells with cleaved caspase-3 was used as a measure for apoptosis. After stimulation, all floating and adherent cells were collected and fixed with 4% paraformaldehyde in phosphate-buffered saline. Subsequently, cells were washed twice with 1% bovine serum albumin in phosphate-buffered saline and once with permeabilizing buffer (phosphate-buffered saline with 0.1% saponin and 0.5% bovine serum albumin). Next, cells were incubated for 20 min with permeabilizing buffer and stained for 1 h with rabbit anti-caspase-3 antibody (1:50; BD Biosciences, Erembodegem, Belgium). Hereafter, cells were washed three times with

permeabilizing buffer and stained for 1 h with AlexaFluor 647-conjugated goat anti-rabbit immunoglobulin (1:100; Molecular Probes, Leiden, The Netherlands). After three more washes with permeabilizing buffer, cells were analyzed by flow cytometry in the FL4 channel.

#### Clonogenic survival assay

Cells were plated at increasing densities in 10-cm dishes (250–12 800 cells per dish). Once attached, cells were exposed to increasing concentrations of etoposide (80–200 ng/ml) or IR (2–10 Gy) and incubated for 14 days. Next, surviving colonies were fixed with 75% MeOH/25% acetic acid and stained with 50% MeOH/10% acetic acid/0.2% Coomassie blue solution. The number of colonies was counted by visual inspection and the relative surviving fraction was determined with the ratio (colonies with dose  $x$  Gy/colonies with dose 0 Gy)  $\times$  1.

#### Immunoblotting

Cells were lysed for 30 min in RIPA lysis buffer (1% Nonidet P-40, 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% deoxycholate, 0.1% SDS, 5 mM EDTA and protease inhibitors) on ice. Cell lysates were subsequently cleared by centrifugation at 13 000 g for 10 min at 4 °C. Of the total cell lysates, samples containing 30 µg total cellular protein were prepared, as determined by BCA assay (Thermo Scientific Pierce, Rockford, IL, USA). Proteins were separated on 4–12% NuPage Bis-Tris gradient gels (Invitrogen) in MES buffer, according to the manufacturer's instructions. Subsequent immunoblotting was performed as described (Tait *et al.*, 2007). Proteins were detected with the following antibodies: sheep anti-p53 (Ab-7) pAb (1:2500; Oncogene Research Products, San Diego, CA, USA), rabbit anti-Bid pAb (1:250; rabbit serum; homemade but available from BD Biosciences), mouse anti-Bcl-2 mAb (1:1000; clone Bcl-2-100, mouse ascites fluid; Sigma-Aldrich), mouse anti-caspase-9 mAb 9508 (1:1000; Cell Signaling Technology, Danvers, MA, USA) and mouse anti-Actin mAb C4, MAB1501R (1:10 000, Chemicon International, Temecula, CA, USA). Second-step antibodies were horseradish peroxidase-conjugated rabbit anti-sheep immunoglobulin (1:7500), swine anti-rabbit immunoglobulin (1:7500) and rabbit anti-mouse immunoglobulin (1:7500, all from Dako A/S, Glostrup, Denmark). The enhanced chemiluminescence kit was from Pierce Biotechnology (Rockford, IL, USA).

#### Conflict of interest

The authors declare no conflict of interest.

#### Acknowledgements

This work was supported by grant NKI 2008-4110 from the Dutch Cancer Society. We thank Dr Atan Gross for MEF cell lines, Rogier Rooswinkel for data analysis, Inge Verbrugge, Bert van de Kooij and Rogier Rooswinkel for discussions and critical reading of the manuscript, and personnel of the flow cytometry facility of the Netherlands Cancer Institute for experimental assistance.

#### References

- Anguissola S, Koehler B, O'Byrne R, Duessmann H, Cannon MD, Murray FE *et al.* (2009). Bid and calpains cooperate to trigger oxaliplatin-induced apoptosis of cervical carcinoma HeLa cells. *Mol Pharmacol* **76**: 998–1010.
- Brown JM, Wouters BG. (1999). Apoptosis, p53, and tumor cell sensitivity to anticancer agents. *Cancer Res* **59**: 1391–1399.
- Chipuk JE, Green DR. (2006). Dissecting p53-dependent apoptosis. *Cell Death Differ* **13**: 994–1002.

- Chou JJ, Li H, Salvesen GS, Yuan J, Wagner G. (1999). Solution structure of BID, an intracellular amplifier of apoptotic signaling. *Cell* **96**: 615–624.
- Cirman T, Oresic K, Mazovec GD, Turk V, Reed JC, Myers RM et al. (2004). Selective disruption of lysosomes in HeLa cells triggers apoptosis mediated by cleavage of Bid by multiple papain-like lysosomal cathepsins. *J Biol Chem* **279**: 3578–3587.
- Erlacher M, Michalak EM, Kelly PN, Labi V, Niederegger H, Coultas L et al. (2005). BH3-only proteins Puma and Bim are rate-limiting for gamma-radiation- and glucocorticoid-induced apoptosis of lymphoid cells *in vivo*. *Blood* **106**: 4131–4138.
- Gross A, Yin XM, Wang K, Wei MC, Jockel J, Milliman C et al. (1999). Caspase cleaved BID targets mitochondria and is required for cytochrome c release, while BCL-XL prevents this release but not tumor necrosis factor-R1/Fas death. *J Biol Chem* **274**: 1156–1163.
- Gross A. (2006). BID as a double agent in cell life and death. *Cell Cycle* **5**: 582–584.
- Guo Y, Srinivasula SM, Druilhe A, Fernandes-Alnemri T, Alnemri ES. (2002). Caspase-2 induces apoptosis by releasing proapoptotic proteins from mitochondria. *J Biol Chem* **277**: 13430–13437.
- Kamer I, Sarig R, Zaltsman Y, Niv H, Oberkovitz G, Regev L et al. (2005). Proapoptotic BID is an ATM effector in the DNA-damage response. *Cell* **122**: 593–603.
- Kaufmann T, Tai L, Ekert PG, Huang DC, Norris F, Lindemann RK et al. (2007). The BH3-only protein bid is dispensable for DNA damage- and replicative stress-induced apoptosis or cell-cycle arrest. *Cell* **129**: 423–433.
- Koehler B, Anguissola S, Concannon CG, Rehm M, Kogel D, Prehn JH. (2008). Bid participates in genotoxic drug-induced apoptosis of HeLa cells and is essential for death receptor ligands' apoptotic and synergistic effects. *PLoS One* **3**: e2844.
- Krajewska M, Zapata JM, Meinhold-Heerlein I, Hedayat H, Monks A, Bettendorf H et al. (2002). Expression of Bcl-2 family member Bid in normal and malignant tissues. *Neoplasia* **4**: 129–140.
- Lam M, Dubyak G, Chen L, Nunez G, Miesfeld RL, Distelhorst CW. (1994). Evidence that BCL-2 represses apoptosis by regulating endoplasmic reticulum-associated Ca<sup>2+</sup> fluxes. *Proc Natl Acad Sci USA* **91**: 6569–6573.
- Letai A, Bassik MC, Walensky LD, Sorcinelli MD, Weiler S, Korsmeyer SJ. (2002). Distinct BH3 domains either sensitize or activate mitochondrial apoptosis, serving as prototype cancer therapeutics. *Cancer Cell* **2**: 183–192.
- Luo X, Budihardjo I, Zou H, Slaughter C, Wang X. (1998). Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell* **94**: 481–490.
- Mandic A, Viktorsson K, Strandberg L, Heiden T, Hansson J, Linder S et al. (2002). Calpain-mediated Bid cleavage and calpain-independent Bak modulation: two separate pathways in cisplatin-induced apoptosis. *Mol Cell Biol* **22**: 3003–3013.
- McDonnell JM, Fushman D, Milliman CL, Korsmeyer SJ, Cowburn D. (1999). Solution structure of the proapoptotic molecule BID: a structural basis for apoptotic agonists and antagonists. *Cell* **96**: 625–634.
- Michalak EM, Villunger A, Adams JM, Strasser A. (2008). In several cell types tumour suppressor p53 induces apoptosis largely via Puma but Noxa can contribute. *Cell Death Differ* **15**: 1019–1029.
- Oberkovitz G, Regev L, Gross A. (2007). Nucleocytoplasmic shuttling of BID is involved in regulating its activities in the DNA-damage response. *Cell Death Differ* **14**: 1628–1634.
- Ploner C, Kofler R, Villunger A. (2008). Noxa: at the tip of the balance between life and death. *Oncogene* **27**(Suppl 1): S84–S92.
- Ren D, Tu H-C, Kim H, Wang GX, Bean GR, Takeuchi O et al. (2010). BID, BIM, and PUMA are essential for activation of the BAX- and BAK-dependent cell death program. *Science* **330**: 1390–1393.
- Sarig R, Zaltsman Y, Marcellus RC, Flavell R, Mak TW, Gross A. (2003). BID-D59A is a potent inducer of apoptosis in primary embryonic fibroblasts. *J Biol Chem* **278**: 10707–10715.
- Sax JK, Fei P, Murphy ME, Bernhard E, Korsmeyer SJ, El Deiry WS. (2002). BID regulation by p53 contributes to chemosensitivity. *Nat Cell Biol* **4**: 842–849.
- Schmitt CA, Rosenthal CT, Lowe SW. (2000). Genetic analysis of chemoresistance in primary murine lymphomas. *Nat Med* **6**: 1029–1035.
- Shelton SN, Shawgo ME, Robertson JD. (2009). Cleavage of Bid by executioner caspases mediates feed forward amplification of mitochondrial outer membrane permeabilization during genotoxic stress-induced apoptosis in Jurkat cells. *J Biol Chem* **284**: 11247–11255.
- Simmen T, Aslan JE, Blagoveshchenskaya AD, Thomas L, Wan L, Xiang Y et al. (2005). PACS-2 controls endoplasmic reticulum-mitochondria communication and Bid-mediated apoptosis. *EMBO J* **24**: 717–729.
- Slee EA, Keogh SA, Martin SJ. (2000). Cleavage of BID during cytotoxic drug and UV radiation-induced apoptosis occurs downstream of the point of Bcl-2 action and is catalysed by caspase-3: a potential feedback loop for amplification of apoptosis-associated mitochondrial cytochrome c release. *Cell Death Differ* **7**: 556–565.
- Strasser A, Harris AW, Jacks T, Cory S. (1994). DNA damage can induce apoptosis in proliferating lymphoid cells via p53-independent mechanisms inhibitable by Bcl-2. *Cell* **79**: 329–339.
- Tait SW, de Vries E, Maas C, Keller AM, D'Santos CS, Borst J. (2007). Apoptosis induction by Bid requires unconventional ubiquitination and degradation of its N-terminal fragment. *J Cell Biol* **179**: 1453–1466.
- Tan KO, Tan KM, Yu VC. (1999). A novel BH3-like domain in BID is required for intramolecular interaction and autoinhibition of pro-apoptotic activity. *J Biol Chem* **274**: 23687–23690.
- Valentijn AJ, Gilmore AP. (2004). Translocation of full-length Bid to mitochondria during anoikis. *J Biol Chem* **279**: 32848–32857.
- Villunger A, Michalak EM, Coultas L, Mullauer F, Bock G, Auserlechner MJ et al. (2003). p53- and drug-induced apoptotic responses mediated by BH3-only proteins puma and noxa. *Science* **302**: 1036–1038.
- Wang X. (2001). The expanding role of mitochondria in apoptosis. *Genes Dev* **15**: 2922–2933.
- Ward MW, Rehm M, Duessmann H, Kacmar S, Concannon CG, Prehn JH. (2006). Real time single cell analysis of Bid cleavage and Bid translocation during caspase-dependent and neuronal caspase-independent apoptosis. *J Biol Chem* **281**: 5837–5844.
- Werner AB, Tait SW, de Vries E, Eldering E, Borst J. (2004). Requirement for aspartate-cleaved bid in apoptosis signaling by DNA-damaging anti-cancer regimens. *J Biol Chem* **279**: 28771–28780.
- Willis SN, Fletcher JI, Kaufmann T, van Delft MF, Chen L, Czabotar PE et al. (2007). Apoptosis initiated when BH3 ligands engage multiple Bcl-2 homologs, not Bax or Bak. *Science* **315**: 856–859.
- Yu J, Zhang L, Hwang PM, Kinzler KW, Vogelstein B. (2001). PUMA induces the rapid apoptosis of colorectal cancer cells. *Mol Cell* **7**: 673–682.
- Zha J, Weiler S, Oh KJ, Wei MC, Korsmeyer SJ. (2000). Post-translational N-myristoylation of BID as a molecular switch for targeting mitochondria and apoptosis. *Science* **290**: 1761–1765.
- Zinkel SS, Hurov KE, Gross A. (2007). Bid plays a role in the DNA damage response. *Cell* **130**: 9–10.
- Zinkel SS, Hurov KE, Ong C, Abtahi FM, Gross A, Korsmeyer SJ. (2005). A role for proapoptotic BID in the DNA-damage response. *Cell* **122**: 579–591.
- Zinkel SS, Ong CC, Ferguson DO, Iwasaki H, Akashi K, Bronson RT et al. (2003). Proapoptotic BID is required for myeloid homeostasis and tumor suppression. *Genes Dev* **17**: 229–239.



This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>