

CORRIGENDUM

RNA-binding motif protein 47 inhibits Nrf2 activity to suppress tumor growth in lung adenocarcinoma

T Sakurai, K Isogaya, S Sakai, M Morikawa, Y Morishita, S Ehata, K Miyazono and D Koinuma

Oncogene (2017) 36, 5083; doi:10.1038/onc.2017.191; published online 12 June 2017

Correction to: *Oncogene* (2016) 35, 5000–5009; doi:10.1038/onc.2016.35; published online 29 February 2016

Since the online publication of this article the authors found that one of the siRNAs used in Figure 2f was not specific for TTF-1/

Nkx2.1. The authors thus wish to replace the figure with the new figure using siRNAs specific for TTF-1/Nkx2.1. The authors apologize for any inconvenience caused, and confirm that the conclusions drawn from this research are not affected by this correction.





Figure 2. (f) H441 cells were transfected with human RBM47 promoter-reporter construct in combination with control or TTF-1 siRNAs (siTTF-1#1 (HSS144278), siTTF-1#3 (HSS144277) and siTTF-1#4 (HSS144279; Thermo Fisher Scientific)). At 48 h after transfection, cells were collected and assayed for luciferase activities. siCT: control siRNA. Averages and standard deviations of the two independent experiments were shown for each condition. The Tukey–Kramer test of the R program (<http://www.r-project.org/>) was used for multiple comparisons of the data. F-test was performed for the equality of variances.

 This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

© The Author(s) 2017