Copyright GPIAG - reproduction prohibited

RM Hopstaken et al.

- Christ-Crain M, Muller B. Biomarkers in respiratory tract infections: diagnostic guides to antibiotic prescription, prognostic markers and mediators. *Eur Respir* J 2007;30(3):556-73. http://dx.doi.org/10.1183/09031936.00166106
- Hopstaken RM, Muris JWM, Knottnerus JA, Kester ADM, Rinkens PELM, Dinant GJ. Contributions of symptoms, signs, erythrocyte sedimentation rate and Creactive protein to a diagnosis of pneumonia in acute lower respiratory tract infection. Br J Gen Pract 2003;53:358-64.
- Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. N Engl J Med 1999;340(6):448-54. http://dx.doi.org/10.1056/NEJM 199902113400607
- 8. Gaini S, Koldkjaer OG, Pedersen C, Pedersen SS. Procalcitonin, lipopolysaccharide-binding protein, interleukin-6 and C-reactive protein in community-acquired infections and sepsis: a prospective study. *Crit Care* 2006; **10**(2):R53. http://dx.doi.org/10.1186/cc4866
- Melbye H, Straume B, Aasebo U, Brox J. The diagnosis of adult pneumonia in general practice. The diagnostic value of history, physical examination and some blood tests. Scand J Prim Health Care 1988;6(2):111-17. http://dx.doi.org/10.3109/02813438809009300
- Zweigner J, Schumann RR, Weber JR. The role of lipopolysaccharide-binding protein in modulating the innate immune response. *Microbes Infect* 2006;8(3):946-52. http://dx.doi.org/10.1016/j.micinf.2005.10.006
- Dahler Eriksen BS, Lauritzen T, Lassen JF, Lund ED, Brandslund I. Near-patient test for C-reactive protein in general practice: assessment of clinical, organizational, and economic outcomes. Clin Chem 1999;45(4):478-85.
- Seamark DA, Backhouse SN, Powell R. Field-testing and validation in a primary care setting of a point-of-care test for C-reactive protein. *Ann Clin Biochem* 2003;40:178-80. http://dx.doi.org/10.1258/000456303763046139

Available online at http://www.thepcrj.org

OBITUARY

Dr Ian Gregg 1925-2009

lan Gregg was well known for his pioneering work in asthma, and its treatment and monitoring in general practice and by the patient at home. He was particularly instrumental in getting the peak flow meter accepted as a valuable indicator of airway resistance change before and during an asthma attack. Perhaps his best known work was in establishing normal peak flow values (Nunn & Gregg, *Br Med J* 1989;**298**:1068-70).

lan was educated at Westminster, and after leaving in 1943 joined the Army. He spent much time in India, which he loved, and became fluent in Urdu. On demobilisation he went up to Wadham College, Oxford, and then qualified at Westminster Hospital Medical School in 1954. House jobs followed at the Westminster, Brompton, and St Stephen's Hospitals.

lan's ambitions were always in general practice, and from 1958 to 1982 he was a principal at practices in Roehampton and later at Kingston-upon-Thames. He continued research as a clinical assistant at the Westminster. In 1962 lan observed the increase in airway resistance in many asthmatic patients without symptoms, and in 1964 published his classic paper on the use of the Wright peak flow meter in general practice (*J Coll Gen Pract* 1964;**7**(2):199-214)

From 1982 until 1987 he was senior lecturer at Southampton University. He travelled widely, lecturing, presenting research, helping and advising industry, and making many friends.

Ian Gregg died on 26 April 2009

John P McNaughton

* Corresponding author: John P McNaughton, Director, Fyne Dynamics Ltd, 1 Horsecroft Place, Harlow, Essex CM19 5BT. Tel: +44(0)1279 423423 Fax: +44(0)1279 454373 E-mail: www.fyne-dynamics.com