

## Neonatal Polycythemia and Hyperviscosity Syndrome

VIRGINIA D. BLACK

*Departments of Pediatrics, Wayne State University, Hutzel Hospital and Children's Hospital of Michigan, Detroit, Michigan U.S.A.*

Infant hematocrit varies with both gestational age and post-natal age. It has been assumed that the rise in hematocrit which occurs from first to third trimester is an adaptation to increase oxygen needs and the relative intrauterine hypoxia. Three decades ago newborns with markedly elevated hematocrits and serious cardiorespiratory distress were described. Often the symptomatic infant was the recipient of a twin-to-twin transfusion. These early case reports are the first documented cases of what is now described as neonatal polycythemia.

In 1966, Baum (1) identified an elevated viscosity in blood from infants who were polycythemic. This elevation in viscosity was believed to decrease peripheral blood flow and hence might explain the pathology associated with neonatal polycythemia. A new condition, neonatal hyperviscosity, was described.

It is common but inaccurate to equate neonatal polycythemia and neonatal hyperviscosity. The definitions of each are specific and, although overlapping, do not exactly define the same population.

Neonatal polycythemia is most commonly described as a venous hematocrit of 65%. Levels as low as 60% or as high as 70% have also been suggested. Variations in sample site or postnatal age compound the difficulty in defining neonatal polycythemia. Linderkamp (3) and Gattl (2) have demonstrated the variability between capillary and venous hematocrit values. Shohat (4) was able to demonstrate dynamic changes in both venous hematocrit and viscosity measurements. By two hours of age there is a linear correlation between cord blood hematocrit and venous hematocrit. Thus, cord blood hematocrit may be a reliable technique for initial screening. However, this technique may miss infants who become polycythemic as a result of acute placental transfusion.

Neonatal hyperviscosity refers to a characteristic of blood which increases with hematocrit. Several other factors also contribute to the viscosity of whole blood. Among these are large plasma proteins and stiffness of the red blood cell membrane. Newborn plasma has a lower viscosity compared to adult plasma since the concentration of plasma proteins is lower in the newborn. Standards for viscosity measurements are available from several authors. Hyperviscosity has been defined as a value more than two standard deviations from mean cord viscosity. Whole blood viscosity is dependent on the shear rate studied. Viscosity measurements are therefore given over several shear rates. Use of the viscosity standards available suggest that 2-6% of newborn infants have hyperviscosity. While hematocrit and viscosity are closely correlated the contribution of other factors to viscosity make it impossible to equate the two conditions. At the higher hematocrit levels all polycythemic infants are hyperviscous. Some infants without polycythemia have abnormal whole blood viscosity. Some polycythemic infants are hyperviscous. In the remainder of this review the clinical signs and outcome associated with the two conditions will be discussed.

Infants with polycythemia and hyperviscosity often present with cardiorespiratory signs which mimic cyanotic congenital heart disease. Tachypnea, tachycardia, cyanosis, and apnea have all been described as well as pulmonary hypertension.

Central nervous system signs are common including the non specific findings of irritability, rapid state changes, vasomotor instability, lethargy, poor tone, and poor feeding. Early reports of polycythemic infants frequently reported seizures. Prospective screening of infants has identified few with neonatal seizures. Obvious central nervous system hemorrhage is also uncommon.

More recently there has become increasing evidence that polycythemia/hyperviscosity is a frequent cause of necrotizing enterocolitis (NEC) when it appears in the term infant. Polycythemic infants undergoing partial plasma exchange with fresh frozen plasma seem particularly vulnerable to developing a spectrum of gastrointestinal injuries. Pneumatosis intestinalis has also been seen in many of these infants with a range of gastrointestinal signs.

Renal failure, thrombocytopenia, and various other hematologic findings have also been reported. Hypoglycemia is commonly associated with polycythemia and may be of greater magnitude than appreciated since the determination of plasma glucose overestimates whole blood glucose. Peripheral blood flow may be severely compromised and has improved following plasmapheresis.

Outcome following neonatal polycythemia/hyperviscosity is largely dependent upon the method used to identify the subjects. Seriously compromised infants as originally described are not the rule. Most infants have symptoms that are less obvious. The most symptomatic infants in early studies had developmental delays and neurologic handicaps. Prospective screening identifies a group of infants for whom the outcome remains controversial. Several authors have failed to show significant impairment at 1-6 years. Our studies suggest that there are clear differences between polycythemic infants and a comparison group at two years of age.

Partial plasma exchange appears to improve sensory outcome. All studies of prospectively identified polycythemic infants suggest that mental development is not significantly affected.

### REFERENCES

1. Baum RS, 1966, Viscous forces in neonatal polycythemia. *J Pediatr* 69: 975
2. Gattl RA, 1967, Hematocrit values for capillary blood in the newborn infant. *J Pediatr* 70: 117
3. Linderkamp O, Versmold HT, Strohhacker I, et al, 1977, Capillary-venous hematocrit differences in newborn infants. *Eur J Pediatr* 127: 9
4. Shohat M, Merlob P, Reisner SH, 1984, Neonatal polycythemia: I. Early diagnosis and incidence relating to time of sampling. *Pediatrics* 73: 7

## Oxygen Radicals and Tissue Damage in the Perinatal Period

OLA D. SAUGSTAD

*Department of Pediatrics, The National Hospital, Oslo, Norway*

Posthypoxic tissue damage could be caused by oxygen free radicals generated by hypoxanthine (Hx)-xanthine oxidase (XO) system. Hx which is accumulated in hypoxia is washed out into the circulation in the posthypoxic reperfusion period. If XO is present the superoxide radical,  $O_2^-$ , is formed:  $Hx + O_2 \xrightarrow{XO} \text{Urate} + O_2^-$ .

We have suggested that this mechanism is responsible, at least partly, for several conditions in perinatal medicine such as bronchopulmonary dysplasia, necrotizing enterocolitis and retrolental fibroplasia. If this hypothesis is correct we are dealing with a common pathogenetic mechanism leading to one disease affecting different organs.

To explore this hypothesis we have mainly studied the lung. In summary we have found: 1) The combination of 100 percent oxygen and intravenous oxygen given 48 h damage the rat lung. The typical changes found are hemorrhage and edema. Furthermore, in spite of a normal lung surfactant profile the surfactant function was abolished indicating that surfactant was inactivated either by peroxidation or by the induction of surfactant inhibitor. 2) XO applied into the airways of guinea pigs acutely induced lung hemorrhage and edema. The lung compliance was dramatically reduced. One single application of XO induced long term effects on the lungs as well. After 2 days thickening of alveolar septae and proliferation of type 2 cells were observed. Furthermore, there was an increased number of alveolar macrophages. 7 and 14 days afterwards the histological changes seemed to have normalized, lung compliance, however, was still significantly lowered. 3) Superoxide dismutase given together with XO protected against XO.

The experimental data show that the hypoxanthine-xanthine oxidase system potentially destroys the lungs under experimental conditions.

## Red Cell Aggregation in the Neonate

MICHAEL RAMPLING

*Department of Physiology and Biophysics, St. Mary's Hospital Medical School, London W2, UK*

The viscometric characteristics of neonatal blood show distinct differences from those of adults (1,2); in particular it is found that at a given haematocrit neonatal blood generally exhibits lower viscosity at all shear rates. In part this is due to the lower plasma viscosity that is also found in the neonate. The effect of this plasma difference can be removed by calculating relative viscosity, and when this is done the high shear rate ( $>5s^{-1}$ ) values of the two types of blood become similar but the low shear rate difference persists. Since it is usually accepted that the elevated low shear rate viscosity of adult blood is due to rouleaux formation, it is to be expected that the latter will be reduced in the neonate and this has been confirmed experimentally.

The cause of the reduced aggregation must lie in the red cell or in the plasma or both. However plasma exchange experiments indicate that adult and neonatal cells aggregate similarly when suspended in adult plasma, while both show reduced aggregation in neonatal plasma (2). Of the aggregating agents in adult blood fibrinogen is by far the most important, so experiments have been performed in which both cell types were suspended in solutions of purified adult fibrinogen. Again they showed similar degrees of rouleaux formation. The conclusion is therefore that the reduced aggregation found in neonatal blood is not due to cellular differences, but rather to variations in the composition of the plasma.