

**CORRECTION**

Correction: Dose-Related Effects of Adjunctive Ketamine in Taiwanese Patients with Treatment-Resistant Depression

Tung-Ping Su^{1,2,3,4}, Mu-Hong Chen^{2,4}, Cheng-Ta Li^{1,2,4}, Wei-Chen Lin^{2,4}, Chen-Jee Hong^{1,2,4}, Ralitzia Gueorguieva^{5,6}, Pei-Chi Tu^{1,2,3}, Ya-Mei Bai^{1,2}, Chih-Ming Cheng² and John H. Krystal^{6,7,8,9}

Neuropsychopharmacology (2019) 44:655–656; <https://doi.org/10.1038/s41386-018-0253-z>

Correction to: *Neuropsychopharmacology* (2017), <https://doi.org/10.1038/npp.2017.94>

Following the publication of this article the authors noted an error in Fig. 5. The units of blood concentration for ketamine and

norketamine should be ng/mL, as per the assay details in Materials and methods section, and not nM/ml as presented in the figure. A revised Fig. 5 is in the correction article. The authors apologize for any inconvenience caused.

¹Division of Psychiatry, Faculty of Medicine, National Yang-Ming University, Taipei, Taiwan; ²Department of Psychiatry, Taipei Veterans General Hospital, Taipei, Taiwan; ³Department of Medical Research, Taipei Veterans General Hospital, Taipei, Taiwan; ⁴Institute of Brain Science, National Yang-Ming University, Taipei, Taiwan; ⁵Department of Biostatistics, School of Public Health, Yale University, New Haven, CT, USA; ⁶Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA; ⁷Department of Neuroscience, Yale University School of Medicine, New Haven, CT, USA; ⁸Yale-New Haven Hospital, New Haven, CT, USA and ⁹VA National Center for PTSD, VA Connecticut Healthcare System, West Haven, CT, USA

Correspondence: T-P. Su (tomsu0402@gmail.com) or John H. Krystal (john.krystal@yale.edu)




Fig. 5