



## Correction: Increasing the bactofection capacity of a mammalian expression vector by removal of the f1 *ori*

Síle A. Johnson<sup>1</sup> · Michael J. Ormsby<sup>1</sup> · Anne McIntosh<sup>1</sup> · Stephen W. G. Tait<sup>2</sup> · Karen Blyth<sup>2</sup> · Daniel M. Wall<sup>1</sup>

Published online: 19 September 2018  
© Springer Nature America, Inc. 2018

Correction to: *Cancer Gene Therapy*; <https://doi.org/10.1038/s41417-018-0039-9>; published online: 13 August 2018

This Article was originally published with one of the panels in Fig. 5a inserted twice (SL-pEGFP). In Fig. 5b

there was also a typo. SL-LacZ should have read SL-pEGFP(-f1). Both 5a and 5b are corrected in both the PDF and HTML versions of the Article.

---

✉ Daniel M. Wall  
Donal.Wall@glasgow.ac.uk

<sup>1</sup> Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary and Life Sciences, Sir Graeme Davies Building, University of Glasgow, Glasgow G12 8TA, UK

<sup>2</sup> Cancer Research UK Beatson Institute, University of Glasgow, Garscube Estate, Switchback Road, Glasgow G61 1BD, UK