



Author Correction: Is antihypertensive treatment based on home blood pressure recommended rather than that based on office blood pressure in adults with essential hypertension? (meta-analysis)

Michihiro Satoh¹ · Toshiki Maeda² · Satoshi Hoshide³ · Takayoshi Ohkubo⁴

Published online: 10 June 2019
© The Japanese Society of Hypertension 2019

Correction to: Hypertension Research

<https://doi.org/10.1038/s41440-019-0221-y>
published online 5 April 2019

The original version of the Supplementary Information associated with this Article contained errors in Supplementary Figs. 1 and 2. The corrected figures are shown below.

Supplementary Fig. 1. Flow diagram of the process of selection of the studies to be included in the meta-analysis for cardiovascular outcome.

Supplementary Fig. 2. Flow diagram of the process of selection of the studies to be included in the meta-analysis for BP change outcome.

These authors contributed equally: Michihiro Satoh, Toshiki Maeda

The original article can be found online at <https://doi.org/10.1038/s41440-019-0221-y>.

Supplementary information The online version of this article (<https://doi.org/10.1038/s41440-019-0280-0>) contains supplementary material, which is available to authorized users.

✉ Takayoshi Ohkubo
tohkubo@med.teikyo-u.ac.jp

¹ Division of Public Health, Hygiene and Epidemiology, Faculty of Medicine, Tohoku Medical and Pharmaceutical University, Sendai, Japan

² Department of Preventive Medicine and Public Health, Faculty of Medicine, Fukuoka University, Fukuoka, Japan

³ Division of Cardiovascular Medicine, Department of Medicine, Jichi Medical University School of Medicine, Shimotsuke, Japan

⁴ Department of Hygiene and Public Health, Teikyo University School of Medicine, Tokyo, Japan




Fig. 1




Fig. 2