














## Author Correction: High oscillator strength interlayer excitons in two-dimensional heterostructures for mid-infrared photodetection

Steven Lukman, Lu Ding , Lei Xu, Ye Tao, Anders C. Riis-Jensen, Gang Zhang, Qing Yang Steve Wu , Ming Yang , Sheng Luo , Chuanghan Hsu, Liangzi Yao , Gengchiao Liang , Hsin Lin , Yong-Wei Zhang, Kristian S. Thygesen , Qi Jie Wang , Yuanping Feng  and Jinghua Teng 

Correction to: *Nature Nanotechnology* <https://doi.org/10.1038/s41565-020-0717-2>, published online 29 June 2020.

In the version of this Article originally published, in the sentence beginning “We also found that the PL intensity...”, the values of the laser fluence (in  $\text{W } \mu\text{m}^{-2}$ ) were incorrect: ‘ $8 \times 10^4$ ’ and ‘ $5 \times 10^5$ ’ should have been ‘ $8 \times 10^{-4}$ ’ and ‘ $5 \times 10^{-5}$ ’, respectively. Furthermore, in the units of the  $x$ -axis labels of Extended Data Fig. 4 and Supplementary Fig. 6, ‘ $10^5$ ’ should have been ‘ $10^{-5}$ ’. The online versions of the Article have been amended and the Supplementary Information file has been replaced.

Published online: 5 February 2021

<https://doi.org/10.1038/s41565-021-00865-9>

© The Author(s), under exclusive licence to Springer Nature Limited 2021