SCIENTIFIC REPORTS

Published online: 12 April 2019

OPEN Author Correction: Reconstruction of cysteine biosynthesis using engineered cysteine-free enzymes

Kosuke Fujishima^{1,2,3}, Kendrick M. Wang⁴, Jesse A. Palmer², Nozomi Abe³, Kenji Nakahigashi^{3,5}, Drew Endy⁴ & Lynn J. Rothschild⁶

Correction to: Scientific Reports https://doi.org/10.1038/s41598-018-19920-y, published online 29 January 2018

This Article contains an error in the order of the figure legends. The legends for Figures 4 and 5 were incorrectly given as those for 5 and 4 respectively.

The legend for Figure 4 should read:

"Growth curve of re-transformed auxotrophic E. coli strains in LB and M9+ glucose media. Each panel represents growth of cysteine-dependent E. coli auxotrophs rescued by wild type enzymes: CysE or CysM (blue), cysteine-free enzymes: CysE-C or CysM-C (orange), cysteine- and methionine-free enzymes: CysE-CM or CysM-CM (yellow), and LacZ α protein expressed from original pUC19 plasmid (gray). Growth curve was monitored every 10 minutes at OD 600 nm using 96-well plate reader. Standard deviations of the growth curves are displayed as calculated from triplicates."

The legend for Figure 5 should read:

"Synthetic cysE and cysM gene transformants display recovery of CysE function without cysteine and methionine and CysM function without cysteine. (A) E. coli $\Delta cysE$ competent cells were transformed with positive control cysE, two cysE variants cysE-C/cysE-CM cloned into the multiple cloning site of pUC19 plasmid, and original pUC19 encoding N-terminal fragment of $lacZ\alpha$ as a negative control. (B) E. coli $\Delta cysM\Delta cysK$ competent cells transformed with positive control cysM, two cysM variants cysM-C/cysM-CM in pUC19 plasmid, and original pUC19 encoding N-terminal fragment of $lacZ\alpha$ as a negative control. Cells were plated on M9 + glucose medium with 0.4 mM IPTG, 50 µg/ml kanamycin, and 100 µg/ml ampicillin and incubated at 30 °C for 72 h."

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2019

¹Earth-Life Science Institute, Tokyo Institute of Technology, Tokyo, 1528550, Japan. ²Universities Space Research Association, NASA Ames Research Center, Moffett Field, California, 94035, USA. ³Institute for Advanced Biosciences, Keio University, Tsuruoka, 9970035, Japan. ⁴Stanford University Department of Bioengineering, Stanford, California, 94305, USA. ⁵Spiber Inc. 234-1 Mizukami, Kakuganji, Tsuruoka, 9970052, Japan. ⁶NASA Ames Research Center, Moffett Field, California, 94035, USA. Kosuke Fujishima and Kendrick M. Wang contributed equally. Correspondence and requests for materials should be addressed to L.J.R. (email: Lynn.J.Rothschild@nasa.gov)