

CORRESPONDENCE

Proximal tibia fracture in a patient with incomplete spinal cord injury associated with robotic treadmill training

Spinal Cord Series and Cases (2016) 2, 16010; doi:10.1038/scsandc.2016.10; published online 7 July 2016

Éimear Smith
*Spinal Cord System of Care, National Rehabilitation Hospital,
Dublin, Ireland*
E-mail: Eimear.Smith@nrh.ie

Filippo *et al.*,¹ report on the T-scores of a 43-year-old female, who sustained a fragility fracture, while using the robotic treadmill, Lokomat (Hocoma, Volketswil, Switzerland). The patient's menopausal status was not outlined. However, at 43 years of age, she is likely to have been pre-menopausal. Is it not widely accepted that for pre-menopausal females and males under 60 years of age z-scores are used in determining fracture risk?^{2,3} Unless, there is uniformity in the reporting of bone mineral density, development of protocols as advocated by the authors, will remain a significant challenge.

COMPETING INTERESTS

The author declares no conflict of interest.

REFERENCES

- 1 Filippo TRM, De Carvalho MCL, Carvalho LB, de Souza DR, Imamura M, Battistella LR. Proximal tibia fracture in a patient with incomplete spinal cord injury associated with robotic treadmill training. *Spinal Cord* 2015; **53**: 875–876.
- 2 International Society for Clinical Densitometry, Official Positions, Adult and Paediatric 2015. www.iscd.org.
- 3 Craven BC, Robertson LA, McGillivray CF, Adachi JD. Detection and treatment of sub-lesional osteoporosis among patients with chronic spinal for injury: proposed paradigms. *Top Spinal Cord Inj Rehabil* 2009; **14**: 1–22.