by encouraging the use of WSWM, but it is another instruction that is given, with minimal input from the literature to provide support for the use of WSWM.

SUMMARY

As mentioned briefly in the first of these articles, the delivery of the advice, instructions and protocols that are provided to patients is as important as the content of what is delivered. Only this holistic approach can ensure that the requirements of consent within the profession are met,⁴³ and patients are given the information they want and need.

As clinicians there is a responsibility to not only realise and be aware of what should be advised/not advised, what instructions to provide and which protocol to follow, but there must also be recognition of where this information has been derived and its relative reliability. A sound knowledge of this information, coupled with a thorough and meticulous history and examination, will aid clinical decision-making, enable appropriate consent to be gained and help support patients who have questions or wish to request information in relation to their healthcare.

- Carrotte P. Endodontic problems. Br Dent J 2005; 198: 127–133.
- Nair P N. On the causes of persistent apical periodontitis: a review. *Int Endod J* 2006; 39: 249–281.
- Chong B S, Rhodes J S. Endodontic surgery. *Br Dent J* 2014; 216: 281–290.
- Eliyas S, Vere J, Ali Z, Harris I. Micro-surgical endodontics. Br Dent J 2014; 216: 169–177.
- Carrotte P. Surgical endodontics. Br Dent J 2005; 198: 71–79.
- Del Fabbro M, Taschieri S, Testori T, Francetti L, Weinstein R L. Surgical vs. non-surgical endodontic re-treatment for periradicular lesions. Cochrane Database Syst Rev 2007; 18: CD005511.
- Chong B S, Pitt Ford T R, Hudson M B. A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root end filling materials in endodontic surgery. Int Endod J 2003; 36: 520–526.
- Saunders W P. A prospective clinical study of periradicular surgery using Mineral Trioxide Aggregate as a root-end filling. J Endod 2008; 34: 660–665.

- Tsesis I, Faivishevsky, Kfir A, Rosen E. Outcome of surgical endodontic treatment performed by a modern technique: a meta-analysis of the literature. J Endod 2009; 35: 1505–1511.
- Setzer F C, Shah S B, Kohli M R, Karabucak B, Kim S. Outcome of surgical endodontic treatment performed by a modern technique: a meta-analysis of the literature – Part 1: Comparison of traditional root-end surgery and endodontic microsurgery. J Endod 2010; 36: 1757–1765.
- Mansoor J. Part 2: Before and after the removal of wisdom teeth. Br Dent J 2015; 218: 279-284.
- High A S, Russell J L. Retrograde root filling using antibiotic-containing, radiopaque, bone cement. J Dent 1989; 17: 241–245.
- Anderssen C, Bernhart T, Honlinger M, Mailath G, Wachter H, Fuchs D. Immune activation after apicoectomy: comparison between patients with and without prophylaxis antibiotic therapy. *Pteridines* 2003; 4: 192–194.
- Lindeboom J A, Frenken J W, Valkenburg P, van den Akker HP. The role of preoperative prophylactic antibiotic administration in periapical endodontic surgery: a randomised, prospective double-blind placebocontrolled study. *Int Endod J* 2005; 38: 887–881.
- Mansoor J. Part 1: Minor oral surgery before and after where is the evidence? Br Dent J 2014; 218: 273–278.
- Scottish Dental Clinical Effectiveness Programme. *Drug prescribing for dentistry*, 2nd ed. NHS Scotland, Dundee Dental Education Centre UK, 2011.
- Faculty of Dental Practitioners (FGDP). Antimicrobial prescribing for general dental practitioners, 2nd ed. London: FGDP, 2012.
- Von Arx T, Hanni S, Jensen S S. Clinical and radiographic assessment of various predictors for healing outcome one year after periapical surgery. J Endod 2007; 33: 123–128.
- National Institute of Clinical Excellence (NICE). Prophylaxis against infective endocarditis, 1st ed. London: NICE, 2008.
- Dental and Eye Care Team; Prescribing Support Unit. Prescribing by dentists, 2010. London: National Information Centre, 2011.
- Martin M V, Nind D. Use of chlorhexidine gluconate for pre-operative disinfection of apicectomy sites. Br Dent J 1987; 162: 459–461.
- Tsesis I, Fuss Z, Lin S, Tilinger G, Peled M. Analysis
 of postoperative symptoms following surgical
 endodontic treatment. *Quintessence Int* 2003; 34:
 756–760.
- Pemberton M N, Gibson J. Chlorhexidine and hypersensitivity reactions in dentistry. Br Dent J 2012;
 213: 547–550.
- Garcia B, Larrazabal C, Penarrocha M, Penarrocha M. Pain and swelling in periapical surgery. A literature update. Med Oral Patol Oral Cir Bucal 2008; 13: E726–E728.
- 25. Penarrocha M, Garcia B, Marti E, Balaguer J. Pain

- and inflammation after periapical surgery in 60 patients. *J Oral Maxillofac Surg* 2006; **64:** 429–433.
- Kvist T, Reit C. Postoperative discomfort associated with surgical and nonsurgical endodontic retreatment. Endod Dent Traumatol 2000; 16: 71–74.
- Garcia B, Larranzabal C, Penarrocha M, Penarrocha M. Pain and swelling in periapical surgery: a literature update. *Med Oral Patol Oral Cir Buccal* 2008; 13: E726–729.
- 28. Sortino F, Cicciù M. Strategies used to inhibit swelling following removal of impacted lower third molar. *Dent Res J (Isfahan)* 2011; **8:** 162–171.
- Gutmann J L. Surgical endodontics: post-surgical care. Endod Topics 2005; 11: 196-205.
- Evans G E, Bishop K, Renton T. Guidelines for surgical endodontics, 2nd ed. Royal College of Surgeons (England). Fac Dent Surg 2012: 5–6.
- Chong B S, Pitt Ford T R. Postoperative pain after root-end resection and filling. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2005; 100: 762–766.
- Rhodes J S. Advanced endodontics: clinical retreatment and surgery, 1st ed. Oxfordshire: Taylor & Francis, 2005.
- Löe H, Schiott C R. The effect of mouthrinses and topical application of chlorhexidine on the development of dental plaque and gingivitis in man. J Periodontal Res 1970; 5: 79–83.
- Harrison J, Jurosky K, Wound healing in tissues of the periodontium following periradicular surgery. I. The incisional wound. *J Endod* 1991; 17: 425–435.
- Parirokh M, Asgary S, Eghbal M J. The effect of different suture removal time intervals on surgical wound healing. *Iran Endod J* 2006; 1: 81–86.
- Gutmann J L, Harrison W H. Flap designs and incisions. *In* Gutmann J L, Harrison W H (eds). *Surgical endodontics*. pp 162–175. St. Louis, Missouri: lshijaku EuroAmerica, Inc., 1994.
- Selvig K, Torabinejad M. Wound healing after mucoperiosteal surgery in the cat. *J Endod* 1996; 22: 507–515.
- Wirthlin M R, Hancock E B, Gaugler R W. The healing of atraumatic and traumatic incisions in the gingivae of monkeys. J Periodontol 1984; 55: 103–113.
- Grung B. Healing of gingival mucoperiosteal flaps after marginal incision in apicoectomy procedures. Int J Oral Surg 1973; 2: 20–25.
- World Health Organization. Post-operative pain relief. 2014. Available at: http://www.who.int/ surgery/publications/Postoppain.pdf (accessed 18 May 14).
- 41. National Institute for Clinical Excellence (NICE) technology appraisal guidance number 1. Guidance on the extraction of wisdom teeth. 2000.
- British Association of Oral Surgeons.
 Apicectomy. 2014. Available at: http://www.baos. org.uk/resources/Apicectomy1.pdf (accessed 3 November 2014).
- General Dental Council. Standards for the dental team. 2013.

Corrigendum

Practice articles (BDJ 2015; 218: 273-278, 279-284)

Pre- and postoperative management techniques. Before and after. Part 1: medical morbidities
Pre- and postoperative management techniques. Before and after. Part 2: the removal of third molars

In the above practice articles, the address for author correspondence was incorrect. It should read: ¹School of Dentistry, University of Manchester, Higher Cambridge Street, Manchester, M15 6FH.