

Invasive cervical resorption and the oro-facial cleft patient: a review and case series

A. O'Mahony,¹ C. McNamara,² A. Ireland,³ J. Sandy³ and J. Puryer^{*3}

In brief

Presents a review of invasive cervical resorption.

Presents a retrospective review of over 500 oro-facial cleft affected patients at St James's Hospital, Dublin.

Summarises the features associated with 14 cases diagnosed with Class 4 invasive cervical resorption.

Reinforces the importance of long-term review of oro-facial cleft affected patients.

Introduction Invasive cervical resorption (ICR) has an unknown aetiology, yet it exhibits very aggressive behaviour compared with typical external root resorption, posing a high risk of tooth loss. **Aim** To investigate the number of patients at the Dublin Cleft Prosthodontic Department with an oro-facial cleft who experienced ICR and to identify any possible aetiological factors. **Materials and method** A retrospective investigation of all oro-facial cleft patients treated at the Dublin Cleft Prosthodontic Department, St James's Hospital, Dublin. All patients' clinical and radiological records were reviewed. Patients where tooth loss became inevitable due to Class 4 ICR were analysed. **Results** From 588 oro-facial cleft patients, 14 (2.38%) patients with ICR were identified. Of these eight (57%) were female and six (43%) were male. Mean age at diagnosis was 28 years (range = 16–49 years). Cleft type: six (42.1%) unilateral cleft lip and palate, eight (57.9%) bilateral cleft lip and palate. Seventeen ICR affected teeth in total, with eleven (65%) maxillary central incisors, two (12%) maxillary lateral incisors, four (23%) maxillary canines, and one (7%) central, lateral and canine affected. Some, (N = 10, 71.4%) presented with ICR resulting in immediate tooth loss. Other patients (N = 4, 28.6%) developed ICR during or following prosthodontic treatment at the Cleft Centre. Tooth loss for this cohort, though not immediate, was inevitable. All had undergone fixed orthodontic appliance treatment and twelve had received dento-alveolar bone grafts. A number (N = 7, 50%) had undergone osteotomy, two (14%) had received night guard vital dental whitening and one had a history of trauma. **Conclusions** ICR, given its aggressive nature and ill-understood aetiology, poses significant treatment challenges. The most severe form of ICR (Class 4) leads inevitably to tooth loss. The slow–moderate progression of ICR may explain the late presentation found in this study, reinforcing the importance of long-term follow-up of this special dental care group.

Introduction

Oro-facial clefting is the most common congenital malformation of the head and neck.¹ Dental anomalies in the anterior and pre-maxillary area are long-recognised in oro-facial clefting, and these include abnormalities in tooth number, size, form, development, position and eruption.^{2–6} Dental

crown anomalies include enamel hypoplasia, microdontia, macrodontia and dilacerations, whilst root malformations may range from delayed development to dilaceration, interrupted formation or microform giving rise to unfavourable crown root ratios.^{3,8} These defects may be attributed to the cleft itself, to genetic factors or to traumatic factors relating to surgery and resultant scar tissue formation.^{2,7}

Invasive cervical resorption (ICR) is a relatively uncommon and often aggressive form of external root resorption that can affect any tooth in the permanent dentition.⁹ ICR is characterised by aggressive invasion of the cervical region of the root by fibro-vascular tissue, which progressively resorbs dentine enamel and cementum leading to progressive destruction of the tooth structure at the zone of connective tissue attachment.^{10–12}

The clinical appearance of teeth affected by ICR varies considerably depending upon the extent of the resorptive process. Some teeth may display an obvious 'pinkish' colour in the tooth crown as the highly vascular resorptive tissue becomes visible through the thin residual enamel,⁹ although more frequently, the condition is only detected radiographically,¹³ often as an incidental finding. Histologically, the pulp remains protected by a thin layer of pre-dentine until late in the process. More advanced lesions may display fibro-osseous characteristics with deposition of ectopic bone-like calcifications, both within the resorbing tissues and directly on the dentine surface.¹⁴

The aetiology of ICR is unknown.^{9,14} Physical or chemical trauma to the cemento-enamel junction region is considered a significant

¹Cleft Pros/Ortho Unit, St James' Hospital, Dublin 8, Ireland;
²HSE Regional Orthodontic Department, St James' Hospital, Dublin 8, Ireland; ³School of Oral and Dental Sciences, Bristol Dental Hospital, Lower Maudlin Street, Bristol, BS1 2LY, United Kingdom

*Correspondence to: Dr James Puryer
Email: james.puryer@bristol.ac.uk

Refereed Paper. Accepted 30 January 2017
DOI: 10.1038/sj.bdj.2017.405

predisposing factor.^{9,14,15} Trauma may be direct or associated with orthodontic tooth movement, periodontal or dentoalveolar surgery, tooth transplantation, bone grafting, segmental maxillary orthognathic surgery, intra-coronal bleaching and tetracycline conditioning of roots.^{10,11,15} Additionally, ICR has been found in association with heat damage to bone or where there has been impairment of blood supply at the cemento-enamel junction.¹⁵ Systemic factors such as viral infections, hormonal abnormalities and renal pathology have also been reported in association with ICR,¹⁵ and there is familial predisposition.¹⁶

It is important to differentiate ICR from internal resorption, and a diagnosis is normally achieved by both clinical and radiographic examination. Long cone periapical views using a parallax technique can be taken in order to follow the outline and continuity of the pulp chamber. The lesion of ICR will appear to move on the films with the X-ray tube angulation.¹⁷ The use of cone beam computed tomography has been advocated¹⁸ as in addition to detecting the presence of a resorptive defect, it will show its extension in three dimensions and whether or not it has invaded the pulp. Treatment of ICR is difficult, and the outcome is often uncertain.^{9,19–21} The main aim of treatment is to completely remove the resorptive tissue with an excavator or slow-speed bur, followed by conditioning of the dentinal walls to remove any remnants of resorptive tissue before restoration. There appears to be no consensus as to the most appropriate restorative material, with glass-ionomer cement, composite resin, amalgam, mineral trioxide aggregate and calcium-enriched mixture cement all having been advocated.^{22–25} To facilitate treatment planning, Heithersay⁹ divided ICR into four distinct clinical types based upon the severity of the lesion (Table 1). The poor survival rate of Class 4 lesions indicate that while the affected tooth may be left to progressively resorb, its extraction and prosthodontic replacement is inevitable.

Approximately 100 oro-facial cleft affected children are born in Ireland annually. These children require comprehensive and complex multidisciplinary care from early infancy through to early adulthood. In addition to a range of essential medical specialties and therapies, dental specialties include paediatric dentistry, orthodontics, maxillofacial surgery and advanced restorative-prosthodontic care. These have significant time and cost implications for the patient and the health system provider. In 2004, an Advanced Restorative Clinic was established at the Cleft Centre, St James's Hospital, Dublin, and since then, care has been provided for over 500 patients.

Aim

The aim of this study was to investigate the number of oro-facial patients at the Dublin Cleft Prosthodontic Department who experienced ICR and to identify any possible aetiological factors.

Method

A retrospective investigation was carried out on all patients treated at the oro-facial Cleft Clinic of the Maxillofacial Department, St James's Hospital, Dublin. All patients treated since the clinic opened in 2004 had their clinical and radiographic records reviewed. The records of ICR patients where tooth loss became inevitable underwent a detailed analysis, and the following parameters were recorded:

- Condition of crown (including enamel hypoplasia and size and form of crown)
- Condition of root (including size, dilaceration, pre-existing external root resorption, vitality and apical form as determined radiologically)
- Gender and ethnicity
- Cleft type, cleft side and if syndromic or non-syndromic
- Surgical history (including bone grafting, implant insertion, fistula repair,

osteotomies, repeat or additional surgery [such as late lip or nasal reconstruction])

- Dental history (including previous orthodontic treatment, crown whitening, restorative or prosthodontic procedures)
- Trauma history (independent of any dental intervention)
- Age at identification and treatment duration before tooth loss became inevitable where ICR developed during prosthodontic operative care.

Results

The records of 588 cleft lip and palate (CLP) patients were reviewed and 14 (2.38%) patients were identified with ICR, of which 8 (53%) were female and 6 (47%) were male. A summary of results is shown in Table 2. All patients were Caucasian and all were non-syndromic CLP. The age range of patients was 16–49 years (mean = 28 years). Cleft types of these patients were identified as six unilateral CLP (three right-sided and three left-sided) and eight bilateral CLP. There were 17 ICR affected teeth in total, and these were all found in the upper arch, of which 11 (65%) were maxillary central incisors, two (12%) were maxillary lateral incisors, and four (23%) were maxillary canines. The majority, (12 [86%]) of patients had one ICR affected tooth, whilst one had two (7%) and another (7%) had three (7%) ICR affected teeth. Bilateral CLP was present in both of the cases with more than one ICR affected teeth. In the single patient with three ICR affected teeth, the maxillary central, lateral and canine on the same side were affected. In all of the patients with unilateral CLP, the ICR affected tooth was on the same side as the cleft. A developmental crown anomaly was found in 12 (85%) patients, and a developmental root anomaly was found in nine (64%) patients. Over half (N = 8, 57%) of patients had a combined crown-root anomaly.

All of the patients had received orthodontic treatment, and two (14%) had received vital night guard dental whitening. Only one patient gave a history of dental trauma. This male patient presented with a non-vital maxillary central incisor prior to the development of ICR. The majority (N = 13, 93%) of patients had undergone surgical bone grafting, seven (50%) had received a maxillary advancement osteotomy, six (43%) had received a surgical fistula repair and four (29%) had received additional surgery for late lip and/or nasal reconstruction. Most (N = 10, 71%) of the

Table 1 Heithersay's classification of ICR⁹

| Class | Description |
|-------|---|
| 1 | A small invasive resorptive lesion near the cervical area with shallow penetration into dentine |
| 2 | A well-defined invasive resorptive lesion that has penetrated close to the coronal pulp chamber but shows little or no extension into the radicular dentine |
| 3 | A deeper invasion of dentine by resorbing tissue not only involving the coronal dentine but also extending into the coronal third of the root |
| 4 | A large invasive resorptive process that has extended beyond the coronal third of the root |

patients presented to the prosthodontic clinic with an existing ICR affected tooth, and all of these patients were symptomless (Figs 1–6). Four patients (29%) developed ICR during or following attendance to the prosthodontic

clinic. These were all symptomless. One of these patients developed ICR immediately after placement of a minimally-prepared resin bonded bridge, with loss of the tooth within a 15 month period (Fig. 4). Another patient

lost two teeth, a maxillary central incisor and maxillary canine, one year and two years respectively following temporary crown preparation. The remaining two patients, who had received prosthodontic care, developed ICR following discharge, losing the affected teeth five years (Fig. 5) and six years (Fig. 6) later.

Discussion

Preserving the integrity of both hard and soft dental tissues is central to oro-facial cleft care. Healthy dental tissues not only benefit and assist the various necessary operative procedures, but are vital to long-term oral health, function, facial aesthetics and quality of life outcomes. It is recognised that surgery,




Fig. 1 (a & b) An example of an ICR affected maxillary canine following extraction; (c & d) An example of an ICR affected maxillary central incisor following extraction

| Table 2 A summary of the 14 patients with Class 4 ICR | | | | | | | | | | | | | | |
|---|-----|----|-----|-----|------------|----|-----|-----|----|----|-------|--------|-------|-------------------|
| Case | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| Gender M:F | F | F | F | M | M | F | M | F | F | M | M | F | M | F |
| Caucasian | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| UCLP Side | ✓ R | – | ✓ R | ✓ L | – | – | ✓ L | ✓ L | | | ✓ R | | | |
| BCLP | – | ✓ | – | – | ✓ | ✓ | – | – | ✓ | ✓ | – | ✓ | ✓ | ✓ |
| Non-syndromic | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| Developmental crown anomaly | ✓ | ✓ | ✓ | – | ✓ | – | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| Developmental root anomaly | ✓ | ✓ | – | ✓ | ✓ | – | ✓ | – | ✓ | ✓ | ✓ | – | – | ✓ |
| Bone graft | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | – | ✓ | ✓ | ✓ | ✓ |
| Osteotomy | – | ✓ | ✓ | ✓ | – | – | – | – | – | ✓ | – | ✓ | ✓ | ✓ |
| Fistula repair | ✓ | ✓ | – | – | – | – | – | – | – | – | ✓ | ✓ | ✓ | ✓ |
| Tooth/teeth ICR affected | 11 | 22 | 13 | 21 | 21, 22, 23 | 13 | 21 | 21 | 21 | 21 | 11 | 21 | 21 | 21, 23 |
| ICR: age at identification | 16 | 21 | 28 | 37 | 27 | 29 | 49 | 26 | 17 | 33 | 29 | 23 | 38 | 23 |
| ICR presentation immediate tooth loss | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | – | ✓ | – | – |
| ICR during/after prosthodontic treatment | – | – | – | – | – | – | – | – | – | – | ✓ | ✓ | ✓ | ✓ |
| Duration to tooth loss | – | – | – | – | – | – | – | – | – | – | 6 yrs | 15 mos | 5 yrs | 21 1 yr, 23 2 yrs |
| Type of prosthodontic treatment | – | – | – | – | – | – | – | – | – | – | *A | *B | *C | *D |
| Orthodontic treatment | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| Tooth whitening | – | – | – | – | – | – | – | – | – | – | – | – | ✓ | ✓ |
| Trauma | – | – | – | – | – | – | – | – | – | – | ✓ | – | – | – |
| Additional surgeries | – | – | – | – | ✓ | – | – | – | – | – | – | ✓ | ✓ | ✓ |
| Implant | – | – | – | – | – | – | – | – | – | – | – | – | ✓ | – |

*A = RCT & bridge preparation *B = minimal preparation resin bonded bridge *C = veneer & implant adjacent *D = crown preparation and temporaries




Fig. 2 (a) Pre-extraction and (b) post-extraction images of a Class 4 ICR affected maxillary canine that suffered general enamel hypoplasia and had been restored with a composite veneer




Fig. 3 (a-c) Case 12 showing pre-oralonasal fistula repair and post-maxillary advancement osteotomy (a) pre-fistula closure; (b) 12 months post fistula closure and placement of a RBB; (c) 24 months post placement of a RBB




Fig. 5 Case 6 showing a Class 4 ICR affected maxillary right canine. Note the 'pinkish' colour at the cervical margin




Fig. 6 Case 13 showing a Class 4 ICR affected maxillary left central incisor. Implant and bone graft had been completed 4 years previously

orthodontic tooth movement, intra-coronal bleaching and prosthodontic operative procedures may have an effect upon dental root

structure.^{8,9} Control of these risk factors is particularly critical in oro-facial cleft-affected patients, given that the available dental tissues,



Fig. 4 Case 11 presenting 6 years following prosthodontic treatment and discharge

both hard and soft, are frequently developmentally compromised.

ICR is an uncommon but aggressive form of external root resorption. Its aetiology is poorly understood but trauma, alone, or in association with dental or surgical operative procedures is a recognised predisposing factor. Treatment of ICR is difficult leading to an uncertain long-term prognosis.⁸⁻¹⁰ The fundamental objectives in managing ICR are to expose the defect, remove the granulation tissue and to seal the defect.^{12,19} Heithersay's clinical classification can provide guidance to clinicians in assessing and managing the affected teeth, and careful case selection is advised in order to achieve a good prognosis.^{9,10} Heithersay recommends treating Class I-3 cases. Management of Class 4 ICR is difficult and has a high risk of failure. Therefore, he considers that Class 4 ICR cases may be left in-situ untreated for as long as they remain asymptomatic.⁹⁻¹¹ Otherwise, extraction is the only viable option, as ultimately was the case for all affected patients in this present study.

In our study, the ICR affected maxillary central, lateral and canine teeth were symptom free at presentation in all cases. Our study also supports previous findings that maxillary central incisor teeth are the most frequently affected and it is thought it is because these teeth are more prone to dental trauma.²⁶ Cleft patients are no more susceptible to trauma than the non-cleft population but incisors are most at risk. All of the unilateral CLP patients had just one ICR affected tooth and this was invariably associated with the cleft side. More than one affected ICR tooth was associated with bilateral CLP patients, and in this study the ICR affected teeth were confined to the cleft side. Common to all subjects was a history of fixed orthodontic treatment.

In patients who developed ICR unexpectedly during prosthodontic care, no common predisposing factor could be determined.

Neither could the duration before tooth loss became inevitable be predicted.

While the severity of the cleft influenced the number of ICR affected teeth, no association with the number of ICR affected teeth and the range or extent of surgical procedures could be found. In addition, no association with age and ICR was found. Interestingly, nine of the patients were over 25 years of age at presentation. Given the recognised improvement and refinements in surgical cleft care techniques in recent years, the possibility of the cohort in the present study being subjected to less conservative surgical procedures could not be discounted. We recognise that we are reporting observations on a small number of cleft patients and that these were from one centre. Nevertheless, the fact that ICR developed late and that this is a difficult condition to treat highlights the need for long-term follow up of cleft patients.

Conclusions

This paper reports on the experience of one specialist Cleft Centre regarding patients with ICR. As with similar Cleft Centres, patients underwent extensive specialised multidisciplinary dental care over a period of many years. This treatment included plastic surgery, orthodontics, prosthodontics, orthognathic surgery and fistula repair – all procedures recognised to have inherent predisposing traumatic risks. Though some patients presented to the

prosthodontic clinic with ICR and the tooth loss was immediately recognised, for others, the onset of ICR at or after prosthodontic care had been completed was unexpected. In neither group could the onset of ICR be predicted. This paper reinforces the importance of long-term review of oro-facial cleft affected patients in order to diagnose and treat possible cases of ICR at an early stage so as to avoid further tooth loss from an already compromised dentition.

1. Shaye D, Liu C C, Tollefson T T. Cleft lip and palate: An evidence-based review. *Facial Plast Surg Clin North Am* 2015; **23**: 357–372.
2. Pioto N R, Costa B, Gomide M R. Dental Development of the Permanent Lateral Incisor in Patients with Incomplete and Complete Unilateral Cleft Lip. *Cleft Palate-Craniofacial J* 2005; **42**: 517–520.
3. Al-Jamal G A, Hass'a A M, Rawashdeh M A. Crown-root ratio of permanent teeth in cleft lip and palate patients. *Angle Orthodontist* 2010; **80**: 1122–1128.
4. Deepti A, Muthu M S, Kumar N S. Root development of permanent lateral incisor in cleft lip and palate children: a radiographic study. *Indian J Dent Res* 2007; **18**: 82–86.
5. Ribero L L, das Neves L T, Costa B, Gomide M R. Dental development of permanent lateral incisor in complete unilateral cleft lip and palate. *Cleft Palate-Craniofacial J* 2002; **39**: 193–196.
6. Solis A, Figueroa A A, Cohen M, Polley J W, Evans C A. Maxillary dental development in complete unilateral alveolar clefts. *Cleft Palate-Craniofacial J* 1998; **35**: 320–328.
7. Zhou W, Li W, Lin J, Liu D, Xie X, Zhang Z. Tooth lengths of the permanent upper incisors in patients with cleft lip and palate determined with cone beam computed tomography. *Cleft Palate-Craniofacial J* 2013; **50**: 88–95.
8. Deepti A, Muthu M S, Kumar N S. Root development of permanent lateral incisor in cleft lip and palate children: a radiographic study. *Indian J Dent Res* 2007; **18**: 82–86.
9. Heithersay G S. Invasive Cervical Resorption. *Endod Topics* 2004; **7**: 73–92.
10. Heithersay G S. Life cycles of traumatized teeth: Long-term observations from a cohort of dental trauma victims. *Aust Dent J* 2016; **61**: 120–127.
11. Patel S, Kanagasingam S, Pitt Ford T. External Cervical Resorption: A Review. *J Endod* 2009; **35**: 616–625.
12. Umer F, Samira A, Khan F R. Conservative Management of Invasive Cervical resorption: A Case Report. *J Dent* 2013; **10**: 289–295.
13. Goon W W Y, Cohen S, Borer R F. External Cervical Root Resorption Following Bleaching. *J Endod* 1986; **12**: 414–418.
14. Kandalgaonkar S D, Gharat L A, Tupsakhare S D, Gabhane. Invasive Cervical Resorption: A Review. *J Int Oral Health* 2013; **5**: 124–130.
15. Yookyung K, Chan-Young L, Byoung-Duck R. Invasive Cervical Resorption: treatment challenges. *Rest Dent Endo* 2012; **37**: 228–231.
16. Neely A L, Gordon S C. A familial pattern of multiple idiopathic root resorption in a father and son: a 22 year follow up. *J Periodontol* 2007; **78**: 367–371.
17. Patel S, Dawood A, Whaites E, Pitt Ford T. New dimensions in endodontic imaging: part 1. Conventional and alternative radiographic systems. *Int Endod J* 2009; **46**: 463–475.
18. Patel S, Dawood A, Pitt Ford T, Whaites E. The potential applications of cone beam computed tomography in the management of endodontic problems. *Int Endod J* 2007; **40**: 818–830.
19. Lo Giudice G, Matarese G, Lizio A, Lo Giudice R, Tumedei M, Zizzari V L, Tete S. Invasive cervical resorption: A case series with 3 year follow-up. *Int J Periodontics Restorative Dent* 2016; **36**: 103–109.
20. Salzano S, Tirone F. Conservative nonsurgical treatment of class 4 invasive cervical resorption: A case series. *J Endod* 2015; **41**: 1907–1912.
21. Royzenblat A, Tordik P A, Goodell G. Cervical Resorption. *Clin Update* 2005; **27**: 1–2.
22. Frank A L. External-internal progressive resorption and its non-surgical correction. *J Endod* 1981; **7**: 473–476.
23. Goldman H M. Spontaneous intermittent resorption of the teeth. *J Am Dent Assoc* 1954; **49**: 522–532.
24. Koh E T, Torabinejad M, Pitt Ford T, Brady K. MTA stimulates a biological response in human osteoblasts. *J Biomed Mater Res* 1997; **37**: 432–439.
25. Asgary S, Nosrat A. Conservative management of Class 4 invasive cervical resorption using calcium enriched mixture cement. *J Endod* 2016; **8**: 1291–1294.
26. Nguyen Q V, Bezemer P D, Habets L, Prahl-Anderson B. A systematic review of the relationship between overjet size and traumatic dental injuries. *Eur J Orthod* 1999; **21**: 503–515.