



OPEN

SUBJECT AREAS:

GEODYNAMICS

GEOLOGY

SEISMOLOGY

TECTONICS

Seismic Evidence for a Geosuture between the Yangtze and Cathaysia Blocks, South China

Chuansong He¹, Shuwen Dong², M. Santosh³ & Xuanhua Chen²

¹Institute of Geophysics, China Earthquake Administration 100081, Beijing, China, ²Chinese Academy of Geological Science, 100037, Beijing, China, ³School of Earth Sciences and Resources, China University of Geosciences, 100083, Beijing, China.

Received
19 December 2012

Accepted
26 June 2013

Published
16 July 2013

Correspondence and
requests for materials
should be addressed to
C.H. (hechuansong@
yahoo.com.cn)

South China, composed of the Yangtze and Cathaysia Blocks and the intervening Jiangnan orogenic belt, has been central to the debate on the tectonic evolution of East Asia. Here we investigate the crustal structure and composition of South China from seismic data employing the *H-k* stacking technique. Our results show that the composition and seismic structure of the crust in the Jiangnan orogenic belt are identical to those of the Cathaysia Block. Our data reveal a distinct contrast in the crustal structure and composition between the two flanks of the Jiujiang-Shitai buried fault. We propose that the Jiujiang-Shitai buried fault defines a geosuture between the Yangtze and Cathaysia Blocks, and that the felsic lower crust of the Cathaysia Block and the Jiangnan orogenic belt may represent fragments derived from the Gondwana supercontinent.

The continental assembly of China, one of the cores of East Asia, is composed of the North China Craton, the Tarim Craton and the South China Block (Figure 1). The South China Block is composed of two sub-blocks, the Yangtze in the NW and Cathaysia in the SE (Figure 1), which collided and amalgamated during the Neoproterozoic, giving rise to the Jiangnan Orogen^{1–10}. Collision zones such as this are critical to our understanding of orogenic process and the evolution of the continents^{11,12}.

The South China continent abuts the western margin of the Pacific plate, and has witnessed plate subduction beneath the China mainland¹³. Since Grabau¹⁴, who first coined the term Cathaysia to describe the geology of southeastern China and part of the coastal region of West Pacific, debate has continued for over 80 years regarding the regional tectonics¹³, and the spatio-temporal evolution of the Cathaysia Block^{14–16}. Several workers consider that the boundary (geosuture) between the Yangtze and Cathaysia Blocks is defined by the Shaoxing-Yichun-Pingxiang fault^{6,17,18}.

In this study, we determine bulk crustal seismic properties and use these to infer the differences in structure and composition across the South China Block. Our study provides new seismic evidence for the precise location of the geosuture between the Yangtze Block and Cathaysia Blocks.

Results

We used the *H-κ* stacking method to determine the average crustal thickness (*H*) and the ratio of *P*- and *S*-wave velocities (*V_p/V_s* ratio, or *κ*) under each station¹⁹. This method treats the crust as a single homogeneous layer, and constrains *H* and *κ* of the crust by searching the most energetic stack of the direct *P_s* phase and multiples such as *PpPs*, *P_sP_s* + *PpS_s* of the Moho according to the predicted delays relative to the incident *P* wave.

Before *H-k* stacking was performed, a reasonable bulk crustal *V_p* and search range of *H* and *k* were adopted. Based on deep seismic sounding investigation, the value of bulk crustal *V_p* in the study area should be approximately 6.3 km s⁻¹²⁰. Therefore, we assume a mean crustal *P*-wave velocity (*V_p*) of 6.3 km s⁻¹ and perform *H-k* stacking^{19,21} (search range: *H* = 25–50 km; *V_p/V_s* = 1.4–2.5). Finally, we obtained 251 estimates on robust crustal thickness and *V_p/V_s* ratios (Table S1). The resulting bulk crustal *V_p/V_s* ratios range from 1.53 to 2.11 with an average of 1.72. The Jiangnan orogenic belt and the Cathaysia Block are characterized by lower *V_p/V_s* ratio of 1.66–1.73, as compared to the *V_p/V_s* ratios of 1.74–1.79 in the SECCLMVZ (Southeast China coastal late Mesozoic volcanic zone). The *V_p/V_s* ratios of 1.76–2.11 in the Yangtze Block in Northwestern part of this study area are even higher (Figure 2, Table S1).

Our study shows that the average crustal thickness of South China is 33 km. However, locally the values vary widely between 25.5 and 56.9 km. Thinner crust of 25.6–32 km is observed in the Jiangnan orogenic belt and the




Figure 1 | The distribution of seismic stations in South China and the general tectonic framework. F1: Shaoxing-Jiangshan-Pingxiang fault, F2: Zhenghe-Dapu fault, F3: Tanlu fault, F4: Xiangfan-Guangji fault, F5: Lu'an fault, F6: Longmenshan fault, F7: Jiujiang-Shitai buried fault²⁷; P1, P2, P3 and P4: profile, blue font: the seismic stations at the profiles; black triangle: seismic stations; SECCLMVZ: Southeast China coastal late Mesozoic volcanic zone. Inset figure (right-upper corner): Distribution of selected event. For each event-station pair, data were selected within the distance ranges of 30°–95° (The figure is generated using Generic Mapping Tool (<http://gmt.soest.hawaii.edu/>) by Chuansong He).

Cathaysia Block, with a relatively flat Moho. A thick crust of 40–56.9 km is observed beneath the Yangtze Block in the northwestern part of the study area (Figure 3, Table S1).

Discussion

For lower-crustal rocks, low σ (<0.26) ($V_p/V_s < 1.75$), intermediate σ ($0.26–0.28$) and high σ (>0.28) ($V_p/V_s > 1.81$) are characteristic of felsic, intermediate and mafic compositions^{21–23,25}, respectively. If the bulk value of the entire crust is $\sigma > 0.28$, the lower crust must have a value of ~ 0.30 and therefore, the low Poisson's ratio ($\sigma = 0.25$) for the Mesozoic-Cenozoic crust is considered to indicate a predominantly felsic composition²¹. The V_p/V_s values are around 1.73 for felsic rocks, whereas for mafic rocks, this value tends to be greater than 1.73^{22–24}. The values of bulk crustal V_p/V_s ratio are slightly less than that of lower crustal V_p/V_s ratio (about 0.02 or so)^{21,22,24,25}. Therefore, we can use bulk crustal V_p/V_s ratio to estimate the lower crust V_p/V_s ratio^{24,26}.

The distribution of bulk crustal V_p/V_s ratio indicates that the Jiangnan orogenic belt and the Cathaysia Block are characterized by felsic lower crust and the Yangtze Block by intermediate and mafic-ultramafic lower crust^{21,25}. Notably, the bulk V_p/V_s ratio in the Jiangnan orogenic belt is lower than 1.70, which might be related to the collisional orogenesis. The similar crustal thickness on both sides of Shaoxing-Jiangshan-Pingxiang fault implies that the mountain root has been completely lost, or that the lower crust delamination beneath the Jiangnan orogenic belt was more intense, leading to the lower V_p/V_s values. The late Mesozoic coastal volcanic zone in Southeast China shows dominantly intermediate lower crust^{21,25}, and the narrow distribution of these strips might reflect deep-seated magmatism associated with tectonic processes.

The distribution of the V_p/V_s presented here indicates that along the boundary of the Jiujiang-Shitai buried fault, there is a marked difference between the Jiangnan orogenic belt, the Cathaysia Block and Yangtze Block. The distribution of crustal thickness also assigns the Jiangnan-Shitai buried fault as the major boundary, with the crustal thickness increasing northwards in the Yangtze Block. The crustal thickness in the Jiangnan orogenic belt and the Cathaysia Block shows minor change laterally with a smooth variation of the Moho²⁸ (Figure 3 and Figure 4). The depth domain profiles from our study at various stations bring out the presence of an important boundary in crustal structure between the BJT and AST (profile 1), HOJ and YOZ (profile 2), TAY and YIY (profile 3), and XNI and DUC stations (profile 4). Notably, this boundary line almost precisely coincides with the Jiangnan-Shitai buried fault. In figure 4, the crust beneath the Jiangnan orogenic belt and the Cathaysia Block along the red lines is characterized by a simple crustal structure. In contrast, the crust above the red lines, relating to the Yangtze Block, is dominantly featured by a complex crustal structure. A relatively large offset, of around 5 km, exists on both sides of the Jiujiang-Shitai buried fault above and below the red (or yellow) lines.

Our results clearly indicate the absence of prominent differences in crustal structure on the two flanks of the Shaoxing-Jiangshan-Pingxiang fault (Figure 2, Figure 3, and Figure 4: above and below the dotted line). However, the Jiangnan-Shitai buried fault obviously defines an important boundary based on crustal composition and structure (Figure 1, Figure 2, Figure 3, Figure 4, Figure 5). Therefore, we propose that the Jiangnan-Shitai buried fault defines the boundary between the Yangtze and Cathaysia Blocks.

Recent studies have proposed that all the East and SE Asian continental terranes/blocks were directly or indirectly derived from the eastern margin of the Gondwana supercontinent²⁹. There is




Figure 2 | Distribution of V_p/V_s ratios in South China. F1: Shaoxing-Jiangshan-Pingxiang fault, F2: Zhenghe-Dapu fault, F3: Tanlu fault, F4: Xiangfan-Guangji fault, F5: Lu'an fault, F6: Longmenshan fault, F7: Jiujiang-Shitai buried fault²⁷; P1, P2, P3 and P4: profile; black triangle: effective data point; SECCLMVZ: Southeast China coastal late Mesozoic volcanic zone (The figure is generated using Generic Mapping Tool (<http://gmt.soest.hawaii.edu/>) by Chuansong He).

increasing evidence to consider the South China block, including the Yangtze Block, as an integral part of East Gondwana in early Palaeozoic, rather than as a discrete continental block in the Palaeo-Pacific or a fragment of Laurentia^{30–33}. If this model is true, the Jiangnan orogenic belt and the Cathaysia Block, with broadly consistent crustal characteristics and thickness, might represent fragments from the Gondwana assembly.

An active continental-margin model, with the subduction of the paleo-Pacific plate beneath the South China Craton, has been invoked in several studies to account for the extensive magmatic zone in southeastern China during the Mesozoic^{3,34–38}. This has led to the popular concept that the Mesozoic low-angle subduction of the Pacific Plate has played a major role in the thinning of the South China Land Mass^{39,40}.

Subduction and collision tectonics may generate local topography on Moho which may survive for a long time^{41,42}. The dipping Moho topography is considered as an indication of the remnants of deep collision and subduction⁴³. Our studies show little lateral variations in crustal thickness and a flat Moho in the Cathaysia block and Jiangnan orogenic belt. These results do not support the model of low-angle subduction of the Pacific Plate and resultant crustal thinning.

Delamination refers to the loss and sinking of the portion of the lower crust and (or) the lowermost lithosphere from the tectonic plate to which it was attached. This can occur when the lower crust and (or) the lower portion of the lithosphere becomes denser than the surrounding mantle. Because of the instability of higher density (e.g. thickening lower crust) material above lower density material, the lower crust and (or) the lower lithosphere separates from the tectonic plate and sinks into the mantle^{44–46}, which results in asthenospheric

upwelling into the space previously occupied by thickened lithosphere. Flow of hot mantle material encounters the base of the thin lithosphere and often results in melting and a new phase of volcanism^{44–46}.

Since the Mesozoic, the South China region has been located at the center of a triangular area surrounded by westward subduction of the Pacific plate (Cenozoic, about ~50 Ma), northward subduction of the India Plate beneath the Eurasia Plate (Cenozoic, about ~50 Ma), and collision of the North and South China blocks along the Central China Orogen (Permian-Triassic, about 290–250 Ma)²⁸. This region (including the Yangtze and Cathaysia blocks) thus marks the frontier of a super-convergent regime. Within the super-convergence domain, the compressional structures in the center of the South China Block are mainly characterized by shortening, thrusting and decollement⁴⁷.

The collision between the Yangtze and Cathaysia Blocks generated a thick lithospheric root in the Hercynian-Indosinian period (409–205 Ma). Subsequently, in Yanshanian (208–135 Ma), delamination and asthenospheric upwelling led to extensive lithospheric extension and thinning^{48,49}. The collision between the Yangtze and the Cathaysia Block in the Triassic as well as crustal detachment of the eastern Yangtze Block might have led to the thickening of lower crust and delamination⁵⁰.

The lower V_p/V_s ratio in the bulk lower crust of the Jiangnan orogenic belt and Cathaysia Block might suggest deep process such as lower crustal delamination, which resulted in the dominantly felsic lower crust^{21,25,26} beneath the Jiangnan orogenic belt and Cathaysia Block. Therefore, we favor lower crustal delamination of the Cathaysia Block and crustal thinning associated with mantle upwelling^{44–46,51} as the most plausible scenario to explain the extensive




Figure 3 | Distribution of crust thickness. F1: Shaoxing-Jiangshan-Pingxiang fault, F2: Zhenghe-Dapu fault, F3: Tanlu fault, F4: Xiangfan-Guangji fault, F5: Lu'an fault, F6: Longmenshan fault, F7: Jiujiang-Shitai buried fault²⁷; P1, P2, P3 and P4: profile; black triangle: effective data point, SECCLMVZ: Southeast China coastal late Mesozoic volcanic zone (The figure is generated using Generic Mapping Tool (<http://gmt.soest.hawaii.edu/>) by Chuansong He).

magmatic activity in this region. A recent *S* receiver function study indicates lithospheric thinning in Southern China⁵², which supports the delamination model for this area^{53,54}. Therefore, we consider that the lower crust/lithosphere delamination⁵⁵ beneath the Cathaysia Block might have led to lithospheric thinning in this area and the surrounding regions.

In summary, we envisage the following processes: (1) tectonic thickening of the crust and lithosphere by convergence; (2) asthenospheric upwelling and infiltration of the lithosphere; (3) subsequent weakening of the lithosphere by this infiltration, resulting in delamination of the lower crust and mantle. The thickened lower crust was eclogitic, hence denser than the underlying mantle. The geosuture between the Yangtze and Cathaysia Blocks is defined by Jiujiang-Shitai buried fault. The Cathaysia Block, dominated by felsic bulk lower crust^{21,24,25}, is probably a fragment of the Gondwana supercontinent. The large offset between the Yangtze and Cathaysia Blocks lends support to the notion of collisional assembly of these two discrete blocks (Figure 6).

Methods

Teleseismic receiver functions⁵⁶ are very sensitive to the *S*-wave velocity beneath the station and have proven to be a useful tool for estimating crustal thicknesses and *V_p/V_s* ratios beneath individual seismic stations^{19,57,58}. The *P*-to-*S* converted phase at the Moho and the first reverberated phases in the crust are generally apparent in the receiver function waveforms, and their relative travel times can then be employed to constrain the crustal thickness and the bulk *V_p/V_s* ratio below the recording station^{21,57}. Deciphering the geological evolution of the Earth's continental crust requires knowledge of its bulk composition and global variability²¹. Average *V_p/V_s* or Poisson's ratio ($\sigma = 0.5[1 - ((V_p/V_s)^2 - 1)^{-1}]$), can be used to complement petrological studies of crustal composition⁵⁸.

This study performs a careful analysis of the receiver functions in South China, in order to characterize the bulk seismic properties of crust with local estimates for the

crustal thickness and *V_p/V_s* ratio^{19,21}. For this purpose, we apply the stacking procedure of Zhu and Kanamori¹⁹ (2000) to 281 seismic stations located in South China (Figure 1).

These stations have been in operation between from July 2008-present. We selected a total of 424 events with magnitude *m_b* ≥ 6.0 recorded by those stations⁵⁹ (Figure 1 and Table S2). For each event-station pair, data were selected within the distance ranges of 30°–95° and initially windowed 15 *s* before and 120 *s* after the *P*-wave pick. Only signals with a good signal-to-noise ratio and a clearly identifiable *P*-wave arrival were used. Data are filtered using a zero-phase Butterworth bandpass filter with corner frequencies of 0.03–3 Hz.

In our study we use a modified frequency domain deconvolution^{56,60}, which is implemented by dividing the spectrum *R*(ω) of the teleseismic *P* waveform by the source spectrum *S*(ω):

$$r(t) = (1 + c) \int \frac{R(\omega)S^*(\omega)}{|S(\omega)|^2 + c\sigma_0^2} e^{-\frac{\omega^2}{4\alpha^2}} e^{i\omega t} d\omega \quad (1)$$

Where *S*^{*}(ω) is the complex conjugate of *S*(ω), the Gaussian-type low-pass filter $e^{-\omega^2/4\alpha^2}$ is added to remove high-frequency noise. The quantity $c\sigma_0^2$ is used to suppress "holes" in the spectrum *S*(ω), thus stabilizing the deconvolution where *r*(*t*) is the radial receiver function. The 1 + *c* factor is used to compensate the amplitude loss due to the water level¹⁹. In this study, the Gaussian factor and water level were set to 3 and 0.01, respectively.

The time separation between *P_s* and *P* can be used to estimate crustal thickness, given the average crustal velocities,

$$H = \frac{t_{P_s}}{\sqrt{\frac{1}{V_s^2} - p^2} - \sqrt{\frac{1}{V_p^2} - p^2}} \quad (2)$$

Where *p* is the ray parameter of the incident wave and the crustal velocity is given, we can obtain the thickness estimation; however, one problem is the trade-off between the thickness and crustal velocities.

$$H = \frac{t_{P_p P_s}}{\sqrt{\frac{1}{V_s^2} - p^2} + \sqrt{\frac{1}{V_p^2} - p^2}} \quad (3)$$




Figure 4 | Profiles in the depth domain. P1: profile 1, P2: profile 2, P3: profile 3, P4: profile 4. Blue lines: the crust-mantle boundary of the Yangtze block, yellow lines: the crust-mantle boundary of the Jiangnan orogenic belt and the Cathaysia block, dotted line: the boundary between the Jiangnan orogenic belt and the Cathaysia block, yellow dotted line: Jiujiang-Shitai buried fault. The red line on all profiles almost precisely coincides with the Jiangnan-Shitai buried fault except for profile 2. The Jiangnan orogenic belt and the Cathaysia Block are characterized by a relatively flat Moho. The large offset between the Yangtze and Cathaysia Blocks lends support to the notion of collisional assembly of these two discrete blocks (The figure is generated using Generic Mapping Tool (<http://gmt.soest.hawaii.edu/>) by Chuansong He).

$$\frac{V_p}{V_s} = \left\{ (1 - p^2 V_p^2) \left[2 \left(\frac{t_{ps} - t_p}{t_{ppps} - t_{ps}} \right) + 1 \right]^2 + p^2 V_p^2 \right\}^{1/2} \quad (4)$$

The trade-off influence can be reduced by using the later phases, which provide additional constraints. The precise crustal H and V_p/V_s ratios can be estimated by (3) and (4)^{19,21}.

The stacking is usually done in the time domain for a cluster of events⁶⁰. We propose a straightforward H - k domain stacking defined as below:

$$s(H, k) = w_1 r(t_1) + w_2 r(t_2) - w_3 r(t_3) \quad (5)$$

Where $r(t)$ is the radial receiver function, t_1, t_2 and t_3 are the predicted P_s , $PpPs$ and $PpPs + PsPs$ arrival time corresponding to crustal thickness H and V_p/V_s ratios (ratio k), as given in (2)–(4). The w_i are weighting factors, and $\sum w_i = 1$. The $s(H, k)$ reaches a maximum when all of these phases are stacked coherently with the correct H and k ²¹. Here, we chose unequal weights (0.6, 0.3 and 0.1) for P_s , $PpPs$ and $PpPs + PsPs$ phases

of the Moho, respectively. Using the Taylor expansion of $s(H, k)$ at the maximum and omitting the higher-order terms, one gets the variances of H and k :

$$\sigma_H^2 = 2\sigma_s \frac{\partial^2 s}{\partial H^2}, \quad (6)$$

$$\sigma_k^2 = 2\sigma_s \frac{\partial^2 s}{\partial k^2}, \quad (7)$$

Where σ_s is the estimated variance of $s(H, k)$ from stacking¹⁹. On the other hand, the study shows that the uncertainty of H is < 0.5 km for a 0.1 km/s uncertainty in V_p ¹⁹.

To quantitatively estimate the uncertainty of our results, we measured error bars for h and κ of the profile 2, which line up with longitude, by taking into account uncertainties (Figure 5). Their uncertainties are estimated using (6) and (7)¹⁹. Here we show two examples of H - k stacking computed at two stations located at the study area (Figure S1).




Figure 5 | Blue dots and error bar represent the best estimates of crustal thickness (h) and V_p/V_s ratios (k) and errors of profile 2 in the H - κ stacking of receiver function. (a) Crustal thickness with error bars. (b) V_p/V_s ratios with error bars. Yellow arrow: Jiujiang-Shitai buried fault. Red arrow: the boundary between the Yangtze and Cathaysia Block from this studies, dotted arrow: the boundary between the Jiangnan orogenic belt and Cathaysia block (The figure is generated using Generic Mapping Tool (<http://gmt.soest.hawaii.edu/>) by Chuansong He).




Figure 6 | Collision model between the Yangtze and Cathaysia Blocks. F1: Jiujiang-Shitai buried fault, F2: Shaoxing-Jiangshan-Pingxiang fault, F3: Longmenshan fault. The geosuture between the Yangtze and the Cathaysia Blocks is defined by Jiujiang-Shitai buried fault. A relatively large offset, of around 5 km, exists on both sides of the Jiujiang-Shitai buried fault. The Cathaysia Block might suggest deep process such as lower crustal delamination, which resulted in the dominantly felsic lower crust and crustal thinning beneath the Jiangnan orogenic belt and Cathaysia Block. The Yangtze Block is characterized by intermediate and mafic-ultramafic lower crust (The figure is generated using Generic Mapping Tool (<http://gmt.soest.hawaii.edu/>) by Chuansong He).

We migrate the receiver functions to depth using the iasp91 velocity model⁶¹. The amplitude itself is proportional to the velocity or, more precisely, the impedance contrast at that location. Based on it, we converted the time domain receiver functions into the depth domain after corrections for the incidence angle effect (correcting the move out of P_s conversions to same incidence or near vertical incidence)⁶².

By stacking all depth domain receiver functions from different backazimuths at each station respectively, we obtain the average receiver function at each station (multi-events to stack their receiver function)⁶⁰. The average (stacked) receiver function can clearly delineate Moho interface which is usually the largest discontinuity⁶³. We then get a rough estimation of the Moho depth variation beneath each station from the average receiver function (Figure S2–S3).

1. Zhao, G. C. & Cawood, P. A. Tectonothermal evolution of the Mayuan Assemblage in the Cathaysia Block: implications for Neoproterozoic collision-related assembly of the South China Craton. *Am. J. Sci.* **299**, 309–339 (1999).
2. Li, L. M. *et al.* U–Pb and Hf isotopic study of zircons from migmatized amphibolites in the Cathaysia Block: Implications for the early Paleozoic peak tectonothermal event in Southeastern China. *Gondwana Res.* **19**, 191–201 (2011).
3. Yan, Y., Hu, X. Q., Lin, G., Santoch, M. & Chan, L. S. Sedimentary provenance of the Hengyang and Mayang basins, SE China, and implications for the Mesozoic topographic change in South China Craton: Evidence from detrital zircon geochronology. *J. Asian Earth Sci.* **41**, 494–503 (2011).
4. Ren, J. S. On the geotectonics of southern China. *Acta Geologica Sinica (English Edition)* **2**, 111–136 (1991).
5. Duan, L., Meng, Q. R., Zhang, C. L. & Liu, X. M. Tracing the position of the South China block in Gondwana: U–Pb ages and Hf isotopes of Devonian detrital zircons. *Gondwana Res.* **19**, 141–149 (2011).
6. Wong, J. *et al.* Zircon U–Pb and Hf isotopic study of Mesozoic felsic rocks from eastern Zhejiang, South China: geochemical contrast between the Yangtze and Cathaysia blocks. *Gondwana Res.* **19**, 244–259 (2011).
7. Li, Z., Li, X., Zhou, H. & Kinny, P. D. Grenvillian continental collision in south China: new SHRIMP U–Pb zircon results and implications for the configuration of Rodinia. *Geology* **30**, 163–166 (2002).
8. Zhou, X. & Zhu, Y. Late Proterozoic collisional orogen and geosuture in southeastern China: petrological evidence. *Chin. J. Geochem.* **12**, 239–251 (1993).
9. Mao, X. L. *et al.* Effective elastic thickness and mechanical anisotropy of South China and surrounding regions. *Tectonophysics* **550–553**, 47–56 (2012).
10. Wang, Y. J. *et al.* Numerical modeling of the formation of Indo-Sinian peraluminous granitoids in Hunan Province: Basaltic underplating versus tectonic thickening. *Sci. China (D)* **45**, 1042–1056 (in Chinese) (2002).
11. Dewey, J. F. Suture zone complexities: a review. In: Toksoz, M. N. *et al.* (Ed.) *Oceanic Ridges and Arcs*: New York, pp. 477–491 (1980).
12. Zhou, X. & Zhu, Y. Late Proterozoic collisional orogen and geosuture in southeastern China: petrological evidence. *Chin. J. Geochem.* **12**, 239–251 (1993).
13. Zhao, B., Zhang, Z., Bai, Z. M., Bada, J. & Zhang, Z. J. Shear velocity and Vp/Vs ratio structure of the crust beneath the southern margin of South China continent. *J. Asian Earth Sci.* **62**, 167–179 (2013).
14. Grabau, A. W. *Stratigraphy of China, Part I*, China Publishing, 1924.
15. Mao, J. W. *et al.* Geodynamic process and metallogeny: History and present research trend, with a special discussion on continental accretion and related metallogeny throughout geological history in South China. *Mineral Deposits* **24**, 193–205 (in Chinese with English abstract) (2005).
16. Wang, Q. C. Preliminary discussion on sedimentary tectonics of the clustered continents of South China. *Acta Sedimentologica Sinica* **27**, 811–817 (in Chinese with English abstract) (2009).

17. Mao, J. R., Tao, K. Y., Xing, G. F., Yang, Z. L. & Zhao, Y. Petrological records of the Mesozoic-Cenozoic mantle plume tectonics in epicontinental area of Southeast China. *Acta Geoscientia Sinica* **20**, 252–258 (in Chinese with English abstract) (1999).
18. Mou, C. L. & Xu, X. S. Sedimentary evolution and petroleum geology in South China during the early Palaeozoic. *Sedimentary Geology and Tethyan Geology* **30**, 24–29 (in Chinese with English abstract) (2010).
19. Zhu, L. & Kanamori, H. Moho depth variation in southern California from teleseismic receiver functions. *J. Geophys. Res.* **105**, 2969–2980 (2000).
20. Deng, Y. F., Li, S. L., Fan, W. M. & Li, J. Crustal structure beneath South China revealed by deep seismic soundings and its dynamics implications. *Chinese J. Geophys.* **54**, 2560–2574 (in Chinese with English abstract) (2011).
21. Zandt, G. & Ammon, C. J. Continental crust composition constrained by measurements of crustal Poissons ratio. *Nature* **374**, 152–154 (1995).
22. Christensen, N. Poisson's ratio and crustal seismology. *J. Geophys. Res.* **101**, 3139–3156 (1996).
23. Chang, S. J. & Baag, C. E. Moho Depth and Crustal Vp/Vs Variation in Southern Korea from Teleseismic Receiver Functions: Implication for Tectonic Affinity. *Bull. Seism. Soc. Am.* **97**, 1621–1631 (2007).
24. Niu, F. & James, D. E. Fine structure of the lowermost crust beneath the Kaapvaal craton and its implications for crustal formation and evolution. *Earth Planet. Sci. Lett.* **200** (1–2), 121–130 (2002).
25. Nair, S. K., Gao, S. S., Liu, K. H. & Silver, P. G. Southern African crustal evolution and composition: Constraints from receiver function studies. *J. Geophys. Res.* **111**, B02304 (2006). doi:10.1029/2005JB003802
26. Thompson, D. A. *et al.* Precambrian crustal evolution: Seismic constraints from the Canadian Shield. *Earth Planet. Sci. Lett.* **297** (3–4), 655–666 (2010). doi:10.1016/j.epsl.2010.07.021
27. Yao, J. L., Shu, L. S., Santosh, M. & Li, J. Y. Geochronology and Hf isotope of detrital zircons from Precambrian sequences in the eastern Jiangnan Orogen: Constraining the assembly of Yangtze and Cathaysia Blocks in South China. *J. Asian Earth Sci.*, (in press) (2012).
28. Chen, Y. *et al.* Crustal structure beneath China from receiver function analysis. *J. Geophys. Res.* **115** (B3) (2010).
29. Duan, L., Meng, Q. R., Wu, G. L., Ma, S. X. & Li, L. Detrital zircon evidence for the linkage of the South China block with Gondwanaland in early Palaeozoic time. *Geol. Mag.* **149**, 1124–1131 (2012).
30. Metcalfe, I. Gondwana dispersion and Asian accretion: Tectonic and palaeogeographic evolution of eastern Tethys. *J. Asian Earth Sci.* (2013) (in press).
31. Wang, H. N. & Zhou, L. Y. A further understanding in geological structure of South China. *Geological Journal of China Universities* **12**, 457–465 (in Chinese with English abstract) (2006).
32. Xu, D. R. *et al.* Petrological, mineralogical and geochemical characteristics of Ordovician volcanic-clastic sedimentary rocks in Bangxi area, Northwest Hainan island, South China: Implications for provenance and tectonic setting. *Geochimica* **36**, 11–26 (in Chinese with English abstract) (2007).
33. Metcalfe, I. Gondwanaland dispersion, Asian accretion and evolution of eastern Tethys. *Aust. J. Earth Sci.* **43**, 605–623 (1996).
34. Charvet, J., Lapiere, H. & Yu, Y. Geodynamic significance of the Mesozoic volcanism of southeastern China. *J. Southeast Asian Earth Sci.* **68**, 387–396 (1994).
35. Lapiere, H., Jahn, B. M., Charvet, J. & Yu, Y. W. Mesozoic magmatism in Zhejiang Province and its relation with the tectonic activities in SE China. *Tectonophysics* **274**, 321–338 (1997).
36. Sewell, R. J. & Campbell, S. D. G. Geochemistry of coeval Mesozoic plutonic and volcanic suites in Hong Kong. *J. Geol. Soc. London.* **154**, 1053–1066 (1997).



37. Zhou, X. M. & Li, W. X. Origin of Late Mesozoic igneous rocks in Southeastern China: implications for lithosphere subduction and underplating of mafic magmas. *Tectonophysics* **326**, 269–287 (2000).
38. Li, Z. X. & Li, X. H. Formation of the 1300 km-wide intracontinental orogen and post-orogenic magmatic province in Mesozoic South China: a flat-slab subduction model. *Geology* **35**, 179–182 (2007).
39. Zheng, Y. F. & Wu, F. Y. Growth and reworking of cratonic lithosphere. *Chinese Sci. Bull.* **54**, 3347–3353 (2009).
40. Zhang, J. J., Zheng, Y. E. & Zhao, Z. E. Geochemical evidence for interaction between oceanic crust and lithospheric mantle in the origin of Cenozoic continental basalts in east-central China. *Lithos* **110**, 305–326 (2009).
41. Cook, F. A., Velden, A., Hall, K. W. & Reberts, B. J. Frozen subduction in Canada's Northwest Territories: lithoprobe deep lithospheric reflection profiling of the western Canadian Shield. *Tectonics* **18**, 1–24 (1999).
42. Balling, N. Deep seismic reflection evidence for ancient subduction and collision zones within the continental lithosphere of northwestern Europe. *Tectonophysics* **329**, 269–300 (2000).
43. Sverning, L. *et al.* Crustal root beneath the highlands of southern Norway resolved by teleseismic receiver functions. *Geophys. J. Int.* **170**, 1129–1138 (2007).
44. Kay, R. W. & Kay, S. M. Delamination and delamination magmatism. *Tectonophysics* **219**, 177–189 (1993).
45. Vlaar, N. J., van Keken, P. E. & van den Berg, A. P. Cooling of the Earth in the Archaean: consequences of pressure-release melting in a hotter mantle. *Earth Planet. Sci. Lett.* **121**, 1–18 (1994).
46. van Thienen, P., van den Berg, A. P. & Vlaar, N. J. Production and recycling of oceanic crust in the early Earth. *Tectonophysics* **386**, 41–65 (2004).
47. Li, S. Z., Santosh, M., Zhao, G. C., Zhang, G. W. & Jin, C. Intracontinental deformation in a frontier of super-convergence: A perspective on the tectonic milieu of the South China Block. *J. Asian Earth Sci.* **49**, 313–329 (2012).
48. Shen, X. M., Zhang, H. X. & Zhang, B. Y. A preliminary study of relationship between metamorphic core complexes and lithospheric thinning over the Mesozoic in South China. *Geotectonica et Metallogenia* **116**, 11–19 (in Chinese with English abstract) (2008).
49. Wang, Y. J., Fan, W. M., Guo, F., Pen, T. P. & Li, C. W. Geochemistry of Mesozoic mafic rocks adjacent to Chenzhou-Linwu fault, South China: Implications for the lithospheric boundary between the Yangze and Cathaysia Block. *In. Geol. Rev.* **45**, 263–286 (2003).
50. Zi, F. *et al.* Shrimp U-Pb zircon geochronology and geochemistry of the Guandian pluton in central Anhui, China: Petrogenesis and geodynamic implications. *Geochimica* **37**, 462–480 (in Chinese with English abstract) (2008).
51. Zheng, T. Y., Zhao, L. & Zhu, R. X. Insight into the geodynamics of cratonic reactivation from seismic analysis of the crust-mantle boundary. *Geophys. Res. Lett.* **35**, L08303, doi:10.1029/2008GL033439 (2008).
52. Sodoudi, F., Yuan, X., Liu, Q., Kind, R. & Chen, J. Lithospheric thickness beneath the Dabie Shan, central eastern China from S receiver functions. *Geophys. J. Int.* **166**, 1363–1367 (2006).
53. Zhang, Q. *et al.* A model of delamination of continental lower crust. *Acta Petrologica Sinica* **22**, 265–276 (in Chinese with English abstract) (2006).
54. Handy, M. R. & Brun, J.-P. Seismicity structure and strength of the continental lithosphere. *Earth Planet. Sci. Lett.* **223**, 427–441 (2004).
55. Levander, A. *et al.* Continuing Colorado plateau uplift by delamination-style convective lithospheric downwelling. *Nature* **472** (7344), 461–465 (2011).
56. Langston, C. A. Structure under Mount Rainier, Washington, inferred from teleseismic body waves. *J. Geophys. Res.* **84**, 4749–4762 (1979).
57. Zandt, G., Myers, S. C. & Wallace, T. C. Crust and mantle structure across the Basin and Range-Colorado Plateau boundary at 37°N latitude and implications for Cenozoic extensional mechanism. *J. Geophys. Res.* **100**, 10529–10548 (1995).
58. Chevrot, S. & van der Hilst, R. D. The Poisson's ratio of the Australian crust: geological and geophysical implications. *Earth planet. Sci. Lett.* **183**, 121–132 (2000).
59. Zheng, X. F., Yao, Z. X., Liang, J. H. & Zheng, J. The role played and opportunities provided by IGP DMC of China National Seismic Network in Wenchuan earthquake disaster relief and researches. *Bull. Seismol. Soc. Am.* **100**(5B), 2866–2872 (2010).
60. Owens, T. J., Zandt, G. & Taylor, S. R. Seismic evidence for ancient rift beneath the Cumberland plateau, Tennessee: A detailed analysis of broadband teleseismic P waveform. *J. Geophys. Res.* **89**, 7783–7795 (1984).
61. Kennett, B. L. N. & Engdahl, E. R. Travel times for global earthquake location and phase identification. *Geophys. J. Int.* **105**, 429–465 (1991).
62. Yuan, X., Ni, J., Kind, R., Sandvol, E. & Mechie, J. Lithospheric and upper mantle structure of southern Tibet from a seismological passive source experiment. *J. Geophys. Res.* **102**, 27491–27500 (1997).
63. Zhu, L. Crustal structure across the San Andreas Fault, southern California from teleseismic converted waves. *Earth planet. Sci. Lett.* **179**, 183–190 (2000).

Acknowledgements

The authors acknowledge SinoProbe-Deep Exploration in China (SinoProbe-08). Waveform data for this study are provided by Data Management Centre of China National Seismic Network at Institute of Geophysics, China Earthquake Administration. This study also contributes to the 1000 Talents Award to M. Santosh from the Chinese Government.

Author contributions

C.S.H. wrote the main manuscript text and S.W.D., M.S. and X.H.C. revised the manuscript. All authors reviewed the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: He, C., Dong, S., Santosh, M. & Chen, X. Seismic Evidence for a Geosuture between the Yangtze and Cathaysia Blocks, South China. *Sci. Rep.* **3**, 2200; DOI:10.1038/srep02200 (2013).



This work is licensed under a Creative Commons Attribution 3.0 Unported license. To view a copy of this license, visit <http://creativecommons.org/licenses/by/3.0>