



## OPEN

Prevalence of *psbA*-containing cyanobacterial podoviruses in the oceanQiang Zheng<sup>1</sup>, Nianzhi Jiao<sup>1</sup>, Rui Zhang<sup>1</sup>, Feng Chen<sup>2</sup> & Curtis A. Suttle<sup>3</sup>SUBJECT AREAS:  
MICROBIAL ECOLOGY  
EVOLUTIONARY ECOLOGY  
BIODIVERSITYReceived  
27 June 2013Accepted  
25 October 2013Published  
13 November 2013Correspondence and  
requests for materials  
should be addressed to  
N.Z.J. (jiao@xmu.edu.  
cn)

<sup>1</sup>State Key Laboratory for Marine Environmental Science, Institute of Marine Microbes and Ecospheres, Xiamen University, Xiamen 361005, People's Republic of China, <sup>2</sup>Institute of Marine and Environmental Technology, University of Maryland Center for Environmental Science, Baltimore, MD 21202, USA, <sup>3</sup>Departments of Earth, Ocean and Atmospheric Sciences, Microbiology and Immunology, Botany, and the Biodiversity Research Centre, The University of British Columbia, Vancouver, British Columbia, Canada.

Podoviruses that infect marine picocyanobacteria are abundant and could play a significant role on regulating host populations due to their specific phage-host relationship. Genome sequencing of cyanophages has unveiled that many marine cyanophages encode certain photosynthetic genes like *psbA*. It appears that *psbA* is only present in certain groups of cyanopodovirus isolates. In order to better understand the prevalence of *psbA* in cyanobacterial podoviruses, we searched the marine metagenomic database (GOS, BATS, HOT and MarineVirome). Our study suggests that 89% of recruited cyanopodovirus scaffolds from the GOS database contained the *psbA* gene, supporting the ecological relevance of the photosynthesis gene for surface oceanic cyanophages. Diversification between Clade A and B are consistent with recent finding of two major groups of cyanopodoviruses. All the data also shows that Clade B cyanopodoviruses dominate the surface ocean water, while Clade A cyanopodoviruses become more important in the coastal and estuarine environments.


Viruses are abundant in the ocean and can influence population dynamics and genetic diversity of their hosts<sup>1–3</sup>. Cyanophage are a specific group of viruses which infect cyanobacteria mainly including *Prochlorococcus* and *Synechococcus*. Many cyanophages have been isolated, and all the known marine cyanophages belong to three phage families: *Myoviridae*, *Siphoviridae* and *Podoviridae*<sup>4–10</sup>. Recent studies showed that cyanopodoviruses might make up 50% of cyanophage community in the sea<sup>11–12</sup>, suggesting that cyanopodoviruses interact actively with cyanobacteria in the marine environment.

Currently, nearly 40 cyanophage genomes have been sequenced, and half of them are cyanomyoviruses. Cyanomyoviruses have a relatively large genome size and acquire many accessory metabolic genes via horizontal gene transfer (HGT), which constitute the large reservoir of genetic diversity pool<sup>13–19</sup>. Five genome sequences of cyanosiphoviruses have been reported with genome size ranging from 30,332 to 105,532 bps<sup>20–21</sup>. Compared to cyanomyoviruses and cyanosiphoviruses, cyanopodoviruses have a relatively conserved genome size ranging from 42,257 to 47,872 bps<sup>11,22–25</sup>.

Genome sequencing of marine cyanophages has shown that many marine cyanophages encode photosynthesis genes. All the isolated cyanomyoviruses and more than half of the isolated cyanopodoviruses were detected to contain the key photosystem II reaction centre gene *psbA* in their genomes<sup>11,13,17–19,23,26–29</sup>, while no *psbA* gene was found among the known cyanosiphoviruses<sup>20,30</sup>. Two recent studies showed that 24 of 39 marine cyanopodovirus isolates contained *psbA*<sup>12</sup> and 8 of 12 sequenced cyanopodovirus genomes encoded *psbA*<sup>13</sup>. In these two studies, the frequency of *psbA*-containing podoviruses was estimated based on isolated cyanophages which could be biased by the host used for isolation. Is it possible to quantify the presence of *psbA* in cyanopodoviruses in the ocean using a culture-independent approach? The metagenomic database is a useful tool, however these datasets in the public domain are also limited and may not represent true community composition.

## Results

In this study, we estimated the relative abundance and distribution of *psbA*-containing podoviruses based on the metagenomic data. Our approach is built on a conserved genomic structure of cyanopodoviruses. Cyanopodovirus genome organization can be divided into three parts: structural genes, nucleotide metabolism related genes and some hypothetical genes regions (Fig. 1)<sup>11,18,22–24</sup>. Both the composition and the arrangement of structural genes are conserved. One gene cluster, the “portal-capsid-tail/fiber”, existed in all cyanopodoviruses, as well as in other T7 phages<sup>31</sup>. Interestingly, the *psbA* gene was commonly located at a fixed position within the



**Figure 1** | The structure and organization of cyanopodoviruses and some scaffolds or contigs.

conserved gene cluster “portal-*psbA*-capsid”<sup>11</sup>. Based on this conserved gene cluster, we searched (BLAST) the GOS scaffold database using portal, capsid assembly, *psbA* and major capsid protein (MCP) genes, and successfully retrieved 79 cyanopodoviral scaffolds from the GOS database.


Among the 79 cyanopodovirus scaffolds, 70 contain *psbA* and 9 have no *psbA*. All the MCP sequences (>200 aa) were used to construct the phylogenetic tree. The MCP based phylogeny separated cyanopodoviruses into two major clades (Clade A and B) (Fig. 2), which is consistent with the phylogenetic relationship based on the DNA polymerase gene<sup>10,12,21,32</sup>. Nearly all cyanopodoviruses in Clade B carry the *psbA* gene whereas none of those in Clade A do (Fig. 2). A recent study also illustrated such *psbA* distribution pattern in cyanopodoviruses<sup>12</sup>.

In the Bermuda (BATS) database, 58 Clade B MCP homologs were recruited, but no Clade A MCP was found (Fig. 3A). We recruited 17 Clade B homologs, but no Clade A homologs from the North Pacific (HOT) database (Fig. 3A). In the GOS database, 729 Clade B MCP homologs and 18 Clade A MCP homologs were found (Fig. 3A). Interestingly, 17 of 18 of reads were recruited from the coastal water. It is likely that most of Clade A like sequences are from the podoviruses infecting marine *Synechococcus*<sup>10,33,34</sup>. In the MarineVirome database, 271 Clade B like MCP sequences and 4 Clade A like MCP sequences were detected (Fig. 3A).


## Discussion

Podoviruses in Clade A could be a transitional group between Clade B and other T7-like non-cyanobacterial podoviruses (Fig. 2). Four scaffolds in Clade B do not contain *psbA*, and the *psbA* gene in these four scaffolds might be lost during the evolution. Interestingly, scaffold JCVI\_SCAF\_109662694693 (in Clade B) contains a high light-induced gene (*hli*), but no *psbA*.

Our analysis suggests that Clade A podoviruses only make up a very small proportion of cyanopodoviruses in the surface ocean. In the open ocean, Clade A podoviruses only account for 0.27% and



**Figure 2** | The neighbor-joining tree based on the MCP sequences. The sequences with red color mean scaffolds or cyanophage genomes without *psbA* genes. Values of >50% are shown, and indicate percentage bootstrap support based on 1000 replicates for distance, maximum parsimony (MP) and minimum evolution (ME) analyses in the order of NJ/MP/ME. Scale bar, 0.1 nucleotide substitution per site.



**Figure 3 | Number and distribution pattern of cyanopodoviral major capsid reads in the database.** A, Read counts of major capsid corresponding to Clade A and B in four metagenomic database, BATS, GOS, HOT and MarineVirome. B, Proportion of reads belonging to Clade A in open ocean and coastal water, respectively.

1.12% of all cyanopodoviruses in the GOS and MarineVirome databases, respectively. In the coastal surface water, Clade A podoviruses can make up 8.02% and 14.29% of total cyanopodoviruses in the GOS and MarineVirome databases, respectively (Fig. 3B). Clade A podoviruses were not detected in the two open ocean stations, BATS and HOT. Clade A mainly consists of the *psbA*-lacking podoviruses which infect marine *Synechococcus*<sup>10–12</sup>. Our study suggests that it may be less important for cyanophages in coastal or estuarine environments to carry the *psbA* gene compared to cyanophages in the open ocean. Sullivan and colleagues also suggested a shorter latent period could explain the lack of *psbA* gene as result of shorter infection duration with no need the help of *psbA*<sup>23</sup>.

The metagenomic recruitment based on the unique portal-capsid structure provides a culture-independent survey on the distribution frequency of *psbA*-carrying cyanopodoviruses. However all of the datasets that were analyzed were mainly derived from the surface ocean. Our analysis suggests: 1) *psbA*-carrying cyanopodoviruses are the dominant cyanopodoviruses in the surface ocean; 2) *Synechococcus* podoviruses become relatively more abundant in the coastal water; 3) *psbA* is more important for oceanic cyanopodoviruses than for their coastal counterparts.

## Methods

**Metagenomics.** Four metagenomic databases were used to search homologs in our study: three from the bacterial fraction: the Global Ocean Survey database (GOS)<sup>35</sup>, the Bermuda database (BATS)<sup>36</sup>, the Hawaii Ocean Time-Series (HOT)<sup>37,38</sup>, and one viral fraction database: the MarineVirome<sup>39</sup>. All databases were obtained from the CAMERA website (<http://camera.calit2.net/index.shtml>).

Based on the cyanopodovirus genomic conserved gene cluster “portal-*psbA*-capsid”, we searched (BLAST) the GOS scaffold database using portal, capsid assembly, *psbA* and major capsid protein (MCP) genes using a reciprocal best-hit BLAST strategy but no e-value cutoff limitation (Fig. 1)<sup>40</sup>. The structural genes (portal, MCP or capsid assembly gene) allowed the identification of cyanopodoviruses via searching against the NCBI non-redundant proteins database.

To analyze the occurrence frequency and geographic pattern of cyanopodoviruses in the ocean, we recruited reads from BATS, GOS, HOT and MarineVirome datasets using all MCP sequences from sequenced cyanopodoviral genomes as published in Labrie’s paper<sup>11,13</sup>. Our approach is similar to the methods described by Zhao et al.<sup>40,41</sup>. Briefly, all homologous reads were recruited from binning by e-value cutoff to avoid potential bias, and then each putative hit was extracted and used as a query to search against the NCBI non-redundant proteins database<sup>42</sup>. Metagenomic sequences returned a best-hit which could be used to confirm the classification, and all identified reads are listed in Table S1. The number of recruited reads was not normalized, because the method for sampling is different among all the sites and doesn’t target the viruses. However, there should be no bias for cyanopodoviruses with or without *psbA* gene using any methods for sampling.

**Phylogenetic analyses.** All the MCP sequences (>200 aa) were used to construct the phylogenetic tree. Sequences were aligned using Clustal X and phylogenetic trees were constructed using the neighbour-joining, minimum-evolution and maximum-parsimony algorithms of MEGA software 3.0<sup>42</sup>. The phylogenetic trees were supported by bootstrap for re-sampling test with 1000 replicates.

- Proctor, L. M. & Fuhrman, J. A. Viral mortality of marine bacteria and cyanobacteria. *Nature* **343**, 60–62 (1990).
- Fuhrman, J. A. Marine viruses and their biogeochemical and ecological effects. *Nature* **399**, 541–548 (1999).
- Suttle, C. A. Viruses in the sea. *Nature* **437**, 356–361 (2005).
- Suttle, C. A. & Chan, A. M. Dynamics and distribution of cyanophages and their effect on marine *Synechococcus* spp. *Appl Environ Microbiol* **60**, 3167–3174 (1994).
- Waterbury, J. B. & Valois, F. W. Resistance to co-occurring phages enables marine *Synechococcus* communities to coexist with cyanophages abundant in seawater. *Appl Environ Microbiol* **59**, 3393–3399 (1993).
- Wilson, W. H., Joint, I. R., Carr, N. G. & Mann, N. H. Isolation and molecular characterization of five marine cyanophages propagated on *Synechococcus* sp. strain WH7803. *Appl Environ Microbiol* **59**, 3736–3743 (1993).
- Lu, J., Chen, F. & Hodson, R. E. Distribution, isolation, host specificity, and diversity of cyanophages infecting marine *Synechococcus* spp. in river estuaries. *Appl Environ Microbiol* **67**, 3285–3290 (2001).
- Marston, M. F. & Sallee, J. L. Genetic diversity and temporal variation in the cyanophage community infecting marine *Synechococcus* species in Rhode Island’s coastal waters. *Appl Environ Microbiol* **69**, 4639–4647 (2003).
- Sullivan, M. B., Waterbury, J. B. & Chisholm, S. W. Cyanophages infecting the oceanic cyanobacterium *Prochlorococcus*. *Nature* **424**, 1047–1051 (2003).
- Wang, K. & Chen, F. Prevalence of highly host-specific cyanophages in the estuarine environment. *Environ Microbiol* **10**, 300–312 (2007).
- Labrie, S. et al. Genomes of marine cyanopodoviruses reveal multiple origins of diversity. *Environ Microbiol* **15**, 1356–1376 (2013).
- Dekel-Bird, N. P. et al. Diversity and evolutionary relationships of T7-like podoviruses infecting marine cyanobacteria. *Environ Microbiol* **15**, 1476–1491 (2013).
- Lindell, D., Sullivan, M. B., Johnson, Z. I., Tolonen, A. C., Rohwer, F. & Chisholm, S. W. Transfer of photosynthesis genes to and from *Prochlorococcus* viruses. *Proc Natl Acad Sci USA* **101**, 11013–11018 (2004).
- Clokic, M. R. J. et al. Transcription of a ‘photosynthetic’ T4-type phage during infection of a marine cyanobacterium. *Environ Microbiol* **8**, 827–835 (2005).
- Clokic, M. R. J., Millard, A. D., Mehta, J. Y. & Mann, N. H. Virus isolation studies suggest short-term variations in abundance in natural cyanophage populations of the Indian Ocean. *J MAR BIOL ASSOC UK* **86**, 499–506 (2006).
- Millard, A., Clokic, M. R. J., Shub, D. A. & Mann, N. H. Genetic organization of the *psbAD* region in phages infecting marine *Synechococcus* strains. *Proc Natl Acad Sci USA* **101**, 11007–11012 (2004).
- Millard, A. D., Zwirgmaier, K., Downey, M. J., Mann, N. H. & Scanlan, D. J. Comparative genomics of marine cyanomyoviruses reveals the widespread occurrence of *Synechococcus* host genes localized to a hyperplastic region: implications for mechanisms of cyanophage evolution. *Environ Microbiol* **11**, 2370–2387 (2009).
- Sullivan, M. B., Coleman, M. L., Weigele, P., Rohwer, F. & Chisholm, S. W. Three *Prochlorococcus* cyanophage genomes: signature features and ecological interpretations. *PLoS Biol* **3**, e144 (2005).
- Sullivan, M. B. et al. Genomic analysis of oceanic cyanobacterial myoviruses compared with T4-like myoviruses from diverse hosts and environments. *Environ Microbiol* **12**, 3035–3056 (2010).
- Sullivan, M. B. et al. The genome and structural proteome of an ocean siphovirus: a new window into the cyanobacterial ‘mobilome’. *Environ Microbiol* **11**, 2935–2951 (2009).



21. Huang, S., Wilhelm, S. W., Jiao, N. & Chen, F. Ubiquitous cyanobacterial podoviruses in the global oceans unveiled through viral DNA polymerase gene sequences. *ISME J* **4**, 1243–1251 (2010).
22. Chen, F. & Lu, J. Genomic sequence and evolution of marine cyanophage P60: a new insight on lytic and lysogenic phages. *Appl Environ Microbiol* **68**, 2589–2594 (2002).
23. Sullivan, M. B., Lindell, D., Lee, J. A., Thompson, L. R., Bielawski, J. P. & Chisholm, S. W. Prevalence and evolution of core photosystem II genes in marine cyanobacterial viruses and their hosts. *PLoS Biol* **4**, e234 (2006).
24. Pope, W. H. *et al.* Genome sequence, structural proteins, and capsid organization of the cyanophage Syn5: a “horned” bacteriophage of marine *Synechococcus*. *J Mol Biol* **368**, 966–981 (2007).
25. Raytcheva, D. A., Haase-Pettingell, C., Piret, J. M. & King, J. A. Intracellular assembly of cyanophage Syn5 proceeds through a scaffold-containing procapsid. *J Virol* **85**, 2406–2411 (2011).
26. Mann, N. H., Cook, A., Millard, A., Bailey, S. & Clokie, M. Marine ecosystems: bacterial photosynthesis genes in a virus. *Nature* **424**, 741–741 (2003).
27. Lindell, D., Jaffe, J. D., Johnson, Z. I., Church, G. M. & Chisholm, S. W. Photosynthesis genes in marine viruses yield proteins during host infection. *Nature* **438**, 86–89 (2005).
28. Lindell, D. *et al.* Genome-wide expression dynamics of a marine virus and host reveal features of co-evolution. *Nature* **449**, 83–86 (2007).
29. Chenard, C. & Suttle, C. A. Phylogenetic diversity of sequences of cyanophage photosynthetic gene *psbA* in marine and freshwaters. *Appl Environ Microbiol* **74**, 5317–5324 (2008).
30. Huang, S., Wang, K., Jiao, N. & Chen, F. Genome sequences of siphoviruses infecting marine *Synechococcus* unveil a diverse cyanophage group and extensive phage-host genetic exchanges. *Environ Microbiol* **14**, 540–558 (2012).
31. Seed, K. D. *et al.* Evidence of a dominant lineage of *Vibrio cholerae*-specific lytic bacteriophages shed by cholera patients over a 10-year period in Dhaka, Bangladesh. *MBio* **2**, e00334–10 (2011).
32. Chen, F. *et al.* Diverse and dynamic populations of cyanobacterial podoviruses in the Chesapeake Bay unveiled through DNA polymerase gene sequences. *Environ Microbiol* **11**, 2884–2892 (2009).
33. Zwirgmaier, K. *et al.* Global phylogeography of marine *Synechococcus* and *Prochlorococcus* reveals a distinct partitioning of lineages among oceanic biomes. *Environ Microbiol* **10**, 147–161 (2008).
34. Scanlan, D. J. *et al.* Ecological genomics of marine picocyanobacteria. *Microbiol Mol Biol Rev* **73**, 249–299 (2009).
35. Rusch, D. B. *et al.* The Sorcerer II Global Ocean Sampling expedition: Northwest Atlantic through Eastern Tropical Pacific. *PLoS Biol* **5**, 398–431 (2007).
36. Ghai, R. *et al.* Metagenome of the Mediterranean deep chlorophyll maximum studied by direct and fosmid library 454 pyrosequencing. *ISME J* **4**, 1154–1166 (2010).
37. Frias-Lopez, J. *et al.* Microbial community gene expression in ocean surface waters. *Proc Natl Acad Sci USA* **105**, 3805–3810 (2008).
38. Coleman, M. L. & Chisholm, S. W. Ecosystem-specific selection pressures revealed through comparative population genomics. *Proc Natl Acad Sci USA* **107**, 18634–18639 (2010).
39. Angly, F. E. *et al.* The marine viromes of four oceanic regions. *PLoS Biol* **4**, e368 (2006).
40. Wilhelm, L., Tripp, H. J., Givan, S., Smith, D. & Giovannoni, S. Natural variation in SAR11 marine bacterioplankton genomes inferred from metagenomic data. *Biology Direct* **2**, 27 (2007).
41. Zhao, Y. *et al.* Abundant SAR11 viruses in the ocean. *Nature* **494**, 357–360 (2013).
42. Kumar, S., Tamura, K. & Nei, M. MEGA3: Integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* **5**, 150–163 (2004).

## Acknowledgments

This work was supported by the 973 Program (2013CB955700, 2011CB808800), the 863 Program (2012AA092003) and the NSFC (41376132). Both QZ and RZ were supported by the Fundamental Research Funds for the Central Universities (2013121051 and 2012121052 respectively). FC was supported by the Xiamen University 111 Program and Hanse-Wissenschaftskolleg Fellowship. Professor John Hodgkiss of The University of Hong Kong is thanked for polishing the English in this manuscript.

## Author contributions

Q.Z., N.J. and C.A.S. conceived and designed the experiments; Q.Z., R.Z. and F.C. analyzed the data. All the authors helped in writing the manuscript, discussed the results and commented on the manuscript.

## Additional information

**Supplementary information** accompanies this paper at <http://www.nature.com/scientificreports>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Zheng, Q., Jiao, N., Zhang, R., Chen, F. & Suttle, C. Prevalence of *psbA*-containing cyanobacterial podoviruses in the ocean. *Sci. Rep.* **3**, 3207; DOI:10.1038/srep03207 (2013).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported license. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0>