



OPEN

SUBJECT AREAS:
PROTEASOME
UBIQUITYLATIONReceived
22 August 2014Accepted
3 December 2014Published
5 January 2015Correspondence and
requests for materials
should be addressed to
N.P.D. (nico.
dantuma@ki.se)* These authors
contributed equally to
this work.† Current address:
Department of
Oncology, CIMA,
University of Navarra,
31008 Pamplona,
Spain.

Cdc48-independent proteasomal degradation coincides with a reduced need for ubiquitylation

Daniela Gödderz*, Christian Heinen*, Francesco P. Marchese†, Tilman Kurz, Klára Acs & Nico P. Dantuma

Department of Cell and Molecular Biology, Karolinska Institutet, S-17177 Stockholm, Sweden.

Ubiquitin fusion degradation (UFD) substrates are delivered at the proteasome by a handover mechanism involving the ubiquitin-selective chaperone Cdc48 and the ubiquitin shuttle factor Rad23. Here, we show that introduction of a 20 amino acid peptide extension not only rendered degradation independent of Cdc48, in line with the model that this chaperone is involved in early unfolding events of tightly folded substrates, but at the same time relieved the need for efficient polyubiquitylation and the ubiquitin shuttle factor Rad23. Removal of the ubiquitylation sites in the N-terminal UFD signal made the degradation of this substrate strictly dependent on the peptide extension and also on Cdc48 and, importantly the presence of a functional ubiquitylation machinery. This suggests that the extension in the absence of N-terminal ubiquitylation sites is not properly positioned to engage the unfoldase machinery of the proteasome. Thus the need for efficient ubiquitylation and Cdc48 in facilitating proteasomal degradation are tightly linked but can be bypassed in the context of UFD substrates by the introduction of an unstructured extension. Our data suggest that polyubiquitin-binding complexes acting upstream of the proteasome, rather than the proteasome itself, can be primary determinants for the level of ubiquitylation required for protein degradation.

Conjugation of ubiquitin chains to lysine residues is a post-translational modification that is best known for its decisive role in proteasomal degradation¹. These chains are conjugated to proteins designated for degradation through the concerted action of ubiquitylation enzymes². In sharp contrast to the topological diversity in ubiquitin chains that can target proteins for proteasomal degradation³, a common feature appears to be the presence of ubiquitin chains that consists of at least four ubiquitin monomers. On the contrary, conjugation of a single ubiquitin, a process known as monoubiquitylation, has been implicated in a number of non-proteolytic ubiquitin-dependent processes, including membrane trafficking, DNA repair, and transcriptional regulation⁴.

An attractive explanation for the dependency of proteasomal degradation of folded proteins on polyubiquitylation is the preferential binding of ubiquitin chains to the 19S regulatory particle of the proteasome⁵. The proteasome, a large proteolytic complex that is responsible for the regulated hydrolysis of proteins, contains at least two intrinsic ubiquitin binding subunits, Rpn10⁶ and Rpn13^{7,8}. Using a panel of synthetically ubiquitylated ubiquitin fusion degradation (UFD) substrates, it has been shown that a chain consisting of four Lys48-linked ubiquitin monomers represents the minimal signal for proteasomal degradation, which correlates with a dramatically increased affinity of the substrate for the proteasome particle⁵. However, the observations of exceptional cases in which monoubiquitylation⁹ or multiple-monoubiquitylation¹⁰ suffices for proteasomal degradation, show that ubiquitin monomers can facilitate interaction with the proteasome. Recently, it has been shown that polypeptides consisting of no more than approximately 150 amino acid residues can be targeted for proteasomal degradation by introduction of a ubiquitin monomer whereas larger proteins require polyubiquitin chains suggesting that the complexity of the substrate, and not the ubiquitin binding sites at the proteasome, may dictate the required minimal length of the polyubiquitin chain¹¹. Moreover, proteasomes are able to degrade denatured proteins *in vitro* suggesting that ubiquitin as such is not a prerequisite for processing of substrates by the proteasome¹².

In addition to proteasome subunits, ubiquitin binding sites are also present in many other proteins involved in ubiquitin-dependent proteasomal degradation. For example, the ubiquitin receptors Rad23 and Dsk2, which shuttle substrates to the proteasome, contain ubiquitin binding domains by which they sequester polyubiquitylated proteins¹³. Other important proteins involved in ubiquitin-dependent degradation are the yeast Cdc48 and its metazoan ortholog p97 (also known as valosin-containing protein), which, like the proteasomal ATPases,



belong to the family of AAA-ATPases¹⁴. Cdc48/p97 forms a homo-hexameric ring that associates with several co-factors linking these ATPases to diverse cellular processes, most importantly proteasomal degradation¹⁵, ubiquitin-dependent extraction from chromatin¹⁶, homotypic membrane fusion¹⁷ and macroautophagy^{18,19}. It has been proposed that Cdc48, in concert with its ubiquitin-binding dimeric co-factor Ufd1/Npl4^{20,21}, is part of a pathway comprising a series of ubiquitin-interacting factors that, in a handover mechanism, escorts substrates to the proteasome²².

Although Cdc48^{Ufd1/Npl4} has originally been identified as a complex that is required for the degradation of cytosolic and nuclear UFD substrates¹⁵, most studies have focused on its role in the proteasomal destruction of proteins derived from the endoplasmic reticulum (ER)²³, a process known as ER-associated degradation (ERAD)²⁴. A large body of evidence suggests that Cdc48/p97 mediates the extraction of ERAD substrates from the ER membrane prior to proteasomal degradation in the cytosol^{25,26}. In contrast, its functional significance in the degradation of soluble proteins is less clear. Taking into consideration the unfoldase activity of Cdc48²⁷, and the fact that it acts upstream of proteasomal degradation²², it is tempting to speculate that this chaperone plays a role in initiating the unfolding of substrates prior to their interaction with the proteasome. Since proteasomes require unstructured initiation sites to engage their intrinsic unfoldase machinery²⁸ and stimulate the ATPase activity of the proteasomal AAA-ATPases²⁹, the primary role of Cdc48/p97^{Ufd1/Npl4} may be the generation of loosely folded structures that can be subsequently used by the proteasome to further unravel the substrate³⁰.

Here we show that the need for efficient UFD-mediated polyubiquitylation and the ubiquitin-dependent chaperone Cdc48 can be simultaneously overcome by introduction of a peptide extension that is sufficiently long to act as an unstructured initiation site. Our data underscore the importance of ubiquitylation not only in properly engaging the proteasome but also in facilitating interaction with upstream factors that prepare substrates for subsequent destruction by the proteasome.

Results

To investigate the role of Cdc48 in the degradation of soluble proteasome substrates in the budding yeast *Saccharomyces cerevisiae*, we generated a set of a commonly used and well-characterized UFD substrate³¹ ubiquitin^{Gly76Val}-green fluorescent protein (Ub-GFP) with or without a C-terminal polypeptide extension of 15 or 20 amino acids derived from the V5-His tag (Fig. 1A). Structural prediction algorithms predict the V5-His extension to be mostly random coil³⁰. Furthermore, proteins whose structures have been solved containing V5-His C-terminal extensions do not show secondary structural elements or were unsolvable for the extension^{32,33}. We compared the steady-state levels of these proteins in yeast strains carrying the wild-type Cdc48 allele or the mutant allele *cdc48-6*. Notably, introduction of a 20 amino acid extension resulted in reduced steady-state levels of this substrate in wild-type yeast (Fig. 1B). All three Ub-GFP fusions accumulated upon administration of proteasome inhibitor although we did not reach the same steady-state level for Ub-GFP-20aa within the 2 hr-treatment with proteasome inhibitor consistent with its half-life being shorter than those of Ub-GFP and Ub-GFP-15aa. Even more striking, we found that while each of these fusions displayed low steady-state levels in the presence of wild-type Cdc48 and accumulated upon administration of the proteasome inhibitor MG132, only the Ub-GFP fusion with a 20 amino acid-long extension was efficiently degraded by proteasomes in the *cdc48-6* strain (Fig. 1B). Analysis of the turnover confirmed that all three fusion proteins were degraded in wild-type yeast and stabilized in the presence of proteasome inhibitor (Fig. 1C), whereas Ub-GFP with a 20 amino acid extension was rapidly degraded even in the absence of functional Cdc48 (Fig. 1D). We conclude that the introduction of a

20 amino acid-long unstructured polypeptide enhances degradation and alleviates the need for Cdc48 in proteasomal degradation of a soluble UFD substrate in yeast.

It has been proposed that the ubiquitin shuttle factor Rad23 is important to facilitate the exchange of substrates from the Cdc48 complex to the proteasome^{22,34}. Interestingly, we found that the Ub-GFP fusion carrying a 20 amino-acid extension was also efficiently degraded by the proteasome in yeast lacking Rad23, whereas fusions lacking the extension or containing a 15 amino acid-long extension were more stable in the *rad23Δ* strain. (Fig. 2A, B). We argued that the lack of Rad23-dependency may reflect a change in the role of ubiquitylation in the degradation of this substrate. Therefore, we next explored whether the degradation of the Ub-GFP fusion was still dependent on the ubiquitin ligase Ufd4, which is responsible for the conjugation of polyubiquitin chains to UFD substrates³⁵. Whereas the degradation of Ub-GFP and Ub-GFP-15aa was strictly dependent on this ubiquitin ligase, degradation of the fusion carrying a 20 amino acid-long extension did not require the presence of Ufd4 (Fig. 2C, D). To rule out any sequence specificity given by the extension, we inserted an unrelated extension derived from mRFP of 15 or 20 amino acids length into Ub-GFP. Only the 20-amino acid-long, but not the 15 amino acid-long, extension abrogated the need for UFD-mediated ubiquitylation (Supplementary Fig. S1).

Our observation that Cdc48/Rad23-independent degradation of UFD substrates coincides with abrogation of the need of UFD-mediated ubiquitylation does not exclude the possibility that ubiquitylation by alternative ubiquitin ligases is still required for the degradation of these substrates. To test this, we analyzed degradation of these fusions in a yeast strain carrying a mutant allele of the ubiquitin activase Uba1, which is an essential gene critical for conjugation of ubiquitin³⁶. Ubiquitylation is strongly compromised in this strain as levels of total ubiquitin conjugates are drastically reduced (Supplementary Fig. S2). Interestingly, a C-terminal extension of 20 amino acids supported degradation of the UFD substrate in the *uba1-ts 26* strain (Fig. 3A). Ubiquitin pull-down experiments showed a strong reduction in Ub-GFP fusions carrying large ubiquitin conjugates in the *uba1-ts 26* strain, which are visible as a smear over the high molecular weight range of the Western blot (Fig. 3B). However, we also noted that there was little change and even a tendency to an increase in Ub-GFP species modified with one, two or three ubiquitin moieties, reflecting multiple monoubiquitylation or short ubiquitin chains that do not support proteasomal degradation of UFD substrates⁵. The observed ubiquitylation was dependent on the presence of an N-terminal ubiquitin moiety since no modified species of GFP-20aa could be detected (data not shown). These data suggest that residual ubiquitylation activity in the conditional *uba1-ts 26* mutant strain still allows the generation of low molecular weight ubiquitin conjugates of Ub-GFP. This minimal ubiquitylation activity was only sufficient to target the Cdc48-independent fusion with a 20 amino acid polypeptide extension for degradation but failed to support degradation of the Cdc48-dependent Ub-GFP and Ub-GFP-15aa. This suggests that proteasomes are able to facilitate degradation of minimal ubiquitylated Ub-GFP fusions as long as their degradation does not require Cdc48 which coincides with a requirement for more robust ubiquitylation. Consistent with the idea that some level of ubiquitylation is still required for proteasomal degradation, we found that purified proteasomes were unable to processively degrade recombinant Ub-GFP carrying a 20 amino acid-long extension *in vitro* (Supplementary Fig. S3A). Interestingly, purified proteasomes were able to process the 15 and 20 amino acid-long C-terminal extensions, albeit with low efficiency, but this was independent of the N-terminal ubiquitin moiety (Supplementary Fig. S3B). This suggests that proteasomes have access to the polypeptide extension but in the absence of ubiquitylation fail to process the GFP moiety of the fusion. A role for residual ubiquitylation in the degradation of this fusion is also in line with an earlier study demonstrating that




Figure 1 | C-terminal peptide extension allows proteasomal degradation of a UFD substrate independent of Cdc48. (A) Schematic representation of the different UFD substrates including the amino acid sequence of the extensions. (B) Western blot analysis showing steady-state levels of Ub-GFP, Ub-GFP-15aa and Ub-GFP-20aa in wild-type or *cdc48-6* yeast strain in the absence or presence of proteasome inhibitor MG132 using GFP antibody. β -actin was detected as loading control. (C) Cycloheximide-chase analysis of Ub-GFP, Ub-GFP-15aa and Ub-GFP-20aa turnover in wild-type yeast in the absence and presence of MG132. Quantification of GFP levels of three independent experiments with error bars representing standard error of the mean is shown. Solid lines, untreated; dashed lines, MG132-treated. (D) Same as B but in a *cdc48-6* mutant yeast strain.

only short unfolded proteins can be targeted for proteasomal degradation by monoubiquitylation¹¹.

To have a closer look at the role of ubiquitylation in the N-terminal UFD signal, we next investigated whether the Lys29 and Lys48 residues in the N-terminal ubiquitin moiety, which are the canonical acceptor sites for polyubiquitin chains in UFD substrates³⁵, were required for proteasomal degradation. Interestingly, we found that whereas Ub-GFP and Ub-GFP-15aa were stabilized when the Lys29 and Lys48 residues had been substituted for arginine residues, pro-

teasomal degradation of the Cdc48-independent Ub-GFP-20aa substrate was not altered (Fig. 3C, D). Also an N-terminal ubiquitin in which all lysine residues had been substituted for arginine residues was still proficient to allow degradation of a GFP fusion (Ub^{K0}-GFP) but only when carrying a 20 amino acid extension (Fig. 3E). The Ub^{K0}-GFP-20aa was stabilized by administration of the inhibitor MG132 confirming that degradation of this fusion is still mediated by the proteasome. This shows that in the absence of lysine residues in the N-terminal UFD signal the degradation of the fusion becomes





Figure 2 | C-terminal peptide extensions abrogate the need for UFD-dependent ubiquitylation. (A) Cycloheximide-chase analysis of Ub-GFP, Ub-GFP-15aa and Ub-GFP-20aa turnover in *rad23Δ* yeast strain. Quantification of GFP levels of three independent experiments with error bars representing standard error of the mean is shown. (B) FACS analysis of GFP fluorescence intensities of wild-type or *rad23Δ* yeast cells expressing Ub-GFP, Ub-GFP-15aa or Ub-GFP-20aa. For each construct, intensities of three transformants were determined. $^{**}P < 0.01$ (Student's *t* test). (C) Cycloheximide-chase analysis of Ub-GFP, Ub-GFP-15aa and Ub-GFP-20aa turnover in *ufd4Δ* yeast strain. Quantification of GFP levels of three independent experiments with error bars representing standard error of the mean is shown. (D) FACS analysis of GFP fluorescence intensities of wild-type and *ufd4Δ* yeast cells expressing Ub-GFP, Ub-GFP-15aa or Ub-GFP-20aa. Experiment was performed as in B. $^{***}P < 0.0001$ (Student's *t* test).

strictly dependent on the C-terminal peptide extension even in wild-type yeast.

The Ub^{K0}-GFP fusion still contains a number of lysine residues in the GFP moiety of the fusion. To explore the role of ubiquitylation in the degradation of this fusion we tested the stability of the Ub^{K0}-GFP-20aa fusion in the *uba1-ts 26* strain. Unlike the Ub-GFP-20aa fusion, we found that the fusion lacking lysine residues in the N-terminal ubiquitin was stabilized in the *uba1-ts 26* strain suggesting that Ub^{K0}-GFP-20aa is still targeted for ubiquitin-dependent degradation but

independent of the UFD pathway (Fig. 4A). Consistent with a possible role of UFD-independent ubiquitylation, we found that this substrate was degraded in the absence of Ufd4, but stabilized in a yeast strain lacking the ubiquitin conjugases Ubc4 and Ubc5, which cooperate with different ubiquitin ligases (Supplementary Fig. S4). The C-terminal extension is responsible for the ubiquitin-dependent degradation of the Ub^{K0}-GFP-20aa fusion since the 20 amino acid extension did indeed destabilize GFP in wild-type yeast although not to the same extent as fusions carrying a UFD signal (Fig. 4B). The





decrease in half-life of GFP was a consequence of ubiquitin-dependent degradation since GFP-20aa was stabilized in the *uba1-ts 26* strain (Fig. 4C).

Interestingly, the requirement for wild-type Uba1 to facilitate degradation of the Ub^{K0}-GFP-20aa (Fig. 4A) and GFP-20aa (Fig. 4C) coincided with a restoration of Cdc48-dependency. This supports the positive correlation between the need for efficient ubiquitylation and Cdc48-dependency, which was also observed for Ub-GFP. It also indicates that in the absence of the N-terminal ubiquitylation site (i.e. N-terminal ubiquitin without lysine residues) the 20aa extension does not suffice to bypass Cdc48. This is not unexpected as it has been shown that the functionality of unstructured initiation sites is determined not only by their length but also by their position relative to the ubiquitylation site³⁷. This suggests that a C-terminal polypeptide extension is properly positioned for the proteasome when combined with an N-terminal degradation signal while in case of an internal ubiquitylation site Cdc48 may be required to generate alternative initiation sites, which re-introduces the need for efficient ubiquitylation.

Discussion

The UFD signal has been commonly used to target proteins-of-interest for ubiquitin-dependent proteasomal degradation³⁸. The identification of a number of proteins that are involved in the degradation of UFD substrates, which are collectively referred to as the UFD pathway³⁵, has given important fundamental insights into the molecular mechanisms underlying proteasomal degradation. For

example, the identification of the ubiquitin chain recognition signal⁵, the discovery of the E4 multiubiquitylation enzyme³⁹, insights into substrate unfolding²⁸, the minimal length of proteasome substrates⁴⁰ and the functional significance of the proteasome binding subunit Rpn13^{7,8} are all exclusively or primarily based on observations with UFD substrates. Also in the current study, it is the unique nature of the UFD substrates that allowed us to uncover a basic principle in proteasomal degradation as it enabled us to directly compare the fate of almost identical substrates carrying a single ubiquitin or, alternatively, polyubiquitin chains.

Our study provides an explanation for the central role of the UFD pathway in the degradation of substrates that are structurally different from the engineered substrates that led to the identification of this pathway^{23,41,42}. We propose that a primary role of the UFD pathway is to assure that folded substrates lacking initiation sites are first delivered to Cdc48 before they are handed over to the proteasome. Although we cannot formally exclude the possibility that the 20 amino acid extension introduces conformational changes in the UFD substrate, that may affect downstream factors such as ubiquitylation, Cdc48 and Rad23 in different ways, we consider this possibility unlikely for various reasons. As pointed out above structural prediction algorithms and structural data on V5-tagged proteins support the idea that the V5 tag is largely unstructured. Moreover, we show that ubiquitylation of Ub-GFP fusions is severely impaired in the *uba1ts-26* strain regardless of the presence of the V5 tag suggesting that the difference in the stability of the Ub-GFP-20aa in this strain is unlikely to be attributed to a difference in the




Figure 4 | Degradation of C-terminally extended substrates lacking an N-terminal ubiquitylation site requires Uba1 and Cdc48. (A) Cycloheximide-chase analysis of Ub^{K0}-GFP-20aa turnover in wild-type, *uba1ts-26* and *cdc48-6* yeast cells. Quantification of GFP levels of three independent experiments with error bars representing standard error of the mean is shown. (B) Cycloheximide-chase analysis of GFP and GFP-20aa turnover in wild-type yeast compared to Ub-GFP-20aa. (C) Cycloheximide-chase analysis of GFP-20aa turnover in wild-type, *uba1ts-26* and *cdc48-6* yeast cells.



ubiquitylation as a consequence of the V5 tag. Finally, we obtained similar results with an unrelated peptide extension consistent with the idea that the presence of a peptide extension and not an intrinsic feature specific for the V5 tag is responsible for proteasomal degradation in the absence of efficient ubiquitylation.

It remains to be seen what fraction of endogenous proteasome substrates is degraded dependent on Cdc48 and/or Rad23 but it is noteworthy that *in silico* predictions suggest that many proteins contain unstructured regions fulfilling at least the length requirement of unstructured initiation sites⁴³. However, whether or not unstructured regions can function as initiation sites is not only length- but also position-dependent^{28,37}, which complicates the prediction of proteins that require Cdc48 for their degradation based on structural features. The N-terminal positioning of an uncleavable ubiquitin moiety in UFD substrates, which is known to dictate the preferential site of unfolding⁴⁴ and may hence confine the location for potential initiation sites, may be the main reason why this pathway has been readily and consistently identified as a common requirement for the degradation of this particular class of substrates.

It is tempting to speculate that short ubiquitin chains or monoubiquitylation may allow proteins to bypass the unfoldase action of Cdc48, which will result either in stable proteins that are not subject to proteasomal degradation (in the case of non-proteolytic ubiquitylation) or will force the proteasome to use pre-existing unfolded structures in substrates (in the case of proteolytic ubiquitylation). In this respect, it is also of interest to point out that monoubiquitylation appears to be specifically implicated in proteasomal processing of several precursor proteins, such as the membrane-tethered transcription factors Mga2 and Spt23²¹, the p105 precursor of the transcription factor NF- κ B⁴⁵, and the ubiquitin ligase bridging factor Def1⁴⁶. In each of these cases it results in selective destruction of the C-terminal part of these proteins which share the presence of low complexity sequences that have been proposed to form loop-like structures from which proteasomal processing is started^{47,48}, in a manner that is mechanistically similar to unstructured initiation sites²⁸. The fact that these processing events are targeted by means of monoubiquitylation may be of functional significance since it may bias the proteasome to pre-existing unstructured initiation sites whereas a polyubiquitin modification may allow Cdc48 to generate alternative initiation sites for degradation that will enable the proteasome to processively degrade the protein.

While our data are in line with the model that localization of a protein at the proteasome can be sufficient to facilitate proteasomal degradation⁴⁹, they suggest that this will only apply to proteins that contain unstructured initiation sites. At least one naturally occurring substrate, ornithine decarboxylase, is degraded independent of ubiquitylation in a manner that strictly depends on the presence of an unstructured region⁵⁰, similar to the substrates used in this study. Degradation of ornithine decarboxylase⁵¹ depends not only on a domain that facilitates binding to the proteasome⁵² but also on the presence of an unstructured region that has to be sufficiently long to initiate degradation⁵⁰. The critical requirement for unstructured regions in protein degradation may also explain why some proteins can functionally operate in close proximity to the proteolytic complex without becoming subject to processive degradation⁵³. Indeed, when interacting with protein complexes, proteasomes tend to preferentially degrade those proteins that carry unstructured initiation sites while leaving folded proteins unharmed⁵⁴, which has been shown to be relevant for the stable nature of ubiquitin receptors that transiently interact with the proteasome^{55,56}.

In conclusion, our study shows that the strict requirement for ubiquitylation in the degradation of proteasome substrates does not necessarily have to be attributed to polyubiquitin-dependent binding to the proteasome but may rather reflect involvement of polyubiquitin-specific proteins acting upstream of the proteasome.

Methods

Yeast strains and media. All experiments with wild-type and mutant strains were performed at 30°C. A detailed list of the yeast strains can be found in the supplementary information. Yeast transformed with episomal pYES2 plasmids encoding the indicated substrates were grown in synthetic medium comprising a mix of glucose, raffinose, and galactose as carbohydrate source, and then in synthetic medium containing galactose until mid-log phase. For proteasome inhibition, synthetic medium with 0.1% proline as sole nitrogen source was used and supplemented with 0.003% SDS 3 h before addition of proteasome inhibitor Z-Leu-Leu-Leu-CHO (MG132; ENZO)⁵⁷. Cultures were incubated for 2h with 50 μ M MG132.

Construction of plasmids. The plasmids were generated by introducing the Ub-GFP open reading frame into the yeast expression vector pYES2 (Invitrogen, 2 μ URA3) under the regulation of the galactose-inducible GAL1 promoter⁵⁸. The C-terminal extensions are derived from the V5 epitope tag (15aa: SCRPL ESRGP FEGKP; 20aa: SCRPL ESRGP FEGKP IPNPL) or the mRFP open reading frame (15aa: GGMS SEDVI KEFMR, 20aa: GGMS SEDVI KEFMR FKVRM) respectively by introducing stop codons at the indicated positions by site-directed mutagenesis (Stratagene). All constructs were verified by DNA sequencing.

Western blot analysis. Yeast expressing the indicated reporter substrate under the GAL1 promoter was grown in galactose-containing medium until mid-log phase. Protein synthesis was terminated by adding glucose (final concentration 2%) and cycloheximide (final concentration 0.1 mg/ml). One OD₆₀₀ unit was harvested at the indicated time points and total protein extracts were prepared by lysis and precipitation in 12.5% trichloroacetic acid. Total yeast lysates were separated by SDS-PAGE (NuPAGE, Invitrogen) followed by transfer to nitrocellulose membranes (PROTRAN; Schleicher & Schuell) or PVDF membrane (Millipore). Western blot analysis was performed as described previously⁵⁶ using the following primary antibodies: a mixed monoclonal antibody specific to GFP (Roche) (1 : 2000), a polyclonal anti-GFP antibody (Ab290, Abcam) (1 : 5000), monoclonal antibodies against β -actin (Abcam) (1 : 5000) or against PGK1 (Life technologies) (1 : 5000).

Flow cytometric analysis. Yeast expressing the indicated reporter substrate was grown in synthetic medium containing galactose until mid-log phase and was subsequently subjected to flow cytometric analysis (FACS caliber, Beckton & Dickinson). CellQuest software was used to analyze the data.

Detection of ubiquitylated proteins. Yeast was co-transformed with an expression plasmid for the substrate of interest and an expression plasmid for His-tagged ubiquitin (gift from Dr. Ben Distel, AMC Amsterdam, The Netherlands). Pulldown of 6xHis-ubiquitin modified proteins was performed as described previously⁵⁶.

- Hershko, A. & Ciechanover, A. The ubiquitin system. *Annu. Rev. Biochem.* **67**, 425–479 (1998).
- Pickart, C. M. Mechanisms underlying ubiquitination. *Annu. Rev. Biochem.* **70**, 503–533 (2001).
- Kravtsova-Ivantsiv, Y. & Ciechanover, A. Non-canonical ubiquitin-based signals for proteasomal degradation. *J. Cell Sci.* **125**, 539–548 (2012).
- Hicke, L. Protein regulation by monoubiquitin. *Nat Rev Mol Cell Biol.* **2**, 195–201 (2001).
- Thrower, J. S., Hoffman, L., Rechsteiner, M. & Pickart, C. M. Recognition of the polyubiquitin proteolytic signal. *EMBO J.* **19**, 94–102 (2000).
- Young, P., Deveraux, Q., Beal, R. E., Pickart, C. M. & Rechsteiner, M. Characterization of two polyubiquitin binding sites in the 26 S protease subunit 5a. *J. Biol. Chem.* **273**, 5461–5467 (1998).
- Husnjak, K. *et al.* Proteasome subunit Rpn13 is a novel ubiquitin receptor. *Nature (London)* **453**, 481–488 (2008).
- Schreiner, P. *et al.* Ubiquitin docking at the proteasome through a novel pleckstrin-homology domain interaction. *Nature (London)* **453**, 548–552 (2008).
- Boutet, S. C., Disatnik, M. H., Chan, L. S., Iori, K. & Rando, T. A. Regulation of Pax3 by proteasomal degradation of monoubiquitinated protein in skeletal muscle progenitors. *Cell* **130**, 349–362 (2007).
- Dimova, N. V. *et al.* APC/C-mediated multiple monoubiquitylation provides an alternative degradation signal for cyclin B1. *Nat. Cell Biol.* **14**, 168–176 (2012).
- Shabek, N. *et al.* The size of the proteasomal substrate determines whether its degradation will be mediated by mono- or polyubiquitylation. *Mol. Cell* **48**, 87–97 (2012).
- Driscoll, J. & Goldberg, A. L. Skeletal muscle proteasome can degrade proteins in an ATP-dependent process that does not require ubiquitin. *Proc. Natl. Acad. Sci. USA* **86**, 787–791 (1989).
- Dantuma, N. P., Heinen, C. & Hoogstraten, D. The ubiquitin receptor Rad23: At the crossroads of nucleotide excision repair and proteasomal degradation. *DNA Repair (Amst)* **8**, 449–460 (2009).
- Sauer, R. T. *et al.* Sculpting the proteome with AAA(+) proteases and disassembly machines. *Cell* **119**, 9–18 (2004).
- Ghislain, M., Dohmen, R. J., Levy, F. & Varshavsky, A. Cdc48p interacts with Ufd3p, a WD repeat protein required for ubiquitin-mediated proteolysis in *Saccharomyces cerevisiae*. *EMBO J.* **15**, 4884–4899 (1996).



16. Dantuma, N. P. & Hoppe, T. Growing sphere of influence: Cdc48/p97 orchestrates ubiquitin-dependent extraction from chromatin. *Trends Cell Biol.* **22**, 483–491 (2012).
17. Patel, S. K., Indig, F. E., Olivieri, N., Levine, N. D. & Latterich, M. Organelle membrane fusion: a novel function for the syntaxin homolog Ufe1p in ER membrane fusion. *Cell* **92**, 611–620 (1998).
18. Tresse, E. *et al.* VCP/p97 is essential for maturation of ubiquitin-containing autophagosomes and this function is impaired by mutations that cause IBMPFD. *Autophagy* **6**, 217–227 (2010).
19. Ju, J. S. *et al.* Valosin-containing protein (VCP) is required for autophagy and is disrupted in VCP disease. *J. Cell Biol.* **187**, 875–888 (2009).
20. Meyer, H. H., Shorter, J. G., Seemann, J., Pappin, D. & Warren, G. A complex of mammalian ufd1 and npl4 links the AAA-ATPase, p97, to ubiquitin and nuclear transport pathways. *EMBO J.* **19**, 2181–2192 (2000).
21. Rape, M. *et al.* Mobilization of processed, membrane-tethered SPT23 transcription factor by CDC48(UFD1/NPL4), a ubiquitin-selective chaperone. *Cell* **107**, 667–677 (2001).
22. Richly, H. *et al.* A series of ubiquitin binding factors connects CDC48/p97 to substrate multiubiquitylation and proteasomal targeting. *Cell* **120**, 73–84 (2005).
23. Neuber, O., Jarosch, E., Volkwein, C., Walter, J. & Sommer, T. Ubx2 links the Cdc48 complex to ER-associated protein degradation. *Nat. Cell Biol.* **7**, 993–998 (2005).
24. Vembar, S. S. & Brodsky, J. L. One step at a time: endoplasmic reticulum-associated degradation. *Nat. Rev. Mol. Cell Biol.* **9**, 944–957 (2008).
25. Ye, Y., Meyer, H. H. & Rapoport, T. A. The AAA ATPase Cdc48/p97 and its partners transport proteins from the ER into the cytosol. *Nature (London)* **414**, 652–656 (2001).
26. Ye, Y., Meyer, H. H. & Rapoport, T. A. Function of the p97-Ufd1-Npl4 complex in retrotranslocation from the ER to the cytosol: dual recognition of nonubiquitinated polypeptide segments and polyubiquitin chains. *J. Cell Biol.* **162**, 71–84 (2003).
27. Rothballer, A., Tzvetkov, N. & Zwickl, P. Mutations in p97/VCP induce unfolding activity. *FEBS Lett.* **581**, 1197–1201 (2007).
28. Prakash, S., Tian, L., Ratliff, K. S., Lehotzky, R. E. & Matouschek, A. An unstructured initiation site is required for efficient proteasome-mediated degradation. *Nat. Struct. Mol. Biol.* **11**, 830–837 (2004).
29. Peth, A., Kukushkin, N., Bosse, M. & Goldberg, A. L. Ubiquitinated proteins activate the proteasomal ATPases by binding to Usp14 or Uch37 homologs. *J. Biol. Chem.* **288**, 7781–7790 (2013).
30. Beskow, A. *et al.* A conserved unfoldase activity for the p97 AAA-ATPase in proteasomal degradation. *J. Mol. Biol.* **394**, 732–746 (2009).
31. Neefjes, J. & Dantuma, N. P. Fluorescent probes for proteolysis: Tools for drug discovery. *Nat. Rev. Drug Discov.* **3**, 58–69 (2004).
32. Hu, Y., Sun, Z., Eaton, J. T., Bouloux, P. M. & Perkins, S. J. Extended and flexible domain solution structure of the extracellular matrix protein anosmin-1 by X-ray scattering, analytical ultracentrifugation and constrained modelling. *J. Mol. Biol.* **350**, 553–570 (2005).
33. Shipley, K. *et al.* Structure of a kinesin microtubule depolymerization machine. *EMBO J.* **23**, 1422–1432 (2004).
34. Medicherla, B., Kostova, Z., Schaefer, A. & Wolf, D. H. A genomic screen identifies Dsk2p and Rad23p as essential components of ER-associated degradation. *EMBO Rep.* **5**, 692–697 (2004).
35. Johnson, E. S., Ma, P. C., Ota, I. M. & Varshavsky, A. A proteolytic pathway that recognizes ubiquitin as a degradation signal. *J. Biol. Chem.* **270**, 17442–17456 (1995).
36. Dohmen, R. J. *et al.* An essential yeast gene encoding a homolog of ubiquitin-activating enzyme. *J. Biol. Chem.* **270**, 18099–18109 (1995).
37. Inobe, T., Fishbain, S., Prakash, S. & Matouschek, A. Defining the geometry of the two-component proteasome degron. *Nat. Chem. Biol.* **7**, 161–167 (2011).
38. Johnson, E. S., Bartel, B., Seufert, W. & Varshavsky, A. Ubiquitin as a degradation signal. *EMBO J.* **11**, 497–505 (1992).
39. Koegl, M. *et al.* A novel ubiquitination factor, E4, is involved in multiubiquitin chain assembly. *Cell* **96**, 635–644 (1999).
40. Verhoef, L. G. *et al.* Minimal length requirement for proteasomal degradation of ubiquitin-dependent substrates. *FASEB J.* **23**, 123–133 (2009).
41. Hoppe, T. *et al.* Regulation of the myosin-directed chaperone UNC-45 by a novel E3/E4-multiubiquitylation complex in *C. elegans*. *Cell* **118**, 337–349 (2004).
42. Janiesch, P. C. *et al.* The ubiquitin-selective chaperone CDC-48/p97 links myosin assembly to human myopathy. *Nat. Cell Biol.* **9**, 379–390 (2007).
43. Tompa, P., Prilusky, J., Silman, I. & Sussman, J. L. Structural disorder serves as a weak signal for intracellular protein degradation. *Proteins* **71**, 903–909 (2008).
44. Lee, C., Schwartz, M. P., Prakash, S., Iwakura, M. & Matouschek, A. ATP-dependent proteases degrade their substrates by processively unraveling them from the degradation signal. *Mol. Cell* **7**, 627–637 (2001).
45. Kravtsova-Ivantsiv, Y., Cohen, S. & Ciechanover, A. Modification by single ubiquitin moieties rather than polyubiquitination is sufficient for proteasomal processing of the p105 NF- κ B precursor. *Mol. Cell* **33**, 496–504 (2009).
46. Wilson, M. D. *et al.* Proteasome-mediated processing of def1, a critical step in the cellular response to transcription stress. *Cell* **154**, 983–995 (2013).
47. Piwko, W. & Jentsch, S. Proteasome-mediated protein processing by bidirectional degradation initiated from an internal site. *Nat. Struct. Mol. Biol.* **13**, 691–697 (2006).
48. Rape, M. & Jentsch, S. Taking a bite: proteasomal protein processing. *Nat. Cell Biol.* **4**, E113–116 (2002).
49. Janse, D. M., Crosas, B., Finley, D. & Church, G. M. Localization to the proteasome is sufficient for degradation. *J. Biol. Chem.* **279**, 21415–21420 (2004).
50. Takeuchi, J., Chen, H. & Coffino, P. Proteasome substrate degradation requires association plus extended peptide. *EMBO J.* **26**, 123–131 (2007).
51. Bercovich, Z., Rosenberg-Hasson, Y., Ciechanover, A. & Kahana, C. Degradation of ornithine decarboxylase in reticulocyte lysate is ATP-dependent but ubiquitin-independent. *J. Biol. Chem.* **264**, 15949–15952 (1989).
52. Murakami, Y. *et al.* Ornithine decarboxylase is degraded by the 26S proteasome without ubiquitination. *Nature (London)* **360**, 597–599 (1992).
53. Heessen, S., Masucci, M. G. & Dantuma, N. P. The UBA2 Domain Functions as an Intrinsic Stabilization Signal that Protects Rad23 from Proteasomal Degradation. *Mol. Cell* **18**, 225–235 (2005).
54. Prakash, S., Inobe, T., Hatch, A. J. & Matouschek, A. Substrate selection by the proteasome during degradation of protein complexes. *Nat. Chem. Biol.* **5**, 29–36 (2009).
55. Fishbain, S., Prakash, S., Herrig, A., Elsasser, S. & Matouschek, A. Rad23 escapes degradation because it lacks a proteasome initiation region. *Nat. Commun.* **2**, 192 (2011).
56. Heinen, C., Acs, K., Hoogstraten, D. & Dantuma, N. P. C-terminal UBA domains protect ubiquitin receptors by preventing initiation of protein degradation. *Nat. Commun.* **2**, 191 (2011).
57. Liu, C., Apodaca, J., Davis, L. E. & Rao, H. Proteasome inhibition in wild-type yeast *Saccharomyces cerevisiae* cells. *Biotechniques* **42**, 158, 160, 162 (2007).
58. Heessen, S., Dantuma, N. P., Tessarz, P., Jellne, M. & Masucci, M. G. Inhibition of ubiquitin/proteasome-dependent proteolysis in *Saccharomyces cerevisiae* by a Gly-Ala repeat. *FEBS Lett.* **555**, 397–404 (2003).

Acknowledgments

We thank Drs. Stefan Jentsch, Jürgen Dohmen, and Ben Distel for reagents and strains, Dr. Steven Bergink for technical advice and members of the Dantuma laboratory for helpful suggestions. This work was supported by the Swedish Research Council (NPD), the European Community Network of Excellence RUBICON (Project no LSHC-CT-2005-018683) (NPD) and the Wenner-Gren Foundation (DG).

Author contributions

D.G., C.H., N.P.D. designed the research; D.G., C.H., F.P.M., T.K., K.A. performed the experiments; D.G., C.H., F.P.M., T.K., K.A., N.P.D. analyzed the data; D.G., C.H., N.P.D. wrote the paper.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Gödderz, D. *et al.* Cdc48-independent proteasomal degradation coincides with a reduced need for ubiquitylation. *Sci. Rep.* **5**, 7615; DOI:10.1038/srep07615 (2015).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>