



OPEN

# Distinct type I and type II toxin-antitoxin modules control *Salmonella* lifestyle inside eukaryotic cells

SUBJECT AREAS:

CELLULAR  
MICROBIOLOGY

MOLECULAR BIOLOGY

Damián Lobato-Márquez<sup>1,2</sup>, Inmaculada Moreno-Córdoba<sup>2</sup>, Virginia Figueroa<sup>1</sup>, Ramón Díaz-Orejas<sup>2</sup> & Francisco García-del Portillo<sup>1</sup>

Received

13 December 2014

Accepted

2 March 2015

Published

20 March 2015

Correspondence and requests for materials should be addressed to R. D.-O. (ramondiaz@cib.csic.es) or F.G.P. (fgportillo@cnb.csic.es)

<sup>1</sup>Centro Nacional de Biotecnología-Consejo Superior de Investigaciones Científicas (CNB-CSIC), Darwin, 3. 28049 Madrid, Spain, <sup>2</sup>Centro de Investigaciones Biológicas-CSIC (CIB-CSIC), Ramiro de Maeztu, 9. 28040 Madrid, Spain.

Toxin-antitoxin (TA) modules contribute to the generation of non-growing cells in response to stress. These modules abound in bacterial pathogens although the bases for this profusion remain largely unknown. Using the intracellular bacterial pathogen *Salmonella enterica* serovar Typhimurium as a model, here we show that a selected group of TA modules impact bacterial fitness inside eukaryotic cells. We characterized in this pathogen twenty-seven TA modules, including type I and type II TA modules encoding antisense RNA and proteinaceous antitoxins, respectively. Proteomic and gene expression analyses revealed that the pathogen produces numerous toxins of TA modules inside eukaryotic cells. Among these, the toxins Hok<sub>ST</sub>, LdrA<sub>ST</sub>, and TisB<sub>ST</sub>, encoded by type I TA modules and T4<sub>ST</sub> and VapC2<sub>ST</sub>, encoded by type II TA modules, promote bacterial survival inside fibroblasts. In contrast, only VapC2<sub>ST</sub> shows that positive effect in bacterial fitness when the pathogen infects epithelial cells. These results illustrate how *S. Typhimurium* uses distinct type I and type II TA modules to regulate its intracellular lifestyle in varied host cell types. This function specialization might explain why the number of TA modules increased in intracellular bacterial pathogens.

Toxin-antitoxin modules (hereafter TA) were discovered in bacteria due to their capacity to stabilize plasmids by interfering with the viability of plasmid-free segregants<sup>1–3</sup>. This phenomenon results from differential stability of the toxin (stable) and the antitoxin (unstable). TA modules are composed of two small genes and classified in five types attending to the antitoxin nature and its mode of action. Antitoxins are either small RNAs (type I and III modules) or proteins (type II, IV and V modules). All known toxins are proteins and exhibit activities ranging from RNAses to DNA gyrase inhibitors<sup>4–6</sup>. TA loci abound in microbial genomes<sup>7–10</sup> and are found in archaea<sup>11</sup> and, in free-living, symbiotic and obligate intracellular bacteria<sup>4,5</sup>.

Toxins encoded by TA modules trigger mainly bacteriostatic effects<sup>12–14</sup>. These toxins are implicated in processes as phage abortive infection<sup>15,16</sup>; survival in response to nutrient starvation<sup>17–20</sup> or to oxidative damage<sup>21,22</sup>; biofilm formation<sup>23–25</sup>; and, tolerance to antimicrobial drugs<sup>14,26</sup>. Growth arrest caused by TA modules leads to selection of persisters, which are rare slow-growing or dormant cells that normally exist in populations of actively growing cells<sup>14,26</sup>. This selection occurs stochastically with no associated heritable genetic alteration.

TA modules have been mostly characterized in bacteria growing in axenic cultures<sup>14,26,27</sup>. In addition, recent studies implicate TA modules in virulence. Thus, uropathogenic *Escherichia coli* and *Haemophilus influenzae* use type II TA modules to colonize and survive in animal organs<sup>28–30</sup>. *Mycobacterium tuberculosis* up-regulates genes encoding type II TA modules inside macrophages<sup>8</sup>. A type II TA module termed *sehAB*, homolog of the *higBA/relBE* type II TA module, is required for survival of *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) in mice<sup>31</sup>. Due to the capacity of TA modules to arrest bacterial growth, recent studies have focused in understanding whether TA modules contribute to formation of dormant cells in chronic and persistent infections<sup>32</sup>. Bacterial pathogens that cause these types of infections contain more TA modules than non-pathogenic species that are related phylogenetically<sup>8,33,34</sup>. Using a macrophage infection model, a recent study reported impaired generation of non-growing *S. Typhimurium* cells in mutants lacking each of the 14 type II TA modules that were tested<sup>32</sup>. An equal contribution of such a large number of TA modules to arrest bacterial growth during infection is, however, intriguing.




*S. enterica* is an intracellular bacterial pathogen associated to persistent infections in humans and livestock<sup>35</sup>. The serovar Typhimurium has been extensively studied in murine models in which the pathogen causes either acute<sup>36</sup> or chronic infections<sup>37,38</sup>. In the animal, *S. Typhimurium* shows limited proliferation inside macrophages<sup>39</sup>. This pathogen also attenuates growth in cultured fibroblasts<sup>40</sup> and in non-phagocytic cells of the intestinal lamina propria<sup>41</sup>. The study reported here includes a comprehensive analysis of *S. Typhimurium* TA modules and shows that a selected group of toxins encoded by these modules might have evolved to control bacterial survival inside host cells. Besides this specialization of functions, our data also implicate for the first time toxins encoded by type I TA modules in promoting pathogen survival in the infected eukaryotic cell.

## Results

**S. Typhimurium has a large number of TA modules.** Our first aim was to identify every putative TA module in the genome of the *S. Typhimurium* virulent strain SL1344 ([http://www.ncbi.nlm.nih.gov/genome/152?genome\\_assembly\\_id=23044](http://www.ncbi.nlm.nih.gov/genome/152?genome_assembly_id=23044)). We used the database described by Fofo et al.<sup>7</sup> and the web resource TADB (<http://bioinformml.sjtu.edu.cn/TADB/>)<sup>9</sup> to search for type I and type II TA modules, respectively. Twenty-four TA loci, accounting for five type I and 19 type II TA modules, were identified (Fig. 1a, Table 1, Supplementary Table S1). Other available tools that predict type II

TA modules such as RASTA (<http://genoweb1.irisa.fr/duals/RASTA-Bacteria/>) did not identify additional hits. PSI-BLAST was also carried out using as queries validated toxins and antitoxins described for TA modules of distinct types (Supplementary Table S2). This PSI-BLAST identified three new putative TA modules, two type I loci (*hok-sok*<sub>ST</sub> and *symER*<sub>ST</sub>) and one type II locus (*pasTI*<sub>ST</sub>) (Fig. 1a, Table 1, Supplementary Table S1). Recent massive analyses performed for type II<sup>10</sup> and type III<sup>42</sup> TA modules were also examined although no new hit was found in the SL1344 genome.

To keep consistently with previous TA nomenclature, we assigned original names to those TA modules displaying high sequence homology to those previously characterized at the functional level in *S. Typhimurium* and other bacteria (e.g. *ccdAB*<sub>ST</sub>, *phd-doc*<sub>ST</sub>, *vapBC*<sub>ST</sub>, *relBE*<sub>ST</sub>, *shpAB*<sub>ST</sub>). Strikingly, most of the genes we identified as encoding putative type I TA modules were not annotated as genes in the *S. Typhimurium* SL1344 genome. This lack of annotation might be related to the marked small size of the toxin and antitoxin genes. Module size ranges from ~50 to ~150 nucleotides in length for the five type I TA modules identified in our study, which are listed in Table 1 with their respective genome coordinates. We also assigned original names to those proteins carrying functional domains of known toxins but not displaying high amino acid sequence similarity with described TA modules. This was the case of proteins encoded by the *relBE3*<sub>ST</sub>, *relBE4*<sub>ST</sub> and *higBA*<sub>ST</sub> loci (Supplementary Table S1). In the case of the type II TA modules



**Figure 1 | TA modules identified in *S. Typhimurium* strain SL1344.** (a) Workflow depicting the *in silico* analyses that led to the identification of 27 putative TA modules in *S. Typhimurium* strain SL1344. The study of Fofo et al.<sup>7</sup> is focused in type I modules whereas that of Shao et al.<sup>9</sup> describes the TADB web resource that compiles type II modules from different organisms; (b) distribution of the 27 TA loci identified encoding seven type I (orange) and 20 type II (red) modules in the chromosome and plasmids of strain SL1344; (c) conservation of the 27 TA loci identified in *S. Typhimurium* SL1344 in the genome of other *Salmonella* species and *S. enterica* serovars. Homolog search was performed using the tBLASTn tool and as query the protein sequences of toxins and antitoxins identified in *S. Typhimurium* SL1344. The results of this search were contrasted with recent data reported by Nuccio and Baumberg<sup>44</sup>, which compared genome sequences of *S. enterica* serovars causing gastrointestinal and extraintestinal pathologies. Abbreviations: STm-SL1344, *S. Typhimurium* SL1344; STm-LT2, *S. Typhimurium* LT2; SNew, *S. Newport* SL254; SHei, *S. Heildeberg* SL476; SSch, *S. Schwarzengrund* CVM19633; SAgo, *S. Agona* SL483; SEnt, *S. Enteritidis* P125109; SPB, *S. Paratyphi B* SP87; STY1, *S. Typhi* CT18; STY2, *S. Typhi* Ty2; SPA1, *S. Paratyphi A* ATCC9150; SPA2, *S. Paratyphi A* AKU\_12601; SDub, *S. Dublin* CT\_02021853; SGal, *S. Gallinarum* 287/91; SCho, *S. Choleraesuis* SC-B67; SPC, *S. Paratyphi C* RKS4594; *S. bongori*-1, strain N268-08; *S. bongori*-2, strain NCTC12419. Colour code: green, both T and A homologs identified; blue, only T homolog identified; yellow: only A homolog identified; black, no homolog identified. Only hits with e-values  $\leq 10^{-5}$  and pairing encompassing a minimum of 30 amino acids were considered significant.



**Table 1 | TA loci identified in the genome of *S. Typhimurium* strain SL1344**

TA locus (*)	TA module type	Coordinates/ gene ID (†)	Identification method/Reference
<i>hok-sok</i> <sub>ST</sub> (‡)	I	41921–42073	PSI-BLAST
<i>ldrB-rdlB</i> <sub>ST</sub>	I	466721–466936	7
<i>ibsB-sibB</i> <sub>ST</sub>	I	2211602–2211658	7
<i>ibsA-sibA</i> <sub>ST</sub>	I	3383044–3383103	7
<i>ldrA-rdlA</i> <sub>ST</sub>	I	3829510–3829724	7
<i>tisB-istR</i> <sub>ST</sub>	I	4019333–4019842	7
<i>symE</i> <sub>ST</sub>	I	SL4454 ( <i>symE</i> <sub>ST</sub> )	PSI-BLAST
<i>relBE</i> <sub>ST</sub>	II	SL1480–SL1479	9
<i>ta1</i> <sub>ST</sub>	II	SL2380–SL2379	9
<i>pasT</i> <sub>ST</sub>	II	SL2659–SL2658	PSI-BLAST
<i>ta2</i> <sub>ST</sub>	II	SL2885–SL2884	9
<i>parDE</i> <sub>ST</sub>	II	SL2935–SL2936	9
<i>vapBC</i> <sub>ST</sub>	II	SL3012–SL3011	9
<i>ta3</i> <sub>ST</sub>	II	SL3437–SL3438	9
<i>dinJ-yafQ</i> <sub>ST</sub>	II	SL3484–SL3483	9
<i>phd-doc</i> <sub>ST</sub>	II	SL3525–SL3524	9
<i>ta4</i> <sub>ST</sub>	II	SL3618–SL3617	9
<i>relBE3</i> <sub>ST</sub>	II	SL3744–SL3743	9
<i>higBA</i> <sub>ST</sub>	II	SL3866–SL3867	9
<i>sehAB</i> <sub>ST</sub>	II	SL3976–SL3977	9
<i>sehCD</i> <sub>ST</sub>	II	SL3979–SL3980	9
<i>ta5</i> <sub>ST</sub>	II	SL4254–SL4253	9
<i>relBE2</i> <sub>ST</sub>	II	SL4379–SL4380	9
<i>shpAB</i> <sub>ST</sub>	II	SL4459–SL4460	9
<i>ccdAB</i> <sub>ST</sub> (‡)	II	PSLT027–PSLT028	9
<i>vapBC2</i> <sub>ST</sub> (‡)	II	PSLT107–PSLT106	9
<i>relBE4</i> <sub>ST</sub> (‡)	II	SLP2_0004–SLP2_0003	9

(\*) Subscript ST is added for clarification purposes to refer exclusively to TA modules characterized functionally in *S. Typhimurium* in this study.

(†) Coordinates and gene ID of *S. Typhimurium* strain SL1344 based on annotations and sequences deposited in NCBI with entries NC\_016810.1 (chromosome), NC\_017720.1 (plasmid 1, pSLT), and NC\_017718.1 (plasmid 2, pCol1B9). Toxin-encoding genes are underlined.

(‡) TA loci mapping in plasmids.

named *sehAB*<sub>ST</sub> and *sehCD*<sub>ST</sub> by De la Cruz et al.<sup>31</sup>, we maintained that terminology although they are highly homologous to the *higBA*/*relBE* modules. Lastly, those unknown TA modules for which we found no homologs in the literature or in databases were referred as “TA-(number)-<sub>ST</sub>”, denoting their first identification in *S. Typhimurium*. We also provide the exact genome coordinates for these novel type II TA modules. Tables 1 and Supplementary Table S1 depict the complete list of TA modules identified in our study and their relevant features.

Most of the 27 TA modules predicted in *S. Typhimurium* strain SL1344 have a chromosomal location (Fig. 1b). Four of the 27 TA loci map in two of the three plasmids that the strain SL1344 bears. A *vapBC*<sub>ST</sub> paralog (here referred as *vapBC2*<sub>ST</sub>) and a *ccdAB*<sub>ST</sub> homolog map in the pSLT virulence plasmid whereas *hok-sok*<sub>ST</sub> and *relBE4*<sub>ST</sub> locate in the pCol1B9 plasmid (Table 1, Fig. 1b). Altogether, these data showed an unsuspected large repertoire of putative TA loci in *S. Typhimurium*, about double of those reported for this pathogen in recent studies<sup>31,32</sup>. This large number of TA modules led us to hypothesize about distinct TA modules that could contribute to the fitness of the pathogen inside eukaryotic cells.

**Acquisition of TA modules has been favoured in pathogenic *Salmonella* species.** The genus *Salmonella* is composed by the pathogenic species *S. enterica* and the non-pathogenic species *S. bongori*<sup>43</sup>. We reasoned that some of the TA modules, if specialised for contributing to the fitness of intracellular bacteria, might show a narrow distribution restricted to pathogenic bacteria. This specialization of TA modules could also occur among to *S. enterica* serovars associated to gastrointestinal or systemic diseases. To test

this, we used as reference the recent study by Nuccio and Baumler<sup>44</sup>, who compared gene content in *S. enterica* serovars that cause either gastrointestinal or systemic pathologies in human and livestock. Those genes encoding TA modules not included in that genome comparative study were used as query in NCBI databases using tBLASTn. Strikingly, up to 17 of the 27 TA loci identified in *S. Typhimurium* SL1344 are absent in the two strains of the non-pathogenic species *S. bongori* with genome sequence available (Fig. 1c). Another prominent feature is the presence of some TA modules in all *S. enterica* serovars whereas other TA modules display a narrow distribution in only a few serovars. This is the case of the chromosomal type I TA module *ibsA-sibA*<sub>ST</sub>, restricted to serovar Typhimurium (Fig. 1c); or, *ibsB-sibB*<sub>ST</sub>, which is present in serovars Typhimurium, Newport, Agona, Typhi, Paratyphi A, Paratyphi B, Paratyphi C and Choleraesuis (Fig. 1c). No TA module was found to be restricted to serovars associated to either gastrointestinal or systemic infections (Fig. 1c). Distribution of the plasmid-encoded TA modules *ccdAB*<sub>ST</sub>, *vapBC2*<sub>ST</sub>, *hok-sok*<sub>ST</sub> and *relBE4*<sub>ST</sub> was limited to *S. Typhimurium* and other serovars causing system infections such as *S. Choleraesuis* and *S. Paratyphi C* (Fig. 1c). Taken together, these data suggest that acquisition of TA modules might have been favoured in *S. enterica* respect the non-pathogenic species *S. bongori*. Some of these TA modules might also have evolved differently among pathogenic *S. enterica* serovars. This scenario is reminiscent to that reported for other bacterial pathogens as *Mycobacterium tuberculosis*, which has more TA modules than environmental non-pathogenic mycobacteria<sup>34</sup>.

### The TA loci identified in *S. Typhimurium* encode multiple toxins with anti-proliferative activity.

Anti-proliferative and neutralizing activities were next tested for predicted toxins and antitoxins, respectively. Genes encoding toxins and antitoxins of type II TA modules and toxins of type I TA modules were cloned in compatible expression vectors suitable to control expression of each component independently (see Methods). These functional assays were performed in the natural host, *S. Typhimurium* strain SL1344. Thirteen out of the 20 type II TA modules predicted in the genome of strain SL1344 behaved as *bona fide* TA modules (Fig. 2a). Thus, toxins encoded by these thirteen type II TA modules decrease bacterial cultivability, which is restored upon antitoxin co-expression (Fig. 2a). Tests in the putative type I TA modules showed that five of the seven predicted toxins, *Hok*<sub>ST</sub>, *IbsA*<sub>ST</sub>, *LdrA*<sub>ST</sub>, *LdrB*<sub>ST</sub>, *TisB*<sub>ST</sub>, impact negatively bacterial growth (Fig. 2b). To our knowledge, these assays are the first demonstrating anti-proliferative activity for some novel toxins predicted here for the first time and for others previously annotated as putative toxins. These novel and validated toxins include *T1*<sub>ST</sub>, *T2*<sub>ST</sub>, *T4*<sub>ST</sub>, *T5*<sub>ST</sub>, *RelE3*<sub>ST</sub>, *RelE4*<sub>ST</sub>, *Hok*<sub>ST</sub>, *IbsA*<sub>ST</sub>, *LdrA*<sub>ST</sub>, *LdrB*<sub>ST</sub> and *TisB*<sub>ST</sub> (Table 1, Fig. 2a, 2b). Of interest, some toxins of the type II modules that behave functional in our assays were not as such in previous studies with *S. Typhimurium* grown in liquid culture<sup>31</sup>. This is the case of *RelE*<sub>ST</sub>, *RelE2*<sub>ST</sub>, *RelE3*<sub>ST</sub>, *SehAB*<sub>ST</sub>, *SehCD*<sub>ST</sub>, and *VapC2*<sub>ST</sub> (Table 1, Fig. 2a). We also found type II TA modules that did not obey the toxicity-neutralization rule, differentiating two groups: i) antitoxins that do not neutralize their cognate toxins, e.g. *A1*<sub>ST</sub> and *RelB3*<sub>ST</sub> (Fig. 2a); and, ii) toxins that do not have clear effects in cultivability as *T3*<sub>ST</sub>, *HigB*<sub>ST</sub>, *CcdB*<sub>ST</sub>, *PasT*<sub>ST</sub>, *VapC*<sub>ST</sub>, *IbsB*<sub>ST</sub>, and *SymE*<sub>ST</sub> (Fig. 2a). Three independent cloning experiments led to identical results. Production of these ‘non-functional’ toxins following inducer addition was confirmed by either Commassie staining or Western blotting (Supplementary Fig. S1). A lack of function that brought our attention was that of *CcdB*<sub>ST</sub>, one of the two type II TA systems that strain SL1344 bears in the virulence pSLT plasmid (Table 1, Fig. 1b). When compared to the *CcdB* of *E. coli* plasmid F, we noticed an R99W amino acid substitution in the *S. Typhimurium* *CcdB*<sub>ST</sub> homolog. Residue R99




is reported to be crucial for toxicity in *E. coli*<sup>45</sup> and an artificial reversion of the R99W mutation in *S. Typhimurium* restores toxin functionality (Supplementary Fig. S2a). Similarly, PasT<sub>ST</sub> of *S. Typhimurium* shows five out of ten changes in the first ten amino acids when compared to the *E. coli* counterpart. These residues are required for toxin activity<sup>28</sup>. Replacement of this region in PasT<sub>ST</sub> by the *E. coli* PasT sequence partially restored anti-proliferative activity (Supplementary Fig. S2b). These results demonstrated the presence in *S. Typhimurium* of an unprecedented large number of active toxins encoded by TA modules. In addition, our data indicate that a few TA modules of *S. Typhimurium* might be diverging and losing some properties respect functional homolog alleles of closely related bacteria.

***S. Typhimurium* produces inside eukaryotic cells bona fide toxins encoded by type II TA modules.** TA modules are in larger numbers in pathogenic versus non-pathogenic *Salmonella* species (Fig. 1). This difference led us to hypothesize that the acquisition of new TA modules could be associated to the emergence of *S. enterica* as an intracellular bacterial pathogen. Given the involvement of TA modules in the generation of growth-arrested cells, we sought to determine whether toxins encoded by these modules control the capacity of *S. Typhimurium* to adapt to distinct intracellular

lifestyles within defined host cell types<sup>32,46</sup>. Production of components of TA modules by intracellular *S. Typhimurium* was confirmed by highly sensitive gel-free proteomics applied to bacteria directly isolated from human fibroblasts, in which intracellular bacteria show limited proliferation. Despite the small size of toxin and antitoxins (average of ~100 amino acids), high-resolution mass spectrometry identified proteins of some functional type II TA modules in intracellular bacteria. These include the toxins T2<sub>ST</sub>, T4<sub>ST</sub>, T5<sub>ST</sub>, RelE<sub>ST</sub>, Doc<sub>ST</sub> and VapC2<sub>ST</sub> and the antitoxins A2<sub>ST</sub>, A5<sub>ST</sub>, ParD<sub>ST</sub> and VapB2<sub>ST</sub> (Table 2). The same proteins were also detected by proteomics in samples prepared from extracellular bacteria (Table 2). This result led us to hypothesize whether pathogen adaptation to a lifestyle of limited proliferation inside fibroblasts is modulated by changes in the relative levels of toxins and antitoxins encoded by TA modules. To this aim, we generated strains bearing 3xFLAG tags in the 3' end of chromosomal and plasmid genes encoding the toxins identified by proteomics. This procedure was not done for the antitoxin-encoding genes as they map upstream of toxin genes and the tagging could result in polar effects. Western blotting assays showed that extracellular bacteria produce rather unequal amounts of toxins T2<sub>ST</sub>, T4<sub>ST</sub>, T5<sub>ST</sub> and VapC2<sub>ST</sub> (Fig. 3a). Noteworthy, the level of these four toxins increases by ~4- to 13-fold in intracellular bacteria isolated from fibroblasts (Fig. 3b). These values were obtained after correcting by both the amount of the toxin produced by extracellular bacteria and those of the housekeeping membrane protein IgaA detected in extra- and intracellular bacteria (Fig. 3b). The toxin VapC2<sub>ST</sub> exhibited the highest increase in intracellular bacteria, an observation in concordance to the proteomic data (Table 2). Altogether, these analyses demonstrate that inside eukaryotic cells *S. Typhimurium* induces the expression of functional toxins encoded by bona fide type II TA modules.

***S. Typhimurium* up-regulates inside eukaryotic cells functional toxins encoded by type I TA modules.** High-resolution mass spectrometry did not identify toxins of type I modules in protein extracts obtained from either extracellular or intracellular bacteria. These toxins are small peptides of 30–50 amino acids in length, difficult to identify even when using highly sensitive proteomic techniques. To determine if these toxins are also produced by intracellular *S. Typhimurium*, we analysed by quantitative RT-PCR (qRT-PCR) those genes encoding toxins of type I modules that showed anti-proliferative activity: *hok*<sub>ST</sub>, *lbsA*<sub>ST</sub>, *ldrA*<sub>ST</sub>, *ldrB*<sub>ST</sub>, and *tisB*<sub>ST</sub> (Fig. 2b). Intracellular bacteria isolated from human fibroblasts up-regulate three of these genes, *hok*<sub>ST</sub>, *ldrA*<sub>ST</sub> and *tisB*<sub>ST</sub> (Fig. 3c). These data indicate that *S. Typhimurium* up-regulates a selected group of type I TA modules in response to the intracellular environment found in fibroblasts in which bacteria undergo limited proliferation.

***S. Typhimurium* regulates inside eukaryotic cells the production of toxins encoded by type II TA modules.** Next, we reasoned that the high levels of toxins encoded by type II TA modules observed in intracellular bacteria could reflect mechanisms that ensure adaptation to an hostile environment that prevents pathogen proliferation. If this was the case, we expected that the relative levels of toxins T2<sub>ST</sub>, T4<sub>ST</sub>, T5<sub>ST</sub> and VapC2<sub>ST</sub> could vary if intracellular bacteria were displaying different growth rates. This hypothesis was tested with the 3xFLAG-tagged strains using HeLa epithelial cells, in which *S. Typhimurium* proliferates massively<sup>47</sup>. Quantification of toxin levels in actively growing bacteria isolated from HeLa cells showed that the amount of T4<sub>ST</sub> and T5<sub>ST</sub> toxins remain constant compared to extracellular bacteria (Fig. 4). In the case of toxin T2<sub>ST</sub>, its production is however down-regulated by intracellular bacteria (Fig. 4). Of interest, levels of the pSLT plasmid-encoded toxin VapC2<sub>ST</sub> increase notoriously (~7–14 fold) in intracellular bacteria, irrespective of the capacity of the pathogen to proliferate



**Figure 2** | Most toxins and antitoxins predicted in *S. Typhimurium* SL1344 are encoded by bona fide functional TA modules. Toxin and antitoxin genes were cloned under  $P_{lac}$  and  $P_{BAD}$  promoters in compatible plasmids. These genes were expressed either independently or combined in the natural host, i.e. *S. Typhimurium* strain SL1344, by plating bacteria onto plates containing IPTG and/or arabinose. Those TA modules highlighted in either red (type II) or orange (type I) boxes correspond to bona fide modules. (a) type II modules; (b) type I modules. Assays were repeated in a minimum of three independent experiments. ‘None’ means no inducer (IPTG or arabinose) added.



**Table 2 |** Toxins and antitoxins of *S. Typhimurium* TA modules detected by mass spectrometry in bacteria isolated from fibroblasts and in bacteria grown in LB medium

Accession Number	Protein (*)	Mass (Da)	Extracellular (†)			Intracellular (‡)		
			Unique peptides	PSMs	Coverage (%)	Unique peptides	PSMs	Coverage (%)
SL1479	<u>RelE<sub>ST</sub></u>	11,070	3	8	29.79	1	4	9.57
SL1480	<u>RelB<sub>ST</sub></u>	9,086	1	2	23.17	-	-	-
SL2379	<u>A1<sub>ST</sub></u>	13,536	4	19	33.33	-	-	-
SL2380	<u>T1<sub>ST</sub></u>	14,966	4	6	34.62	-	-	-
SL2658	<u>PasI<sub>ST</sub></u>	10,769	3	6	31.25	-	-	-
SL2659	<u>PasI<sub>ST</sub></u>	17,725	2	3	13.29	-	-	-
SL2884	<u>A2<sub>ST</sub></u>	10,673	2	5	28.13	2	2	31.25
SL2885	<u>T2<sub>ST</sub></u>	19,059	2	4	13.14	2	4	20.00
SL2936	<u>ParD<sub>ST</sub></u>	10,369	-	-	-	1	1	7.69
SL3011	<u>VapC<sub>ST</sub></u>	14,930	-	-	-	1	1	21.21
SL3012	<u>VapB<sub>ST</sub></u>	14,276	2	3	13.60	1	2	6.40
SL3437	<u>T3<sub>ST</sub></u>	22,767	11	27	68.00	8	27	47.00
SL3484	<u>DinJ<sub>ST</sub></u>	9,442	2	4	24.42	-	-	-
SL3524	<u>DocG<sub>ST</sub></u>	13,580	1	2	7.38	1	2	10.66
SL3617	<u>T4<sub>ST</sub></u>	17,693	2	5	22.36	1	3	8.07
SL3744	<u>RelB3<sub>ST</sub></u>	11,754	4	6	36.45	-	-	-
SL3976	<u>SehB<sub>ST</sub></u>	15,915	1	2	7.75	-	-	-
SL3979	<u>SehC<sub>ST</sub></u>	12,326	1	1	8.33	-	-	-
SL4253	<u>A5<sub>ST</sub></u>	10,972	1	2	9.28	2	4	23.71
SL4254	<u>T5<sub>ST</sub></u>	17,636	1	1	7.98	1	6	7.98
SL4460	<u>ShpB<sub>ST</sub></u>	11,368	2	3	39.00	-	-	-
SL4459	<u>ShpA<sub>ST</sub></u>	11,289	2	2	22.11	-	-	-
SLP2_0003	<u>RelB4</u>	10,176	1	1	7.87	-	-	-
SLP2_0004	<u>RelE4<sub>ST</sub></u>	10,888	1	4	15.05	-	-	-
PSLT028	<u>CcdB<sub>ST</sub></u>	11,581	3	13	40.59	3	8	40.59
PSLT107	<u>VapB2<sub>ST</sub></u>	8,636	1	2	10.53	2	3	22.37
PSLT106	<u>VapC2<sub>ST</sub></u>	14,879	2	6	21.21	3	7	29.55

(\*) Subscript ST is added for clarification purposes to refer exclusively to TA protein characterized functionally in *S. Typhimurium* (see Fig. 2). Toxins are underlined.

(†) Protein extracts prepared from bacteria grown in LB medium to stationary phase at 37 °C in non-shaking conditions. Two independent experiments were performed. PSM, peptide spectrum match.

(‡) Protein extracts were prepared from non-growing intracellular bacteria at 24 h post-infection of BJ5ta human fibroblasts. Three independent experiments were performed. PSM, peptide spectrum match.

inside the infected cell (Figs. 3b and 4). These data indicate that *S. Typhimurium* regulates differently the production of some toxins encoded by type II TA modules in varied host cell types.

**Only selected groups of type I and type II TA modules impact *S. Typhimurium* fitness inside eukaryotic cells.** The significance of the enhanced toxin production by *S. Typhimurium* inside fibroblasts compared to epithelial cells was examined with isogenic mutants lacking the respective toxins. We inactivated those *bona fide* TA modules displaying anti-proliferative activity (Fig. 2) and that were induced according to RT-qPCR analyses or detected by proteomics (Table 2, Fig. 3). A total of ten toxin-defective mutants were tested in human fibroblasts, in which *S. Typhimurium* show limited proliferation. Control experiments with these toxin mutants discarded polar effects in the genetic constructions (Supplementary Fig. S3). None of them showed a discernable phenotype in the invasion of the eukaryotic cell lines used (Supplementary Fig. S4) or the growth rate in microbiological media (Supplementary Fig. S5). In contrast, the absence of the three type I TA modules tested, Hok<sub>ST</sub>, TisB<sub>ST</sub> and LdrA<sub>ST</sub>, or the toxins T4<sub>ST</sub> and VapC2<sub>ST</sub>, encoded by type II TA modules, impacted negatively the survival of intracellular bacteria inside fibroblasts (Fig. 5a). These toxins were also required for survival of intracellular bacteria in the unrelated rat fibroblast cell line NRK-49F (Supplementary Fig. S6). The  $\Delta$ vapC2<sub>ST</sub> mutant displayed the strongest phenotype with a decrease of up to ~80% in the intracellular survival rate compared to wild-type bacteria (Fig. 5a and Supplementary Fig. S6).


We further tested a multiple mutant lacking the five toxins required for survival of intracellular bacteria ( $\Delta$ hok-sok<sub>ST</sub>  $\Delta$ tisB-istR<sub>ST</sub>  $\Delta$ ldrA-ldrA<sub>ST</sub>  $\Delta$ ta4<sub>ST</sub>  $\Delta$ vapBC2<sub>ST</sub>). This mutant showed a phenotype undistinguishable from individual mutants (Fig. 5a).

These data indicate that this set of toxins could be acting in a coordinated manner to prepare bacteria to cope with host defences. Alternatively, some of these toxins might operate in similar if not identical targets or processes, explaining why we did not observe additive effects. Noteworthy, when the same set of mutants was tested in HeLa epithelial cells, only the  $\Delta$ vapC2<sub>ST</sub> mutant and the quintuple mutant showed a clear phenotype. These two mutants proliferate intracellularly with lower growth rates than wild-type bacteria (Fig. 5b). Taken together, these results differentiated toxins of TA modules that impact positively the fitness of intracellular bacteria showing limited proliferation (case of Hok<sub>ST</sub>, TisB<sub>ST</sub>, LdrA<sub>ST</sub>, and T4<sub>ST</sub>) from other toxins such as VapC2<sub>ST</sub>, required for fitness of *S. Typhimurium* inside the host cell irrespective of the growth rate.


## Discussion

The data reported here increase significantly the current list of TA modules known for *S. Typhimurium* and provide experimental evidence for a vast number of *bona fide* TA modules, more than those recently characterized or predicted in databases<sup>9,31,32,48</sup>. We identified 27 TA modules and 20 toxins encoded by these modules have anti-proliferative activity in its natural host, *S. Typhimurium*. This vast repertoire of 'functional' toxins, which includes members of type I and type II modules, contrasts with the only four functional toxins recognized by De la Cruz et al. based on diminished growth rate when overexpressed in *E. coli*<sup>31</sup>. The proportion of functional toxins characterized here in *S. Typhimurium* is also higher than the reported for *M. tuberculosis*, in which 30 of the 88 predicted and individually-tested TA modules were proved to be functional<sup>8</sup>.

Our study also uncovered apparent 'non-functional' TA modules. Seven toxin candidates did not affect *S. Typhimurium* growth (T3<sub>ST</sub>,




**Figure 3** | *S. Typhimurium* up-regulates inside fibroblasts *bona fide* toxins encoded by type I and type II TA modules. *S. Typhimurium* was chromosomally tagged with 3xFLAG epitope in genes encoding toxins of type II TA modules detected by proteomics (see Table 2). Protein extracts were prepared from intracellular bacteria at 24 h post-infection of BJ5Ta human fibroblasts and extracellular bacteria grown to stationary phase in LB medium. (a) toxins of type II modules detected in extracellular bacteria. Note the distinct relative levels of the toxins examined; (b) synthesis of toxins of type II modules in intracellular bacteria. Note that the four toxins shown (T2<sub>ST</sub>, T4<sub>ST</sub>, T5<sub>ST</sub>, and VapC2<sub>ST</sub>) are produced at higher relative levels inside the eukaryotic cells. The inner membrane IgaA was used as loading control. Numbers below the toxin bands correspond to the relative values determined by densitometry, referred to values measured in extracellular bacteria and corrected by those obtained for the loading control, the IgaA protein; (c) quantitative RT-PCR (qRT-PCR) assays showing the relative expression of five genes encoding toxins of type II modules. Note that three of them displayed significant induction in intracellular bacteria. Data are the means and standard deviations from three independent experiments. \*,  $P = 0.01$  to 0.05; \*\*,  $P = 0.01$  to 0.001, by one-way ANOVA with Dunnett's post-test.



**Figure 4** | Up-regulation of toxins encoded by type II modules by *S. Typhimurium* is restricted to VapC2<sub>ST</sub> when growing inside HeLa epithelial cells. 3xFLAG-tagged *S. Typhimurium* strains were used to infect human HeLa epithelial cells for 16 h. At this time, protein extracts were prepared from intracellular bacteria and tested for toxin protein by Western blotting assay. Numbers below the toxin bands correspond to the relative values determined by densitometry, referred to values measured in extracellular bacteria and corrected by those obtained for the loading control, the IgaA protein. Note that among the four toxins tested only VapC2<sub>ST</sub> is produced by bacteria at higher relative levels inside the eukaryotic cells. The Western blots shown are representative of a total of three independent experiments.

CcdB<sub>ST</sub>, HigB<sub>ST</sub>, PasT<sub>ST</sub>, VapC<sub>ST</sub>, IbsB<sub>ST</sub> and SymE<sub>ST</sub>). Additionally, two putative type II antitoxins did not neutralize their cognate toxins partners (A1<sub>ST</sub> and RelB3<sub>ST</sub>). We confirmed that these proteins were efficiently produced from the respective plasmids, so other factors should explain such negative outcome. For CcdB<sub>ST</sub> and PasT<sub>ST</sub>, which have active orthologs in *E. coli*, site-directed mutagenesis showed that discrete differences in sequence compromise activity in *S. Typhimurium*. This was an unexpected finding that opens new questions regarding the evolution and deterioration (or specialization to other yet unknown functions) of certain TA modules in otherwise rather genetically close bacteria. Lastly, the possibility that a chromosomally-encoded antitoxin could counteract plasmid-encoded toxin cannot be discarded. However, this case seems unlikely given the marked difference in the stability and stoichiometry of the two components. For the 'inactive antitoxins' A1<sub>ST</sub> and RelB3<sub>ST</sub>, a possible alternative is that other components are needed for proper antitoxin function, as it was shown for the HigA antitoxin of *M. tuberculosis*<sup>49</sup>. These two proteins could also not correspond to true antitoxins even considering they map in proximity to the toxin gene. A precedent exists in the *mazEF* TA module of *Myxococcus xanthus*, in which antitoxin and toxin genes map apart one from the other<sup>50</sup>.

To our knowledge, the activity of defined TA modules in controlling growth of an intracellular pathogen inside non-phagocytic host cell types was not investigated at the cellular level before this study. The data obtained in fibroblasts uncover a selected class of type I and type II TA modules (Hok<sub>ST</sub>, TisB<sub>ST</sub>, LdrA<sub>ST</sub>, T4<sub>ST</sub> and VapC2<sub>ST</sub>) that are required for *S. Typhimurium* to survive within this host cell type. Our data provide new information with respect to the recent study of Helaine et al., which analysed type II TA modules in the *S. Typhimurium*-macrophage infection model<sup>32</sup>. First, we demonstrate that toxins encoded by type I TA modules are also as important as those of type II TA modules for the fitness of intracellular *S. Typhimurium*. Second, unlike the work of Helaine et al., which



**Figure 5 | Distinct type I and type II TA modules impact fitness of intracellular *S. Typhimurium* in fibroblasts and epithelial cells.** Shown are the survival rates of *S. Typhimurium* mutants lacking the indicated type I (grey bars) and type II (white bars) TA modules. (a) ratio of viable intracellular bacteria at 24 h and 2 h post-infection in human BJ5ta fibroblasts; (b) ratio of viable intracellular bacteria at 16 h and 2 h post-infection in human HeLa epithelial cells. “ $\Delta$ 5” refers to the  $\Delta$ hok-sok<sub>ST</sub>  $\Delta$ tisB-istR<sub>ST</sub>  $\Delta$ ldrA-ldrA<sub>ST</sub>  $\Delta$ ta4<sub>ST</sub>  $\Delta$ vapBC2<sub>ST</sub> mutant. Data are the means and standard deviations from three independent experiments. \*,  $P = 0.01$  to  $0.05$ ; \*\*,  $P = 0.01$  to  $0.001$ ; \*\*\*,  $P < 0.001$ , by one-way ANOVA with Dunnett’s post-test.

implicated all type II TA modules that they tested (14 modules) in persister cell formation inside macrophages<sup>32</sup>, our data support a clear specialization of functions among distinct type II TA modules. Differences in the host cell type that is invaded by the pathogen might dictate a critical function for distinct sets of TA modules. Thus, among the type II TA modules tested, only the toxins VapC2<sub>ST</sub> and T4<sub>ST</sub> impact fitness of *S. Typhimurium* inside fibroblasts. VapC2<sub>ST</sub> is predicted to have RNase activity due to the functional domain conservation with its paralog toxin VapC<sub>ST</sub><sup>51</sup>, which behaved as a ‘non-functional’ toxin in our assays (Fig. 2). Regarding T4<sub>ST</sub>, a novel toxin encoded by a TA module discovered in this study, it bears a Gcn5-related acetyl transferase (GNAT) domain. This domain is present in members of a protein superfamily that utilizes acyl coenzyme A (CoA) as donor for the acylation of lysine residues<sup>52</sup>. These enzymes have a large variety of substrates, from nascent endogenous proteins to histones or antibiotics<sup>53</sup>. In *S. Typhimurium*, the action of a GNAT domain-containing protein termed Pat has been related to the control of carbon utilization and metabolic flux via acetylation of several metabolic enzymes<sup>54</sup>. The decreased survival inside fibroblasts experienced by the mutant lacking T4<sub>ST</sub> supports an important role of metabolic readjustment(s) in ensuring the fitness of intracellular *S. Typhimurium* under conditions of limited growth. This assumption agrees with genome-wide profiling data obtained in this pathogen under identical infection conditions<sup>41</sup>, which showed up-regulation of metabolic genes involved in adaptation to microaerophilic conditions.

TisB<sub>ST</sub>, LdrA<sub>ST</sub> and Hok<sub>ST</sub> form the subset of toxins encoded by type I TA modules important for fitness of intracellular *S. Typhimurium*. In *E. coli* these three toxins have been related to dissipation of transmembrane potential and ATP depletion<sup>55–57</sup>. A ‘controlled’ action of these toxins in intracellular bacteria could potentially assist *S. Typhimurium* for acquiring the required state of metabolic dormancy in fibroblasts. This model fits with the increased expression of genes encoding Phage-shock proteins (Psp) observed in intracellular *S. Typhimurium*<sup>41,58</sup>. The bacterial

response involving Psp proteins is triggered as a consequence of stress reducing the energy status of the cell. It is tempting to hypothesize about these Psp proteins as elements balancing the limited damage induced by the toxins encoded by TA modules. This idea could be evaluated in future studies.

Among the five toxins that ensure *S. Typhimurium* fitness inside fibroblasts, LdrA<sub>ST</sub>, Hok<sub>ST</sub>, and VapC2<sub>ST</sub> are absent in the non-pathogenic species *S. bongori*. Hok<sub>ST</sub> and VapC2<sub>ST</sub> are encoded by plasmids that are absent in *S. bongori*, however LdrA<sub>ST</sub> is encoded by a chromosomal gene. Noteworthy, *ldrA<sub>ST</sub>* is conserved in all *S. enterica* serovars (Fig. 1), which suggests that acquisition of this particular TA module might be occurred during the evolution and speciation of *S. enterica* as pathogen. Another conclusion of our study involves the different toxins encoded by TA modules that ensure fitness of intracellular *S. Typhimurium* in distinct host cell types. Thus, the behaviour of actively growing *S. Typhimurium* located HeLa epithelial cells is altered exclusively in the absence of the VapC2<sub>ST</sub> toxin. This is an interesting observation since all toxins tested were shown by Western blotting to be produced by intracellular *S. Typhimurium* in both fibroblasts and epithelial cells, although at varied relative amounts. Based on these findings, not only the type of toxins but also the proportion at which are produced by the pathogen inside the eukaryotic cell might balance the final outcome in terms of persistence or active proliferation. This assumption also considers the importance that the coordination among members of such ‘toxin-cocktail’ might have. Lack of a particular toxin in such specialized group could lead to de-regulation of the rest and, as a consequence, to a negative impact on the fitness of intracellular bacteria. This assumption is consistent with two recent studies that demonstrate cross-regulation among distinct TA modules of *E. coli*<sup>59,60</sup>.

Intracellular *S. Typhimurium* is exposed within the phagosome to stresses like acidic pH, oxidative and nitrosative reactive molecules and, high osmolarity<sup>61</sup>. This fact could explain the apparent higher content of TA modules in bacterial pathogens that infect eukaryotic cells compared to environmental bacteria. Nonetheless, we did not



observe any additive effect following the absence of all those toxins of type I and type II modules that impact positively the fitness of intracellular bacteria. This finding could be in concordance with the model supporting a coordinated action of distinct toxin groups. The loss of this hypothetical coordination due the absence of a particular toxin might not change in quantitative terms when additional toxins are lacking. Precedents for this phenomenon exist in the literature. Thus, the two VapBC TA modules encoded by *H. influenzae* contribute to virulence<sup>29</sup>. However, the phenotype of a double mutant lacking both VapBC modules is indistinguishable from those of the individual mutants.

In summary, the data shown here support a marked specialization of defined TA modules for regulating the intracellular lifestyle of bacterial pathogens. These TA modules would be involved in ensuring pathogen intracellular survival, which is critical for the progression of the infection. It is also worthy to note that, to our knowledge, the data shown here implicate for the first time two plasmid-encoded TA modules, *hok-sok*<sub>ST</sub> and *vapBC2*<sub>ST</sub>, in infection. Of interest in future studies will be also to characterize the regulators that alter the relative levels of toxins encoded by some of these TA modules when the pathogen resides inside the eukaryotic cell.

## Methods

**Bacterial strains and plasmids used in this study.** The *S. enterica* serovar Typhimurium and *E. coli* strains together with plasmids used in this study are listed in Supplementary Table S3.

**Identification of TA modules in *S. Typhimurium* SL1344.** Complete nucleotide sequences of the *S. Typhimurium* SL1344 chromosome and the three plasmids carried by this strain were downloaded from NCBI. The respective entries are: NC\_016810.1 (chromosome), NC\_017720.1 (plasmid 1, pSLT), and NC\_017718.1 (plasmid 2, pCol1B9) and NC\_017719.1 (plasmid 3, pRSF1010). A total of 27 TA modules were identified (information compiled in Supplementary Table S1). The TADB resource (<http://bioinfo-mml.sju.edu.cn/TADB/>)<sup>9</sup> and the catalogue published by Fojo et al.<sup>7</sup>, were used to identify type II and type I TA modules, respectively. In the case of the TADB resource, the WU-BLAST 2.0 tool was selected. To upload genome sequences into these resources in an operative manner, the *S. Typhimurium* SL1344 chromosomal sequence was divided in four sections. Three inter-fragment sequences were created to detect possible operons overlapping borders between fragments of consecutive fragments. Plasmids sequences were uploaded un-fragmented. BLOSUM62 was selected as comparison matrix, default parameters for cut-off score and word length, and an e-value of 0.01. To avoid spurious hits, searches were manually cured. Thus, hits were discarded when having more than 650 bp in length, not forming part of an operon (establishing a maximum distance of 150 bp between toxin and antitoxin genes), or previously characterized as genes non-related to TA modules. In a second survey, several already identified toxin and antitoxin sequences (including TA modules of type I, II, III, IV and V) were used as queries to a PSI-BLAST search against genome of *S. Typhimurium* SL1344 (Supplementary Table S2). PSI-BLAST was executed as previously described<sup>62</sup>.

**Cloning of *S. Typhimurium* toxin and antitoxin genes.** The expression vectors pACYC184<sup>63</sup> and pFUS2<sup>64</sup> were used to clone and express toxin and antitoxin genes with the following modifications. A *P*<sub>lac</sub> promoter was obtained from plasmid pNDM220<sup>65</sup> as a *Z*raI/*B*amHI restriction fragment and introduced in the *B*amHI site of pACYC184. An optimized Shine-Dalgarno sequence<sup>66</sup> and a polylinker containing *Z*raI/*S*peI restriction sites (between *B*amHI and *S*all sites), were also introduced. Additionally, *lacIQ* repressor was cloned between *P*siI/*E*coRV sites to ensure proper transcription control. The same Shine-Dalgarno and polylinker were cloned behind the *P*<sub>BAD</sub> promoter of plasmid pFUS2, between *E*coRI and *K*pnl restriction sites. Candidate toxin and antitoxin genes were cloned in either pACYC184- or pFUS2-based expression vectors. Toxin and antitoxin genes were amplified using Pfu polymerase (Promega) using *S. Typhimurium* genomic DNA as template and primers containing *Z*raI/*S*peI restriction sites (Supplementary Table S4). The fragments were ligated (T4 DNA ligase, Promega) 30 min at room temperature (24°C) and transformed in *S. Typhimurium* SL1344 competent cells. All inserts were sequenced to verify a correct construction of the recombinant plasmid.

**Construction of *S. Typhimurium* mutants defective in TA modules.** For disruption of toxin-antitoxin genes, the deletion method described by Maisonneuve et al.<sup>67</sup>, was followed.

**Construction of *S. Typhimurium* recombinant strains expressing 3x-FLAG-tagged toxins.** 3x-FLAG tagging was performed at the 3'-end of toxin genes in their respective chromosomal or plasmidic locations using the procedure of Uzau et al.<sup>68</sup>. Oligonucleotide primers used in these procedures are listed in Supplementary Table S4.

**Functional assays to detect toxin and antitoxin activities.** Toxicity and neutralization assays were developed based on expression of toxin and antitoxin proteins from the compatible vectors pACYC184 and pFUS2 bearing *P*<sub>lac</sub> or *P*<sub>BAD</sub> promoters, respectively. Induction of the desirable protein was carried out with IPTG (isopropyl β-D-1-thiogalactopyranoside) or arabinose. Most toxins were expressed in pFUS2 and its cognate antitoxins in pACYC184. Corresponding pair components of TA modules were co-expressed in *S. Typhimurium* SL1344. Bacteria were grown overnight in LB medium containing the corresponding antibiotic for each vector and supplemented with 0.4% glucose for those cases in which the toxin gene was cloned in pFUS2. This culture was diluted 1 : 100 in fresh LB medium and bacteria grown up to an optical density (OD<sub>600</sub>) of 0.3. At this time, serial dilutions made in phosphate buffered saline (PBS) pH 7.0 were plated onto solid media containing toxin and antitoxin inducers (IPTG and/or arabinose). IPTG and arabinose concentrations were adjusted to optimize visualization of toxin and antitoxin activities.

**Distribution of TA modules among *Salmonella* species and serovars.** To determine the phylogenetic distribution of the 27 TA modules identified in *S. Typhimurium* SL1344, we first examined the comparative genomic study recently reported by Nuccio and Bäumlér<sup>44</sup>. This study analyzed 17 genomes of different *S. enterica* serovars mostly associated to intestinal or extraintestinal infections. In addition, TBLASTN was used: i) to determine the distribution of those TA modules identified in our study that were not listed in the Nuccio and Bäumlér compilation; and, ii) to compare the distribution of all 27 identified TA modules in *S. Typhimurium* strain SL1344 with those present two strains of the non-pathogenic species *S. bongori* whose genome sequence is available (strains N268-08 and NCTC12419). For TBLASTN, only hits with e-values of ≤ 10<sup>-5</sup> with a length of 30 amino acids or over were considered significant.

**Bacterial infection of fibroblasts and epithelial cells.** Foreskin human BJ-5ta fibroblasts (ATCC CRL-4001) and HeLa epithelial cells (ATCC CCL-2) were used. These two cell lines were propagated and infected as described<sup>69</sup>. Number of viable intracellular bacteria was determined by plating (in triplicate) serial dilutions of the cell lysate at the different post-infection times. Proliferation index was calculated as the ratio of colony forming units (cfu) counted at 24 h (fibroblasts) or 16 h (epithelial cells) versus 2 h post-infection. Assays were performed in a minimum of three independent repetitions.

**Isolation of intracellular bacteria for proteomic and Western analyses.** Protocols optimized for the isolation of intracellular *S. Typhimurium* from infected fibroblasts to obtain bacterial RNA and protein have been described elsewhere<sup>44,46</sup>. These large-scale infections were designed with a minimum of 12 BioDish-XL 500-cm<sup>2</sup> plates (ref. 351040, BD Biosciences) per bacterial strain and post-infection time.

**Toxin/antitoxin expression in intracellular bacteria monitored by Western immunoblot.** Isogenic *S. Typhimurium* strains carrying a 3xFLAG tag in defined chromosomal toxin genes were constructed (Supplementary Table S3). These tagged strains were used to infect human BJ-5ta fibroblasts or HeLa epithelial as described<sup>69</sup>. Protein extracts prepared from intracellular bacteria were resolved in SDS-PAGE using 10% polyacrylamide gels and processed for Western blot assays with anti-FLAG antibody (ref. F3165, Sigma-Aldrich). Levels of the *S. Typhimurium* inner membrane protein IgaA were monitored as loading control.

**Proteomic analysis in extra- and intracellular bacteria.** Proteome comparison was performed in bacteria grown overnight in LB medium at 37°C and intracellular bacteria collected at 24 h post-infection from human BJ-5ta fibroblasts. Protein extracts were prepared as described<sup>44,46</sup>, run in SDS-PAGE using 15% acrylamide gels and stained with 'Colloidal blue staining kit' (Invitrogen). These extracts were prepared as three biological replicates for intracellular bacteria and two replicates for extracellular bacteria grown to stationary phase. Samples were run by SDS-PAGE, stained and collected as 1-mm width gel slices. These slices were cut in eight pieces, washed in 50 mM ammonium bicarbonate and 50% acetonitrile (ACN) and dehydrated with ACN (100%). Gel pieces were rehydrated in 50 mM ammonium bicarbonate with 12.5 ng/μl trypsin and incubated overnight at 30°C. Tryptic peptides were extracted at 37°C using ACN 100% and 0.5% trifluoroacetic acid (TFA), dried, cleaned using ZipTip (Millipore) and reconstituted in 5 μl 0.1% formic acid/2% ACN. This peptide mix was loaded into a C18-A1 ASY-Column 2 cm precolumn (Thermo Scientific EASY-Column), eluted with a Biosphere C18 column (75 μm inner diameter, 15 cm long, 3 μm particle size) (NanoSeparations), and finally separated using a 150 min gradient with the last 100 min from 2–35% buffer B. Composition of buffer A was: 0.1% formic acid/2% ACN and that of buffer B: 0.1% formic acid in ACN. Flow-rate was 250 nL/min on a nano EASY-nLC (Thermo Scientific) coupled to a nanoelectrospray ion source (Thermo Scientific). Mass spectra were acquired on the LTQ-Orbitrap Velos (Thermo Scientific, San Jose, CA) in the positive ion mode. Full-scan MS spectra (m/z 300–2000) were acquired in the Orbitrap with a target value of 1,000,000 at a resolution of 30,000 at m/z 400. For internal mass calibration, the 445.120025 ion for lock mass was used. Charge state screening was enabled and precursors with charge state unknown or 1 were excluded. After the survey scan, the ten most intense precursor ions were selected for CID-HCD MS/MS fragmentation. Peptide identification was performed in both CID (Collision Induced Dissociation) and HCD (Higher-Energy Collision Induced Dissociation) spectra. For CID fragmentation the target value was set to 10,000 and normalized collision energy to 35%. For HCD, target value was set to 50,000 and collision energy





was set to 45%. Dynamic exclusion was applied during 30 s. MS data were analyzed with Proteome Discoverer (version 1.3.0.339) (Thermo Fisher, San Jose, CA) using standardized workflows with Sequest search engine. MS/MS spectra were searched against an in-house created FASTA database with the sequences of the toxin and antitoxin protein identified *in silico* in *S. Typhimurium* strain SL1344 (Table 1, Supplementary Table S1). Search parameters included a maximum of two missed cleavages allowed, carbamidomethylation of cysteines as a fixed modification, and oxidation of methionine as variable modifications. Precursor and fragment mass tolerance were set to 10 ppm and 0.8 Da, respectively. Identified peptides were filtered by false discovery rate of 0.01 using Percolator.

**Real-time quantitative PCR.** Large scale infection of human BJ-5ta fibroblasts was performed as described<sup>41</sup>. Total RNA (prokaryotic and eukaryotic RNA) was isolated by addition of 1 ml of Trizol reagent (Invitrogen) following indications of manufacturer and co-precipitated with 20 µg of glycogen (Roche). RNA samples were treated with DNaseI for 1 hour at 37°C (Turbo DNA-free kit Ambion/Applied Biosystems). cDNA libraries were constructed from 1 or 2 µg of RNA obtained from extracellular or intracellular bacteria, respectively (High-capacity cDNA archive kit, Applied Biosystems). For qPCR, the Power Sybr Green PCR master mix (Applied Biosystems) was used in a 10 µl final volume and reactions were carried out in an ABI Prism 7500 equipment following standard reaction conditions according to manufacturer's recommendations. Each cDNA sample was run in triplicate and expression data from each condition were obtained from three independent experiments.

**Statistical analyses and densitometry.** Statistical significance was analyzed with GraphPad Prism v5.0b software (GraphPad Inc.) using one-way analysis of variance (ANOVA) with Dunnett's multiple comparison post-test. A *P* value ≤ 0.05 was considered significant. Densitometry on toxins bands obtained in the Western blotting assays was performed using ImageJ, made available to the public by National Institute of Health, USA.

- Ogura, T. & Hiraga, S. Mini-F plasmid genes that couple host cell division to plasmid proliferation. *Proc. Natl. Acad. Sci. USA* **80**, 4784–4788 (1983).
- Bravo, A., de Torrontegui, G. & Diaz, R. Identification of components of a new stability system of plasmid R1, ParD, that is close to the origin of replication of this plasmid. *Mol. Gen. Genet.* **210**, 101–110 (1987).
- Gerdes, K., Rasmussen, P. B. & Molin, S. Unique type of plasmid maintenance function: postsegregational killing of plasmid-free cells. *Proc. Natl. Acad. Sci. USA* **83**, 3116–3120 (1986).
- Gerdes, K. in *Prokaryotic toxin-antitoxins* (Springer, 2013).
- Hayes, F. & Van Melderen, L. Toxins-antitoxins: diversity, evolution and function. *Crit. Rev. Biochem. Mol. Biol.* **46**, 386–408 (2011).
- Wang, X. *et al.* A new type V toxin-antitoxin system where mRNA for toxin GhoT is cleaved by antitoxin GhoS. *Nat. Chem. Biol.* **8**, 855–861 (2012).
- Fozo, E. M. *et al.* Abundance of type I toxin-antitoxin systems in bacteria: searches for new candidates and discovery of novel families. *Nucleic Acids Res.* **38**, 3743–3759 (2010).
- Ramage, H. R., Connolly, L. E. & Cox, J. S. Comprehensive functional analysis of *Mycobacterium tuberculosis* toxin-antitoxin systems: implications for pathogenesis, stress responses, and evolution. *PLoS Genet.* **5**, e1000767 (2009).
- Shao, Y. *et al.* TADB: a web-based resource for Type 2 toxin-antitoxin loci in bacteria and archaea. *Nucleic Acids Res.* **39**, D606–611 (2011).
- Sberro, H. *et al.* Discovery of functional toxin/antitoxin systems in bacteria by shotgun cloning. *Mol. Cell* **50**, 136–148 (2013).
- Yamaguchi, Y., Park, J. H. & Inouye, M. Toxin-antitoxin systems in bacteria and archaea. *Annu. Rev. Genet.* **45**, 61–79 (2011).
- Yamaguchi, Y. & Inouye, M. Regulation of growth and death in *Escherichia coli* by toxin-antitoxin systems. *Nat. Rev. Microbiol.* **9**, 779–790 (2011).
- Jensen, R. B. & Gerdes, K. Programmed cell death in bacteria: proteic plasmid stabilization systems. *Mol. Microbiol.* **17**, 205–210 (1995).
- Maisonneuve, E. & Gerdes, K. Molecular mechanisms underlying bacterial persisters. *Cell* **157**, 539–548 (2014).
- Fineran, P. C. *et al.* The phage abortive infection system, ToxIN, functions as a protein-RNA toxin-antitoxin pair. *Proc. Natl. Acad. Sci. USA* **106**, 894–899 (2009).
- Hazan, R. & Engelberg-Kulka, H. *Escherichia coli* mazEF-mediated cell death as a defense mechanism that inhibits the spread of phage P1. *Mol. Genet. Genomics* **272**, 227–234 (2004).
- Hazan, R., Sat, B. & Engelberg-Kulka, H. *Escherichia coli* mazEF-mediated cell death is triggered by various stressful conditions. *J. Bacteriol.* **186**, 3663–3669 (2004).
- Buts, L., Lah, J., Dao-Thi, M. H., Wyns, L. & Loris, R. Toxin-antitoxin modules as bacterial metabolic stress managers. *Trends Biochem. Sci.* **30**, 672–679 (2005).
- Aguirre-Ramirez, M., Ramirez-Santos, J., Van Melderen, L. & Gomez-Eichelmann, M. C. Expression of the F plasmid ccd toxin-antitoxin system in *Escherichia coli* cells under nutritional stress. *Can. J. Microbiol.* **52**, 24–30 (2006).
- Gerdes, K., Christensen, S. K. & Lobner-Olesen, A. Prokaryotic toxin-antitoxin stress response loci. *Nat. Rev. Microbiol.* **3**, 371–382 (2005).
- Kim, Y. *et al.* *Escherichia coli* toxin/antitoxin pair MqsR/MqsA regulate toxin CspD. *Environ. Microbiol.* **12**, 1105–1121 (2010).
- Frampton, R., Aggio, R. B., Villas-Boas, S. G., Arcus, V. L. & Cook, G. M. Toxin-antitoxin systems of *Mycobacterium smegmatis* are essential for cell survival. *J. Biol. Chem.* **287**, 5340–5356 (2012).
- Kim, Y., Wang, X., Ma, Q., Zhang, X. S. & Wood, T. K. Toxin-antitoxin systems in *Escherichia coli* influence biofilm formation through YjgK (TabA) and fimbriae. *J. Bacteriol.* **191**, 1258–1267 (2009).
- Muranaka, L. S., Takita, M. A., Olivato, J. C., Kishi, L. T. & de Souza, A. A. Global expression profile of biofilm resistance to antimicrobial compounds in the plant-pathogenic bacterium *Xylella fastidiosa* reveals evidence of persister cells. *J. Bacteriol.* **194**, 4561–4569 (2012).
- Wang, X. & Wood, T. K. Toxin-antitoxin systems influence biofilm and persister cell formation and the general stress response. *Appl. Environ. Microbiol.* **77**, 5577–5583 (2011).
- Helaine, S. & Kugelberg, E. Bacterial persisters: formation, eradication, and experimental systems. *Trends Microbiol.* **22**, 417–424 (2014).
- Schuster, C. F. & Bertram, R. Toxin-antitoxin systems are ubiquitous and versatile modulators of prokaryotic cell fate. *FEMS Microbiol. Lett.* **340**, 73–85 (2013).
- Norton, J. P. & Mulvey, M. A. Toxin-antitoxin systems are important for niche-specific colonization and stress resistance of uropathogenic *Escherichia coli*. *PLoS Pathog.* **8**, e1002954 (2012).
- Ren, D., Walker, A. N. & Daines, D. A. Toxin-antitoxin loci *vapBC-1* and *vapXD* contribute to survival and virulence in nontypeable *Haemophilus influenzae*. *BMC Microbiol.* **12**, 263 (2012).
- Ren, D., Kordis, A. A., Sonenshine, D. E. & Daines, D. A. The *ToxA-vapA* toxin-antitoxin locus contributes to the survival of nontypeable *Haemophilus influenzae* during infection. *PLoS One* **9**, e91523 (2014).
- De la Cruz, M. A. *et al.* A toxin-antitoxin module of *Salmonella* promotes virulence in mice. *PLoS Pathog.* **9**, e1003827 (2013).
- Helaine, S. *et al.* Internalization of *Salmonella* by macrophages induces formation of nonreplicating persisters. *Science* **343**, 204–208 (2014).
- Georgiades, K. & Raoult, D. Genomes of the most dangerous epidemic bacteria have a virulence repertoire characterized by fewer genes but more toxin-antitoxin modules. *PLoS One* **6**, e17962 (2011).
- Sala, A., Bordes, P. & Genevaux, P. Multiple toxin-antitoxin systems in *Mycobacterium tuberculosis*. *Toxins* **6**, 1002–1020 (2014).
- Ruby, T., McLaughlin, L., Gopinath, S. & Monack, D. *Salmonella*'s long-term relationship with its host. *FEMS Microbiol. Rev.* **36**, 600–615 (2012).
- Tsolis, R. M. *et al.* Of mice, calves, and men. Comparison of the mouse typhoid model with other *Salmonella* infections. *Adv. Exp. Med. Biol.* **473**, 261–274 (1999).
- Crawford, R. W. *et al.* Gallstones play a significant role in *Salmonella* spp. gallbladder colonization and carriage. *Proc. Natl. Acad. Sci. USA* **107**, 4353–4358 (2010).
- Monack, D. M., Bouley, D. M. & Falkow, S. *Salmonella typhimurium* persists within macrophages in the mesenteric lymph nodes of chronically infected Nramp1<sup>+/+</sup> mice and can be reactivated by IFNγ neutralization. *J. Exp. Med.* **199**, 231–241 (2004).
- Mastroeni, P., Grant, A., Restif, O. & Maskell, D. A dynamic view of the spread and intracellular distribution of *Salmonella enterica*. *Nat. Rev. Microbiol.* **7**, 73–80 (2009).
- García-del Portillo, F. *Salmonella* intracellular proliferation: where, when and how? *Mol. Microbiol. Infect.* **3**, 1305–1311 (2001).
- Núñez-Hernández, C. *et al.* Genome expression analysis of nonproliferating intracellular *Salmonella enterica* serovar Typhimurium unravels an acid pH-dependent PhoP-PhoQ response essential for dormancy. *Infect. Immun.* **81**, 154–165 (2013).
- Blower, T. R. *et al.* Identification and classification of bacterial Type III toxin-antitoxin systems encoded in chromosomal and plasmid genomes. *Nucleic Acids Res.* **40**, 6158–6173 (2012).
- Fookes, M. *et al.* *Salmonella bongori* provides insights into the evolution of the Salmonellae. *PLoS Pathog.* **7**, e1002191 (2011).
- Nuccio, S. P. & Baumler, A. J. Comparative analysis of *Salmonella* genomes identifies a metabolic network for escalating growth in the inflamed gut. *MBio* **5**, e00929–00914 (2014).
- Bahassi, E. M., Salmon, M. A., Van Melderen, L., Bernard, P. & Couturier, M. F. plasmid CcdB killer protein: *ccdB* gene mutants coding for non-cytotoxic proteins which retain their regulatory functions. *Mol. Microbiol.* **15**, 1031–1037 (1995).
- Núñez-Hernández, C., Alonso, A., Pucciarelli, M. G., Casadesu, J. & García del Portillo, F. Dormant intracellular *Salmonella enterica* serovar Typhimurium discriminates among *Salmonella* pathogenicity island 2 effectors to persist inside fibroblasts. *Infect. Immun.* **82**, 221–232 (2014).
- Malik-Kale, P., Winfree, S. & Steele-Mortimer, O. The bimodal lifestyle of intracellular *Salmonella* in epithelial cells: replication in the cytosol obscures defects in vacuolar replication. *PLoS One* **7**, e38732 (2012).
- Slattery, A., Victorsen, A. H., Brown, A., Hillman, K. & Phillips, G. J. Isolation of highly persistent mutants of *Salmonella enterica* serovar typhimurium reveals a new toxin-antitoxin module. *J. Bacteriol.* **195**, 647–657 (2013).
- Bordes, P. *et al.* SecB-like chaperone controls a toxin-antitoxin stress-responsive system in *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. USA* **108**, 8438–8443 (2011).
- Nariya, H. & Inouye, M. MazF, an mRNA interferase, mediates programmed cell death during multicellular *Myxococcus* development. *Cell* **132**, 55–66 (2008).



51. Cook, G. M. *et al.* Ribonucleases in bacterial toxin-antitoxin systems. *Biochim. Biophys. Acta* **1829**, 523–531 (2013).
52. Vetting, M. W. *et al.* Structure and functions of the GNAT superfamily of acetyltransferases. *Arch. Biochem. Biophys.* **433**, 212–226 (2005).
53. Spange, S., Wagner, T., Heinzl, T. & Kramer, O. H. Acetylation of non-histone proteins modulates cellular signalling at multiple levels. *Int. J. Biochem. Cell Biol.* **41**, 185–198 (2009).
54. Thao, S. & Escalante-Semerena, J. C. Biochemical and thermodynamic analyses of *Salmonella enterica* Pat, a multidomain, multimeric N(epsilon)-lysine acetyltransferase involved in carbon and energy metabolism. *MBio* **2**, e00216–11 (2011).
55. Wagner, E. G. & Unoson, C. The toxin-antitoxin system *tisB-istRI*: Expression, regulation, and biological role in persister phenotypes. *RNA Biol.* **9**, 1513–1519 (2012).
56. Yamaguchi, Y., Tokunaga, N., Inouye, M. & Phadtare, S. Characterization of LdrA (long direct repeat A) protein of *Escherichia coli*. *J. Mol. Microbiol. Biotechnol.* **24**, 91–97 (2014).
57. Steif, A. & Meyer, I. M. The hok mRNA family. *RNA Biol.* **9**, 1399–1404 (2012).
58. Eriksson, S., Lucchini, S., Thompson, A., Rhen, M. & Hinton, J. C. Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*. *Mol. Microbiol.* **47**, 103–118 (2003).
59. Kasari, V., Mets, T., Tenson, T. & Kaldalu, N. Transcriptional cross-activation between toxin-antitoxin systems of *Escherichia coli*. *BMC Microbiol.* **13**, 45 (2013).
60. Winther, K. S. & Gerdes, K. Ectopic production of VapCs from Enterobacteria inhibits translation and trans-activates YoeB mRNA interferase. *Mol. Microbiol.* **72**, 918–930 (2009).
61. Malik-Kale, P. *et al.* *Salmonella* - at home in the host cell. *Front. Microbiol.* **2**, 125 (2011).
62. Leplae, R. *et al.* Diversity of bacterial type II toxin-antitoxin systems: a comprehensive search and functional analysis of novel families. *Nucleic Acids Res.* **39**, 5513–5525 (2011).
63. Chang, A. C. & Cohen, S. N. Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. *J. Bacteriol.* **134**, 1141–1156 (1978).
64. Lemonnier, M. & Lane, D. Expression of the second lysine decarboxylase gene of *Escherichia coli*. *Microbiology* **144**, 10 (1998).
65. Gotfredsen, M. & Gerdes, K. The *Escherichia coli relBE* genes belong to a new toxin-antitoxin gene family. *Mol. Microbiol.* **29**, 1065–1076 (1998).
66. Nieto, C. *et al.* The chromosomal *relBE2* toxin-antitoxin locus of *Streptococcus pneumoniae*: characterization and use of a bioluminescence resonance energy transfer assay to detect toxin-antitoxin interaction. *Mol. Microbiol.* **59**, 1280–1296 (2006).
67. Maisonneuve, E., Shakespeare, L. J., Jorgensen, M. G. & Gerdes, K. Bacterial persistence by RNA endonucleases. *Proc. Natl. Acad. Sci. USA* **108**, 13206–13211 (2011).
68. Uzzau, S., Figueroa-Bossi, N., Rubino, S. & Bossi, L. Epitope tagging of chromosomal genes in *Salmonella*. *Proc. Natl. Acad. Sci. USA* **98**, 15264–15269 (2001).
69. Aiastui, A., Pucciarelli, M. G. & Garcia-del Portillo, F. *Salmonella enterica* serovar typhimurium invades fibroblasts by multiple routes differing from the entry into epithelial cells. *Infect. Immun.* **78**, 2700–2713 (2010).

## Acknowledgments

We are grateful to Josep Casadesús (University of Seville, Spain) for strains and critical comments and Elizabeth Diago-Navarro for her initial involvement in this project and for the critical reading of the manuscript. We also thank the technical assistance of Pablo García and Diana Barroso (CNB-CSIC) and Alicia Rodríguez-Bernabé. The proteomics analysis LC-MS/MS by LTQ Orbitrap Velos was carried out by María I. Fernández López-Lucendo in the Proteomics and Genomics Facility (CIB-CSIC), a member of ProteoRed-ISCI network. This study was supported by grants BIO2013-46281-P (to F.G.-d.P.), BFU2011-25939 (to R. D.-O.), and CSD2008/00013 (INTERMODS, Consolider Program) (to F.G.-d.P. and R. D.-O.), funded by the Spanish Ministry of Economy and Competitiveness. D. L.-M. was supported by a F.P.U. fellowship from the Spanish Ministry of Education, Culture and Sports.

## Author contributions

D.L.-M., R.D.-O. and F.G.-d.P. conceived and designed the experiments for this study. D.L.-M., I.M.C. and V.F. performed the experiments. F.G.-d.P. wrote the manuscript. All authors discussed the data and made comments on the manuscript.

## Additional information

**Supplementary information** accompanies this paper at <http://www.nature.com/scientificreports>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Lobato-Márquez, D., Moreno-Córdoba, I., Figueroa, V., Díaz-Orejas, R. & García-del Portillo, F. Distinct type I and type II toxin-antitoxin modules control *Salmonella* lifestyle inside eukaryotic cells. *Sci. Rep.* **5**, 9374; DOI:10.1038/srep09374 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>