# SCIENTIFIC REPORTS

### **OPEN**

SUBJECT AREAS: CRYOSPHERIC SCIENCE ATMOSPHERIC SCIENCE

> Received 2 November 2014

Accepted 13 February 2015

> Published 9April 2015

Correspondence and requests for materials should be addressed to K.K. (kawamura@ lowtem.hokudai.ac.jp) or S.C.K. (shichang. kang@lzb.ac.cn)

## Penetration of biomass-burning emissions from South Asia through the Himalayas: new insights from atmospheric organic acids

Zhiyuan Cong<sup>1,2,3</sup>, Kimitaka Kawamura<sup>2</sup>, Shichang Kang<sup>3,4</sup> & Pingqing Fu<sup>5</sup>

<sup>1</sup>Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China, <sup>2</sup>Institute of Low Temperature Science, Hokkaido University, Sapporo, 060-0819, Japan, <sup>3</sup>CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100101, China, <sup>4</sup>State Key Laboratory of Cryospheric Sciences, Cold and Arid Regions Environmental and Engineering Research Institute, CAS, Lanzhou 730000, China, <sup>5</sup>LAPC, Institute of Atmospheric Physics, CAS, Beijing 100029, China.

High levels of carbonaceous aerosol exist over South Asia, the area adjacent to the Himalayas and Tibetan Plateau. Little is known about if they can be transported across the Himalayas, and as far inland as the Tibetan Plateau. As important constituents of aerosols, organic acids have been recognized as unique fingerprints to identify the atmospheric process. Here we measured dicarboxylic acids and related compounds in aerosols on the northern slope of Mt. Everest (Qomolangma, 4276 m a.s.l.). Strong positive correlations were observed for dicarboxylic acids with biomass burning tracers, levoglucosan and K<sup>+</sup>, demonstrating that this area was evidently affected by biomass burning. The seasonal variation pattern of dicarboxylic acids is consistent with OC and EC, being characterized by a pronounced maximum in the pre-monsoon season. Molecular distributions of dicarboxylic acids and related compounds (malonic acid/ succinic acid, maleic acid/fumaric acid) further support this finding. We suggest that the local meteorological conditions and regional atmospheric flow process could facilitate the penetration of the carbonaceous aerosols from South Asia throughout the Himalayas. With the consideration of the darkening force of carbonaceous aerosols, our finding has important implication for this climate-sensitive area, where the glacier melting supplies water for billions of people downstream.

ith vast mountain glaciers (ca. 10<sup>5</sup> km<sup>2</sup>), the Himalayas and Tibetan Plateau (HTP) present a unique and sensitive area under the regime of climate change<sup>1,2</sup>. Meltwater from those glaciers supplies major Asian rivers, such as the Indus River, Ganges River, Yarlung Tsangpo (Brahmaputra), Yangtze River and Yellow River<sup>3,4</sup>. At the same time, serious air pollution (Atmospheric Brown Clouds, ABC) is widely spread on the Indo Gangetic Plain (IGP)<sup>5</sup>. To evaluate the influence of the brown cloud over IGP on the high Himalayas, several continuous observations have been implemented under the framework of the UNEP-ABC project<sup>6</sup>. Those research clearly demonstrated that the ABC could affect the high altitudes, and even be lifted to more than 10 km in height<sup>7</sup>. However, most current works were confined to the southern slope of the Himalayas and little is known about the northern slope, especially the spread and transport mechanism of these air pollutants<sup>8</sup>. Actually, if the pollutants could be transported across the Himalayas, the gentle surface of the Tibetan Plateau will favor their further spread to the north, where more glaciers locate<sup>9</sup>. The deposition and accumulation of carbonaceous particles on the snow/ice surface may dramatically decrease the snow albedo, thereby resulting in increased glacier melt<sup>10-13</sup>.

As major constituents of atmospheric aerosols, organic acids can originate either from the primary process such as biomass burning and traffic emission<sup>14</sup> or the secondary formation by gas-to-particle conversion from various precursors<sup>15,16</sup>. Due to the hygroscopicity and the capability to act as cloud condensation nuclei (CCN), organic acids in aerosols are of great importance in the climate system<sup>17</sup>. Moreover, several specific compounds and their concentration ratios have been well recognized as unique fingerprints to identify the sources, transport and reaction processes of atmospheric aerosols. Given the importance of organic acids, they were widely studied in different locations such as urban<sup>18</sup>, continental background<sup>19,20</sup>, ocean<sup>21</sup> and the Arctic<sup>22</sup>. However, such

research is very scarce in the HTP. In ice core samples from Mt. Everest, Kang<sup>23</sup> found the oxalate concentration level during the 1950s–1980s was three times the background value of the 19<sup>th</sup> century values. This tripling was attributed to the anthropogenic production.

In this study, dicarboxylic acids and related compounds in aerosols were measured for the first time on Mt. Everest, a high-altitude site on the northern slope of the Himalayas (28.36°N, 86.95°E; 4276 m above sea level), from August 2009 to July 2010. The major purpose of this research is to reveal the transport process of atmospheric pollutants across the Himalayas by exploiting the sourceindicating faction of organic acids.

#### Results

**Molecular distribution of dicarboxylic acids and related compounds.** Table 1 shows the average concentration and ranges of various dicarboxylic and related compounds in aerosols from Mt. Everest. Annual average concentrations of dicarboxylic acids, oxoacids and  $\alpha$ -dicarbonyls were 109, 7.39 and 0.85 ng m<sup>-3</sup>, accounting for 7.45%, 0.39% and 0.06% of total organic carbon (OC), respectively. The contribution of dicarboxylic acids and related compounds to aerosol OC could be up to 15.5%, indicating that organic acids are major components in the high mountain aerosols. The average concentration of dicarboxylic acids in Mt. Everest aerosol is two to four times higher than that (64 ng m<sup>-3</sup>) reported in Alert, Arctic<sup>24</sup>, and that (30 ng m<sup>-3</sup>) in Syowa, Antarctica<sup>25</sup>. The value found in this study is comparable with data from some remote marine sites, such as 139 ng m<sup>-3</sup> at Chichi-jima Island in the western North Pacific<sup>26</sup>, but much lower than those observed in Asian cities. For example, high levels of dicarboxylic acids were reported in New Delhi  $(2330 \text{ ng m}^{-3})^{27}$  and Chennai (612 ng m<sup>-3</sup>)<sup>28</sup>, India, and Hong Kong (692 ng m<sup>-3</sup>)<sup>29</sup>. Furthermore, in this study the average concentration of total dicarboxylic acids in the pre-monsoon period (235 ng m<sup>-3</sup>) is significantly higher than during other seasons (63.1 ng m<sup>-3</sup> in monsoon, 58.4 ng m<sup>-3</sup> in post-monsoon and 69.7 ng m<sup>-3</sup> in winter, Table S1). The high loading of dicarboxylic

Table 1 | Concentrations of dicarboxylic acids, oxocarboxylic acids and  $\alpha$ -dicarbonyls (ng m<sup>-3</sup>) in the aerosols from QOMS station, the north slope of the Himalayas

	Concentrations (ng m <sup>-3</sup> )			
	Mean	S.D.	Min	Max
Dicarboxylic acids				
Oxalic, C <sub>2</sub>	59.8	73.4	7.64	426
Malonic, C <sub>3</sub>	7.30	7.37	0.76	39.7
Succinic, C <sub>4</sub>	13.7	13.13	2.55	72.1
Glutaric, C5	2.49	2.22	0.50	12.7
Adipic, C <sub>6</sub>	1.23	1.34	0.15	7.43
Pimelic, C <sub>7</sub>	0.54	0.98	0.03	5.35
Sebacic, C <sub>8</sub>	0.35	0.61	0.00	2.91
Azelaic, C <sub>9</sub>	1.38	1.01	0.42	6.75
Decanedioic, C <sub>10</sub>	0.28	0.50	0.00	2.78
Undecanedioic, C <sub>11</sub>	0.26	0.53	0.00	3.55
Dodecanedioc, C <sub>12</sub>	0.11	0.10	0.00	0.51
Methylmalonic, iC <sub>4</sub>	0.36	0.33	0.00	1.97
Methylsuccinic, iC <sub>5</sub>	1.43	1.66	0.20	10.0
2-Methylalutaric, iC <sub>6</sub>	0.24	0.16	0.05	0.85
Maleic, M	2.98	2.71	0.78	16.4
Fumaric, F	0.72	0.62	0.22	3.34
Methylmaleic, mM	2.20	2.68	0.56	14.2
Phthalic. Ph	9 47	6 14	2 27	33.2
Isophthalic, iPh	0.23	0.17	0.06	0.87
Terenbthalic tPh	1 55	1 43	0.32	8.07
Malic bC	0.15	0.10	0.02	0.52
Oxomalonic kC.	1 18	1 31	0.16	6.52
4-Oxonimelic kC-	0.67	1.00	0.00	3 00
Subtotal	109	117	16.00	677
Contribution to OC (%)	7.45	2.52	1.71	14.7
Oxocarboxylic acids				
Pyruvic, Pyr	0.83	0.87	0.04	4.33
Glyoxylic (2-oxoethanoic), $\omega C_2$	2.09	3.97	0.00	19.6
3-Oxopropanoic, $\omega C_3$	0.44	0.54	0.00	2.46
4-Oxobutanoic, $\omega C_4$	1.62	1.76	0.16	9.87
5-Oxopentanoic, $\omega C_5$	0.17	0.20	0.02	1.21
<b>7-Oxoheptanoic</b> , $\omega C_7$	0.77	0.86	0.00	4.49
8-Oxooctanoic, $\omega C_8$	1.04	1.23	0.00	5.72
9-Oxononanoic, $\omega C_{o}$	0.43	0.60	0.00	3.28
Subtotal	7.39	10.0	0.22	50.9
Contribution to OC (%)	0.39	0.20	0.03	0.79
α-Dicarbonyls				
Glyoxal, Gly	0.40	0.56	0.03	2.68
Methylglyoxal, mGly	0.45	0.56	0.00	2.45
Subtotal	0.85	1.12	0.03	5.12
Contribution to OC (%)	0.06	0.05	0.01	0.29

acids in the pre-monsoon season in the Mt. Everest region was likely related to the source strength rather than photochemical oxidation. According to a previous study, a positive linear correlation exists between dicarboxylic acids and air temperature<sup>30</sup>. However, the maximum air temperature at Mt. Everest was observed in the summer monsoon season.

Seasonal average molecular distributions of dicarboxylic acids, oxoacids and  $\alpha$ -dicarbonyls in the aerosols are presented in Figure 1. Although large differences occurred among seasons, oxalic acid (C<sub>2</sub>) was detected as the most abundant dicarboxylic acid species, followed by succinic (C<sub>4</sub>) and malonic acid (C<sub>3</sub>) in all seasons. Oxalic acid was commonly detected as the predominant species in various environments worldwide. However, a clear pattern that C<sub>4</sub>

being more abundant than  $C_3$ , a typical signal of biomass burning emission<sup>31,32</sup>, was found in the Mt. Everest aerosols. Therefore, our finding indicates the importance of biomass burning influences in this area, the point to be further discussed below.

#### Discussion

Primary versus secondary contribution reflected by ratios of dicarboxylic acids. Because succinic acid (C<sub>4</sub>) tends to be degraded into malonic acid (C<sub>3</sub>), the ratio of C<sub>3</sub>/C<sub>4</sub> has widely been used as an indicator to evaluate the photochemical production of dicarboxylic acids<sup>18,21</sup>. The C<sub>3</sub>/C<sub>4</sub> ratios in the Mt. Everest aerosols ranged from 0.11 to 0.81 with an average of 0.51, which is similar to some urban sites (such as 0.6 in New Delhi)<sup>27</sup>, while much lower than those from



Figure 1 | Molecular distributions of dicarboxylic acids and related compounds in different seasons at Mt. Everest (QOMS station), the northern slope of the Himalayas.





Figure 2 | Relationship between the concentrations of total dicarboxylic acids and (a) Elemental Carbon (EC), (b) Levoglucosan and (c) water-soluble potassium (K<sup>+</sup>).

continental and remote marine sites. For example, the  $C_3/C_4$  ratio from Qinghai Lake, northern TP is 2.2 due to the photochemical production of  $C_3$  from  $C_4^{33}$ . Even higher ratios are found over the Atlantic Ocean  $(2.1-3.4)^{21}$  and the equatorial central Pacific (up to  $10)^{34}$ . Low  $C_3/C_4$  ratios in this study show that the aerosols from Mt. Everest were relatively fresh, again indicating that the direct influence from primary emission source is more important than photochemical oxidation.

The unsaturated dicarboxylic acid, maleic acid (M, *cis* configuration), is formed by the degradation of aromatic hydrocarbons (e.g. toluene and benzene). Under the solar radiation, it could be further isomerized to its *trans* isomer, fumaric acid (F), through photochemical processes<sup>30</sup>. Therefore, the ratios of M/F can be applied to assess the aging of aerosols, i.e, lower M/F ratio means higher photochemical aging. In the present study, the M/F ratios (1.55–8.16, average 4.44) are much higher than those at marine sites (e.g. the North Pacific, 0.06–1.3, average 0.26)<sup>34</sup>. While the M/F ratios of Mt. Everest aerosols are similar to those reported at urban sites (New Delhi, 2.0–3.6; Beijing, 2.3)<sup>27,35</sup> and sites intensively impacted by biomass burning (Mt. Tai, China, 2.0; Rondonia, Amazonia, 2.8)<sup>31,32</sup>. The high M/F ratios found in this study also suggest that isomerization of maleic acid to fumaric acid by photochemical trans-





Figure 3 | CALIPSO retrieved backscatter signal at 532 nm (upper panel) and aerosol sub-type information (lower panel) on 17 April 2010. The Himalayas and southern TP (marked with circles) are covered by a thick aerosol layer, suggesting that air pollutants could be uplifted to more than 6 km high in altitude. CALIPSO profiles were obtained from (http://www-calipso.larc.nasa.gov).

formation is not significant at Mt. Everest. This finding confirms that Mt. Everest aerosols are rather fresh, without being substantially aged.

Source attribution - the influence of biomass burning. Dicarboxylic acids have various sources, including primary sources such as biomass burning, vehicular exhausts and cooking, and secondary sources i.e. atmospheric photooxidation of organic precursors. Since EC is only emitted by combustion sources (fossil fuel and/or biomass burning), it has been used frequently as a conservative tracer for primary combustion-generated OC. In this study, the dicarboxylic acids exhibit a strong correlation with EC ( $R^2 = 0.77$ , Fig. 2a), which demonstrates that primary combustion sources are the predominant contributor to dicarboxylic acids, while secondary photochemical production is negligible. Levoglucosan is a specific tracer of biomass burning, because it can only be produced through the pyrolysis of cellulose during the combustion process<sup>36</sup>. For the aerosols from Mt. Everest, a strong positive correlation between levoglucosan and dicarboxylic acids was found (Fig. 2b), with a correlation coefficient (R<sup>2</sup>) of 0.83. Furthermore, dicarboxylic acids also closely correlated with another biomass burning tracer, water-soluble potassium (K<sup>+</sup>) (Fig. 2c) (Note here, only the samples with K<sup>+</sup> concentration above the detection limits were used for the correlation analysis.). Therefore, these results suggested that among the various primary combustion sources, biomass burning rather than fossil fuel (coal combustion or vehicular exhaust) is the prevalent source of dicarboxylic acids at Mt. Everest. Especially in the pre-monsoon season, the higher ratios of levoglucosan to EC emphasize the importance of biomass burning influence (Fig. S1). Our findings differs from a previous study at Qinghai Lake (3200 m a.s.l.) over the northern TP (similar background conditions to this study), where dicarboxylic acids are found to be mainly derived from photochemical oxidation due to strong radiation<sup>33</sup>.

**Potential source region and transport mechanism.** Because molecular distributions and ratios of organic acids in the Mt. Everest aerosols exhibited a strong influence from biomass burning, we further checked such emission strength for different seasons using the active fire spots from MODIS (FIRMS, https://earthdata.nasa.





Figure 4 | Illustration of the aerosol transport mechanism from the lowlands of South Asia to across the Himalayas by the mountain/valley wind system (generated by Z.Y. Cong).

gov/firms). Results showed that in the pre-monsoon period there were a great number of agricultural burning and forest fires along the southern Himalayan foothills and Northern IGP (Figure S2). Similarly, a recent work (BC and O<sub>3</sub>) conducted at the Nepal Climate Observatory-Pyramid station (NCO-P) on the southern slope of the Himalayas also pointed out the importance of open fire emission from that area<sup>37</sup>. Our finding is also in agreement with the viewpoint of Vadrevu et al.38, that the pre-monsoon period (especially April) is the major fire season in the lowland of the southern Himalayas. The meteorological regime of this region is characterized by humid air masses from the Indian Ocean in the summer monsoon season and strong westerlies in other seasons (Fig. S3). Therefore, the wind system could facilitate the transport of air pollutants from South Asia to the Himalayas, regardless the shift of air circulation pattern among seasons. The aerosol vertical distribution achieved by Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) retrievals demonstrates that smoke plume could reach beyond 6 km in altitude, which is higher than most of the mountain valleys in the Himalayas. An example of such pollution phenomenon can be observed on 17 April 2010 (Fig. 3), which clearly demonstrates that the Himalayas and the southern TP (marked with circles) are covered by a thick polluted aerosol layer, which apparently originated from South Asia.

In addition to large-scale atmospheric circulation, the local orography may also play an important role in air pollutant transport. The mountain/valley wind system in the southern Himalayas is characterized by an evident up-valley wind in daytime with a maximum in the afternoon, which delivers substantial pollutants from South Asian lowland to higher altitude (e.g. NCO-P)<sup>6,39</sup>. In contrast, a predominant down-valley wind occurs on the northern slope of the Himalayas with peaks in the afternoon, because the downward "glacier wind" produced by the vast snow/ice cover in the northern Himalayas can overcome the normal up-valley air flow in daytime40,41. Therefore, acting as efficient channels of south-to-north air flow, the mountain valleys could allow the air pollutants to easily penetrate throughout the Himalayas (Fig. 4). A previous work<sup>42</sup> has revealed a trans-Himalayan pollution episode from Khumbu Valley, Nepal to the Tibetan Plateau (Rongbuk Valley) based on the simultaneous observation of condensation nucleus. When carbonaceous aerosols emitted from South Asia are transported to the far north (i.e. inland on the TP) and eventually deposited and accumulated on glacier surfaces, undoubtedly, they will change the energy balance of glaciers<sup>43</sup>.

#### Method

**Research site and sampling.** During August 2009 to July 2010, total suspended particle (TSP) samples (n = 50) were collected weekly using a medium-volume

sampler with pre-combusted quartz filters at Mt. Everest (Qomolangma Station for Atmospheric and Environmental Observation and Research, 28.36°N, 86.95°E; 4276 m above sea level). Given the remote location and very sparse local population, QOMS is an ideal place to monitor the atmospheric environment in the Himalayas<sup>44</sup>. According to meteorological measurements, the seasonality at QOMS was divided into pre-monsoon, monsoon, post-monsoon and winter (Table S2).

After the water extraction and butyl ester derivatization, dicarboxylic acids (C<sub>2</sub>-C<sub>12</sub>), oxocarboxylic acids (C<sub>2</sub>-C<sub>9</sub>) and  $\alpha$ -dicarbonyls (glyoxal and methylglyoxal) (Fig. S4) in the aerosol samples were determined using gas chromatography with a flame ionization detector (GC-FID) following the modified analytical methods established by Kawamura and Ikushima<sup>18</sup> and Kawamura<sup>45</sup>.

The concentration of OC and EC was measured using an OC/EC analyzer following the IMPROVE-A protocol. Levoglucosan was determined by GC/MS after the extraction of the samples with a methanol/methylene chloride mixture followed by BSTFA derivatization<sup>46</sup>. The water-soluble ionic species were determined using an ion chromatograph (761 Compact IC, Metrohm). Details about the analytical procedures for dicarboxylic acids, oxocarboxylic acids,  $\alpha$ -dicarbonyls, OC, EC, and K<sup>+</sup>, as well as quality assurance and control are presented in the Supplementary Materials. All the concentrations of dicarboxylic acids and related compounds, carbonaceous and ionic components in this study are field-blank corrected.

- 1. Kang, S. *et al.* Review of climate and cryospheric change in the Tibetan Plateau. *Environ. Res. Lett.* **5**, 015101 (2010).
- 2. Yao, T. et al. Third Pole Environment (TPE). Environ. Dev. 3, 52-64 (2012).
- Lutz, A., Immerzeel, W., Shrestha, A. & Bierkens, M. Consistent increase in High Asia's runoff due to increasing glacier melt and precipitation. *Nat. Clim. Change* 4, 587–592 (2014).
- Schaner, N., Voisin, N., Nijssen, B. & Lettenmaier, D. P. The contribution of glacier melt to streamflow. *Environ. Res. Lett.* 7, 034029 (2012).
- Ramanathan, V. *et al.* Atmospheric brown clouds: Impacts on South Asian climate and hydrological cycle. *Proc. Natl. Acad. Sci. USA* 102, 5326–5333 (2005).
- Bonasoni, P. *et al.* Atmospheric Brown Clouds in the Himalayas: first two years of continuous observations at the Nepal Climate Observatory-Pyramid (5079 m). *Atmos. Chem. Phys.* 10, 7515–7531 (2010).
- Lawrence, M. G. Atmospheric science: Asia under a high-level brown cloud. Nat. Geosci. 4, 352–353 (2011).
- Rahul, P. R. C. *et al.* Double blanket effect caused by two layers of black carbon aerosols escalates warming in the Brahmaputra River Valley. *Sci. Rep.* 4, DOI:10.1038/srep03670 (2014).
- Gardner, A. S. et al. A Reconciled Estimate of Glacier Contributions to Sea Level Rise: 2003 to 2009. Science 340, 852–857 (2013).
- Qian, Y., Flanner, M. G., Leung, L. R. & Wang, W. Sensitivity studies on the impacts of Tibetan Plateau snowpack pollution on the Asian hydrological cycle and monsoon climate. *Atmos. Chem. Phys.* 11, 1929–1948 (2011).
- 11. Menon, S. et al. Black carbon aerosols and the third polar ice cap. Atmos. Chem. Phys. 10, 4559–4571 (2010).
- Lau, W. K. M., Kim, M.-K., Kim, K.-M. & Lee, W.-S. Enhanced surface warming and accelerated snow melt in the Himalayas and Tibetan Plateau induced by absorbing aerosols. *Environ. Res. Lett.* 5, 025204 (2010).
- 13. Ming, J. et al. Darkening of the mid-Himalaya glaciers since 2000 and the potential causes. *Environ. Res. Lett.* 7, 014021 (2012).
- Narukawa, M., Kawamura, K., Takeuchi, N. & Nakajima, T. Distribution of dicarboxylic acids and carbon isotopic compositions in aerosols from 1997 Indonesian forest fires. *Geophys. Res. Lett.* 26, 3101–3104 (1999).
- Claeys, M. et al. Formation of secondary organic aerosols through photooxidation of isoprene. Science 303, 1173–1176 (2004).

- Kawamura, K., Kasukabe, H. & Barrie, L. A. Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in arctic aerosols: One year of observations. *Atmos. Environ.* **30**, 1709–1722 (1996).
- 17. Zhang, R. *et al.* Atmospheric new particle formation enhanced by organic acids. *Science* **304**, 1487–1490 (2004).
- Kawamura, K. & Ikushima, K. Seasonal changes in the distribution of dicarboxylic acids in the urban atmosphere. *Environ. Sci. Technol.* 27, 2227–2235 (1993).
- van Pinxteren, D., Neusüß, C. & Herrmann, H. On the abundance and source contributions of dicarboxylic acids in size-resolved aerosol particles at continental sites in Central Europe. *Atmos. Chem. Phys.* 14, 3913–3928 (2014).
- Legrand, M. *et al.* Origin of C2-C5 dicarboxylic acids in the European atmosphere inferred from year-round aerosol study conducted at a west-east transect. *J. Geophys. Res. -Atmos* 112, DOI:10.1029/2006JD008019 (2007).
- Fu, P., Kawamura, K., Usukura, K. & Miura, K. Dicarboxylic acids, ketocarboxylic acids and glyoxal in the marine aerosols collected during a round-the-world cruise. *Mar. Chem.* 148, 22–32 (2013).
- 22. Kawamura, K., Kasukabe, H. & Barrie, L. A. Secondary formation of water-soluble organic acids and  $\alpha$ -dicarbonyls and their contributions to total carbon and water-soluble organic carbon: Photochemical aging of organic aerosols in the Arctic spring. *J. Geophys. Res.* -*Atmos* **115**, doi:10.1029/2010JD014299 (2010).
- 23. Kang, S., Dahe, Q., Mayewski, P. A. & Wake, C. P. Recent 180 Year Oxalate (C<sub>2</sub>O<sub>4</sub><sup>2</sup>) Records Recovered from the Mount Everest Ice Core: Some Environmental Implications. *J. Glaciol.* 47, 155–156 (2001).
- Kawamura, K., Imai, Y. & Barrie, L. A. Photochemical production and loss of organic acids in high Arctic aerosols during long-range transport and polar sunrise ozone depletion events. *Atmos. Environ.* 39, 599–614 (2005).
- Kawamura, K. *et al.* Water soluble dicarboxylic acids and related compounds in Antarctic aerosols. J. Geophys. Res. -Atmos 101, 18721–18728 (1996).
- Mochida, M. *et al.* Seasonal variation and origins of dicarboxylic acids in the marine atmosphere over the western North Pacific. *J. Geophys. Res. -Atmos* 108, DOI:10.1029/2002JD002355 (2003).
- Miyazaki, Y. *et al.* Dicarboxylic acids and water-soluble organic carbon in aerosols in New Delhi, India, in winter: Characteristics and formation processes. *J. Geophys. Res. - Atmos* 114, DOI:10.1029/2009JD011790 (2009).
- Pavuluri, C. M., Kawamura, K. & Swaminathan, T. Water-soluble organic carbon, dicarboxylic acids, ketoacids, and α-dicarbonyls in the tropical Indian aerosols. J. Geophys. Res. -Atmos 115, DOI:10.1029/2009JD012661 (2010).
- Ho, K. F. *et al.* Dicarboxylic acids, ketocarboxylic acids and dicarbonyls in the urban roadside area of Hong Kong. *Atmos. Environ.* **40**, 3030–3040 (2006).
  Sempéré, R. & Kawamura, K. Trans-hemispheric contribution of C2C10 α, ω-
- Sempéré, R. & Kawamura, K. Trans-hemispheric contribution of C2C10 α, ωdicarboxylic acids, and related polar compounds to water-soluble organic carbon in the western Pacific aerosols in relation to photochemical oxidation reactions. *Glob. Biogeochem. Cycle* 17, DOI:10.1029/2002GB001980 (2003).
- Kawamura, K. *et al.* High abundances of water-soluble dicarboxylic acids, ketocarboxylic acids and α-dicarbonyls in the mountaintop aerosols over the North China Plain during wheat burning season. *Atmos. Chem. Phys.* 13, 8285–8302 (2013).
- 32. Kundu, S. *et al.* Molecular distributions of dicarboxylic acids, ketocarboxylic acids and  $\alpha$ -dicarbonyls in biomass burning aerosols: implications for photochemical production and degradation in smoke layers. *Atmos. Chem. Phys.* **10**, 2209–2225 (2010).
- Meng, J. *et al.* Atmospheric oxalic acid and related secondary organic aerosols in Qinghai Lake, a continental background site in Tibet Plateau. *Atmos. Environ.* 79, 582–589 (2013).
- Kawamura, K. & Sakaguchi, F. Molecular distributions of water soluble dicarboxylic acids in marine aerosols over the Pacific Ocean including tropics. *J. Geophys. Res. - Atmos* 104, 3501–3510 (1999).
- 35. Ho, K. F. et al. Dicarboxylic acids, ketocarboxylic acids, α-dicarbonyls, fatty acids, and benzoic acid in urban aerosols collected during the 2006 Campaign of Air Quality Research in Beijing (CAREBeijing-2006). J. Geophys. Res. -Atmos 115, DOI:10.1029/2009JD013304 (2010).
- Simoneit, B. R. T. Biomass burning a review of organic tracers for smoke from incomplete combustion. *Appl. Geochem.* 17, 129–162 (2002).

- Putero, D. *et al.* Influence of open vegetation fires on black carbon and ozone variability in the southern Himalayas (NCO-P, 5079 m a.s.l.). *Environ. Pollut.* 184, 597–604 (2014).
- Vadrevu, K. P. et al. Vegetation fires in the himalayan region Aerosol load, black carbon emissions and smoke plume heights. Atmos. Environ. 47, 241–251 (2012).
- Marinoni, A. *et al.* Aerosol mass and black carbon concentrations, a two year record at NCO-P (5079 m, Southern Himalayas). *Atmos. Chem. Phys.* 10, 8551–8562 (2010).
- Zou, H. *et al.* Local wind system in the Rongbuk Valley on the northern slope of Mt. Everest. *Geophys. Res. Lett.* 35, DOI:10.1029/2008GL033466 (2008).
- Chen, X., Su, Z., Ma, Y. & Sun, F. Analysis of Land-Atmosphere Interactions over the North Region of Mt. Qomolangma (Mt. Everest). *Arct. Antarct. Alp. Res.* 44, 412–422 (2012).
- Hindman, E. E. & Upadhyay, B. P. Air pollution transport in the Himalayas of Nepal and Tibet during the 1995–1996 dry season. *Atmos. Environ.* 36, 727–739 (2002).
- 43. Qian, Y. *et al.* Light-absorbing particles in snow and ice: Measurement and modeling of climatic and hydrological impact. *Adv. Atmos. Sci.* 32, 64–91 (2015).
  44. Cara, Z. Y. *et al.* Elemental and the last of the measurement of the second seco
- 44. Cong, Z. Y. et al. Elemental and individual particle analysis of atmospheric aerosols from high Himalayas. Environ. Monit. Assess. 160, 323-335 (2010).
- 45. Kawamura, K. Identification of C2-C10  $\omega$ -oxocarboxylic acids, pyruvic acid, and C2-C3  $\alpha$ -dicarbonyls in wet precipitation and aerosol samples by capillary GC and GC/MS. *Anal. Chem.* **65**, 3505–3511 (1993).
- Fu, P. et al. Organic molecular compositions and temporal variations of summertime mountain aerosols over Mt. Tai, North China Plain. J. Geophys. Res. -Atmos 113, DOI:10.1029/2008JD009900 (2008).

#### Acknowledgments

This study is supported by National Natural Science Foundation of China (41271073, 41075089 and 41225002), Strategic Priority Research Program-Climate Change: Carbon Budget and Relevant Issues (XDA05100105), CAS, and also by the Japanese Society for the Promotion of Science (24221001). P.Q.F. appreciates the financial support from the "One Hundred Talents" program of CAS. We thank Zhongyan WANG and Yaoming MA for their sampling support at QOMS, Eri TACHIBANA, Kaori ONO, Daisuke SAWAOKA and Shaopeng GAO for the help in chemical analysis. The MODIS FIRMS, CALIPSO of NASA and HYSPLIT (ARL) of NOAA are also acknowledged.

#### Author contributions

All authors designed the research framework, Z.Y.C. conducted the fieldwork, Z.Y.C., P.Q. F. and K.K. performed the laboratory analysis and data interpretation. Z.Y.C., K.K. and S.C. K contributed to the manuscript preparation.

#### Additional information

Supplementary information accompanies this paper at http://www.nature.com/ scientificreports

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Cong, Z., Kawamura, K., Kang, S. & Fu, P. Penetration of biomass-burning emissions from South Asia through the Himalayas: new insights from atmospheric organic acids. *Sci. Rep.* **5**, 9580; DOI:10.1038/srep09580 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/