

SCIENTIFIC REPORTS



OPEN

Concurrence of oral and genital human papillomavirus infection in healthy men: a population-based cross-sectional study in rural China

Received: 23 June 2015
Accepted: 28 September 2015
Published: 27 October 2015

Fangfang Liu^{1,*}, Dong Hang^{2,*}, Qiuju Deng^{1,*}, Mengfei Liu¹, Longfu Xi^{3,4}, Zhonghu He¹, Chaoting Zhang¹, Min Sun¹, Ying Liu¹, Jingjing Li¹, Yaqi Pan¹, Tao Ning¹, Chuanhai Guo¹, Yongmei Liang¹, Ruiping Xu⁵, Lixin Zhang⁵, Hong Cai¹ & Yang Ke¹

Human papillomavirus (HPV) infection, a primary cause of genital cancer, is also related to the increasing incidence of oropharyngeal cancer among young men. Relatively little is known about the concurrence of oral and genital infection among healthy individuals. Oral and genital swab exfoliated cells were collected simultaneously from 2566 men in rural China. Using general primer-mediated (SPF1/GP6+) PCR and sequencing, HPV testing results were obtained from 2228 men with both valid oral and genital specimens (β -globin-positive). The prevalence of HPV infection was 6.7% in the oral cavity and 16.9% for the external genitalia. Among 43 men (1.9%, 43/2228) with oral-genital coinfection, 60.5% (26/43) harbored an identical HPV type at both sites. The risk of oral HPV infection was higher among men with genital infection than among uninfected men (11.4% vs. 5.7%, Adjusted OR = 2.3, 95% CI: 1.6–3.4). In addition, having multiple lifetime sexual partners was a significant risk for oral-genital HPV coinfection (Adjusted OR = 2.6, 95% CI: 1.0–7.0; 2 partners vs. 1 partner). These findings provide a basis for further understanding the natural history and transmission dynamics of oral HPV infection.

Human papillomavirus (HPV) infection is an etiologic factor for cervical, anal and penile cancer¹. More recent evidence shows that HPV also causes a subset of head and neck cancer including 20–90% of oropharyngeal cancer². In 2008, approximately 85,000 cases of oropharyngeal cancer were diagnosed globally, with more than 22,000 cases being HPV positive². Notably, the incidence of HPV-related oropharyngeal cancer is significantly increasing, particularly in young men^{3,4}. The prevalence of oral HPV infection in healthy individuals ranges from 1.3% to 9.2% worldwide, with HPV-16 being the most prevalent oncogenic type⁵. Previous studies have suggested that individuals with high-risk sexual behaviors (e.g. practicing oral sex and having multiple sexual partners) are at a greater risk for oral HPV infection^{5–9}.

However, whether oral HPV infection is associated with genital infection remains largely unknown, especially among men. Previous studies investigating the association of oral and genital HPV infection among men were based largely on particular high-risk groups such as university students, sexually transmitted disease clinic patients, HIV-infected men, or homosexual men^{10–16}. These studies, mostly limited

¹Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Laboratory of Genetics, Peking University Cancer Hospital & Institute, Beijing, China. ²Department of Epidemiology and Biostatistics, School of Public Health, Nanjing Medical University, Nanjing, China. ³Department of Epidemiology, School of Public Health, University of Washington, Seattle, WA, USA. ⁴Department of Pathology, School of Medicine, University of Washington, Seattle, WA, USA. ⁵Anyang Cancer Hospital, Anyang, Henan Province, People's Republic of China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to H.C. (email: drhcai@gmail.com) or Y.K. (email: keyang@bjmu.edu.cn)

in sample size, have indicated that concurrent oral-genital HPV infections are rare and have reported inconsistent results regarding type-specific association of HPV infection at these two sites^{10–16}.

Concurrence of oral and genital HPV infection has rarely been investigated in a general male population. Previously, we reported an overall oral HPV prevalence of 6% in 5410 25–65 year old healthy males and females in rural China¹⁷. The intent of this study is to evaluate the association of HPV DNA simultaneously present in collected male external genital and oral specimens from this general population and identify relevant risk factors.

Results

Participant characteristics. Among 2721 eligible subjects, 2228 (81.9%) participants showed valid HPV DNA results for both oral and genital specimens. The 493 individuals not included in analysis were either non-responding ($n = 155$) or were negative for beta-globin ($n = 338$). These excluded subjects were more likely to work outside of the local area, show loss of teeth, have a history of oral disease, and report oral sex practices, but were less likely to have a smoking history (Supplementary Table S1). The median age of the participants was 43 years (range, 25–65 years), and half of these individuals were engaged in farming (45.3%) (Table 1). Most subjects were married or cohabiting (94.7%), and had a level of education of less than 9 years total (88.6%). Both cigarette smoking (65.9%) and alcohol consumption (36.4%) were common. The proportion of individuals who reported having lost >10 teeth or who reported a history of oral disease were 8.3% and 1.8% respectively. With respect to sexual behaviors, 81.7% reported occasionally or never washing external genitalia before sex, 14.6% reported having ≥ 2 lifetime sexual partners, and 3.9% had a history of oral sex practices.

HPV detection and type-specific concordance. Among 2228 men, 376 (16.9%) and 149 (6.7%) were positive for genital and oral HPV infection, respectively (Table 1). Compared with oral infections, the spectrum of HPV types on the external genitalia (43 types) was more diverse and included all 11 HPV types which were detected in oral specimens. A predominance of non-oncogenic HPV types was observed both in the oral cavity and on the external genitalia. HPV-16 was the most prevalent type of oncogenic HPV (oral cavity: 0.45%; external genitalia: 2.83%), and HPV-3 was the most prevalent type of non-oncogenic HPV (oral cavity: 4.53%; external genitalia: 3.37%) (Supplementary Table S2).

The prevalence of oral HPV infection among men without genital HPV infection was 5.7%. Of the men with HPV detected on the genitalia, 11.4% also had an oral HPV infection. Among 43 men (1.9%, 43/2228) positive for HPV in both oral and genital specimens, 37.2% (16/43) had no types in common, 62.8% (27/43) were concordant for at least one type, and 60.5% (26/43) harbored exactly the same HPV type at both sites, including 18 men concordant for HPV-3, 4 men concordant for HPV-10, 3 men concordant for HPV-57, and 1 man concordant for HPV-29 (Table 2). The Monte-Carlo simulation showed that the observed events of positive concordance significantly exceeded the expected value (27 vs. 0.6, $P < 0.001$) (Supplementary Table S3).

Association between oral and genital HPV infections. Overall, the prevalence of oral HPV infection was higher among men with genital infection than among those without genital HPV infection (11.4% vs. 5.7%, Adjusted OR = 2.3, 95% CI: 1.6–3.4) (Table 3). In the type-specific analysis, the presence of one specific HPV type in oral specimens was associated with the presence of the same type in genital specimens (Adjusted OR = 55.7, 95% CI: 36.1–86.0).

Risk factor analysis. Men with multiple lifetime sexual partners were found to be at a higher risk for oral-genital HPV coinfection (Adjusted OR = 2.6, 95% CI: 1.0–7.0; 2 partners vs. 1 partner), as was the case for genital HPV infection (Table 1).

Discussion

This population-based investigation of the concordance of HPV infections in the oral cavity and on the external genitalia is to our knowledge the first such investigation to be undertaken among Chinese men. Although oral-genital coinfection in healthy men was a rare event in this study (1.9%), over 60.0% of these concurrent infections were completely type-specific concordant, and oral HPV prevalence was significantly higher among men with genital HPV infection than among those without. These data provide a basis for further understanding the natural history and transmission dynamics of oral HPV infection.

Prevalence of oral HPV was significantly lower than on the external genitalia in healthy adult Chinese men, similar to what has been observed in other populations⁶. The distinct microenvironment of the oral cavity including the existence of the abundant antimicrobial agents in the saliva may in part explain the overall lower prevalence of infection and less diversity in oral HPV types^{18,19}. Given the low prevalence of oral HPV infection (6.7%), it is not surprising that concurrent oral-genital HPV infection (1.9%) was uncommon in this study. Despite this rarity of coinfection, a relatively high level of type-specific concordance (62.8%, 27/43) was observed among concurrent oral and genital infections, in keeping with some previous studies^{10,16}. It is of note that coinfection was only observed for non-oncogenic HPV in this population. The absence of concurrent infections with oncogenic HPV may be due to the lower frequency of oncogenic HPV infection in the oral cavity (0.5%), in agreement with the relatively low incidence of HPV-related oropharyngeal cancer²⁰.

| Variables | No. (%) | Oral cavity (N = 2228) | | | External genitalia (N = 2228) | | | Oral cavity and external genitalia (N = 43) ^a | | |
|---|--------------|------------------------|--------------------------------|-----------------------------------|-------------------------------|--------------------------------|-----------------------------------|--|--------------------------------|-----------------------------------|
| | | % | Crude OR (95% CI) ^b | Adjusted OR (95% CI) ^b | % | Crude OR (95% CI) ^b | Adjusted OR (95% CI) ^b | % | Crude OR (95% CI) ^b | Adjusted OR (95% CI) ^b |
| Total | 2228 (100.0) | 6.7 | | | 16.9 | | | 1.9 | | |
| Age, years | | | | | | | | | | |
| 25–35 | 515 (23.1) | 6.4 | 1.0 | 1.0 | 14.6 | 1.0 | 1.0 | 1.6 | 1.0 | 1.0 |
| 36–45 | 792 (35.6) | 6.2 | 1.0 (0.6–1.5) | 1.0 (0.6–1.6) | 19.3 | 1.4 (1.0–1.9) | 1.6 (1.2–2.2) | 2.4 | 1.6 (0.7–3.6) | 1.7 (0.7–4.0) |
| 46–55 | 458 (20.6) | 6.8 | 1.1 (0.6–1.8) | 1.2 (0.7–2.1) | 16.4 | 1.1 (0.8–1.6) | 1.4 (0.9–2.0) | 2.2 | 1.4 (0.6–3.6) | 1.4 (0.5–3.9) |
| 56–65 | 463 (20.8) | 7.8 | 1.2 (0.8–2.0) | 1.3 (0.7–2.3) | 15.8 | 1.4 (0.8–1.6) | 1.4 (0.9–2.1) | 1.3 | 0.8 (0.3–2.4) | 1.0 (0.3–3.5) |
| <i>P</i> value for trend ^d | | | 0.340 | 0.363 | | 0.972 | 0.251 | | 0.686 | 0.963 |
| Education level | | | | | | | | | | |
| Illiteracy, <1 year | 120 (5.4) | 10.0 | 1.0 | 1.0 | 18.3 | 1.0 | 1.0 | 1.7 | 1.0 | 1.0 |
| Primary school, 1–6 years | 618 (27.7) | 7.3 | 0.7 (0.4–1.4) | 0.7 (0.3–1.4) | 16.4 | 0.9 (0.5–1.5) | 0.9 (0.5–1.5) | 1.0 | 0.6 (0.1–2.9) | 0.6 (0.1–3.2) |
| Junior high school, 7–9 years | 1236 (55.5) | 6.2 | 0.6 (0.3–1.1) | 0.6 (0.3–1.2) | 17.2 | 0.9 (0.6–1.5) | 0.9 (0.5–1.6) | 2.5 | 1.5 (0.4–6.4) | 1.5 (0.3–7.3) |
| Senior high school or above, >9 years | 254 (11.4) | 6.4 | 0.6 (0.3–1.3) | 0.6 (0.3–1.4) | 15.9 | 0.8 (0.5–1.5) | 0.8 (0.4–1.5) | 1.6 | 1.0 (0.2–5.3) | 0.9 (0.1–5.8) |
| <i>P</i> value for trend ^d | | | 0.150 | 0.222 | | 0.828 | 0.666 | | 0.228 | 0.332 |
| Marital status | | | | | | | | | | |
| Married or cohabiting | 2110 (94.7) | 6.7 | 1.0 | 1.0 | 16.6 | 1.0 | 1.0 | 1.9 | 1.0 | 1.0 |
| Not married, divorced, separated or widowed | 118 (5.3) | 5.9 | 0.9 (0.4–1.9) | 0.7 (0.3–1.6) | 22.0 | 1.4 (0.9–2.2) | 1.5 (1.0–2.4) | 2.5 | 1.4 (0.4–4.4) | 1.6 (0.5–5.5) |
| Type of employment | | | | | | | | | | |
| Farming at home | 1010 (45.3) | 6.3 | 1.0 | 1.0 | 15.7 | 1.0 | 1.0 | 1.9 | 1.0 | 1.0 |
| Working in local area | 366 (16.4) | 6.3 | 1.0 (0.6–1.6) | 1.1 (0.6–1.8) | 16.7 | 1.1 (0.8–1.5) | 1.0 (0.7–1.5) | 1.4 | 0.7 (0.3–1.9) | 0.7 (0.2–1.8) |
| Working outside local area | 662 (29.7) | 7.1 | 1.1 (0.8–1.7) | 1.2 (0.8–1.9) | 16.9 | 1.1 (0.8–1.4) | 1.0 (0.8–1.4) | 2.1 | 1.1 (0.6–2.3) | 1.0 (0.5–2.1) |
| Other | 190 (8.5) | 7.9 | 1.3 (0.7–2.3) | 1.5 (0.8–2.8) | 23.2 | 1.6 (1.1–2.4) | 1.5 (1.0–2.2) | 2.6 | 1.4 (0.5–3.8) | 1.2 (0.4–3.5) |
| Cigarette smoking ^c | | | | | | | | | | |
| Never | 760 (34.1) | 6.8 | 1.0 | 1.0 | 14.6 | 1.0 | 1.0 | 2.0 | 1.0 | 1.0 |
| Ever | 1468 (65.9) | 6.6 | 1.0 (0.7–1.4) | 0.9 (0.7–1.4) | 18.1 | 1.3 (1.0–1.6) | 1.3 (1.0–1.6) | 1.9 | 1.0 (0.5–1.8) | 1.0 (0.5–2.0) |
| Alcohol consumption ^c | | | | | | | | | | |
| Never | 1418 (63.6) | 7.1 | 1.0 | 1.0 | 15.7 | 1.0 | 1.0 | 1.9 | 1.0 | 1.0 |
| Ever | 810 (36.4) | 6.1 | 0.8 (0.6–1.2) | 0.8 (0.6–1.2) | 19.0 | 1.3 (1.0–1.6) | 1.1 (0.9–1.4) | 2.0 | 1.0 (0.6–1.9) | 1.0 (0.5–1.8) |
| History of oral disease ^c | | | | | | | | | | |
| Never | 2188 (98.2) | 6.5 | 1.0 | 1.0 | 17.0 | — | — | 2.0 | — | — |
| Ever | 40 (1.8) | 15.0 | 2.5 (1.0–6.1) | 2.8 (1.1–7.1) | 10.0 | — | — | 0.0 | — | — |
| Number of missing teeth | | | | | | | | | | |
| None | 382 (17.2) | 7.1 | 1.0 | 1.0 | 18.0 | — | — | 2.1 | 1.0 | 1.0 |
| 1–10 | 1639 (73.6) | 6.5 | 0.9 (0.6–1.4) | 0.9 (0.6–1.4) | 16.6 | — | — | 1.9 | 0.9 (0.4–2.0) | 1.0 (0.4–2.1) |
| >10 | 185 (8.3) | 7.0 | 1.0 (0.5–2.0) | 0.8 (0.4–1.7) | 15.7 | — | — | 2.2 | 1.0 (0.3–3.5) | 1.6 (0.4–6.2) |
| <i>P</i> value for trend ^d | | | 0.890 | 0.551 | | — | — | | 0.965 | 0.694 |
| Wash external genitalia before sex | | | | | | | | | | |
| Occasionally or never | 1820 (81.7) | 6.9 | 1.0 | 1.0 | 16.5 | 1.0 | 1.0 | 1.9 | 1.0 | 1.0 |
| Often or every time | 408 (18.3) | 5.6 | 0.8 (0.5–1.3) | 0.9 (0.5–1.4) | 18.6 | 1.2 (0.9–1.5) | 1.2 (0.9–1.6) | 2.2 | 1.2 (0.6–2.5) | 1.2 (0.5–2.5) |
| Oral sex practices | | | | | | | | | | |
| Never | 2141 (96.1) | 6.5 | 1.0 | 1.0 | 16.9 | 1.0 | 1.0 | 2.0 | 1.0 | 1.0 |
| Ever | 87 (3.9) | 10.3 | 1.6 (0.8–3.4) | 1.7 (0.8–3.5) | 16.1 | 0.9 (0.5–1.7) | 0.8 (0.4–1.5) | 1.1 | 0.6 (0.1–4.3) | 0.6 (0.1–4.9) |
| Lifetime number of sexual partners | | | | | | | | | | |
| 0–1 | 1903 (85.4) | 6.7 | 1.0 | 1.0 | 14.8 | 1.0 | 1.0 | 1.7 | 1.0 | 1.0 |
| 2 | 115 (5.2) | 7.0 | 1.0 (0.5–2.2) | 1.1 (0.5–2.3) | 25.2 | 1.9 (1.2–3.0) | 1.9 (1.2–3.0) | 4.3 | 2.6 (1.0–6.7) | 2.6 (1.0–7.0) |
| Continued | | | | | | | | | | |

| Variables | No. (%) | Oral cavity (N = 2228) | | | External genitalia (N = 2228) | | | Oral cavity and external genitalia (N = 43) ^a | | |
|---------------------------------------|-----------|------------------------|--------------------------------|-----------------------------------|-------------------------------|--------------------------------|-----------------------------------|--|--------------------------------|-----------------------------------|
| | | % | Crude OR (95% CI) ^b | Adjusted OR (95% CI) ^b | % | Crude OR (95% CI) ^b | Adjusted OR (95% CI) ^b | % | Crude OR (95% CI) ^b | Adjusted OR (95% CI) ^b |
| ≥3 | 210 (9.4) | 6.7 | 1.0 (0.6–1.8) | 0.9 (0.5–1.7) | 31.0 | 2.6 (1.9–3.5) | 2.7 (1.9–3.7) | 2.4 | 1.4 (0.5–3.6) | 1.5 (0.6–4.0) |
| <i>P</i> value for trend ^d | | | 0.975 | 0.918 | | <0.001 | <0.001 | | 0.241 | 0.197 |

Table 1. Prevalence of HPV infection by anatomic site and selected characteristics in men of rural Anyang, China, 2009–2011. Abbreviation: HPV, human papillomavirus; OR, odds ratio; CI, confidence interval. “—” denotes that the indicated variables were not included in the logistic regression models. ^aData for the oral cavity and external genitalia category include the subset of men with HPV in both sites. ^bAdjusted ORs and 95% CIs were calculated by multivariate logistic regression models including all listed variables. ^cCigarette smoking was defined as consuming an average of one cigarette or more per day for ≥12 months, and alcohol consumption was defined as drinking Chinese liquor at least twice per week for ≥12 months. A history of oral disease was defined as self-reported oral ulcers, gum disease, or chronic oral inflammation in the preceding 12 months. ^d*P* values for trends were derived by logistic regression analyses taking categorical variables as continuous variables.

| HPV status | No. (%) | Genital HPV types | Oral HPV types |
|-----------------------------|-------------|-------------------|-----------------|
| Genital–/Oral– ^a | 1746 (78.4) | None | None |
| Genital+/Oral– ^a | 333 (15.0) | Types not shown | Types not shown |
| Genital–/Oral+ ^a | 106 (4.8) | Types not shown | Types not shown |
| Genital+/Oral+ ^a | 43 (1.9) | | |
| Full concordance | 26 (1.1) | | |
| | 18 | HPV-3 | HPV-3 |
| | 4 | HPV-10 | HPV-10 |
| | 3 | HPV-57 | HPV-57 |
| | 1 | HPV-29 | HPV-29 |
| Partial concordance | 1 (0.0) | HPV-3 and 16 | HPV-3 |
| No concordance | 16 (0.7) | | |
| | 4 | HPV-16 | HPV-3 |
| | 1 | HPV-18 | HPV-3 |
| | 1 | HPV-33 | HPV-3 |
| | 1 | HPV-43 | HPV-3 |
| | 1 | HPV 84 | HPV-3 |
| | 1 | HPV-91 | HPV-3 |
| | 1 | HPV-30 | HPV-10 |
| | 1 | HPV-33 | HPV-16 |
| | 1 | HPV-3 | HPV-75 |
| | 1 | HPV-18 | HPV-75 |
| | 1 | HPV-66 | HPV-75 |
| | 1 | HPV-16 and 73 | HPV-3 |
| | 1 | HPV-43 and 45 | HPV-3 |

Table 2. Concordance and discordance of type-specific HPV types detected in paired oral and genital specimens from individual male participants in rural Anyang, China, 2009–2011. Abbreviation: HPV, human papillomavirus. ^aKappa = –1.460; Percent agreement = 80.3%; Percent positive agreement = 8.9%; McNemar chi-square = 117.380, *P* < 0.001; Pearson chi-square = 16.345, *P* < 0.001.

Among concurrent non-oncogenic HPV types found in this population, HPV-3, which has been rarely reported elsewhere, was the most dominant type. Since antibody against HPV-3 L1 was also frequently detected in serum from the same cohort²¹, it is possible that this type is relatively more prevalent in China. Apart from differences in populations, the frequent identification of HPV-3 may be partly explained by the use of different HPV testing methods. Compared to hybridization with pre-assigned

| Analysis | HPV status of external genitalia | HPV status of oral cavity | | Total | Crude OR (95% CI) ^a | Adjusted OR (95% CI) ^b |
|----------------------------|----------------------------------|---------------------------|----------|-------|--------------------------------|-----------------------------------|
| | | Negative | Positive | | | |
| Overall | | | | | | |
| | Negative | 1746 | 106 | 1852 | 1.0 | 1.0 |
| | Positive | 333 | 43 | 376 | 2.1 (1.5, 3.1) | 2.3 (1.6–3.4) |
| | Total | 2079 | 149 | 2228 | | |
| Type-specific ^c | | | | | | |
| | Negative | 95247 | 405 | 95652 | 1.0 | 1.0 |
| | Positive | 125 | 27 | 152 | 50.8 (33.1, 77.8) | 55.7 (36.1, 86.0) |
| | Total | 95372 | 432 | 95804 | | |

Table 3. Association of HPV infection in paired oral and genital specimens from individual male participants in rural Anyang, 2009–2011. Abbreviation: HPV, human papillomavirus; OR, odds ratio; CI, confidence interval. ^aCrude ORs and 95% CIs were calculated by univariate logistic regression analyses with generalized estimating equations (GEE). ^bAdjusted ORs and 95% CIs were calculated by multivariate logistic regression analyses with GEE including age, education level, marital status, type of employment, cigarette smoking, alcohol consumption, history of oral disease, number of missing teeth, frequency of washing genitalia before sex, oral sex practices, and lifetime number of sexual partners. ^cNumber of observations for type-specific analysis = 43 (total number of types detected among oral and genital specimens) × 2228 (number of individuals).

probes, the method adopted in this study using general primer mediated PCR followed by sequencing may maximize identification of the spectrum of HPV types including HPV-3. As a member of Alpha-2 clade, HPV-3 is typically found in benign forms of epidermodysplasia verruciformis (EV) and can induce flat, wart-like lesions over the body^{22,23}. However, the pathological consequences of HPV-3 infection in oral mucosa remain unknown and warrant further investigation.

In this study, significantly higher prevalence of type-specific HPV in oral specimens was found among men positive for genital HPV of the same type, which is consistent with some but not all previous surveys^{10,14,16}. The biological mechanism of this correlation is unclear. However, it may reflect the fact that HPV infections at these two sites in the same individual interact with each other and thus share similarities in their natural history of infection. Oral sex with a genital HPV-positive partner, or oral-oral contact with an oral HPV-positive partner may give rise to concurrent infection with this observed type specificity. In addition to partner to partner transmission, it is also possible that HPV infection at one site can be acquired by autoinoculation from the other site (e.g. via fingers) in the same individual¹¹. In support of the clinical significance of the association between oral and genital HPV infection, a study using the Surveillance, Epidemiology, and End Results (SEER) cancer database of United States found that while there was no elevated risk of HPV-unrelated cancer (e.g. prostate or bladder) following anogenital cancer, the incidence of a second primary oral cavity/pharyngeal cancer was significantly increased following an index genital cancer²⁴. In order to develop more efficient and cost-effective strategies for control of HPV-associated cancer, further research is warranted to pinpoint the major route(s) of oral-genital HPV transmission.

Many studies have suggested that both oral and genital HPV infections are mainly sexually transmitted^{7,25}. However, studies regarding the risk factors for concurrence of oral-genital HPV infections are few in number. As expected, we found that individuals with multiple lifetime sexual partners have a higher likelihood of coinfection. Nevertheless, we did not find a significant association of oral sex and oral-genital coinfection. This result is similar with that in our previous study of oral HPV infection¹⁷. The possibility of self-reporting bias which might lead to exposure misclassification and thus result in dilution of risk estimates to the null cannot be excluded. However, efforts were made to minimize this bias (e.g. conducting interviews in private settings). Another possible explanation is insufficient statistical power, as oral sex is rare among rural Chinese population whose sexual activity is considerably more conservative as compared with westerners. Recent studies have observed an absence of any relationship of oral sex and oral HPV infection^{6,26}, implying there are alternative modes of transmission (nonsexual routes). Thus, more adequately-powered studies are needed to clarify these associations.

This study has limitations. First, the nonparticipation and exclusion of participants because of specimen inadequacy may undermine the validity for generalizability of study results to some extent. Second, self-reported sexual behavior features may be subject to reporting bias. Additionally, despite a large population sample size, the low prevalence of oral-genital coinfection may limit our ability to identify some associations. This low prevalence of coinfection may also have rendered subgroup analysis of carcinogenic HPV impossible. Finally, due to the cross-sectional nature of this study, temporal relationship cannot be inferred. A longitudinal cohort study with rigorous follow-up is needed to evaluate the relationship of HPV acquisition at different anatomic sites and its relevant factors.

In conclusion, our data suggested that HPV infections of the oral cavity and the male genitalia in the same individual were type-specifically correlated. Concurrence of oral-genital infections was associated with multiple sexual partners. Given that HPV-related oropharyngeal cancer appears to be on the increase, improved insight into the relationship of HPV infection at the oral cavity and other anatomical sites may help to inform and direct future prevention efforts, such as HPV vaccination.

Materials and Methods

Study population. A population-based prospective cohort study of esophageal cancer and its associated determinants has been published previously²⁷. The current investigation was carried out in 6 of the 9 villages composing the parent cohort study in 2009–2011. Eligibility criteria were as follows: 1) male permanent residents in the target villages; 2) 25–65 years of age; 3) no prior diagnosis of cancer, mental illness, or cardiovascular disease; and 5) no history of HBV, HCV, or HIV infection. All participants provided written informed consent, and the study was approved by the Institutional Review Board of the School of Oncology, Peking University. The methods were carried out in accordance with the approved guidelines.

Specimen and data collection. Exfoliated cells from oral cavity and genitalia were collected by an experienced clinician using saline-moistened cotton swabs, as described previously^{17,25}. Oral specimens were collected from the buccal mucosa, palate, top and bottom of the tongue, the inner upper and lower lips, and the gingival surfaces by rubbing five times at each site. Exfoliated cells from the male genital shaft, glans penis, coronal sulcus, and scrotum were collected in the same manner. The cells collected on these swabs were then rinsed in 0.9% saline solution. After centrifugation at 3000 g for 10 minutes at 4 °C, the supernatant was decanted and the cell pellet was frozen at –70 °C until DNA was extracted.

A one-on-one computer-aided interview was carried out by a trained interviewer of male gender. Information regarding demographic characteristics, cigarette consumption (defined as 1 cigarette or more each day for at least 1 year), alcohol consumption (defined as use of Chinese liquor twice each week or more for at least 1 year), characteristics of sexual behavior (history of oral sex practices and lifetime number of sexual partners), and self-reported history of oral disease in the preceding 12 months (e.g. ulcers, gum disease, or chronic inflammation) was collected. Numbers of missing teeth were also recorded.

Laboratory procedure. DNA was extracted and purified on a Biomek 3000 automated workstation with the E.Z.N.A.™ Mag-Bind Tissue DNA Kit (Omega Bio-Tek, Inc.). The β -globin gene was evaluated in all purified DNA samples (forward primer, 5'-GAAGAGCCAAGGACAGGTAC-3'; reverse primer, 5'-CAACTTCATCCAGGTTACC-3'). β -globin PCR mixture contained 50 ng template DNA, 0.2 μ M each primer, 0.2 mM deoxynucleoside triphosphate, 1 \times PCR buffer containing 1.5 mM MgCl₂, and 0.8 U HotStarTaq polymerase (Qiagen, Germany). After a 15-min enzyme activation at 95 °C, 40 cycles of amplification (94 °C for 40 s, 58 °C for 40 s, and 72 °C for 40 s) and a final extension at 72 °C for 10 min were carried out. Detection of HPV DNA was performed for the β -globin positive samples by amplifying a fragment of 184 bp in the L1 gene using a highly sensitive PCR primer set (SPF1/GP6+). The PCR was performed in a reaction mixture containing 50 ng template DNA, 0.4 μ M each primer, 0.15 mM deoxynucleoside triphosphate, 1 \times PCR buffer containing 1.5 mM MgCl₂, and 0.8 U HotStarTaq polymerase. PCR conditions were as follows: preheating for 15 min at 95 °C, 45 cycles of amplification (94 °C for 40 s, 49 °C for 50 s, and 72 °C for 30 s), and a final extension at 72 °C for 10 min. PCR products of HPV positive specimens were purified and then directly sequenced using an ABI 3730XL DNA Analyzer²⁵. A broad spectrum of HPV types was identifiable using SPF1/GP6+-mediated PCR and sequencing method (Oncogenic: HPV-16, 18, 26, 31, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68; non-oncogenic: HPV-2, 3, 6, 7, 10, 11, 27, 29, 30, 32, 33, 37, 40, 42, 43, 44, 53, 54, 55, 57, 61, 62, 67, 69, 70, 72, 73, 74, 75, 77, 81, 82, 83, 84, 87, 89, 90, 91, 94, and 127)^{22,25,28}. Samples were considered to be HPV positive if any of these types were detected. Rigorous quality control procedures were implemented to avoid potential contamination as previously described^{17,29}.

Statistical analysis. Kappa value, percent agreement, and percent positive agreement were calculated to assess the concordance of HPV infection in paired oral and genital specimens from the same individuals. The McNemar's test was used to evaluate the difference in the positivity of HPV infection in the oral cavity and on the external genitalia. To evaluate type-specific positive concordance, the analyzed observations were restricted to those positive for at least one site (excluding observations which were negative for HPV infection at both oral cavity and external genitalia) and a Monte-Carlo simulation method (1000 iterations) was used for comparisons of observed concordant events and expected value^{30,31}.

To identify independent determinants for HPV infection, exposure variables that showed statistical significance in univariate logistic regression analysis, together with factors previously reported to be relevant (e.g., age, gender, smoking, and sexual behaviors), were included in multivariate logistic regression models (Variance inflation factor was used to examine whether variables in multivariate models were collinear). To examine linear trends, ordered categorical variables were treated as continuous covariates in the regression analysis.

For type-specific analysis to assess whether the presence of one specific HPV type in the oral cavity was associated with the presence of the same type in the genitalia, generalized estimating equations with an exchangeable correlation structure were used to account for possible correlations of different types of HPV. For groups of HPV types (i.e. any HPV type), infection of each type among the analyzed group for each individual was treated as one observation in the type-specific analysis. For instance, for any HPV type, each subject could have 43 HPV-type outcomes (a total of 43 HPV types were detected in this population).

All tests were two-sided and differences with $P < 0.05$ were considered statistically significant. Statistical analyses were conducted using Stata 11.0 (Stata Corp.).

References

- Cogliano, V. *et al.* Carcinogenicity of human papillomaviruses. *Lancet Oncol* **6**, 204 (2005).
- de Martel, C. *et al.* Global burden of cancers attributable to infections in 2008: a review and synthetic analysis. *Lancet Oncol* **13**, 607–615 (2012).
- Chaturvedi, A. K. *et al.* Worldwide trends in incidence rates for oral cavity and oropharyngeal cancers. *J Clin Oncol* **31**, 4550–4559 (2013).
- Torre, L. A. *et al.* Global cancer statistics, 2012. *CA Cancer J Clin.* **65**, 87–108 (2015).
- Bui, T. C., Markham, C. M., Ross, M. W. & Mullen, P. D. Examining the Association between Oral Health and Oral HPV Infection. *Cancer Prev Res (Phila)* **6**, 917–924 (2013).
- Kreimer, A. R. *et al.* The epidemiology of oral HPV infection among a multinational sample of healthy men. *Cancer Epidemiol Biomarkers Prev.* **20**, 172–182 (2011).
- Gillison, M. L. *et al.* Prevalence of oral HPV infection in the United States, 2009–2010. *JAMA* **307**, 693–703 (2012).
- Pickard, R. K., Xiao, W., Broutian, T. R., He, X. & Gillison, M. L. The prevalence and incidence of oral human papillomavirus infection among young men and women, aged 18–30 years. *Sex Transm Dis.* **39**, 559–566 (2012).
- Lang Kuhs, K. A. *et al.* Prevalence of and risk factors for oral human papillomavirus among young women in Costa Rica. *J Infect Dis.* **208**, 1643–1652 (2013).
- Dahlstrom, K. R. *et al.* Sexual Transmission of Oral Human Papillomavirus Infection among Men. *Cancer Epidemiol Biomarkers Prev.* **23**, 2959–2964 (2014).
- Hernandez, B. Y. *et al.* Transmission of human papillomavirus in heterosexual couples. *Emerg Infect Dis.* **14**, 888–894 (2008).
- Xavier, S. D. *et al.* Prevalence of human papillomavirus (HPV) DNA in oral mucosa of men with anogenital HPV infection. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **108**, 732–737 (2009).
- van Doornum, G. J. *et al.* Prevalence of human papillomavirus infections among heterosexual men and women with multiple sexual partners. *J Med Virol.* **37**, 13–21 (1992).
- Widdice, L. E. *et al.* Human papillomavirus concordance in heterosexual couples. *J Adolesc Health* **47**, 151–159 (2010).
- Sirera, G. *et al.* High prevalence of human papillomavirus infection in the anus, penis and mouth in HIV-positive men. *AIDS* **20**, 1201–1204 (2006).
- Videla, S. *et al.* Natural history of human papillomavirus infections involving anal, penile, and oral sites among HIV-positive men. *Sex Transm Dis.* **40**, 3–10 (2013).
- Hang, D. *et al.* Oral Human Papillomavirus Infection and Its Risk Factors among 5,410 Healthy Adults in China, 2009–2011. *Cancer Epidemiol Biomarkers Prev* **23**, 2101–2110 (2014).
- Fakhry, C. *et al.* Comparison of the immune microenvironment of the oral cavity and cervix in healthy women. *Cytokine* **64**, 597–604 (2013).
- Termine, N. *et al.* Low rate of oral human papillomavirus (HPV) infection in women screened for cervical HPV infection in Southern Italy: A cross-sectional study of 140 immunocompetent subjects. *J Med Virol* **81**, 1438–1443 (2009).
- Kreimer, A. R. Prospects for prevention of HPV-driven oropharynx cancer. *Oral Oncol* **50**, 555–559 (2014).
- Zhang, C. *et al.* Seroprevalence of 10 human papillomavirus types in the general rural population of Anyang, China: a cross-sectional study. *Sex Transm Infect* (2015). doi:10.1136.
- de Villiers, E. M., Fauquet, C., Broker, T. R., Bernard, H. U. & zur Hausen, H. Classification of papillomaviruses. *Virology* **324**, 17–27 (2004).
- Leiding, J. W. & Holland, S. M. Warts and all: human papillomavirus in primary immunodeficiencies. *J Allergy Clin Immunol* **130**, 1030–1048 (2012).
- Sikora, A. G., Morris, L. G. & Sturgis, E. M. Bidirectional association of anogenital and oral cavity/pharyngeal carcinomas in men. *Arch Otolaryngol Head Neck Surg* **135**, 402–405 (2009).
- He, Z. *et al.* Human Papillomavirus Genital Infections among Men, China, 2007–2009. *Emerg Infect Dis.* **19**, 992–995 (2013).
- Kreimer, A. R. *et al.* Incidence and clearance of oral human papillomavirus infection in men: the HIM cohort study. *Lancet* **382**, 877–887 (2013).
- Liu, F. *et al.* The Anyang Esophageal Cancer Cohort Study: study design, implementation of fieldwork, and use of computer-aided survey system. *Plos One* **7**, e31602 (2012).
- Huang, S. L. *et al.* Comparison between the Hybrid Capture II Test and an SPF1/GP6 + PCR-based assay for detection of human papillomavirus DNA in cervical swab samples. *J Clin Microbiol* **44**, 1733–1739 (2006).
- He, Z. *et al.* Evaluation of quality control procedures in an oesophageal cancer cohort study in Anyang, China. *J Hosp Infect* **76**, 336–339 (2010).
- Burchell, A. N., Tellier, P. P., Hanley, J., Coutlee, F. & Franco, E. L. Human papillomavirus infections among couples in new sexual relationships. *Epidemiology* **21**, 31–37 (2010).
- Zeger, S. L. & Liang, K. Y. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics* **42**, 121–130 (1986).

Acknowledgements

We thank Dr. Michael A. McNutt for editing and correction of this manuscript. This work was supported by Natural Science Foundation of China [30872937, 81502855, 30930102]; “973” Project of National Ministry of Science and Technology Grant [2011CB504300, 2012CB910800]; Charity Project of National Ministry of Health [201202014, 200902002]; and Natural Science Foundation of Beijing [7100001].

Author Contributions

Y.K. and H.C. were involved in the design and supervision of the survey. F.L., D.H., Q.D., M.L., L.X., Z.H., C.Z., M.S., Y.L., J.L., Y.P., T.N., C.G., Y.L., R.X. and L.Z. were involved in conducting the field work and laboratory testing of specimens. F.L., D.H., and Q.D. performed the statistical analyses and wrote the manuscript text. All authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Liu, F. *et al.* Concurrence of oral and genital human papillomavirus infection in healthy men: a population-based cross-sectional study in rural China. *Sci. Rep.* **5**, 15637; doi: 10.1038/srep15637 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>