SCIENTIFIC REPORTS

OPEN

Received: 13 December 2014 Accepted: 16 November 2015 Published: 15 December 2015

Responses of photosynthetic parameters to drought in subtropical forest ecosystem of China

Lei Zhou¹, Shaoqiang Wang¹, Yonggang Chi², Qingkang Li¹, Kun Huang¹ & Quanzhou Yu¹

The mechanism underlying the effect of drought on the photosynthetic traits of leaves in forest ecosystems in subtropical regions is unclear. In this study, three limiting processes (stomatal, mesophyll and biochemical limitations) that control the photosynthetic capacity and three resource use efficiencies (intrinsic water use efficiency (iWUE), nitrogen use efficiency (NUE) and light use efficiency (LUE)), which were characterized as the interactions between photosynthesis and environmental resources, were estimated in two species (*Schima superba* and *Pinus massoniana*) under drought conditions. A quantitative limitation analysis demonstrated that the drought-induced limitation of photosynthesis in *Schima superba* was primarily due to stomatal limitation, whereas for *Pinus massoniana*, both stomatal and non-stomatal limitations generally exhibited similar magnitudes. Although the mesophyll limitation represented only 1% of the total limitation in *Schima superba*, it accounted for 24% of the total limitations for *Pinus massoniana*. Furthermore, a positive relationship between the LUE and NUE and a marginally negative relationship or trade-off between the NUE and iWUE were observed in the control plots. However, drought disrupted the relationships between the resource use efficiencies. Our findings may have important implications for reducing the uncertainties in model simulations and advancing the understanding of the interactions between ecosystem functions and climate change.

Water deficit is the primary factor that limits ecosystem productivity in most terrestrial biomes¹. The physiological responses of trees to drought (i.e., carbon uptake) are directly related to vegetation growth², ecosystem productivity^{3,4}, frequency of fires^{5,6} and tree mortality^{7,8}. The subtropical region experiences frequent seasonal droughts⁹ that result in declines in terrestrial carbon sequestration¹⁰. However, the mechanism underlying the effects of drought on the carbon uptake of subtropical ecosystems at the leaf level remains unclear¹¹.

The carbon uptake of forest ecosystems is driven by leaf photosynthesis, the responses of which to drought are mediated by three physiological processes. First, stomatal closure is recognized as the main driver of the photosynthetic response to water stress by limiting CO₂ diffusion from the atmosphere to the substomatal cavities to slow photosynthesis^{12,13}. Second, the mesophyll conductance (g_m) may rapidly decrease, thereby limiting CO₂ diffusion from the substomatal cavities to the chloroplast stroma during water stress^{14,15}. Finally, photosynthesis may be limited by biochemical processes in long-lasting, severe droughts, resulting in decreased photosynthetic enzyme activity (i.e., the maximum rate of Rubisco carboxylation, V_{cmax}), ribulose-1,5-bisphophate (RuBP) regeneration capacity (i.e., the maximum rate of photosynthetic electron transport, J_{max}) and triose-phosphate utilization (TPU)¹⁶⁻¹⁸. As a result, drought stress directly influences CO₂ diffusion and/or the biochemical process of photosynthesis, which in turn limits the net CO₂ assimilation rate (A_n). For example, Maseda and Fernandez (2006) found that the rapid closure of stomata during water stress resulted in a decline in transpiration and the A_n ¹⁹. Increasing evidence has shown that mesophyll conductance is finite²⁰ and plays an important role in limiting the photosynthetic capacity¹². Additionally, drought-stressed plants exhibit significant reductions in V_{cmax} , J_{max} and *TPU* relative to plants with sufficient water²¹, indicating that biochemical processes dramatically inhibit photosynthesis during long-term severe droughts. These apparent discrepancies may arise from the fact that photosynthesis induced

¹Key Laboratory of Ecosystem Network Observation and Modelling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China. ²State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China. Correspondence and requests for materials should be addressed to S.W. (email: sqwang@igsnrr.ac.cn)

| | | Schima superba | | | | Pinus massoniana | | | |
|----------------|-------------------|---------------------|--|---|-------------------------------------|---------------------|---|--|-------------------------------------|
| Treatments | SWC (g g^{-1}) | SLA $(m^2 kg^{-1})$ | $C_{\rm area} ({ m g}{ m C}{ m m}^{-2})$ | $N_{\rm area}({ m g}{ m N}{ m m}^{-2})$ | Leaf C/N ratio (g g ⁻¹) | SLA $(m^2 kg^{-1})$ | $C_{\rm area}({ m g}{ m C}{ m m}^{-2})$ | $N_{\rm area} ({ m g}{ m N}{ m m}^{-2})$ | Leaf C/N ratio (g g ⁻¹) |
| Control | 0.21 ± 0.01 | 10.81 ± 0.58 | 47.01 ± 2.19 | 1.67 ± 0.09 | 28.27 ± 0.62 | 6.88 ± 0.24 | 75.84 ± 3.10 | 2.25 ± 0.09 | 33.86 ± 1.03 |
| Drought | 0.13 ± 0.01 | 10.85 ± 0.32 | 46.27 ± 1.35 | 1.54 ± 0.04 | 30.08 ± 0.57 | 7.52 ± 0.41 | 70.50 ± 4.68 | 2.10 ± 0.13 | 33.90 ± 1.37 |
| <i>p</i> value | 0.000 | 0.959 | 0.777 | 0.191 | 0.044 | 0.262 | 0.351 | 0.351 | 0.982 |

Table 1. Soil water content and leaf traits of *Schima superba* and *Pinus massoniana* grown in control and drought plots. Note: The drought effects on the soil water content (SWC), specific leaf area (SLA), C concentration (C_{area}), N concentration (N_{area}) and Leaf C/N ratio were analyzed for *Schima superba* and *Pinus massoniana* using an independent sample T-test. Significant values (P < 0.05) are shown bold (Mean \pm SE, N=12).

by drought stress is not limited by a single process. Instead, the combined effect of the stomatal, mesophyll, and biochemical limitations simultaneously regulates the decrease in photosynthesis in response to water stress^{21–23}. However, a quantitative limitation analysis is needed to separate the three physiological processes in subtropical climatic zones^{12,22,24}.

Drought not only decreases the leaf photosynthetic rate but also regulates the interaction of plant carbon uptake and environmental resources, which is termed the resource use efficiency. Three resource use efficiencies (water use efficiency (WUE), carbon gain at the expense of water loss; nitrogen use efficiency (NUE), carbon gain per nitrogen content; and light use efficiency (LUE), carbon gain per available light quantum flux density) are important functional parameters that intimately couple the uptake of carbon with the major growth limiting factors (water, nitrogen and light). In general, the leaf-level WUE has been reported to increase during soil water stress^{25,26}, which suggests that stomata closure in response to H₂O flux is more sensitive than the response to carbon flux²⁷. Apart from increasing the WUE, stomatal closure during drought stress has an effect on photosynthesis but no effect on leaf nitrogen, leading to a decline in the NUE^{26,27}. During drought stress periods, the LUE generally decreases with increasing drought intensity; indeed, no change was observed in the electron transport rate under mild and moderate water stress²⁸, or the electron transport rate declined to a lesser extent than the net CO₂ assimilation rate²⁹. The changes in single resource use efficiency induced by drought have been well documented^{25,26,28}, but the trade-off among the multiple resource use efficiencies of plants requires investigation.

Based on a recent integration of eddy covariance observations, subtropical forests in the East Asian region exhibit a high carbon dioxide uptake rate $(362 \text{ g C m}^{-2} \text{ year}^{-1})$ compared with Asian tropical and temperate forests³⁰. A model simulation indicated that drought caused the net exchange of carbon in the subtropical forests in Southern China to decrease by 63% and 47% in 2003 and 2004, respectively³¹. Despite the ecological importance of this region, the carbon uptake response of subtropical forests to drought is poorly constrained. To address this knowledge gap, a rainfall exclusion experiment was established in 2010 using two gradients of soil moisture content and three replicates. The three limiting processes (i.e., stomatal limitation, mesophyll limitation and biochemical limitation) that control photosynthesis and the three resource use efficiencies (iWUE, NUE and LUE) that represent the interaction of photosynthesis and environmental resources were estimated after a 3-year drought. We focused on (1) testing the sensitivity of the photosynthetic characteristics of *Schima superba* and *Pinus massoniana* in subtropical regions of China during an experimental drought and (2) determining the changes in the iWUE, NUE and LUE in response to water stress and identifying whether a trade-off existed among resource use efficiencies.

Results

The response of soil water content and leaf chemical characteristics to drought. The experimental drought significantly reduced the soil water content by 38% (t = 9.840; P < 0.0001) (Table 1). In general, the leaf traits (i.e., SLA, C concentration, N concentration and C/N ratio) for each species were not affected by the drought based on an independent sample T-test, except that a significant difference in the C/N ratio for *Schima superba* (P = 0.044) occurred between the control and drought plots (Table 1).

The response of the carbon assimilation process to drought. Based on an independent sample T-test, significant effects of drought on the A_n of *Schima superba* (t = 3.080, P = 0.005) and *Pinus massoniana* (t = 3.769, P = 0.001) were observed, which showed significant reductions during drought treatment (Fig. 1a,e). However, no significant effects of soil moisture on the R_d were observed (both P > 0.05) for either species (Fig. 1b,f). For *Schima superba* and *Pinus massoniana*, the responses of the A_g to the soil water treatments were significant (t = 3.134, P = 0.005 and t = 3.867, P = 0.001) and resulted in significant decreases in the drought plots compared with the control plots (Fig. 1c,g). Although there were no significant differences in R_d/A_g between drought treatments for each species, a general increasing trend from the control to the drought plots was indicated (Fig. 1d,h). No significant differences in drought resistance for A_n , R_d , A_g , and R_d/A_g were observed between *Schima superba* and *Pinus massoniana* (Fig. 1i–l).

The response of the CO₂ diffusion process to drought. Drought produced a 42% decrease in the g_s (t = 2.709, P = 0.013) for *Schima superba* (Fig. 2a), whereas the effect of drought on the g_s for *Pinus massoniana* was not significantly different between the control and drought plots (P > 0.05) (Fig. 2d). The responses of the g_m to drought were not significantly different for either species, based on an independent sample T-test (both P > 0.05) (Fig. 2b,e). However, significant decreases in the g_{tot} were observed (*Schima superba*: t = 2.618, P = 0.016; *Pinus massoniana*: t = 3.583, P = 0.002) in the drought plots relative to the control plots (Fig. 2c,f). The drought



Figure 1. The effect of drought on the carbon assimilation process for the two species. (a,e) The A_n (net CO_2 assimilation rate, μ mol $CO_2 m^{-2} s^{-1}$), (b,f) R_d (day respiration, μ mol $CO_2 m^{-2} s^{-1}$), (c,g) A_g (gross CO_2 assimilation, μ mol $CO_2 m^{-2} s^{-1}$) and (d,h) ratio of R_d and A_g in the control and drought plots for *Schima* superba (a–d) and *Pinus massoniana* (e–h) are shown. The drought resistance of (i) A_n , (j) R_d , (k) A_g and (l) R_d/A_g in *Schima superba* and *Pinus massoniana* is indicated. ANOVA: *P < 0.05; **P < 0.01; and ***P < 0.001.



Figure 2. The effect of drought on the CO₂ diffusion process for the two species. (a,d) The g_s (stomatal conductance, mmol CO₂ m⁻² s⁻¹), (**b**,**e**) g_m (mesophyll conductance, mmol CO₂ m⁻² s⁻¹) and (**c**,**f**) g_{tot} (the total conductance, mmol CO₂ m⁻² s⁻¹) in the control and drought plots of *Schima superba* (**a**-**c**) and *Pinus massoniana* (**d**-**f**) are shown. The drought resistance of (**g**) g_s , (**h**) g_m and (**i**) g_{tot} in *Schima superba* and *Pinus massoniana* is indicated. ANOVA: *P < 0.05; **P < 0.01; and ***P < 0.001.

resistance of *Pinus massoniana* with regard to the g_s appeared to be considerably higher than that of *Schima superba* (P = 0.065), whereas no significant differences in drought resistance related to the g_m and g_{tot} were found between the species (Fig. 2g–i).

The response of biochemical processes to drought. The effects of drought on V_{cmax} and J_{max} were not significantly different between the control and drought plots for *Schima superba* and *Pinus massoniana* (all P > 0.05, Figure S1a-b, e-f). For *Schima superba*, a significant increasing trend in $J_{\text{max}}/V_{\text{cmax}}$ (t = -2.229, P = 0.036) was observed from the control to the drought plots, whereas no significant effect of drought on $J_{\text{max}}/V_{\text{cmax}}$ was observed



Figure 3. The quantitative limitation of A_n (net CO₂ assimilation rate, μ mol CO₂ m⁻² s⁻¹) for Schima superba (Ss) and Pinus massoniana (Pm) during drought stress. The stomatal limitation (S_L , red area), mesophyll conductance limitation (MC_L , yellow area) and biochemical limitation (B_L , green area) are shown. The total diffusional limitation ($D_L = S_L + MC_L$, blue area) and the non-stomatal limitation ($NS_L = MC_L + B_L$, sky blue area) are also indicated.

for *Pinus massoniana* (P > 0.05). No significant differences in drought resistance were observed for V_{cmax} , J_{max} , TPU and J_{max}/V_{cmax} between *Schima superba* and *Pinus massoniana* (Figure S1i-I).

Quantitative limitation analysis. For *Schima superba*, the values of S_L , MC_L and B_L accounted for 87%, 1% and 12% of the limitations, respectively (Fig. 3). The contributions of the stomatal (S_L) and non-stomatal limitations ($NS_L = MC_L + B_L$) represented approximately seven-eighths and one-eighth of the total limitation, respectively. The role of the total diffusional limitation ($D_L = S_L + MC_L$) was more important than that of the biochemical (B_L) limitation.

For *Pinus massoniana*, the S_L , MC_L and B_L values were equal to 54%, 24%, and 22% of the limitations, respectively. The stomatal (S_L) and non-stomatal limitations (NS_L) generally showed a similar magnitude. The contributions of the total diffusional limitation (D_L) and biochemical limitation (B_L) represented approximately two-thirds and one-third of the total limitation, respectively.

Resource use efficiency. A general increasing trend the in iWUE from the control to the drought plots was observed for *Schima superba* (P = 0.056), whereas significant declines in the iWUE due to the drought treatments (P = 0.021) were observed for *Pinus massoniana* (Fig. 4a,d). The effects of drought on the NUE of *Schima superba* and *Pinus massoniana* exhibited significant reductions from the control to drought plots (Fig. 4b,e). No significant differences were observed in the LUE for *Schima superba* and *Pinus massoniana* during drought treatments (all P > 0.05, Fig. 4c,f). The drought resistance of the iWUE exhibited significant differences between species, whereas no significant differences in drought resistance related to the LUE and NUE were observed between species (Fig. 4g-i).

To determine the trade-off in resource use efficiency, simple linear regressions of LUE vs. NUE and NUE vs. iWUE were performed (Fig. 5). A significant positive correlation was found between the LUE and NUE in the control plots (y = 7.29x - 0.06, $R^2 = 0.57$, P < 0.0001). However, poor correlations between the LUE and NUE were found for all species in the drought plots ($R^2 = 0.06$, P = 0.237). As a result, the regression slopes of the LUE and NUE were different between the control and drought treatments (P = 0.034). A marginally negative relationship between the NUE and iWUE was observed for all species in the control plots (y = -0.06x + 0.09, $R^2 = 0.12$, P = 0.092), whereas the correlations between the NUE and iWUE were weak ($R^2 = 0.02$, P = 0.481) in the drought plots. However, no significant difference was observed in the regression slopes of the NUE and iWUE for the control and drought treatments (P = 0.179).

Discussion

Quantitative limitation analysis of photosynthesis in response to drought. As expected, drought stress significantly decreased the leaf photosynthesis of the dominate species (Schima superba and Pinus massoniana) in subtropical forests. Our study reported that an approximately 37% decrease in the A_n in the drought plots was related to a decrease of approximately 38% in the SWC. The pattern of decreasing A_n with drought was similar to the pattern observed in forests under field conditions^{24,32,33}. However, the degree to which drought affected the A_n did not significantly vary between Schima superba and Pinus massoniana (Fig. 1i). Two independent experiments on mesic and xeric species from diverse hydroclimates in Australia and Europe indicated interspecific differences in the drought response²³. The velocity of the photosynthetic changes in response to water stress imposition were faster in evergreen forests than in semi-deciduous forests, although the declines in photosynthetic rate were similar in magnitude³⁴. Although water stress is known to reduce the photosynthetic rate, the processes responsible for the key limitations are still a matter of debate³⁵. Previous studies have demonstrated that the photosynthetic reaction to water stress is dominated by only one of the three physiological processes (stomatal conductance, mesophyll conductance and biochemical processes)^{36–39}. Increasing evidence had shown that the combined effect of the stomatal, mesophyll, and biochemical limitations simultaneously regulates the decline in photosynthesis in response to drought. Thus, a quantitative limitation analysis of the changes in the photosynthetic rate in response to water stress was necessary^{12,40,41}. Our quantitative limitation analysis demonstrated that the drought-induced limitation of photosynthesis in *Schima superba* was mainly due to the stomatal limitation (87%), whereas the stomatal (S_1 ,



Figure 4. The effect of drought on the resource use efficiency for the two species. (a,d) The iWUE (the water use efficiency, μ mol CO₂ mmol H₂O⁻¹), (b,e) NUE (the nitrogen use efficiency, μ mol CO₂ mol N⁻¹) and (c,f) LUE (the light use efficiency, μ mol CO₂ μ mol Photons⁻¹) in the control and drought plots of *Schima superba* (a-c) and *Pinus massoniana* (d-f) are shown. The drought resistance of the (g) iWUE, (h) NUE, and (i) LUE in *Schima superba* and *Pinus massoniana* is indicated. ANOVA: **P* < 0.05; ***P* < 0.01; and ****P* < 0.001.

54%) and non-stomatal limitations (NS_L , 46%) for *Pinus massoniana* generally showed similar magnitudes. A study of temperate deciduous forests reported a maximum value of 50% for the S_L during drought stress³⁹. A study of tropical evergreen forests (*Campsiandra laurifolia*, *Symmeria paniculata*, *Acosmium nitens* and *Eschweilera tenuifolia*) indicated that the S_L accounted for 30–39% of the limitations in the dry season (March 2004)⁴². Therefore, the magnitude of the photosynthetic reaction to water stress for the three types of limitations varied between species.

Mesophyll conductance is typically absent in gas exchange measurements, which are assumed to be infinite⁴³. However, studies have demonstrated that changes in mesophyll conductance in response to stress and limit photosynthesis are an important physiological process^{12,44}. For a variety of climate zones and species, the MC_L is responsible for approximately 14-30% of the limitations^{12,22-24}. Similarly, Grassi and Magnani (2005) found a maximum value of 14% for the MC_L for an ash forest under seasonal drought¹². Another rainfall exclusion experiment in a *Quercus ilex* forest showed a maximum value of 20% for the MC_L^{22} . Our quantitative limitation analysis showed that the MC_L was responsible for only 1% of the total limitation of *Schima superba*, although this limitation accounted



Figure 5. The relationships between the LUE (light use efficiency, μ mol CO₂ μ mol Photons⁻¹) and NUE (nitrogen use efficiency, μ mol CO₂ mol N⁻¹) (**a**) or the NUE and iWUE (water use efficiency, μ mol CO₂ mmol H₂O⁻¹) (**b**) in the control and drought plots. *Ss* and *Pm* represent *Schima superba* and *Pinus massoniana*, respectively. The dotted line indicates the linear regression for the control plots.

for 24% of the total limitation for *Pinus massoniana*. Therefore, it is important to include mesophyll conductance into any detailed study of the gas exchange response to drought and the processed model.

An explanation for the discrepancies in the $MC_{\rm L}$ between both species in the subtropical region may be related to the different phylogenetic clades (gymnosperms vs. angiosperms)⁴⁵. The SLA was reported to strongly separate gymnosperms from angiosperms based on 305 North American woody species that spanned boreal to subtropical climates⁴⁶. The $g_{\rm m}$ can be influenced by changes in leaf anatomical characteristics, such as the thickness of leaf/ mesophyll cell walls/chloroplasts and the stomata density⁴⁷⁻⁴⁹. Gymnosperms have a lower SLA value⁵⁰, lower mesophyll porosity, thicker mesophyll cell wall⁵¹, and lower $g_{\rm m}^{51,52}$ than angiosperms. In our study, the gymnosperm species (*Pinus massoniana*) had lower SLA and $g_{\rm m}$ values than angiosperms, which might contribute to the high $MC_{\rm L}$ in evergreen conifers.

The balance between J_{max} and V_{cmax} . The J_{max} and V_{cmax} relationship represents the resource allocation between the two photosynthetic cycles: electron transport and the Calvin cycle⁵³. In the biochemically-based photosynthesis model, V_{cmax} was scaled to J_{max} based on the hypothesis that the average J_{max}/V_{cmax} ratio was 2.1⁵⁴. However, some studies have demonstrated that the J_{max} to V_{cmax} ratio is not constant but varies with temperature⁵⁵, leaf nitrogen⁵⁶, and species⁵⁷. The underlying processes responsible for the changes in the J_{max}/V_{cmax} ratio due to water stress are still a matter of debate. We found that the J_{max}/V_{cmax} ratio varied considerably among drought treatments in *Schima superba* (Fig. S1d), which was consistent with previous studies^{24,58}. The hypothesis that droughts modify the balance between RuBP carboxylation and regeneration was supported by our study (i.e., the resource allocation between the two photosynthetic cycles (the Calvin cycle and electron transport) was changed). Current ecosystem models are less capable of accounting for climate extremes (warming, nitrogen deposition or drought), which attributes to model calibration using data collected under standard climate conditions⁵⁹. The findings from manipulation experiments could reduce the uncertainties of model parameter estimates and the predictions made by the models.

The trade-off between resource use efficiencies (iWUE, NUE and LUE). In the present study, we found a significant positive correlation between the LUE and NUE for all species in the control plots (Fig. 5a). Generally, the plants tended to obtain the maximum NUE with an increase in the LUE⁶⁰. However, a poor correlation was observed between the LUE and NUE in the drought plots, indicating that drought stress might have weakened the relationship between the LUE and NUE in these plots (Fig. 5a). The slopes of the regression lines (LUE vs. NUE) from the control and drought plots were significantly different (P=0.034). Furthermore, the NUE but not the LUE was significantly decreased in the drought treatments for each species (Fig. 4b,c,e,f). These results suggested that the responses of the LUE and NUE to drought were uncoupled. The use of resources (light and nitrogen) for plants may also be influenced by other factors, such as water stress.

Additionally, a marginally negative correlation was observed between the NUE and iWUE in the control plots for the two species studied here, which provided evidence for the existence of a trade-off between the uses of resources. An increase in the NUE together with a decrease in the iWUE indicated that compromises existed between the iWUE and NUE. This finding was consistent with previous studies of a large number of plants species⁶¹⁻⁶³. An increase in the efficiency of the use of one resource can lead to a reduction in the efficiency of use of another resource⁶⁰, suggesting that maximization of resource use efficiency depends on the most limited resources³³. This trade-off may help enforce the species distribution across moisture and nutrient gradients⁶².



Figure 6. Rainfall exclusion experiments at the QYZ station: (a) Control plot and (b) Drought plot. Photo credit: Lei Zhou.

Methods

Experimental site. The experimental site was located in the Qianyanzhou (QYZ) Forest Experimental Station (26°44'N, 115°03'E) in Jiangxi Province of South China and belongs to Chinaflux. The mean annual air temperature is approximately 17.9 °C, and the highest and lowest daily temperatures recorded are 39.5 and -5.8 °C, respectively⁶⁴. The annual average precipitation is 1,489 mm and mainly occurs from March to June (52% of total). The annual air temperature and annual precipitation in 2012 were 18.5 °C and 1741 mm, respectively. These characteristics are typical of the prevailing subtropical monsoon climate. The soil mainly consists of the red soil type formed principally from red sandstone, sand gravel or mudstone and river alluvial deposits. The main forest types include Masson's pine (*Pinus massoniana*), slash pine (*Pinus elliottii*), Chinese fir (*Cunninghamia lanceolata*), and Schima (*Schima superba*). *Pinus massoniana* and *Schima superba* are the pioneer species and the dominant species in this region, respectively. Both species were mixed as conifer-broadleaf forests, with a tree density of stems of 700 ha⁻¹.

Rainfall exclusion experiment. The rainfall exclusion experiment was initiated in January 2010. Rainfall was withheld for the entire year with no changes in other meteorological variables (Fig. 6). We used transparent polyester film placed at a height of 2 m on the trees to partially exclude throughfall drops onto the floor and allow light penetration. Trenches were dug around the perimeters of the plots to reduce the lateral inflow of water from the surrounding forest into the plots. Three rainfall exclusion plots $(12 \text{ m} \times 12 \text{ m})$ were used. Three control plots were established in the same environment. An average of three *Schima superba* and five *Pinus massoniana* were growing in each plot. The height and diameter at breast height for each species in the control and drought plots are listed in Table S1. The plots were established in theW middle of the forests, and we attempted to select trees in the middle of the plots.

CO₂ response curve. We measured the foliar gas exchange of two species (*Schima superba* and *Pinus mas*soniana) using a portable photosynthesis system (LI-6400, LI-COR Inc., USA) during the growing season (early September) of 2012 (Figure S2). Measurements were performed with two portable photosynthesis systems on clear sunny days between 8:00 a.m. to 2:00 p.m. Twelve individuals (4 individuals per plot) for each species were measured in each treatment (control and drought plots). All measurements were performed on fully expanded leaves with no signs of senescence or immaturity. Branches with sun-facing leaves were excised from the middle of the crown with the help of a lopper (3 m) affixed to a bamboo shoot (approximately 10 m) (Figure S3) and then immediately stored in a bottle with fresh water. The gas exchange measurements were conducted soon after branch excision. The time period from branch excision to the completion of the measurement was typically less than 30 min. Typical A_n/C_i curves (A_n versus the calculated intercellular CO₂ concentrations, C_i) were measured at the ambient CO_2 concentration (C_a) (ranging from 50 to 1400 μ mol mol⁻¹). The C_a was lowered stepwise from 400 to $50\,\mu\text{mol}\,\text{mol}^{-1}$ and then increased again from 50 to $1400\,\mu\text{mol}\,\text{mol}^{-1}$, with a total of 10 points. Photosynthesis was induced for 10 min at the saturating photosynthetically active photon flux density (PPFD) (1500 µmol photons $m^{-2} s^{-1}$) at a given leaf temperature (25 °C). The CO₂ concentrations in the cuvette were controlled using an injector system (LI-6400-01, LI-COR Inc.) that used a CO₂ mixer and compressed CO₂ cartridges. The PPFD was provided by the red/blue LED light source built into the foliar cuvette (LI-6400-02B, LI-COR Inc.) and was calibrated against an internal photodiode. The average value of the air temperature on the measurement days was 24.1 °C. The leaf temperature in the cuvette, which was controlled by the thermoelectric cooling elements of the Li-6400, was 25 °C. The cuvette was sealed with plasticine to prevent leakage. We placed twelve needles of Pinus massoniana side by side into a 2×3 cm sealed cuvette. The cuvette was sealed with plasticine to prevent leakage^{65,66}.

 A_n/C_i curves were fitted to estimate the V_{cmax} , TPU and g_m using spreadsheet-based software by minimizing the root mean square error (RMSE) of each curve⁶⁷. The g_s (mmolH₂O m⁻² s⁻¹) was initially measured with the Li-6400. Due to the differences in diffusion coefficients between water vapor and CO₂, the stomatal conductance to H₂O was 1.6-fold higher than the stomatal conductance to CO₂⁶⁸. To achieve consistency with the g_m , we converted the $g_{s,w}$ to $g_{s,c}$. The A_n measured at the 400 µmol mol⁻¹ CO₂ concentration and 25 °C leaf temperature from each A_n/C_i curve was used to track the differences between treatments for each species. The total conductance (g_{tot}) was calculated from the sum of the g_s and g_m .

Light response curve. The light response curves of *Schima superba* and *Pinus massoniana* were measured using the LI-6400 after each A_n/C_i curve measurement. The PPFD was sequentially lowered from 1800 to 0μ mol m⁻² s⁻¹, with a total of 14 points. During each measurement, the CO₂ concentration was maintained at 400 μ mol mol⁻¹, and the leaf temperature was maintained at 25 °C.

A non-rectangular hyperbola model⁶⁹ solved for its negative root was used to describe the light response curves. In our study, the leaf maximum apparent quantum yield of CO₂ uptake (AQY, μ mol CO₂ m⁻²s⁻¹) and R_d (μ mol CO₂ m⁻²s⁻¹) were derived from the light response curve. The gross CO₂ assimilation (A_g) was calculated by adding R_d to A_n.

Soil water content and leaf chemical characteristics. The gravimetric soil water content (SWC) was measured at the depth of 0–20 cm in the field. Soil samples from each plot were placed into aluminum boxes and then dried in an oven at 106 °C for 24 h. The SWC in this study was expressed as follows:

$$SWC(gg^{-1}) = (W_1 - W_2)/W_2$$
 (1)

where W_1 is the sample fresh weight, and W_2 is the sample dry weight. Finally, eight soil moisture contents were averaged to represent the water content of each plot.

The foliage covered in the cuvette during the gas exchange measurements was used to measure the leaf C and N concentrations (C_{area} and N_{area}) with respect to area. Foliage was removed from branches after the gas exchange measurements, and then the area was measured with an area meter (LI-3100, Li-Cor Inc.). Foliage samples were dried at 65 °C for 48 h, and the leaf characteristics were measured with a CN analyzer. The leaf C/N ratio with respect to area was calculated using the leaf C and N concentrations. The specific leaf area (SLA) was calculated based on the measurements of the leaf area and dry mass.

Data analyses. The relative limitation to assimilation imposed by the stomatal conductance (S_L), mesophyll conductance (MC_L) and biochemical processes (B_L) were separated using the approach¹² proposed by Grassi and Magnani (2005). This approach makes it possible to compare relative limitations to assimilation, which partitions photosynthetic limitations into components related to stomatal conductance, mesophyll conductance, and leaf biochemical characteristics. The non-stomatal limitation (NS_L) was defined as the sum of the contribution due to the mesophyll conductance and leaf biochemistry ($NS_L = MC_L + B_L$). The total diffuse limitation (D_L) was the sum of the stomatal and mesophyll conductance components ($D_L = S_L + MC_L$).

The leaf-level intrinsic WUE (iWUE, μ mol CO₂ mmol H₂O⁻¹) was expressed as the ratio of the net CO₂ assimilation rate versus the stomatal conductance. The NUE (μ mol CO₂ mol N⁻¹) for leaf photosynthesis was defined as the ratio of the net photosynthesis rate to the leaf nitrogen content. The AQY derived from the light response curve was the proxy for the leaf-level LUE (μ mol CO₂ μ mol Photons⁻¹)⁷⁰ in this study.

The drought resistance in this study was expressed as the ratio of the variables in drought plots to those in control plots^{71,72} (i.e., variable_drought/variable_control). Values closer to 1 imply greater drought resistance.

We performed independent sample T-tests with a 95% confidence level to examine the differences in each variable among treatments. Previously, the homogeneity of variables was evaluated with Levene's test (P < 0.05). If the homogeneity test failed, the variables were log-transformed or sin-transformed prior to analysis. Regression models were used to determine the relationship between different resource use efficiencies. The general linear model (GLM) was used to test the significance of the slopes of the linear regression among resource use efficiencies. All statistical analyses were performed using SPSS Version 14.0 (SPSS Inc. Chicago, IL, USA).

References

- 1. Nemani, R. R. *et al.* Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science* **300**, 1560–1563 (2003).
- Pasho, E., Camarero, J. J., de Luis, M. & Vicente-Serrano, S. M. Impacts of drought at different time scales on forest growth across a wide climatic gradient in north-eastern Spain. Agr. Forest Meteorol. 151, 1800–1811 (2011).
- 3. Ciais, P. et al. Europe-wide reduction in primary productivity caused by the heat and drought in 2003. Nature 437, 529–533 (2005).
- 4. Zhao, M. S. & Running, S. W. Drought-Induced reduction in global terrestrial net primary production from 2000 through 2009. *Science* **329**, 940–943 (2010).
- Nepstad, D. C., Tohver, I. M., Ray, D., Moutinho, P. & Cardinot, G. Mortality of large trees and lianas following experimental drought in an Amazon forest. *Ecology* 88, 2259–2269 (2007).
- 6. Phillips, O. L. et al. Drought Sensitivity of the Amazon Rainforest. Science 323, 1344-1347 (2009).
- Adams, H. D. et al. Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under globalchange-type drought. Proc. Natl. Acad. Sci. USA 106, 7063–7066 (2009).
- Anderegg, W. R. L., Kane, J. M. & Anderegg, L. D. L. Consequences of widespread tree Mortality triggered by drought and temperature stress. Nat. Clim. Change 3, 30–36 (2013).
- Sun, X. M., Wen, X. F., Yu, G. R., Liu, Y. F. & Liu, Q. J. Seasonal drought effects on carbon sequestration of a mid-subtropical planted forest of southeastern China. Sci. China Ser. D 49, 110–118 (2006).
- Wen, X. F., Wang, H. M., Wang, J. L., Yu, G. R. & Sun, X. M. Ecosystem carbon exchanges of a subtropical evergreen coniferous plantation subjected to seasonal drought, 2003–2007. *Biogeosciences* 7, 357–369 (2010).
- 11. Domingues *et al.* Seasonal patterns of leaf-level photosynthetic gas exchange in an eastern Amazonian rain forest. *Plant Ecol. Divers.* 7, 189–203 (2014).
- 12. Grassi, G. & Magnani, F. Stomatal, mesophyll conductance and biochemical limitations to photosynthesis as affected by drought and leaf ontogeny in ash and oak trees. *Plant Cell Environ* **28**, 834–849 (2005).
- Egea, G., Verhoef, A. & Vidale, P. L. Towards an improved and more flexible representation of water stress in coupled photosynthesisstomatal conductance models. Agr. Forest Meteorol. 151, 1370–1384 (2011).
- Flexas, J., Ribas-Carbo, M., Diaz-Espejo, A., Galmes, J. & Medrano, H. Mesophyll conductance to CO₂: current knowledge and future prospects. *Plant Cell Environ.* 31, 602–621 (2008).
- Flexas, J. et al. Mesophyll diffusion conductance to CO₂: An unappreciated central player in photosynthesis. Plant Sci. 193–194, 70–84 (2012).

- Bota, J., Medrano, H. & Flexas, J. Is photosynthesis limited by decreased Rubisco activity and RuBP content under progressive water stress? New Phytol. 162, 671–681 (2004).
- 17. Keenan, T. *et al.* Improved understanding of drought controls on seasonal variation in Mediterranean forest canopy CO_2 and water fluxes through combined *in situ* measurements and ecosystem modelling. *Biogeosciences* **6**, 1423–1444 (2009).
- Galmés, J., Flexas, J., Savé, R. & Medrano, H. Water relations and stomatal characteristics of Mediterranean plants with different growth forms and leaf habits: responses to water stress and recovery. *Plant Soil* 290, 139–155 (2007).
- 19. Maseda, P. H. & Fernandez, R. J. Stay wet or else: three ways in which plants can adjust hydraulically to their environment. *J. Exp Bot.* 57, 3963–3977 (2006).
- 20. Loreto, F., Harley, P. C., Dimarco, G. & Sharkey, T. D. Estimation of Mesophyll Conductance to CO₂ Flux by 3 Different Methods. *Plant Physiol.* **98**, 1437–1443 (1992).
- Martin-StPaul, N. K. *et al.* Photosynthetic sensitivity to drought varies among populations of Quercus ilex along a rainfall gradient. *Funct. Plant Biol.* 39, 25–37 (2012).
- 22. Limousin, J. M., Misson, L., Lavoir, A. V., Martin, N. K. & Rambal, S. Do photosynthetic limitations of evergreen Quercus ilex leaves change with long-term increased drought severity? *Plant Cell Environ.* **33**, 863–875 (2010).
- Zhou, S. X., Medlyn, B., Sabaté, S., Sperlich D. & Prentice, I. C. Short-term water stress impacts on stomatal, mesophyll and biochemical limitations to photosynthesis differ consistently among tree species from contrasting climates. *Tree Physiol.* 34, 1035–1046. (2014).
- Wilson, K. B., Baldocchi, D. D. & Hanson, P. J. Spatial and seasonal variability of photosynthetic parameters and their relationship to leaf nitrogen in a deciduous forest. *Tree Physiol.* 20, 565–578 (2000).
- Delucia, E. H. & Heckathorn, S. A. The Effect of Soil Drought on Water-Use Efficiency in a Contrasting Great-Basin Desert and SierranMontane Species. *Plant Cell Environ.* 12, 935–940 (1989).
- Reich, P. B., Walters, M. B. & Tabone, T. J. Response of Ulmus-Americana Seedlings to Varying Nitrogen and Water Status. 2. Water and Nitrogen Use Efficiency in Photosynthesis. *Tree Physiol.* 5, 173–184 (1989).
- Warren, C. R. & Adams, M. A. Internal conductance does not scale with photosynthetic capacity: implications for carbon isotope discrimination and the economics of water and nitrogen use in photosynthesis. *Plant Cell Environ.* 29, 192–201 (2006).
- 28. Flexas, J., Bota, J., Escalona, J. M., Sampol, B. & Medrano, H. Effects of drought on photosynthesis in grapevines under field conditions: an evaluation of stomatal and mesophyll limitations. *Funct. Plant Biol.* **29**, 461–471 (2002).
- Flexas, J., Escalona, J. M. & Medrano, H. Water stress induces different levels of photosynthesis and electron transport rate regulation in grapevines. *Plant Cell Environ.* 22, 39–48 (1999).
- Yu, G. R. et al. High carbon dioxide uptake by subtropical forest ecosystems in the East Asian monsoon region. Proc. Natl. Acad. Sci. USA 111, 4910–4915 (2014).
- 31. Gu, F. X. *et al.* Drought effects on carbon exchange in a subtropical coniferous plantation in China. *J. Plant Ecol.* **32**, 1041–1051 (2008) (in chinese with english abstract).
- 32. Ni, B. R. & Pallardy, S. G. Stomatal and Nonstomatal Limitations to Net Photosynthesis in Seedlings of Woody Angiosperms. *Plant Physiol.* **99**, 1502–1508 (1992).
- Xu, L. K. & Baldocchi, D. D. Seasonal trends in photosynthetic parameters and stomatal conductance of blue oak (Quercusdouglasii) under prolonged summer drought and high temperature. *Tree Physiol.* 23, 865–877 (2003).
- 34. Galle, A., Florez-Sarasa, I., Aououad, H. E. & Flexas, J. The Mediterranean evergreen *Quercus ilex* and the semi-deciduous *Cistus albidus* differ in their leaf gas exchange regulation and acclimation to repeated drought and re-watering cycles. *J. Exp. Bot.* **62**, 5207–5216 (2011).
- Flexas, J. & Medrano, H. Drought-inhibition of photosynthesis in C-3 plants: Stomatal and non-stomatal limitations revisited. Ann. Bot. 89, 183–189 (2002).
- Cornic, G. [Drought stress and high light effects on leaf photosynthesis]Photoinhibition of photosynthesis [Baker, N. R. & Bowyer, J. R. (ed.)] [297–313] (Bios Scientific Publishers, Oxford, 1994).
- Lawlor, D. W. [The effects of water deficit on photosynthesis] Environment and Plant metabolism [Smirnoff, N. (ed.)] [129–160] (Bios Science Publishers, Oxford, 1995).
- 38. Chaves, M. M. et al. How plants cope with water stress in the field. Photosynthesis and growth. Ann. Bot. 89, 907-916 (2002).
- Wilson, K. B., Baldocchi, D. D. & Hanson, P. J. Quantifying stomatal and non-stomatal limitations to carbon assimilation resulting from leaf aging and drought in mature deciduous tree species. *Tree Physiol.* 20, 787–797 (2000).
- Diaz-Espejo, A., Nicolás, E. & Enrique Fernández, J. E. Seasonal evolution of diffusional limitations and photosynthetic capacity in olive under drought. *Plant Cell Environ.* 30, 922–933 (2007).
- Varone, L. et al. Stomatal and non-stomatal limitations to photosynthesis in seedlings and saplings of Mediterranean species preconditioned and aged in nurseries: Different response to water stress. Environ. Exp. Bot. 75, 235–247 (2012).
- Herrera, A., Tezara, W., Mari'n, O. & Rengifo, E. Stomatal and non-stomatal limitations of photosynthesis in trees of a tropical seasonally flooded forest. *Physiol. Plantarum* 134, 41–48 (2008).
- Farquhar, G. D., Caemmerer, S. V. & Berry, J. A. A Biochemical-Model of Photosynthetic Co₂ Assimilation in Leaves of C-3 Species. *Planta* 149, 78–90 (1980).
- Niinemets, U., Cescatti, A., Rodeghiero, M. & Tosens, T. Leaf internal diffusion conductance limits photosynthesis more strongly in older leaves of Mediterranean evergreen broad-leaved species. *Plant Cell Environ.* 28, 1552–1566 (2005).
- Flexas, J. et al. Stomatal and mesophyll conductances to CO₂ in different plant groups: Underrated factors for predicting leaf photosynthesis responses to climate change? Plant Sci. 226, 41–48 (2014).
- 46. Stahl, U. *et al.* Whole-plant trait spectra of North American woody plant species reflect fundamental ecological strategies. *Ecosphere* **4**, 128. http://dx.doi.org/10.1890/ES13-00143.1 (2013).
- 47. Tholen, D. & Zhu, X. G. The mechanistic basis of internal conductance. Plant Physiol. 156, 90-105 (2011).
- Tosens T., Niinemets Ü., Vislap V., Eichelmann H. & Castro-Díez P. Developmental changes in mesophyll diffusion conductance and photosynthetic capacity under different light and water availabilities in Populus tremula: how structure constrains function. *Plant Cell Environ.* 35, 839–856 (2012).
- Scafaro, A. P., von Caemmerer, S., Evans, J. R. & Atwell, B. J. Temperature response of mesophyll conductance in cultivated and wild Oryza species with contrasting mesophyll cell wall thickness. *Plant Cell Environ.* 34, 1999–2008 (2011).
- Lusk, C. H., Wright, I. J. & Reich, P. B. Photosynthetic differences contribute to competitive advantage of evergreen angiosperms over evergreen conifers. New Phytol. 160, 329–336 (2003).
- De Lucia E. H., Whitehead, D. & Clearwater, M. J. The relative limitation of photosynthesis by mesophyll conductance in co-occurring species in a temperate rainforest dominated by the conifer Dacrydiumcupressinum. *Funct. Plant Biol.* 30, 1197–1204 (2003).
- Tomás *et al.* Importance of leaf anatomy in determining mesophyll diffusion conductance to CO₂ across species: quantitative limitations and scaling up by models. J. Exp. Bot. 64, 2269–2281 (2013).
- Walker, A. P. et al. The relationship of leaf photosynthetic traits-Vcmax and Jmax-to leaf nitrogen, leaf phosphorus, and specific leaf area: a meta-analysis and modeling study. Ecol. Evol. 4, 3218–3235 (2014).
- Wullschleger, S. D. Biochemical limitations to carbon assimilation in C3 plants a retrospective analysis of the A/ci curves from 109 species. J. Exp. Bot. 44, 907–920 (1993).
- 55. Čhi, Y. *et al.* Acclimation of foliar respiration and photosynthesis in response to experimental warming in a temperate steppe in northern China. *PLOS One* **8**, e56482, doi: 10.1371/journal.pone.0056482 (2013).

- Yamori, W., Nagai, T. & Makino A. The rate-limiting step for CO₂ assimilation at different temperatures is influenced by the leaf nitrogen content in several C3 crop species. *Plant Cell Environ.* 34, 764–777 (2011).
- Miao, Z. W., Xu, M., Lathrop, R. G. & Wang, Y. F. Comparison of the A-Cc curve fitting methods in determining maximum ribulose 1.5-bisphosphate carboxylase/oxygenase carboxylation rate, potential light saturated electron transport rate and leaf dark respiration. *Plant Cell Environ.* 32, 109–122 (2009).
- Medlyn, B. E. et al. Temperature response of parameters of a biochemically based model of photosynthesis. II. A review of experimental data. Plant Cell Environ. 25, 1167–1179 (2002).
- Misson, L., Limousin, J. M., Rodriguez, R. & Letts, M. G. Leaf physiological responses to extreme droughts in Mediterranean Quercus ilex forest. *Plant Cell Environ.* 33, 1898–1910 (2010).
- 60. Hirose, T. & Bazzaz, F. A. Trade-off between light- and nitrogen-use efficiency in canopy photosynthesis. Ann. Bot. 82, 195–202 (1998).
- Field, C., Merino, J. & Mooney, H. A. Compromises between Water-Use Efficiency and Nitrogen-Use Efficiency in 5 Species of California Evergreens. Oecologia 60, 384–389 (1983).
- Patterson, T. B., Guy, R. D. & Dang, Q. L. Whole-plant nitrogen- and water-relations traits, and their associated trade-offs, in adjacent muskeg and upland boreal spruce species. *Oecologia* 110, 160–168 (1997).
- Chen, S. P., Bai, Y. F., Zhang, L. X. & Han, X. G. Comparing physiological responses of two dominant grass species to nitrogen addition in Xilin River Basin of China. *Environ. Exp. Bot.* 53, 65–75 (2005).
- 64. Wang, S. Q., Liu, J. Y., Zhang, C., Yi, C. X. & Wu, W. X. Effects of afforestation on soil carbon turnover in China's subtropical region. *J. Geogr. Sci*, 21, 118–134 (2011).
- Tjoelker, M. G., Oleksyn, J., Lorenc-Plucinska, J. & Reich, P. B. Acclimation of respiratory temperature responses in northern and southern populations of *Pinus banksiana*. New Phytol. 181, 218–229 (2009).
- Tjoelker, M. G., Oleksyn, J., Reich, P. B. & Zytkowiak, R. Coupling of respiration, nitrogen, and sugars underlies convergent temperature acclimation in *Pinus banksiana* across wide-ranging sites and populations. *Global Change Biol.* 14, 782–797 (2008).
- Sharkey, T. D., Bernacchi, C. J., Farquhar, G. D. & Singsaas E. L. Fitting photosynthetic carbon dioxide response curves for C-3 leaves. Plant Cell Environ. 30, 1035–1040 (2007).
- Damour, G., Simonneau, T., Cochard, H. & Urban, L. An overview of models of stomatal conductance at the leaf level. *Plant Cell Environ.* 33, 1419–1438 (2010).
- Prioul, J. L. & Chartier, P. Partitioning of transfer and carboxylation components of intracellular resistance to photosynthetic CO₂ fixation: a critical analysis of the methods used. Ann. Bot. 41, 789–800 (1977)
- 70. Milcu, A. et al. Functional diversity of leaf nitrogen concentrations drives grassland carbon fluxes. Ecol Lett. 17, 435–444 (2014).
- 71. Tilman, D. & Downing, J. A. Biodiversity and Stability in Grasslands. Nature 367, 363-365 (1994).
- 72. Lan, Z. C. & Bai, Y. F. Testing mechanisms of N-enrichment-induced species loss in a semiarid Inner Mongolia grassland: critical thresholds and implications for long-term ecosystem responses. *Philos. T. R. SOC. B.* **367**, 3125–3134 (2012).

Acknowledgements

This study was supported by the National Natural Science Foundation of China (Grant No. 41401110 and 31400393), Strategic Priority Research Program of the CAS (Grant No. XDA05050602 and XDA05050702), the Key Project in the National Science & Technology Pillar Program of China (Grant No. 2013BAC03B00), the National Natural Science Foundation of China (Grant No. 30970484), and the National Key Technologies R & D Program (Grant NO. 2012BAC11B05).

Author Contributions

L.Z., Y.G.C. and Q.K.L. conceived and designed the experiment. L.Z., S.Q.W. and Y.G.C. analyzed the data and wrote the paper. L.Z., Y.G.C., K.H. and Q.Z.Y. performed the field investigation and sample analyses and contributed to the draft of the manuscript.

Additional Information

Supplementary information accompanies this paper at http://www.nature.com/srep

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhou, L. *et al.* Responses of photosynthetic parameters to drought in subtropical forest ecosystem of China. *Sci. Rep.* **5**, 18254; doi: 10.1038/srep18254 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/