

SCIENTIFIC REPORTS

OPEN

Corrigendum: Soot superaggregates from flaming wildfires and their direct radiative forcing

Rajan K. Chakrabarty, Nicholas D. Beres, Hans Moosmüller, Swarup China, Claudio Mazzoleni, Manvendra K. Dubey, Li Liu & Michael I. Mishchenko

Scientific Reports 4:5508; doi: 10.1038/srep05508; published online 01 July 2014; updated on 10 March 2016

The original version of this Article contained an error in the Abstract.

“We estimate that SAs contribute, per unit optical depth, up to 35% less atmospheric warming than freshly-emitted ($D_f \approx 1.8$) aggregates, and $\approx 90\%$ more warming than the volume-equivalent spherical soot particles simulated in climate models.”

now reads:

“At 550 nm wavelength, we estimate that SAs contribute, per unit optical depth, up to 35% less atmospheric warming than freshly-emitted ($D_f \approx 1.8$) aggregates, and $\approx 90\%$ more warming than the volume-equivalent spherical soot particles simulated in climate models.”

Additionally, the original version of this Article contained an error in the Methods section.

“The parameter β is the up-scatter fraction, which is a function of asymmetry parameter g as follows:”

now reads:

“The parameter β is the up-scatter fraction, which is a function of asymmetry parameter g using the Henyey-Greenstein phase function as follows:”

These errors have been corrected in the HTML version of the Article but have not been corrected in the PDF version of the Article.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>