

SCIENTIFIC REPORTS



OPEN

On Wiener polarity index of bicyclic networks

Jing Ma¹, Yongtang Shi¹, Zhen Wang² & Jun Yue³

Received: 16 October 2015

Accepted: 02 December 2015

Published: 11 January 2016

Complex networks are ubiquitous in biological, physical and social sciences. Network robustness research aims at finding a measure to quantify network robustness. A number of Wiener type indices have recently been incorporated as distance-based descriptors of complex networks. Wiener type indices are known to depend both on the network's number of nodes and topology. The Wiener polarity index is also related to the cluster coefficient of networks. In this paper, based on some graph transformations, we determine the sharp upper bound of the Wiener polarity index among all bicyclic networks. These bounds help to understand the underlying quantitative graph measures in depth.

In order to decide whether a given network is robust, a way to quantitatively measure network robustness is needed. Intuitively robustness is all about back-up possibilities, or alternative paths, but it is a challenge to capture these concepts in a mathematical formula. During the past years a lot of robustness measures have been proposed¹. Network robustness research is carried out by scientists with different backgrounds, like mathematics, physics, computer science and biology. As a result, quite a lot of different approaches to capture the robustness properties of a network have been undertaken. All of these approaches are based on the analysis of the underlying graph—consisting of a set of vertices connected by edges of a network^{1–6}.

One such category is the distance-based descriptors which include Wiener index, Harary index, etc. The use of Wiener index and related type of indices dates back to the seminal work of Wiener in 1947⁷. Wiener introduced his celebrated index to predict the physical properties, such as boiling point, heats of isomerization and differences in heats of vaporization, of isomers of paraffin by their chemical structures. Wiener index has since inspired many distance-based descriptors in Chemometrics. These include Harary index⁸, hyper Wiener index^{9,10}, Wiener polynomial¹¹, Balaban index¹², Wiener polarity index⁷ and information indices^{13–15}. These indices, or commonly called descriptors, play significant roles in quantitative structure-activity relationship/quantitative structure-property relationship (QSAR/QSPR) models. It is known that the Wiener type indices depend both on a network's number of nodes and its topology. For more results, we refer to^{16,17}.

Let $G = (V, E)$ be a connected simple graph. The distance between two vertices u and v in G , denoted by $d_G(u, v)$, is the length of a shortest path between u and v in G . The Wiener polarity index of a graph $G = (V, E)$, denoted by $W_p(G)$, is the number of unordered pairs of vertices $\{u, v\}$ of G such that $d_G(u, v) = 3$, i.e.,

$$W_p(G) = |\{\{u, v\} | d(u, v) = 3, u, v \in V(G)\}|. \quad (1)$$

The name “Wiener polarity index” is introduced by Harold Wiener⁷ in 1947. Wiener himself conceived the index only for acyclic molecules and defined it in a slightly different – yet equivalent – manner. In the same paper, Wiener also introduced another index for acyclic molecules, called *Wiener index* or *Wiener distance index* and defined by $W(G) = \sum_{\{u, v\} \subseteq V} d_G(u, v)$. Wiener⁷ used a linear formula of W and W_p to calculate the boiling points t_B of the paraffins, i.e., $t_B = aW + bW_p + c$, where a , b and c are constants for a given isomeric group. The Wiener index $W(G)$ is popular in chemical literatures. For more results on Wiener index, we refer to the survey paper¹⁸ written by Dobrynin, Entringer and Gutman, and some recent papers^{19–23}.

The Wiener polarity index is used to demonstrate quantitative structure-property relationships in a series of acyclic and cycle-containing hydrocarbons by Lukovits and Linert²⁴. Hosoya in²⁵ found a physical-chemical interpretation of $W_p(G)$. Du, Li and Shi²⁶ described a linear time algorithm APT for computing the Wiener polarity index of trees, and characterized the trees maximizing the Wiener polarity index among all trees of given order. From then on, the Wiener polarity index started to attract the attention of a remarkably large number of mathematicians and so many results appeared. The extremal Wiener polarity index of (chemical) trees with given

¹Center for Combinatorics and LPMC-TJKLC, Nankai University, Tianjin, 300071, China. ²Interdisciplinary Graduate School of Engineering Sciences, Kyushu University, Kasuga-koen, Kasugashi, Fukuoka, Japan. ³School of Mathematical Sciences, Shandong Normal University, Jinan 250014, Shandong, China. Correspondence and requests for materials should be addressed to Y.S. (email: shi@nankai.edu.cn)




Figure 1. The three types of bicyclic graphs.

different parameters (e.g. order, diameter, maximum degree, the number of pendants, etc.) were studied, see^{27–33}. Moreover, the unicyclic graphs minimizing (resp. maximizing) the Wiener polarity index among all unicyclic graphs of order n were given in³⁴. There are also extremal results on some other graphs, such as fullerenes, hexagonal systems and cactus graph classes, we refer to^{35–37}. Observe that the Wiener polarity index is also related to the cluster coefficient of networks.

Results

The main contributions of this paper can be summarized as follows:

- We provide a formula of the Wiener polarity index of bicyclic networks, from which the value of the index can be computed easily.
- We introduce three graph transformations, which can be used to increase the values of Wiener polarity index. These transformations can help to find more extremal values for other classes of molecular networks.
- We determine the maximum value of the Wiener polarity index of bicyclic networks and characterize the corresponding extremal graphs.

Now let us introduce some notations. Let $N_G(v)$ be the neighborhood of v , and $d_G(v) = |N_G(v)|$ denote the degree of vertex v . For $i = 2, 3, \dots$, we call $N_G^i(v) = \{u \in V(G) \mid d(u, v) = i\}$ the i th neighborhood of v . If $d_G(v) = 1$, then we call v a *pendant vertex* of G . Let $g(C_x)$ be the length of cycle C_x in graph G , P_i denote a path with length i . For all other notations and terminology, not given here, see e.g.³⁸.

Let B be a bicyclic graph. Suppose $C_p = v_1v_2 \dots v_pv_1$ and $C_q = u_1u_2 \dots u_qu_1$ are two cycles in B with l ($l \geq 0$) common vertices. Without loss of generality, we label the vertices of C_p in the clockwise direction, and the vertices of C_q in the inverse clockwise direction. If $l = 0$, then there is one unique path P connecting C_p and C_q , which starts with v_1 and ends with u_1 . We call this kind of bicyclic graph **type I** (see Fig. 1). If $l = 1$, then C_p and C_q have exactly one common vertex $v_1(u_1)$. We call this kind of bicyclic graphs **type II** (see Fig. 1). If $l \geq 2$, then B contains exactly three cycles. The third cycle is denoted by C_z , where $z = p + q - 2l + 2$. Without loss of generality, assume that $p \leq q \leq z$ and $l - 2 \leq p - 2 \leq q - 2$. The two cycles C_p and C_q have more than one common vertex $v_1(u_1), \dots, v_l(u_l)$. We call this kind of bicyclic graphs **type III** (see Fig. 1). In the following section, we use B, C_p, C_q, v_i ($1 \leq i \leq p$), u_j ($1 \leq j \leq q$), l as defined above, except as noted.

Let $C'_{3,3}(s_1, s_2, s_3; t_1, t_2, t_3)$ be the bicyclic graph of type I, where $P = v_1u_1$ and $s_1 + s_2 + s_3 + t_1 + t_2 + t_3 = n - 6$. Especially, we denote this kind of graphs by $C^*_{3,3}$, if $t_1 = t_2 = t_3 = 0$, $0 \leq s_i - s_j \leq 2$ ($i = 2, 3$), $|s_2 - s_3| \leq 1$. For a graph $G = (V, E)$ and $P_l = v_1v_2 \dots v_{l+1}$, we can construct a new graph H by identifying v_1 with $v \in G$, denoted by $H: = G + P_l$, and we say P_l is *incident* to vertex v .

Theorem 0.1. Let B_1 be a bicyclic graph in type I and $|V(B_1)| = n$ ($n \geq 6$), B_1^* be the desired graph attaining the maximum Wiener polarity index.

- (1) If $n = 6$, then $B_1^* = C'_{3,3}(0,0,0;0,0,0)$, and $W_p(B_1) = W_p(B_1^*) = 4$;
- (2) If $n = 7$, then $B_1^* \cong C'_{3,3}(1,0,0;0,0,0)$, and $W_p(B_1) \leq W_p(B_1^*) = 6$;
- (3) If $n = 8$, then $B_1^* \cong C'_{3,3}(1,0,0;1,0,0)$, $C'_{3,3}(1,0,0;0,0,0) + P_1$, where P_1 is incident to the pendant vertex of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 9$;
- (4) If $n = 9$, then $B_1^* \cong C'_{3,3}(2,0,0;1,0,0)$, $C'_{3,3}(1,0,0;1,0,0) + P_1$, where the path P_1 is incident to the pendant vertex of v_1 , $C'_{3,3}(2,0,0;0,0,0) + P_1$, where the path P_1 is incident to one pendant vertex of v_1 , $C'_{3,3}(1,0,0;0,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertex of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 12$;
- (5) If $n = 10$, then $B_1^* \cong C'_{3,3}(2,0,0;2,0,0)$, $C'_{3,3}(2,0,0;1,0,0) + P_1$, where the path P_1 is incident to one pendant vertex of v_1 , $C'_{3,3}(2,0,0;0,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 16$;
- (6) If $n = 11$, then $B_1^* \cong C'_{3,3}(3,0,0;2,0,0)$, $C'_{3,3}(2,0,0;2,0,0) + P_1$, where the path P_1 is incident to one pendant vertex of v_1 , $C'_{3,3}(2,0,0;1,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertices of v_1 , $C'_{3,3}(2,0,0;0,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 20$;
- (7) If $n = 12$, then $B_1^* \cong C'_{3,3}(3,0,0;3,0,0)$, $C'_{3,3}(3,0,0;2,0,0) + P_1$, where P_1 is incident to one pendant vertex of v_1 , $C'_{3,3}(3,0,0;1,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertices of v_1 , $C'_{3,3}(3,0,0;0,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 25$;

- (8) If $n = 13$, then $B_1^* \cong C_{3,3}^* C_{3,3}'(4,0,0;3,0,0), C_{3,3}'(3,0,0;3,0,0) + P_1$, where P_1 is incident to one pendent vertex of $v_1, C_{3,3}'(4,0,0;2,0,0) + P_1$, where P_1 is incident to one pendent vertex of $v_1, C_{3,3}'(4,0,0;1,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertices of $v_1, C_{3,3}'(3,0,0;2,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertices of $v_1, C_{3,3}'(4,0,0;0,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertices of $v_1, C_{3,3}'(3,0,0;1,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertices of $v_1, C_{3,3}'(4,0,0;0,0,0) + P_1 + P_1 + P_1 + P_1$, where the four paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 30$;
- (9) If $n = 14$, then $B_1^* \cong C_{3,3}^* C_{3,3}'(4,0,0;4,0,0), C_{3,3}'(4,0,0;3,0,0) + P_1$, where P_1 is incident to one pendent vertex of $v_1, C_{3,3}'(4,0,0;2,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertices of $v_1, C_{3,3}'(4,0,0;1,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertices of $v_1, C_{3,3}'(4,0,0;0,0,0) + P_1 + P_1 + P_1 + P_1$, where the four paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 36$;
- (10) If $n \geq 15$, then $B_1^* \cong C_{3,3}^*$, and $W_p(B_1) \leq W_p(B_1^*)$. \square

Let $C_{3,3}''(s_1, s_2, s_3; t_1, t_2, t_3)$ be the bicyclic graph in type II, where $s_1 + s_2 + s_3 + t_1 + t_2 + t_3 = n - 5$, and $s_1 = t_1$. When n is large enough, it can be easily checked that the graph maximizing the Wiener polarity index is $B_2^* = C_{3,3}''(s_1, s_2, s_3; s_1, 0, 0)$ (see support information).

Theorem 0.2. Let B_2 be a bicyclic graph in type II and $|V(B_2)| = n (\geq 5)$, B_2^* be the desired graph attaining the maximum Wiener polarity index.

- (1) If $n = 5$, then $B_2^* = B_2 = C_{3,3}''(0,0,0;0,0,0)$, and $W_p(B_2) = 0$;
- (2) If $n = 6$, then $B_2^* \cong C_{3,3}''(0,1,0;0,0,0), C_{3,4}''(0,0,0;0,0,0)$, and $W_p(B_2) \leq W_p(B_2^*) = 2$;
- (3) If $n = 7$, then $B_2^* \cong C_{3,3}''(0,1,1;0,0,0), C_{3,4}''(0,0,0;1,0,0)$, and $W_p(B_2) \leq W_p(B_2^*) = 5$;
- (4) If $n = 8$, then $B_2^* \cong C_{3,3}''(0,1,2;0,0,0), C_{3,4}''(0,0,0;1,0,1)$, and $W_p(B_2) \leq W_p(B_2^*) = 8$;
- (5) For $n \geq 9$, let $s_1 + s_2 + s_3 = 3k + r (r \in \{0,1,2\})$.

If $r = 0$, then $B_2^* \cong C_{3,3}''(k - 2, k + 1, k + 1; k - 2, 0, 0), C_{3,3}''(k - 1, k, k + 1; k - 1, 0, 0)$, and $W_p(B_2) \leq W_p(B_2^*) = 3k^2 + 4k + 1$;

If $r = 1$, then $B_2^* \cong C_{3,3}''(k - 1, k + 1, k + 1; k - 1, 0, 0)$, and $W_p(B_2) \leq W_p(B_2^*) = 3k^2 + 6k + 3$;

If $r = 2$, then $B_2^* \cong C_{3,3}''(k - 1, k + 1, k + 2; k - 1, 0, 0)$, and $W_p(B_2) \leq W_p(B_2^*) = 3k^2 + 8k + 5$. \square

Let $C_{3,3}'''(s_1, s_2, s_3; t_1, t_2, t_3)$ be the bicyclic graph in type III, where $s_1 + s_2 + s_3 + t_1 + t_2 + t_3 = n - 4$, $s_1 = t_1, s_2 = t_1$ and $l = 1$. Let $C_{3,4}'''(s_1, s_2, s_3; t_1, t_2, t_3, t_4)$ be the bicyclic graph in type III, where $s_1 + s_2 + s_3 + t_1 + t_2 + t_3 + t_4 = n - 5, s_1 = t_1, s_2 = t_1$ and $l = 1$. When n is large enough, it can be checked that the graph maximizing the Wiener polarity index is $B_3^* = C_{3,4}'''(s_1, s_2, s_3; s_1, s_2, 0, 0)$.

Theorem 0.3. Let B_3 be a bicyclic graph in type III and $|V(B_3)| = n (\geq 4)$, B_3^* be the desired graph attaining the maximum Wiener polarity index.

- (1) If $n = 4$, then $B_3^* = B_3 = C_{3,3}'''(0,0,0;0,0,0)$, and $W_p(B_3) = 0$;
- (2) If $n = 5$, then $B_3^* \cong C_{3,3}'''(0,0,1;0,0,0)$, and $W_p(B_3) \leq W_p(B_3^*) = 1$;
- (3) If $n = 6$, then $B_3^* \cong C_{3,3}'''(0,0,0;0,0,0) + P_2$, where P_2 is incident to vertex v_1 or v_3 , and $W_p(B_3) \leq W_p(B_3^*) = 3$;
- (4) If $n = 7$, then $B_3^* \cong C_{3,3}'''(1,0,0;1,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertex of v_1 , and $W_p(B_3) \leq W_p(B_3^*) = 6$;
- (5) If $n = 8$, then $B_3^* \cong C_{3,3}'''(1,0,0;1,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertex of v_1 , and $W_p(B_3) \leq W_p(B_3^*) = 9$;
- (6) If $n = 9$, then $B_3^* \cong C_{3,3}'''(1,0,0;1,0,0) + P_1 + P_1 + P_1 + P_1$, where the four paths P_1 are incident to the pendant vertex of $v_1, C_{3,3}'''(2,0,0;2,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_3) \leq W_p(B_3^*) = 12$;
- (7) If $n = 10$, then $B_3^* \cong C_{3,3}'''(2,0,0;2,0,0) + P_1 + P_1 + P_1 + P_1$, where the four paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_3) \leq W_p(B_3^*) = 16$;
- (8) If $n = 11$, then $B_3^* \cong C_{3,3}'''(2,0,0;2,0,0) + P_1 + P_1 + P_1 + P_1 + P_1$, where the five paths P_1 are incident to the pendant vertices of $v_1, C_{3,3}'''(3,0,0;3,0,0) + P_1 + P_1 + P_1 + P_1$, where the four paths P_1 are incident to the pendant vertices of $v_1, C_{3,4}'''(2,2,2;2,2,0,0), C_{3,4}'''(1,2,3;1,2,0,0)$, and $W_p(B_3) \leq W_p(B_3^*) = 20$;
- (9) For $n \geq 12$, let $s_1 + s_2 + s_3 = 3k + r (r \in \{0,1,2\})$.

If $r = 0$, then $B_3^* \cong C_{3,4}'''(k - 1, k - 1, k + 2; k - 1, k - 1, 0, 0), C_{3,4}'''(k - 1, k, k + 1; k - 1, k, 0, 0)$, and $W_p(B_3) \leq W_p(B_3^*) = 3k^2 + 2k + 1$;

If $r = 1$, then $B_3^* \cong C_{3,4}'''(k, k, k + 1; k, k, 0, 0), C_{3,4}'''(k - 1, k, k + 2; k - 1, k, 0, 0)$, and $W_p(B_3) \leq W_p(B_3^*) = 3k^2 + 4k + 2$;

If $r = 2$, then $B_3^* \cong C_{3,4}'''(k, k, k + 2; k, k, 0, 0)$, and $W_p(B_3) \leq W_p(B_3^*) = 3k^2 + 6k + 4$. \square

Theorem 0.4. Let B be a bicyclic graph of order $n (\geq 4)$, B^* be the bicyclic graph with the maximum polarity index among all bicyclic graphs.

- (1) If $n = 4$, then $B^* = B = C_{3,3}'''(0,0,0;0,0,0)$, and $W_p(B) = 0$;
- (2) If $n = 5$, then $B^* \cong C_{3,3}'''(0,0,1;0,0,0)$, and $W_p(B) \leq W_p(B^*) = 1$;

- (3) If $n = 6$, then $B^* \cong C'_{3,3}(0,0,0;0,0,0)$, and $W_p(B) \leq W_p(B^*) = 4$;
- (4) If $n = 7$, then $B_1^* \cong C'_{3,3}(1,0,0;0,0,0), C''_{3,3}(1,0,0;1,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertex of v_1 and $W_p(B_1) \leq W_p(B_1^*) = 6$;
- (5) If $n = 8$, then $B^* \cong C'_{3,3}(1,0,0;1,0,0), C''_{3,3}(1,0,0;0,0,0) + P_1$, where P_1 is incident to one pendant vertex of v_1 , $C''_{3,3}(1,0,0;1,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertex of v_1 , and $W_p(B) \leq W_p(B^*) = 9$;
- (6) If $n = 9$, then $B^* \cong C'_{3,3}(2,0,0;1,0,0), C'_{3,3}(1,0,0;1,0,0) + P_1$, where the path P_1 is incident to the pendant vertex of v_1 , $C'_{3,3}(2,0,0;0,0,0) + P_1$, where the path P_1 is incident to one pendant vertex of v_1 , $C'_{3,3}(1,0,0;0,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertex of v_1 , $C''_{3,3}(0,2,2;0,0,0), C''_{3,3}(1,0,0;1,0,0) + P_1 + P_1 + P_1 + P_1$, where the four paths P_1 are incident to the pendant vertex of v_1 , $C''_{3,3}(2,0,0;2,0,0) + P_1 + P_1 + P_1$, where the three paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 12$;
- (7) If $n = 10$, then $B^* \cong C'_{3,3}(2,0,0;2,0,0), C'_{3,3}(2,0,0;1,0,0) + P_1$, where the path P_1 is incident to one pendant vertex of v_1 , $C'_{3,3}(2,0,0;0,0,0) + P_1 + P_1$, where the two paths P_1 are incident to the pendant vertices of v_1 , $C''_{3,3}(0,2,3;0,0,0), C''_{3,3}(2,0,0;2,0,0) + P_1 + P_1 + P_1 + P_1$, where the four paths P_1 are incident to the pendant vertices of v_1 , and $W_p(B_1) \leq W_p(B_1^*) = 16$;
- (8) For $n \geq 11$, let $s_1 + s_2 + s_3 = 3k + r$ ($r \in \{0,1,2\}$).

If $r = 0$, then $B^* \cong C''_{3,3}(k - 2, k + 1, k + 1; k - 2, 0, 0)$, $C''_{3,3}(k - 1, k, k + 1; k - 1, 0, 0)$, and $W_p(B) \leq W_p(B^*) = 3k^2 + 4k + 1$;
 If $r = 1$, then $B^* \cong C''_{3,3}(k - 1, k + 1, k + 1; k - 1, 0, 0)$, and $W_p(B) \leq W_p(B^*) = 3k^2 + 6k + 3$;
 If $r = 2$, then $B^* \cong C''_{3,3}(k - 1, k + 1, k + 2; k - 1, 0, 0)$, and $W_p(B) \leq W_p(B^*) = 3k^2 + 8k + 5$. \square

Discussion

Quantifying the structure of complex networks is still intricate because the structural interpretation of quantitative network measures and their interrelations have not yet been explored extensively. In this paper, we studied sharp upper bounds for the Wiener polarity index among all bicyclic networks, by using some transformations. The graphs attaining these bounds are also characterized. The proof techniques use structural properties of the graphs under consideration and it may be intricate to extend the techniques when using more general graphs.

An interesting thing is that the Wiener polarity index is related to a pure mathematical problem: counting the number of subgraphs of a graph. This counting problem is a basic problem in mathematics but much more complicated. For example, Alon and Bollobás provide some results on this topic, e.g.^{39–41}.

As a future work, we will consider the extremal problems of the Wiener polarity index for general networks and also some special networks. Furthermore, we would like to explore advanced structural properties of the Wiener polarity index, and relations between the Wiener polarity index and some other topological indices. On the other hand, it would be interesting to investigate the applications of Wiener polarity index in characterizing the structure properties of complex networks and studying algorithm theory and computational complexity. For instance, one can consider the possibility of using the Wiener polarity index or other distance measures to study other very interesting algorithms, like the google algorithm in complex networks^{42,43}.

Methods

First we introduce some operations on bicyclic graphs, then we give the corresponding lemmas which state that the Wiener polarity index is not decreasing after applying these operations on bicyclic graphs.

Let B be a bicyclic graph. As we have claimed, suppose $C_p = v_1v_2 \dots v_p v_1$ and $C_q = u_1u_2 \dots u_q u_1$ are two cycles. If both $T_B[v_i] (1 \leq i \leq p)$ and $T_B[u_j] (1 \leq j \leq q)$ are stars, then we denote such a bicyclic graph by $C_{p,q}(s_1, \dots, s_p; t_1, \dots, t_q)$, where s_i and t_j represent the number of pendant vertices of v_i and u_j , respectively.

We define **Operation I** as follows. Let $T_B[v]$ denote a hanging tree on vertex v of a bicyclic graph B with $p \geq 4$, $q \geq 4$, where v is on the cycle of B . Among all hanging trees, suppose $vc_1 \dots c_{r-1}c_r$ is one of the longest paths from the root v to a leaf c_r in $T_B[v]$. If $r \geq 2$, then after deleting the edge vc_1 from B , we obtain a bicyclic graph A and a tree T such that $v \in A$ and $c_1 \in T$. Let B' denote the bicyclic graph obtained from A and T by identifying c_1 and v' (a neighbor of v on the cycle of B) and adding a new hanging leaf vx to v .

We define **Operation II** as follows. Let B be a bicyclic graph with $p = 3$, $T_B[v_i]$ be a hanging tree rooted at v_i ($i = 1, 2, 3$). Let $v_i c_1 \dots c_{r-1}c_r$ be one of the longest paths from the root v_i to a leaf c_r of the hanging tree $T_B[v_i]$.

For $r \geq 3$, we define a new graph B^* as follows:

$$B^* = \begin{cases} B - c_{r-1}c_r + c_{r-3}c_r, & \text{if } r > 3, \\ B - c_{r-1}c_r + v_i c_r, & \text{if } r = 3. \end{cases} \tag{2}$$

For $r = 2$, the operation differs on the three types of bicyclic graphs.

- (1) For bicyclic graphs in type I, we let

$$B^* = \begin{cases} B - c_1c_2 + c_2w_1, & \text{if } v_i = v_1, \\ B - c_1c_2 + c_2v_1, & \text{if } v_i = v_2 \text{ or } v_3, \end{cases} \tag{3}$$

where $w_1 \in N_B(v_i)$ is on the path $v_1 \dots u_1$.

(2) For bicyclic graphs in type II, by considering the value of q , there are two cases.

Case 1. $q \geq 4$. In this case, let

$$B^* = \begin{cases} B - c_1c_2 + c_2v_1, & \text{if } v_i \neq v_1, \\ B - c_1c_2 + c_2u_2, & \text{if } v_i = v_1, \end{cases} \tag{4}$$

where $v_i (i \in \{1,2,3\})$ is the root vertex mentioned above.

Case 2. $q = 3$ and $|V(B)| \geq 9$. Here we let $C_q = v_1v_4v_5v_1$. We define an operation as follows: delete $T_B[v_i] \setminus v_i$ and add a copy of $T_B[v_i]$ to v_j by identifying v_j and v_i' which is a copy of v_i . We call this operation “move $T_B[v_i]$ to v_j ”. By considering the number of vertices on the cycles of B with hanging trees, there are two subcases.

Subcase 2.1. There is only one vertex $v_i (i \in \{1,2,3,4,5\})$ with a hanging tree. Let $N_B(v_i) = \{c_1^1, \dots, c_1^a\}$, $N_B^2(v_i) = \{c_2^1, \dots, c_2^b\}$.

For the case $v_i = v_1$, we apply operations as follows. If $b \geq 4$, then move c_2^1, c_2^1 to v_2 and $c_2^j (3 \leq j \leq b)$ to v_3 ; if $b = 3$, then move c_2^1, c_2^2 to v_2 and c_3^1, c_1^1 to v_3 ; if $b = 2$, then move c_2^1, c_2^2 to v_2 and c_1^1 to v_3 ; if $b = 1$, then move c_2^1 to v_2 and c_1^1 to v_3 . The new graph is denoted by B' .

For the case $v_i = v_2$, we construct a new graph $B^* = B - c_1c_2 + c_2v_3$.

Subcase 2.2. There are at least two vertices $v_s, v_t (s, t \in \{1,2,3,4,5\})$ with hanging trees. In this subcase, let $B^* = B - c_1c_2 + c_2v_k$, where $v_k \in N_B(v_s) \cap N_B(v_t)$.

(3) For the bicyclic graphs in type III. By considering the value of q , there are two cases.

Case 1. $q \geq 4$. In this case, we can apply Operation 1 on C_q .

Case 2. $q = 3$ and $|V(B)| \geq 12$. Here let $C_q = v_1v_2v_4v_1$. We can move $T_B[v_4]$ to v_3 to get a new graph B' satisfying $W_p(B') = W_p(B)$. By considering the number of vertices on the cycles of B' with hanging trees, there are two subcases.

Subcase 2.1. There exists only one vertex, say $v_i (i \in \{1,2,3\})$, which has a hanging tree. Firstly, move $T_B[v_i]$ to v_3 (denote the new graph by B''), delete a vertex in $N_{B''}^2(v_3)$ and meanwhile subdivide edge v_1v_4 (denote the new graph by B'''); secondly, move all the other vertices in $N_{B''}^2(v_3)$ to v_1 (denote the new graph by B''''); thirdly, if $d_{B''''}(v_1) \geq 5$, then just move one pendant vertex of v_1 to v_2 ; if $d_{B''''}(v_1) = 4$, then move one pendant vertex of v_3 to v_2 .

Subcase 2.2. There exist two vertices, say $v_i, v_j (i, j \in \{1,2,3\})$, which have hanging trees. If $i = 1$ and $j = 2$, then move $T_B[v_2]$ to v_3 . Now we can only consider the case $i = 1$ and $j = 3$.

If there exists $c_2 \in N_{B'}^2(v_3)$, then delete c_2 and subdivide the edge v_1v_4 (denote the new graph by B''). Now return to the situation in Case 1.

If and $d_{B''}(v_3) \geq 4$, then delete a vertex $c_1 \in N_{B''}^2(v_1)$ and subdivide the edge v_1v_4 . Now return to Case 1. For the situation that $d_{B''}(v_3) = 3$, delete a vertex $c_2 \in N_{B''}^2(v_1)$ and subdivide the edge v_1v_4 , move all pendant vertices in $N_{B''}^2(v_1)$ to v_2 , at last move one pendant vertex of v_1 or v_2 to v_3 .

Subcase 2.3. There exist three vertices which have hanging trees. By deleting some pendant vertex in $N_{B'}^2(v_i)$, where $i \in \{1,2,3\}$, and meanwhile subdividing the edge v_1v_4 , we return to the situation in Case 1.

The final graph obtained after the above operation is denoted by B' .

We define **Operation III** as follows. Let B be a bicyclic graph. If $d_B(v) = 2$, then let $B^* = B - vv' - vv'' + v'v'' + vx$, where $v', v'' \in N_B(v), x \in V(B)$. We call such an operation *smooth* v to x .

We define **Operation IV** as follows. Let B be a bicyclic graph, where $T_B[v_i] (1 \leq i \leq p)$ and $T_B[u_j] (1 \leq j \leq q)$ are both stars. Denote the set of the pendant vertices of $v_i(u_j)$ by $V_i(U_j)$.

For bicyclic graphs in type I, we will take the following two steps.

Step 1. For C_p and $i \in \{3, \dots, p - 1\}$, if i is odd, then move V_i to v_1 and smooth v_i to v_2 ; if i is even, then move V_i to v_2 and smooth v_i to v_1 . For C_q and $j \in \{3, \dots, q - 1\}$, if j is odd, then move U_j to u_1 and smooth u_j to u_2 ; if j is even, then move U_j to u_2 and smooth u_j to u_1 . Therefore, we obtain a graph $B' = C_{3,3}(s_1, s_2, s_3; t_1, t_2, t_3)$ with a unique path P connecting C_p and C_q . Let the set of hanging leaves of u_1, u_2, u_q be U'_1, U'_2, U'_q , respectively.

Step 2. Let $P = v_1w_1 \dots w_t u_1, W_k = T_{B'}[w_k] (1 \leq k \leq t)$.

If k is odd, then move W_k to v_3 and smooth w_k to v_2 ; if k is even, then move W_k to v_2 and smooth w_k to v_3 .

If t is odd, then move U'_1 to v_2, U'_2 to v_1, U'_q to v_3 ; if t is even and $t \geq 2$, then move U'_1 to v_3, U'_2 to v_1 , and U'_q to v_2 , respectively; if $t = 0, |V(B')| \geq 9$ and $d(v_2) = d(v_3) = 2$, let $N_{B'}(v_1) = \{a_1, \dots, a_s\}$ and $N_{B'}(u_1) = \{b_1, \dots, b_t\}$, then for the situation that $b = 1$, move b_1 to v_2 and move a_1 to v_3 , for the situation that $b \geq 2$, move b_1 to v_2 and move b_2, \dots, b_t to v_3 ; if $t = 0$ and $d(v_i) = 2, d(v_j) > 2 (i, j \in \{1,2\})$, then move U'_1 to v_i, U'_2 to v_1, U'_q to v_j , respectively.

Finally, we get a new graph $B'' = C_{3,3}(s'_1, s'_2, s'_3; 0,0,0)$ and there is a unique path $P = v_1u_1$ connecting C_p and C_q .

For bicyclic graphs in type II, we also give two steps as follows.

Step 1. For C_p and $i \in \{3, \dots, p - 1\}$, if i is odd, then move V_i to v_1 and smooth v_i to v_2 ; if i is even, then move V_i to v_2 and smooth v_i to v_1 . For C_q and $j \in \{3, \dots, q - 1\}$, if j is odd, then move U_j to u_1 and smooth u_j to u_2 ; if j is even, then move U_j to u_2 and smooth u_j to u_1 . Thus we get a graph $B' = C_{3,3}(s_1, s_2, s_3; t_1, t_2, t_3)$ with $s_1 = t_1$. Let the set of hanging leaves of u_1, u_2, u_q be U'_1, U'_2, U'_q , respectively.

Step 2. By moving U'_2 to v_2, U'_q to v_p , we have $B'' = C_{3,3}(s_1, s'_2, s'_3; t_1, 0, 0)$ with $s_1 = t_1$.

For bicyclic graphs in type III, the operation is defined as follows. Recall that we use $l (\geq 1)$ to denote the number of common vertices of C_p and C_q , and without loss of generality, assume $l - 2 \leq p - 2 \leq q - 2$.

(1) If $p \geq 3$ and $q \geq 4$, then we will take the following three steps.

Step 1. For $i \in \{3, \dots, p-1\}$, $j \in \{3, \dots, q-1\}$. If i is odd, then move V_i to v_1 ; if i is even, then move V_i to v_2 ; if j is odd, then move U_j to v_1 ; if j is even, then move U_j to v_2 ; move U_q to v_p .

Step 2. If $l = 2$ or 3 , smooth vertices v_{l+1}, \dots, v_{p-1} to v_1 and v_2 alternately.

If $l \geq 4$, then we first smooth vertices v_3, \dots, v_{l-1} to v_1 and v_2 alternately; then smooth vertices v_{l+1}, \dots, v_{p-1} to v_1 and v_2 alternately.

After applying this operation, we get a new graph B' with cycles $C_p, C_{q'}$ and C_z . Let l' be the number of common vertices of C_p and $C_{q'}$, l' ($l' = 3$ or 4) be the number of vertices of the smallest cycle of B' , then we have $l' = 2$ or $l' = 3$. Now relabel the vertices on C_p and $C_{q'}$ of B' , and we have $C_{p'} = v_1 \dots v_{p'} v_1$ and $C_{q'} = u_1 \dots u_{q'} u_1$.

Step 3. Considering the value of l' , there are two cases.

Case 1. $l' = 2$.

We just smooth $u_{l'+2} \dots u_{q'-2}$ to v_1 and v_2 alternately, and smooth $u_{q'-1}$ to v_p . The new graph obtained is denoted by $B^* = C_{3,4}(s_1, s_2, s_3; s_1, s_2, 0, 0)$.

Case 2. $l' = 3$, $|V(B')| \geq 6$ and $B' = C_{4,q}(s_1, s_2, 0, s_4; t_1, t_2, 0, \dots, 0)$ with $s_1 = t_1$.

Let $B'' = B' - v_3 v_4 + v_2 v_4$. If $q' \geq 5$, then smooth v_3 to vertex v_1 , smooth $u_5, \dots, u_{q'-2}$ to v_1 and v_2 alternately and smooth $u_{q'-1}$ to v_4 ; if $q' = 4$ and $d_{B''}(v_4) \geq 3$, then we do nothing; if $q' = 4$ and $d_{B''}(v_4) = 2$, then move the pendant vertices of v_2 to v_4 .

Finally, we get the desired graph $B^* = C_{3,4}(s'_1, s'_2, s'_3; s'_1, s'_2, 0, 0)$.

(2) If $p = 3$ and $q = 3$, by Operation II on B and its resultant graphs repeatedly, we have a new graph $B' = C_{3,3}(s'_1, s'_2, s'_3; s'_1, s'_2, t'_3)$. Move the pendant vertices of u_3 to v_3 , we obtain $B'' = C_{3,3}(s''_1, s''_2, s''_3; s''_1, s''_2, 0)$.

References

- Sydney, A., Scoglio, C., Schumm, P. & Kooij, R. Elasticity: topological characterization of robustness in complex networks. *IEEE/ACM Bionetics* (2008).
- Boccaletti, S., Latora, V., Moreno, Y., Chavez, M. & Hwang, D. Complex networks: structure and dynamics. *Physics Reports* **424**, 175–308 (2006).
- da F. Costa, L., Rodrigues, F. & Travieso, G. Characterization of complex networks: A survey of measurements. *Adv. Phys.* **56**, 167–242 (2007).
- Dorogovtsev, S. & Mendes, J. Evolution of networks. *Adv. Phys.* **51**, 1079–1187 (2002).
- Ellens, W. & Kooij, R. Graph measures and network robustness. *arXiv:1311.5064v1 [cs. DM]* (2013).
- Kraus, V., Dehmer, M. & Emmert-Streib, F. Probabilistic inequalities for evaluating structural network measures. *Inform. Sciences* **288**, 220–245 (2014).
- Wiener, H. Structural determination of paraffin boiling points. *J. Amer. Chem. Soc.* **69**, 17–20 (1947).
- Azari, M. & Iranmanesh, A. Harary index of some nano-structures. *MATCH Commun. Math. Comput. Chem.* **71**, 373–382 (2014).
- Feng, L. & Yu, G. The hyper-wiener index of cacti. *Utilitas Math.* **93**, 57–64 (2014).
- Feng, L., Liu, W., Yu, G. & Li, S. The hyper-wiener index of graphs with given bipartition. *Utilitas Math.* **95**, 23–32 (2014).
- Eliasi, M. & Taeri, B. Extension of the wiener index and wiener polynomial. *Appl. Math. Lett.* **21**, 916–921 (2008).
- Balaban, A. Topological indices based on topological distance in molecular graphs. *Pure Appl. Chem.* **55**, 199–206 (1983).
- Cao, S., Dehmer, M. & Shi, Y. Extremality of degree-based graph entropies. *Inform. Sciences* **278**, 22–33 (2014).
- Chen, Z., Dehmer, M. & Shi, Y. A note on distance-based graph entropies. *Entropy* **10**, 5416–5427 (2014).
- Dehmer, M., Emmert-Streib, F. & Grabner, M. A computational approach to construct a multivariate complete graph invariant. *Inform. Sciences* **260**, 200–208 (2014).
- Dehmer, M. & Ilić, A. Location of zeros of wiener and distance polynomials. *PLoS One* **7**(3), e28328 (2012).
- Tian, D. & Choi, K. Sharp bounds and normalization of wiener-type indices. *PLoS One* **8**(11), e78448 (2013).
- Dobrynin, A., Entringer, R. & Gutman, I. Wiener index of trees: theory and applications. *Acta Appl. Math.* **66**, 211–249 (2001).
- da Fonseca, C., Ghebleh, M., Kanso, A. & Stevanovic, D. Counterexamples to a conjecture on wiener index of common neighborhood graphs. *MATCH Commun. Math. Comput. Chem.* **72**, 333–338 (2014).
- Knor, M., Luzar, B., Skrekovski, R. & Gutman, I. On wiener index of common neighborhood graphs. *MATCH Commun. Math. Comput. Chem.* **72**, 321–332 (2014).
- Lin, H. Extremal wiener index of trees with given number of vertices of even degree. *MATCH Commun. Math. Comput. Chem.* **72**, 311–320 (2014).
- Skrekovski, R. & Gutman, I. Vertex version of the wiener theorem. *MATCH Commun. Math. Comput. Chem.* **72**, 295–300 (2014).
- Soltani, A., Iranmanesh, A. & Majid, Z. A. The multiplicative version of the edge wiener index. *MATCH Commun. Math. Comput. Chem.* **71**, 407–416 (2014).
- Lukovits, I. & Linert, W. Polarity-numbers of cycle-containing structures. *J. Chem. Inform. Comput. Sci.* **38**, 715–719 (1998).
- Hosoya, H. & Gao, Y. Mathematical and chemical analysis of wiener's polarity number. In Rouvray, D. & King, R. (eds.) *Topology in Chemistry—Discrete Mathematics of Molecules*, 38–57 (Elsevier, 2002). Horwood, Chichester.
- Du, W., Li, X. & Shi, Y. Algorithms and extremal problem on wiener polarity index. *MATCH Commun. Math. Comput. Chem.* **62**, 235–244 (2009).
- Deng, H. On the extremal wiener polarity index of chemical trees. *MATCH Commun. Math. Comput. Chem.* **66**, 305–314 (2011).
- Deng, H. & Xiao, H. The maximum wiener polarity index of trees with k pendants. *Appl. Math. Lett.* **23**, 710–715 (2010).
- Deng, H., Xiao, H. & Tang, F. On the extremal wiener polarity index of trees with a given diameter. *MATCH Commun. Math. Comput. Chem.* **63**, 257–264 (2010).
- Liu, B., Hou, H. & Huang, Y. On the wiener polarity index of trees with maximum degree or given number of leaves. *Comput. Math. Appl.* **60**, 2053–2057 (2010).
- Liu, M. & Liu, B. On the wiener polarity index. *MATCH Commun. Math. Comput. Chem.* **66**, 293–304 (2011).
- Ma, J., Shi, Y. & Yue, J. On the extremal wiener polarity index of unicyclic graphs with a given diameter. In Gutman, I. (ed.) *Topics in Chemical Graph Theory*, vol. Mathematical Chemistry Monographs No.16a, 177–192 (University of Kragujevac and Faculty of Science Kragujevac, 2014). Horwood, Chichester.
- Ma, J., Shi, Y. & Yue, J. The wiener polarity index of graph products. *Ars Combin.* **116**, 235–244 (2014).
- Hou, H., Liu, B. & Huang, Y. On the wiener polarity index of unicyclic graphs. *Appl. Math. Comput.* **218**, 10149–10157 (2012).
- Behmarama, A., Yousefi-Azari, H. & Ashrafi, A. Wiener polarity index of fullerenes and hexagonal systems. *Appl. Math. Lett.* **25**, 1510–1513 (2012).

36. Deng, H. & Xiao, H. The wiener polarity index of molecular graphs of alkanes with a given number of methyl groups. *J. Serb. Chem. Soc.* **75**, 1405–1412 (2010).
37. Hua, H., Faghani, M. & Ashrafi, A. The wiener and wiener polarity indices of a class of fullerenes with exactly $12n$ carbon atoms. *MATCH Commun. Math. Comput. Chem.* **71**, 361–372 (2014).
38. Bondy, J. A. & Murty, U. S. R. (eds.) *Graph Theory* (Springer–Verlag, 2008). Berlin.
39. Alon, N. On the number of certain subgraphs contained in graphs with a given number of edges. *Israel J. Math.* **53**, 97–120 (1986).
40. Bollobás, B. & Sarkar, A. Paths in graphs. *Studia Sci. Math. Hungar.* **38**, 115–137 (2001).
41. Bollobás, B. & Tyomkyn, M. Walks and paths in trees. *J. Graph Theory* **70**, 54–66 (2012).
42. Paparo, G. D. & Martin-Delgado, M. A. Google in a quantum network. *Sci. Rep.* **2**, 444 (2012).
43. Paparo, G. D., Müller, M., Comellas, F. & Martin-Delgado, M. A. Quantum google in a complex network. *Sci. Rep.* **3**, 2773 (2013).

Acknowledgements

We wish to thank the referee for valuable comments and helpful suggestions. This work was supported by National Natural Science Foundation of China and PCSIRT, China Postdoctoral Science Foundation.

Author Contributions

All authors designed the research. J.M., Y.S., Z.W. and J.Y. contributed equally to conducting the research and doing simulations. J.M., Y.S., Z.W. and J.Y. contributed to the main idea. All authors wrote the paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Ma, J. *et al.* On Wiener polarity index of bicyclic networks. *Sci. Rep.* **6**, 19066; doi: 10.1038/srep19066 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>