

# SCIENTIFIC REPORTS



OPEN

## Effect of simulated tillage on microbial autotrophic CO<sub>2</sub> fixation in paddy and upland soils

Received: 13 December 2014

Accepted: 18 December 2015

Published: 22 January 2016

Tida Ge<sup>1,2,\*</sup>, Xiaohong Wu<sup>1,3,\*</sup>, Qiong Liu<sup>1,4</sup>, Zhenke Zhu<sup>1,4</sup>, Hongzhao Yuan<sup>1,4</sup>, Wei Wang<sup>1</sup>, A. S. Whiteley<sup>4,5</sup> & Jinshui Wu<sup>1,4</sup>

Tillage is a common agricultural practice affecting soil structure and biogeochemistry. To evaluate how tillage affects soil microbial CO<sub>2</sub> fixation, we incubated and continuously labelled samples from two paddy soils and two upland soils subjected to simulated conventional tillage (CT) and no-tillage (NT) treatments. Results showed that CO<sub>2</sub> fixation (<sup>14</sup>C-SOC) in CT soils was significantly higher than in NT soils. We also observed a significant, soil type- and depth-dependent effect of tillage on the incorporation rates of labelled C to the labile carbon pool. Concentrations of labelled C in the carbon pool significantly decreased with soil depth, irrespective of tillage. Additionally, quantitative PCR assays revealed that for most soils, total bacteria and *cbbL*-carrying bacteria were less abundant in CT versus NT treatments, and tended to decrease in abundance with increasing depth. However, specific CO<sub>2</sub> fixation activity was significantly higher in CT than in NT soils, suggesting that the abundance of *cbbL*-containing bacteria may not always reflect their functional activity. This study highlights the positive effect of tillage on soil microbial CO<sub>2</sub> fixation, and the results can be readily applied to the development of sustainable agricultural management.

Autotrophic bacteria in terrestrial ecosystems can partially compensate for increasing atmospheric CO<sub>2</sub> concentration, predicted to reach 450–600 ppm by 2050<sup>1</sup>. These bacteria have the capacity to fix CO<sub>2</sub> and are widely distributed in agricultural soils<sup>2–4</sup>. Among the six pathways developed by microbial autotrophs for CO<sub>2</sub> fixation<sup>5</sup>, autotrophic bacteria predominantly use the Calvin–Benson–Bassham cycle. This pathway depends on the activity of ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO), encoded by the *cbbL* gene<sup>6</sup>. Recently, environmental studies based on *cbbL* gene detection have shown that autotrophic bacteria are sensitive to agricultural management practices such as fertiliser treatment, land use alteration, and cropping systems<sup>2,7–9</sup>. Changes in soil physical, chemical, and biological properties caused by different management practices are reported to affect the abundance, diversity, and activity of CO<sub>2</sub>-fixing autotrophic bacteria and their associated fixation rates<sup>2,4,7–9</sup>.

Conventional tillage (CT), including ploughing and disking, has been the dominant agricultural practice for the past century of crop production<sup>10,11</sup>. Such management practices are excellent for loosening soil, which improves surface soil compaction, thus repressing annual weeds and benefiting precise seeding<sup>12,13</sup>. However, the intensive mechanical disturbance of soil structures introduced by CT practices is accompanied by surface soil erosion, a reduction in soil aggregate stability, and the acceleration of soil organic matter decomposition<sup>14,15</sup>. On-going changes in CT soil properties (e.g. porosity, bulk density, and organic carbon concentration) will affect water, gas, and nutrient diffusion, potentially triggering changes in soil bacterial communities<sup>16,17</sup>. Many studies have shown that CT practices negatively affect soil bacterial populations, often resulting in a decrease in community abundance, diversity, and activity when compared to no-tillage (NT) management<sup>18–21</sup>. However, this phenomenon is not observed in all reports, indicating that the underlying mechanisms driving the changes within CT soil bacterial communities are likely linked to a wide range of factors, including soil texture and depth

<sup>1</sup>Changsha Research Station for Agricultural and Environmental Monitoring & Key Laboratory of Agro-ecological Processes in Subtropical Region, Institute of Subtropical Agriculture, Chinese Academy of Sciences, Hunan, 410125, China. <sup>2</sup>State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese Academy of Sciences, Jiangsu, 210008, China. <sup>3</sup>Faculty of Life Science and Technology, Central South University of Forestry and Technology, Changsha, 410004, China. <sup>4</sup>ISA-CAS and UWA Joint Laboratory for Soil Systems Biology, Hunan, 410125, China. <sup>5</sup>School of Earth and Environment, The University of Western Australia, Crawley, WA 6009, Australia. \*These authors contributed equally to this work. Correspondence and requests for materials should be addressed to T.G. (email: gtd@isa.ac.cn) or J.W. (email: jswu@isa.ac.cn)

of tillage<sup>20,21</sup>. Despite intensive studies concerning the impact of different tillage practices on soil bacterial communities, we still have limited knowledge about the ecological functions of specific microbial communities under various tillage managements. Our current understanding is that soil autotrophic bacteria do play a central role—modulated by tillage practices—in mitigating atmospheric CO<sub>2</sub> emission<sup>4,22,23</sup>, but little data exist to clarify that role. Now, however, advances in molecular microbial ecology allow us to investigate the function of soil autotrophic bacteria by quantifying *cbbL* gene abundances<sup>24</sup> and their associated CO<sub>2</sub> fixation rates under different tillage managements. This method fills a major knowledge gap in clarifying the effects of tillage upon important global C sequestration processes.

The objective of this study was to evaluate the effect of different tillage practices on soil autotrophic bacterial populations and their CO<sub>2</sub> assimilation rates at varying soil depths. Soils were sieved to experimentally generate conventional tillage treatment soils (CT), and intact soil cores without sieving were collected as the corresponding no-till treatment soils (NT). Using continuous labelling with <sup>14</sup>CO<sub>2</sub>, we quantified the carbon fixed by soil autotrophs (<sup>14</sup>C-SOC), the distribution of newly assimilated carbon in the soil microbial biomass carbon pool (<sup>14</sup>C-MBC), and the dissolved organic carbon pool (<sup>14</sup>C-DOC) at different depths (0–1, 1–5, and 5–17 cm) of both CT and NT soils. Real-time quantitative PCR analysis was also conducted to assess how the abundance of autotrophic bacteria changed in response to tillage. We hypothesised that the mechanical disturbance from CT practices would decrease soil autotrophic bacterial abundance, leading to lower rates of CO<sub>2</sub> fixation when compared with NT soils.

## Results

**Soil autotrophic bacteria CO<sub>2</sub> fixation rate.** The CT treatment significantly increased the <sup>14</sup>C-SOC over the 110-day incubation period. The amount of <sup>14</sup>C-SOC was, on average, 87% higher in CT soils when compared with NT soils at depths of 0–1 cm, and 210% higher at 1–5 cm (Fig. 1; Table 1). At 5–17 cm, the <sup>14</sup>C-SOC concentration was 141% greater in P1 (paddy) soils under CT relative to the NT treatment, and no <sup>14</sup>C-SOC content was detected under NT treatments of three other soils (Fig. 1). Generally, different types of soils responded differently to tillage treatments: CT treatment had a greater impact on upland soils than on paddy soils (Fig. 1; Table 1). Under both CT and NT treatments, the overall <sup>14</sup>C-SOC concentrations decreased with increasing soil depth, with deeper soil layers being more sensitive to tillage practices (Fig. 1; Table 1). ANOVA analyses revealed no significant interactive effect of soil type, soil depth, and soil tillage on the measured <sup>14</sup>C-SOC content (Table 1).


**The incorporation of <sup>14</sup>C into MBC and DOC.** The incorporation rates of autotrophically fixed <sup>14</sup>C into microbial biomass carbon (MBC) and dissolved organic carbon (DOC) were modulated by soil tillage (Fig. 2, 3; Table 1). Larger amounts of <sup>14</sup>C-MBC were recovered from CT soils than from NT soils, but the difference was not significant in P2 (0–1 cm) and U1 (upland soil; 0–1 cm) (Fig. 2; Table 1). No significant interaction between soil type, soil depth, and soil tillage was observed to affect <sup>14</sup>C-MBC concentration. Compared with NT treatments, CT also significantly increased <sup>14</sup>C-DOC concentration in the 0–1 cm depth by an average of 33%, whereas the effect of tillage on <sup>14</sup>C-DOC contents at greater depths was highly dependent on soil type (Fig. 3; Table 1). Under both CT and NT treatments, larger amounts of <sup>14</sup>C-MBC and <sup>14</sup>C-DOC were observed in paddy soils when compared with upland soils at 0–1 cm and 1–5 cm (Fig. 2), with a significant soil type × soil tillage interaction (Table 1). Both <sup>14</sup>C-MBC and <sup>14</sup>C-DOC contents decreased with increasing soil depth in CT and NT soils, although a significant soil depth × soil tillage interaction was only observed with <sup>14</sup>C-DOC (Figs 2 and 3, Table 1). Additionally, a significant interactive effect on <sup>14</sup>C-DOC concentration was observed among soil type, soil depth, and soil tillage.

**Bacterial and *cbbL* gene abundance.** The bacterial (*16S rRNA*) and *cbbL* gene abundance differed across tillage managements, as well as across soil type and depth (Table 2). Lower bacterial and *cbbL* gene abundances were observed in CT treatments when compared with NT, but only in P1, P2, and U2 soils. In contrast, higher *cbbL* gene abundance was observed in U1 soil under CT and NT treatments (Table 2). Soil type exhibited a significant effect on bacterial and *cbbL* gene abundance, with paddy soils generally being more susceptible to tillage than upland soils (Tables 1 and 2). In both CT and NT soils, *cbbL*-carrying bacteria decreased with increasing soil depths in paddy soils, but increased with depth in upland soils, with no significant interaction being observed among soil type, soil depth, and soil tillage (Table 1).

**The specific CO<sub>2</sub> fixation activity of autotrophic bacteria.** The CT treatments enhanced the specific CO<sub>2</sub> fixation activity compared to NT treatments, irrespective of soil type and depth, with the increase being significant in all soils except U1 (Tables 1 and 2). Significant soil type × soil tillage and soil depth × soil tillage interactions were observed, suggesting that soil tillage affects CO<sub>2</sub> fixation activity differently depending on the specific soil type or depth (Tables 1 and 2).

## Discussion

We were able to clearly detect <sup>14</sup>C in soils of both the CT and NT treatments, indicating that CO<sub>2</sub> fixation had taken place (Fig. 1). Mechanistically, both biotic and abiotic processes could be responsible for the recovery of <sup>14</sup>C labelled pools in soils. For example, Miltner *et al.*<sup>25</sup> documented that after 81 days of incubation, 0.83 μmol·g<sup>-1</sup> soil of <sup>14</sup>CO<sub>2</sub> was fixed to biologically active soil, with 96% bound as organic compounds. However, approximately 0.02 μmol·g<sup>-1</sup> soil of <sup>14</sup>CO<sub>2</sub> was detected in the fumigated control soil, which did not have microbial activity, and 91% of this fixed <sup>14</sup>C was bound as carbonates. Similar results were observed in studies with longer incubation time or different tracers (e.g. <sup>13</sup>CO<sub>2</sub>)<sup>26,27</sup>. In this study, we were able to remove all <sup>14</sup>C bound as carbonates via the concentrated H<sub>2</sub>SO<sub>4</sub>–H<sub>3</sub>PO<sub>4</sub> treatment, allowing us to conclude that observed CO<sub>2</sub> fixation was mainly from biological processes. Within these processes, we discounted heterotrophic fixation as a primary route because



**Figure 1.** The  $^{14}\text{C}$ -SOC concentrations recovered at different depths (0–1 cm, 1–5 cm, and 5–17 cm) in conventional tillage (CT) and no-till (NT) soils after 110 days of incubation. Error bars indicate the standard error of the mean ( $n = 4$ ). \* indicates significant differences between CT and NT soils at  $P < 0.05$ ; nd, not detectable.

our previous study did not detect fixed  $^{14}\text{C}$  in dark-incubated soils<sup>4</sup>. Therefore, we concluded that microbial autotrophs are the primary source of carbon fixation.

Crucially, significantly higher amounts of  $^{14}\text{C}$  fixation occurred in the CT treatments compared with the NT treatments (Fig. 1). However, tillage effects on the abundances of total bacteria and *cbbL*-carrying bacteria were variable among soils. Both decreases and increases of 16S rRNA and *cbbL* gene numbers were observed under CT treatment. Negative effects of tillage on soil microbial abundance were found in P1, P2, and U2 soils, irrespective of soil depth. These data support previous studies showing that tillage negatively affects the abundances of diverse functional groups such as denitrifiers and nitrifying microorganisms<sup>28–30</sup>. For example, tillage reduced denitrifier populations in loam soils with wheat/fallow rotation<sup>28</sup>, and after 22 years of conventional tillage, the abundance of ammonia-oxidising bacteria was found to decrease in a subtropical rice-based ecosystem<sup>29</sup>. These studies have

Factors	<sup>14</sup> C-SOC (mg·kg <sup>-1</sup> dry soil)		<sup>14</sup> C-MBC (mg·kg <sup>-1</sup> dry soil)		<sup>14</sup> C-DOC (mg·kg <sup>-1</sup> dry soil)		<i>cbbL</i> gene copies (10 <sup>9</sup> copies g <sup>-1</sup> dry soil)		Specific CO <sub>2</sub> fixation activity (10 <sup>-7</sup> g per copy)	
	F	P	F	P	F	P	F	P	F	P
Soil type	421.27	<0.001	112.54	<0.001	1385.00	<0.001	64.09	<0.001	19.25	<0.001
Soil depth	450.65	<0.001	113.57	<0.001	359.68	<0.001	4.43	0.017	6.59	0.013
Soil tillage	47.6	<0.001	8.93	0.005	103.04	<0.001	67.65	<0.001	18.26	<0.001
Soil type × Soil depth	284.3	<0.001	67.72	<0.001	229.25	<0.001	6.23	0.004	22.48	<0.001
Soil type × Soil tillage	11.12	0.002	4.79	0.034	21.97	<0.001	54.98	<0.001	8.11	0.001
Soil depth × Soil tillage	5.06	0.029	0.11	0.742	82.30	<0.001	4.58	0.038	5.33	0.007
Soil type × Soil depth × Soil tillage	0.07	0.795	0	0.980	22.14	<0.001	4.00	0.052	0.02	0.883

**Table 1. Results of ANOVA investigating the effects of soil type, depth, tillage, and their interactions on <sup>14</sup>C-SOC concentration, <sup>14</sup>C-MBC concentration, <sup>14</sup>C-DOC concentration, *cbbL* gene copies, and specific CO<sub>2</sub> fixation activity of autotrophic bacteria.** DOC, dissolved organic carbon; MBC, microbial biomass carbon; SOC, soil organic carbon. *F* and *P* values are results of the ANOVA, with the following factors: soil type (paddy soil, upland soil), soil depth (0–1 cm, 1–5 cm, and 5–17 cm), and soil tillage (CT soil, NT soil).


suggested several underlying mechanisms to explain tillage effects on microbial numbers, including the disruption of soil aggregates by CT that alters nutrient availability and intensifies carbon source preemption<sup>28–30</sup>. The suppression of total bacterial and *cbbL*-carrying bacterial abundance in the P1, P2, and U2 soils of this study may have been caused by similar mechanisms.

Interestingly, we found that CT exerted the opposite effect on bacterial abundance in U1 soil (Table 2), which is a vegetable soil subject to frequent tillage, in contrast to the other three soils. Tillage frequency influences the threshold rates of aggregate turnover; when the thresholds are passed, organic carbon is actually retained in, rather than released from, tilled soil aggregates<sup>31</sup>. Thus, we propose that the CT treatment of U1 soil probably surpassed the threshold aggregate turnover rate, resulting in the accumulation of soil organic carbon. Additionally, we found that total P was almost twice as high in U1 soil than in the other three soils (Table 2). Phosphorus is essential to microbial growth and its availability is highly related to tillage practices<sup>32</sup>. Therefore, CT treatment of U1 soil may create favourable growth conditions for soil microbial communities by improving P and organic substrate availability<sup>33</sup>, resulting in a higher abundance of *cbbL*-containing bacteria.

The present results showed that the amount of key players in CO<sub>2</sub> fixation (*cbbL*-carrying bacteria) was significantly lower for CT than NT in most soils, but microbially assimilated <sup>14</sup>C was markedly higher under CT conditions than under NT, indicating that the population size of *cbbL*-containing bacteria may not necessarily reflect their functional activities under changing soil conditions. This disparity in abundance and activity may be due to the enhancement of carbon fixation under tillage management. Previous research has demonstrated that mechanical disturbance from conventional tillage can largely modify soil physical properties such as bulk density and porosity<sup>34–37</sup>. For instance, Gruber *et al.*<sup>34</sup> reported that ploughed soils exhibited a lower bulk density than no-till soil (1.1 g cm<sup>-3</sup> versus 1.3 g cm<sup>-3</sup>), and tillage also lowered total porosity compared with no-till conditions<sup>35</sup>. These differences in soil porosity and soil bulk density affect the contact between *cbbL*-carrying bacteria and their substrate, <sup>14</sup>CO<sub>2</sub><sup>36–37</sup>, improving soil gas diffusivity<sup>37</sup>. In turn, more <sup>14</sup>CO<sub>2</sub> is likely to be supplied, thus stimulating the CO<sub>2</sub> assimilation activity of *cbbL*-bearing bacteria. Our data support this hypothesis because CT soils exhibit greater specific CO<sub>2</sub> fixation activity than NT soils (Table 2). Previous research has also demonstrated that tillage practices enhance light transmittance<sup>38</sup> and hydraulic conductivity<sup>39</sup>, as well as create novel ecological niches<sup>40</sup>. All of these factors substantially enhance the CO<sub>2</sub> fixation activity of *cbbL*-bearing bacteria<sup>4,8</sup>, leading to our observation of greater fixed <sup>14</sup>C under CT treatment. Moreover, we can assume that tillage management, in altering soil properties, may also affect alternative CO<sub>2</sub> fixation pathways that are sensitive to such changes, thereby contributing to the differences in soil autotrophic microbial CO<sub>2</sub> fixation between CT and NT treatments. We hope that future analyses will resolve these possibilities.

Generally, we observed a consistent trend across both CT and NT treatments, where the microbial fixed <sup>14</sup>C concentrations are higher in paddy soils than upland soils, although this pattern was less obvious at deeper soil depths (Fig. 1, Table 1). We believe this result was caused by the anaerobic conditions in flooded paddy soils, which were flooded with 1–2 cm of sterile water during incubation. The anaerobic paddy soils, in contrast with the aerobic environment of upland soils, may have provided favourable anaerobic niches that promoted the activity of autotrophic CO<sub>2</sub> fixation bacteria and slowed the decomposition of newly fixed <sup>14</sup>C<sup>8,22</sup>.

In this study, fixed <sup>14</sup>C amounts also varied across different soil depths for both CT and NT treatments (Fig. 1, Table 1). Specifically, conventional tillage had a more pronounced effect on deeper soils than on topsoil, increasing the differences in CO<sub>2</sub> fixation rate at 1–5 cm and 5–17 cm compared with 0–1 cm. Previous studies have shown that photoautotrophs are the main contributors to surface soil CO<sub>2</sub> fixation, whereas chemoautotrophs may be involved in CO<sub>2</sub> assimilation in deeper soil layers by using inorganic compounds as electron donors<sup>23</sup>. Without tillage, soils are more compact and less porous, conditions that will slow <sup>14</sup>CO<sub>2</sub> and H<sub>2</sub> diffusivity, as well as inorganic substrate transfer, down the soil profile. Relative to tilled soil, this inhibition is more pronounced at deeper depths for un-tilled soil<sup>41</sup>. Because <sup>14</sup>CO<sub>2</sub>, H<sub>2</sub>, and inorganic compounds are important electron donors for chemoautotrophic bacterial CO<sub>2</sub> fixation, variation in the vertical stratification of electron donors down the soil profile across CT and NT treatments will likely exert differential effects on chemoautotrophic bacterial activity<sup>23</sup>. Moreover, our previous study indicated that a portion of the fixed <sup>14</sup>C in deep soil layers probably originated




**Figure 2.** The  $^{14}\text{C}$ -MBC concentrations recovered at different depths (0–1 cm, 1–5 cm, and 5–17 cm) in conventional tillage (CT) and no-till (NT) soils after 110 days of incubation. Error bars indicate the standard error of the mean ( $n = 4$ ). \* indicates a significant differences between CT and NT soils,  $P < 0.05$ ; nd, not detectable.

from the downward translocation of microbially assimilated C at the soil surface<sup>23</sup>. Tillage-induced changes in soil structure, such as increases in bulk density and decreases in porosity, may therefore benefit the transfer of microbially assimilated C down the soil profile, providing an explanation for the differential impact of tillage at varying soil depths.

## Methods

**Soil sampling.** The experiment was carried out with two paddy soils (P1, P2) and two upland soils (U1, U2) from different regions of Hunan Province, in the subtropical region of China. The properties of the soils before incubation are shown in Table 3. These soils cover the typical land use types in this area: P1 and P2 are from double-rice plantations, U1 is a vegetable plantation base, and U2 receives upland-crop rotation. Soil sampling was conducted in November 2010 after the final harvest of crops. For each site, two sets of soil samples, termed CT treatment soil and NT treatment soil, were prepared.



**Figure 3.** The  $^{14}\text{C}$ -DOC concentrations recovered at different depths (0–1 cm, 1–5 cm, and 5–17 cm) in conventional tillage (CT) and no-till (NT) soils after 110 days of incubation. Error bars indicate the standard error of the mean ( $n = 4$ ). \* indicates significant differences between CT and NT soils,  $P < 0.05$  level; nd, not detectable.

Intact soil cores (i.e. NT treatment soils) were collected directly by inserting four polyvinyl chloride (PVC) containers (10 cm diameter, 20 cm height) approximately 17 cm into soils. The PVC pipes were immediately sealed with a fitted end cap after being extracted from the sampling site and transferred to the laboratory.

To establish the CT treatment soil at the corresponding site, soil cores were randomly sampled using a stainless steel auger and homogenised with mixing. After plant residues and stones were removed, the mixed soil was air-dried and then sieved through a  $< 5$  mm mesh. Before commencing the  $^{14}\text{C}$ -CO $_2$  labelling experiment, the air-dried soils were rewetted using distilled water (P1 and P2, flooding; U1 and U2, 45% water holding capacity [WHC]), and all soils were equilibrated for 2 weeks to stabilise microbial activity. Finally, four CT treatment soils were obtained by packing the pre-incubated sieved soils into PVC containers (10 cm diameter, 20 cm height) to a depth of approximately 17 cm, equivalent to the depth of the NT treatment soil. Four replicates of each soil type (P1, P2, U1, U2) were prepared for each treatment.



Soil depth	Soil type	Bacterial (16S rRNA) Abundance (10 <sup>11</sup> copies g <sup>-1</sup> dry soil)		<i>cbbL</i> Abundance (10 <sup>9</sup> copies g <sup>-1</sup> dry soil)		Specific CO <sub>2</sub> fixation activity (10 <sup>-7</sup> g per copy)	
		CT soil	NT soil	CT soil	NT soil	CT soil	NT soil
0–1 cm	P1	0.86 ± 0.01 b	1.58 ± 0.01 a	0.58 ± 0.05 b	1.65 ± 0.08 a	18.10 ± 1.49 a	5.04 ± 0.33 b
	P2	0.59 ± 0.06 b	1.86 ± 0.02 a	0.38 ± 0.11 b	2.98 ± 0.50 a	31.25 ± 6.01 a	3.33 ± 0.44 b
	U1	0.41 ± 0.05 a	0.27 ± 0.03 b	0.26 ± 0.06 a	0.13 ± 0.01 a	10.85 ± 1.82 a	10.10 ± 1.17 a
	U2	0.14 ± 0.02 b	0.39 ± 0.04 a	0.03 ± 0.01 b	0.30 ± 0.13 a	10.40 ± 1.75 a	0.42 ± 0.00 b
1–5 cm	P1	0.73 ± 0.05 b	1.28 ± 0.07 a	0.36 ± 0.01 b	1.11 ± 0.23 a	6.09 ± 0.16 a	0.94 ± 0.02 b
	P2	0.36 ± 0.04 b	1.27 ± 0.13 a	0.07 ± 0.01 b	1.16 ± 0.19 a	29.36 ± 2.84 a	0.57 ± 0.00 b
	U1	0.37 ± 0.04 a	0.30 ± 0.01 a	0.33 ± 0.04 a	0.13 ± 0.01 b	1.87 ± 0.33 a	1.49 ± 0.00 a
	U2	0.12 ± 0.01 b	0.29 ± 0.02 a	0.16 ± 0.06 a	0.39 ± 0.14 a	1.81 ± 0.41 a	0.02 ± 0.00 b
5–17 cm	P1	0.74 ± 0.06 b	1.08 ± 0.07 a	0.34 ± 0.06 b	0.99 ± 0.08 a	1.24 ± 0.44 a	0.01 ± 0.00 b
	P2	0.32 ± 0.04 b	0.86 ± 0.06 a	0.06 ± 0.01 b	0.66 ± 0.04 a	–	–
	U1	0.38 ± 0.04 a	0.31 ± 0.04 a	0.52 ± 0.08 a	0.13 ± 0.00 b	0.30 ± 0.00 a	–
	U2	0.11 ± 0.01 b	0.27 ± 0.02 a	0.16 ± 0.03 b	0.34 ± 0.04 a	0.51 ± 0.00 a	–

**Table 2.** The abundance and specific CO<sub>2</sub> fixation activity of autotrophic bacteria at different depths of CT and NT soils. Different lower case letters represent significant differences ( $P < 0.05$ ) in CT and NT soils.

Soil	Location	Land use	pH	SOC (g·kg <sup>-1</sup> )	Total N (g·kg <sup>-1</sup> )	Total P (g·kg <sup>-1</sup> )	Clay content (%)	CEC (cmol·kg <sup>-1</sup> )
P1	Ganshan	Double-rice	6.15	21.89	2.64	1.05	39.76	11.79
P2	Pantang	Double-rice	5.66	20.93	2.81	0.70	33.19	13.16
U1	Huangxing	Vegetable	6.60	18.73	2.57	3.65	26.12	13.69
U2	Pantang	Rape-corn	4.40	6.19	1.39	0.75	31.38	11.05

**Table 3.** Basic study site information and corresponding soil physicochemical characteristics. CEC, cation exchange capacity; SOC, soil organic carbon.

**Incubation experiment design.** Microcosms of soils under CT and NT treatments were placed into an airtight growth chamber (80 × 250 × 120 cm). Soils were incubated for 110 days with <sup>14</sup>C-CO<sub>2</sub> produced by the reaction between Na<sub>2</sub><sup>14</sup>CO<sub>3</sub> (at a radioactivity of 1.65 × 10<sup>4</sup> Bq·mL<sup>-1</sup>) and HCl (2 M). The concentration of <sup>14</sup>C-CO<sub>2</sub> in the incubation system was maintained at about 350 μL·L<sup>-1</sup>. The concentration of CO<sub>2</sub> in the growth chamber was monitored with an infrared CO<sub>2</sub> sensor (GasCard NG, 6132 A, Guangzhou, China). During the incubation period, lamps generating artificial light (intensity: 500 mmol photons m<sup>-2</sup>·s<sup>-1</sup> PAR) were open from 08:00 AM to 08:00 PM each day. Day/night temperatures were set at 31 ± 1 °C/24 ± 1 °C, respectively, and relative atmospheric humidity was held at 80–90%. Paddy soils were incubated by flooding with a 1–2 cm water layer while upland soils were kept drained (45% WHC) during the incubation period. Soil moisture was also determined at harvest and was nearly identical to the soil water content at the beginning of the experiment (data not shown). At the end of the incubation period, each soil column in the PVC container was sectioned from the top into three intervals (0–1 cm, 1–5 cm, and 5–17 cm). The sectioned soils were stored separately in three parts. One part was dried for the determination of <sup>14</sup>C-SOC content and another portion was immediately used to measure <sup>14</sup>C-MBC. The remaining part was stored at –70 °C for molecular ecological analysis. Soil moisture content was measured by oven-drying the soil at 105 °C immediately after sampling.

**Soil property analysis.** Soils were air-dried and sieved (2 mm) for SOC and total nitrogen measurements, which were performed with dry combustion using a macro elemental analyser (Vario MAX C/N, Elementar, Germany). Total phosphorus was measured using the Mo-Sb colorimetric method<sup>42</sup>. Soil pH was measured in suspension using a soil:H<sub>2</sub>O ratio of 1:2.5 (w/v). Soil clay content was determined using the pipette method<sup>43</sup>, and cation exchange capacity was measured according to the procedure detailed by Thomas<sup>44</sup>.

**Soil <sup>14</sup>C radioactivity analysis.** To remove inorganic carbon (such as CaCO<sub>3</sub>) from soil samples, 3.0 mL 2.5 M HCl was added and mixed with 1.50 g of soil (sieved with a mesh < 0.149 mm) (v:w = 2:1) in Dolphin tubes for 24 hours. Then, prior to measuring <sup>14</sup>C-SOC, aliquots were washed twice with 3.0 mL H<sub>2</sub>O to remove any remaining HCl. Post-washing, 1.50 g of the concentrated, HCl-treated dried soil was transferred to a flask containing K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> (0.2 M, 20 mL) and concentrated H<sub>2</sub>SO<sub>4</sub>–H<sub>3</sub>PO<sub>4</sub> (v:w = 5:1). This mixture was digested at 165 °C for 8 min under continually replenished pure O<sub>2</sub> and for 10 min without O<sub>2</sub> thereafter<sup>22</sup>. The evolved CO<sub>2</sub> was trapped with NaOH (0.4 M, 40 mL), and the <sup>14</sup>C radioactivity was measured using an automated liquid scintillation counter (LS-6500, Beckman, Germany). The measurement of <sup>14</sup>C-MBC was performed with the fumigation-extraction method, and the amount of <sup>14</sup>C-DOC was determined using K<sub>2</sub>SO<sub>4</sub> extracts of non-fumigated soil<sup>38</sup>. Finally, <sup>14</sup>C-SOC, <sup>14</sup>C-MBC, and <sup>14</sup>C-DOC concentrations were calculated according to the procedure described by Ge *et al.*<sup>22</sup>, with additional details available in our previous reports<sup>4,22,23</sup>.

**Soil DNA extraction.** The extraction of soil microbial DNA was performed in triplicate using a FastDNA Spin Kit, following manufacturer protocol (BIO101, Qbiogene Inc., Carlsbad, CA). The DNA extracts were resuspended in sterilised water for quality and quantity checks. The amount of extracted DNA was determined with a spectrophotometer (Nanodrop ND-1000, PeqLab, Germany), and the quality was evaluated using 1% agarose gel electrophoresis.

**Bacterial (16S rRNA) and *cbbL* gene abundance analysis.** Quantification of the bacterial (16S rRNA) gene and the *cbbL* gene was performed with real-time quantitative PCR, using the primers described by Yuan *et al.*<sup>8</sup> and Wu *et al.*<sup>23</sup> respectively. Gene copy numbers were quantified in triplicate using the primers 799F (5'-ACCMGGATTAGATACCCCKG-3') and 1492R (5'-ACGGTTACCTTG-TTACGACTT-3')<sup>8</sup> for 16S rRNA, as well as the primers K2f (5'-ACCAYCAAAGCCSAAGCTSGG-3') and V2r (5'-GCCTTCSAGCTTGCCSACCRC-3')<sup>24</sup> for *cbbL*. The quantification followed previously described procedures<sup>23</sup> using a *cbbL* cloned standard for constructing standard curves. Briefly, the target *cbbL* gene fragment was amplified from extracted DNA and the generated PCR amplicons ligated into the pGEM-T Easy Vector. The vectors were then transformed into *Escherichia coli* DH5 $\alpha$  competent cells following manufacturer protocol (Promega, Mannheim, Germany). Randomly chosen white colonies were sequenced, and positive clones with target inserts were used for plasmid DNA extraction. Ten-fold serial dilutions of plasmid DNA were subjected to quantitative PCR in triplicate to establish the standard curve. Negative controls without template DNA were run in parallel with template DNA for the soil *cbbL* genes. The reaction was performed in 10  $\mu$ L reaction mixtures containing: 5 ng template DNA, 5  $\mu$ L SYBR Premix Extaq (Takara Bio Inc., Shiga, Japan), and 0.1  $\mu$ M of each primer, following previously described thermal cycling conditions<sup>23</sup>. The copy numbers of the 16S rRNA and bacterial *cbbL* gene in the reaction mixture were automatically calculated using SDS 2.3 software within the Real-Time PCR System, with reference to the standard curve generated for each run.

**Specific CO<sub>2</sub> fixation activity.** The CO<sub>2</sub> fixation activity per *cbbL* copy was defined as the specific CO<sub>2</sub> fixation activity of the autotrophic bacteria and was calculated by dividing the observed <sup>14</sup>C-SOC concentration by the detected *cbbL* gene copy number.

**Statistical analysis.** All data are expressed as means with standard errors. Differences in <sup>14</sup>C radioactivity and *cbbL* gene abundance between CT and NT soils were tested using independent samples *t* tests. The effects of soil type, soil depth, soil disturbance, and their interactions on autotrophic bacterial CO<sub>2</sub> uptake rate, bacterial abundance, and specific CO<sub>2</sub> fixation activity were evaluated using univariate ANOVA. Significance for all tests was set at *P* < 0.05, and analyses were performed in SPSS 13.0 for Windows (IBM, Armonk, NY, USA).

## References

- Moser, S. C. Communicating climate change: history, challenges, process and future directions. *WIREs Clim Change* **1**, 31–53. (2010).
- Selesi, D., Schmid, M. & Hartmann, A. Diversity of green-like and red-like ribulose-1,5-bisphosphate carboxylase/oxygenase large-subunit genes (*cbbL*) in differently managed agricultural soils. *Appl. Environ. Microbiol.* **71**, 175–184 (2005).
- Videmsek, U. *et al.* Abundance and Diversity of CO<sub>2</sub>-fixing Bacteria in Grassland Soils Close to Natural Carbon Dioxide Springs. *Microb. Ecol.* **58**, 1–9 (2009).
- Yuan, H., Ge, T., Chen, C., O'Donnell, A. G. & Wu, J. Microbial autotrophy plays a significant role in the sequestration of soil carbon. *Appl. Environ. Microbiol.* **78**, 2328–2336 (2012a).
- Fuchs, G. Alternative pathways of carbon dioxide fixation: insights into the early evolution of life? *Annu. Rev. Microbiol.* **65**, 631–658 (2011).
- Tabita, F. Microbial ribulose 1,5-bisphosphate carboxylase/oxygenase: A different perspective. *Photosynth. Res.* **60**, 1–28 (1999).
- Yuan, H. *et al.* Long-term field fertilization alters the diversity of autotrophic bacteria based on the ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO) large-subunit genes in paddy soil. *Appl. Microbiol. Biotechnol.* **95**, 1061–1071 (2012b).
- Yuan, H. *et al.* Effect of land use on the abundance and diversity of autotrophic bacteria as measured by ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO) large subunit gene abundance in soils. *Biol. Fert. Soils* **49**, 609–616 (2013).
- Tolli, J. & King, G. M. Diversity and structure of bacterial chemolithotrophic communities in pine forest and agroecosystem soils. *Appl. Environ. Microbiol.* **71**, 8411–8418 (2005).
- Bayer, C., Mielniczuk, J., Amado, T., Martin-Neto, L. & Fernandes, S. Organic matter storage in a sandy clay loam Acrisol affected by tillage and cropping systems in southern Brazil. *Soil Till. Res.* **54**, 101–109 (2000).
- Ding, X., Zhang, B., Zhang, X., Yang, X. & Zhang, X. Effects of tillage and crop rotation on soil microbial residues in a rainfed agroecosystem of northeast China. *Soil Till. Res.* **114**, 43–49 (2011).
- Capelle, C., Schrader, S. & Brunotte, J. Tillage-induced changes in the functional diversity of soil biota - A review with a focus on German data. *Eur. J. Soil Biol.* **50**, 165–181 (2012).
- Holland, J. The environmental consequences of adopting conservation tillage in Europe: reviewing the evidence. *Agric. Ecosyst. Environ.* **103**, 1–25 (2004).
- White, P. & Rice, C. Tillage effects on microbial and carbon dynamics during plant residue decomposition. *Soil Sci. Soc. Am. J.* **73**, 138–145 (2009).
- Bhattacharyya, R., Tuti, M., Kundu, S., Bisht, J. & Bhatt, J. Conservation tillage impacts on soil aggregation and carbon pools in a sandy clay loam soil of the Indian Himalayas. *Soil Sci. Soc. Am. J.* **76**, 617–627 (2012).
- Kahlon, M., Lal, R. & Ann-Varughese, M. Twenty two years of tillage and mulching impacts on soil physical characteristics and carbon sequestration in Central Ohio. *Soil Till. Res.* **126**, 151–158 (2013).
- Ji, B., Zhao, Y., Mu, X. & Li, C. Effects of tillage on soil physical properties and root growth of maize in loam and clay in central China. *Plant Soil Environ.* **59**, 295–302 (2013).
- Jiang, X., Wright, A., Wang, X. & Liang, F. Tillage-induced changes in fungal and bacterial biomass associated with soil aggregates: A long-term field study in a subtropical rice soil in China. *Appl. Soil Ecol.* **48**, 168–173 (2011).
- Wang, J. *et al.* Effects of tillage and residue management on soil microbial communities in North China. *Plant Soil Environ.* **58**, 28–33 (2012).
- Cookson, W., Murphy, D. & Roper, M. Characterizing the relationships between soil organic matter components and microbial function and composition along a tillage disturbance gradient. *Soil Biol. Biochem.* **40**, 763–777 (2008).



21. Moebius-Clune, D., Moebius-Clune, B., van Es, H. & Pawłowska, T. Arbuscular mycorrhizal fungi associated with a single agronomic plant host across the landscape: community differentiation along a soil textural gradient. *Soil Biol. Biochem.* **64**, 191–199 (2013).
22. Ge, T. *et al.* Microbial phototrophic fixation of atmospheric CO<sub>2</sub> in China subtropical upland and paddy soils. *Geochim. Cosmochim. AC.* **113**, 70–78 (2013).
23. Wu, X. *et al.* Changes in bacteria CO<sub>2</sub> fixation with depth in agricultural soils. *Appl. Microbiol. Biot.* **98**, 2309–2319 (2014).
24. Nanba, K., King, G. & Dunfield, K. Analysis of facultative lithotroph distribution and diversity on volcanic deposits by use of the large subunit of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Appl. Environ. Microb.* **71**, 2245–2253 (2004).
25. Miltner, A., Richnow, H.-H., Kopinke, F.-D. & Kästner, M. Incorporation of carbon originating from CO<sub>2</sub> into different compounds of soil microbial biomass and soil organic matter. *Isot. Environ. Health.* **41**, 135–140 (2005a).
26. Miltner, A., Richnow, H.-H., Kopinke, F.-D. & Kästner, M. Assimilation of CO<sub>2</sub> by soil microorganisms and transformation into soil organic matter. *Org. Geochem.* **35**, 1015–1024 (2004).
27. Miltner, A. *et al.* Non-phototrophic CO<sub>2</sub> fixation by soil microorganisms. *Plant Soil* **269**, 193–203 (2005b).
28. Doran, J., Elliott, E. & Paustian, K. Soil microbial activity, nitrogen cycling, and long-term changes in organic carbon pools as related to fallow tillage management. *Soil Till. Res.* **49**, 3–18 (1998).
29. Li, S., Jiang, X., Wang, X. & Wright, A. Tillage effects on soil nitrification and the dynamic changes in nitrifying microorganisms in a subtropical rice-based ecosystem: A long-term field study. *Soil Till. Res.* **150**, 132–138 (2015).
30. Tatti, E. *et al.* Tillage management and seasonal effects on denitrifier community abundance, gene expression and structure over Winter. *Microbial Ecol.* doi: 10.1007/s00248-015-0591-x (2015).
31. Plante, A. & McGill, W. Soil aggregate dynamics and the retention of organic matter in laboratory-incubated soil with differing simulated tillage frequencies. *Soil Till. Res.* **66**, 79–92 (2002).
32. Rodrigues, M. *et al.* Legacy phosphorus and no tillage agriculture in tropical oxisols of the Brazilian savanna. *Sci. Total Environ.* (2014).
33. Ziadi, N. *et al.* Long-term tillage and synthetic fertilization affect soil functioning and crop yields in a corn-soybean rotation in eastern Canada. *Can. J. Soil. Sci.* **94**, 365–376 (2014).
34. Gruber, S., Möhring, J. & Claupein, W. On the way towards conservation tillage-soil moisture and mineral nitrogen in a long-term field experiment in Germany. *Soil Till. Res.* **115**, 80–87 (2011).
35. Franzluebbers, A. Water infiltration and soil structure related to organic matter and its stratification with depth. *Soil Till. Res.* **66**, 197–205 (2002).
36. Abid, M. & Lal, R. Tillage and drainage impact on soil quality: I. Aggregate stability, carbon and nitrogen pools. *Soil Till. Res.* **100**, 89–98 (2008).
37. Yonemura, S., Yokozawa, M., Shirato, Y., Nishimura, S. & Nouchi, I. Soil CO<sub>2</sub> concentrations and their implications in conventional and no-tillage agricultural fields. *J. Agric. Meteorol.* **65**, 141–149 (2009).
38. Campiglia, E., Radicetti, E. & Mancinelli, R. Cover crops and mulches influence weed management and weed flora composition in strip-tilled tomato (*solanum lycopersicum*). *Weed Res.* **55**, 416–425 (2015).
39. Castellini, M. & Ventrella, D. Impact of conventional and minimum tillage on soil hydraulic conductivity in typical cropping system in Southern Italy. *Soil Till. Res.* **124**, 47–56 (2012).
40. Kumaresan, D., Stralis-Pavese, N., Abell, G., Bodrossy, L. & Murrell, J. Physical disturbance to ecological niches created by soil structure alters community composition of methanotrophs. *Environ. Microbiol. Rep.* **3**, 613–621 (2011).
41. Yonemura, S. *et al.* Soil respiration, N<sub>2</sub>O, and CH<sub>4</sub> emissions from an Andisol under conventional-tillage and no-tillage cultivation for 4 years. *Biol. Fert. Soils* **50**, 63–74 (2014).
42. Ding, L. J. *et al.* Mobilisation of inorganic phosphorus induced by rice straw in aggregates of a highly weathered upland soil. *J. Sci. Food Agric.* **92**(5): 1073–1079 (2012).
43. Gee, G.W. & Bauder, J. W. Particle-size analysis. In: A. Klute, editor, Methods of soil analysis. Part 1. 2nd ed. Agron. Monogr. 9. ASA and SSSA, Madison, WI. p. 383–411 (1986).
44. Thomas, G. W. Exchangeable cations. In: Page AL, Miller RH, Keeney DR eds. Methods of Soil Analysis, part 1, 2nd edn. American Society Agronomy, Soil Science Society of America, Madison, p. 159–165(1982).

## Acknowledgements

This study was supported financially by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB15020401), the National Natural Science Foundation of China (41271279; 41503081), the Research Fund of State Key Laboratory of Soil and Sustainable Agriculture, Nanjing Institute of Soil Science, CAS (Y412201410), the Research Fund of Key Laboratory of Agro-ecological Processes in Subtropical Region, ISA, CAS (ISA2015101).

## Author Contributions

T.G. designed experiments, conducted <sup>14</sup>C-SOC, <sup>14</sup>C-DOC, and <sup>14</sup>C-MBC measurements, performed data analyses, and wrote the manuscript. X.W. performed 16S rRNA and *ccbL* abundance quantification and wrote the manuscript. Q.L. and H.Y. extracted DNA and took soil samples. Z.Z. and W.W. performed <sup>14</sup>C-CO<sub>2</sub> labelling experiments. A.S. and J.W. analysed data and revised the manuscript. All authors reviewed the manuscript.

## Additional Information

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Ge, T. *et al.* Effect of simulated tillage on microbial autotrophic CO<sub>2</sub> fixation in paddy and upland soils. *Sci. Rep.* **6**, 19784; doi: 10.1038/srep19784 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>