

# SCIENTIFIC REPORTS



OPEN


## Deficiency in DGCR8-dependent canonical microRNAs causes infertility due to multiple abnormalities during uterine development in mice

Yeon Sun Kim<sup>1</sup>, Hye-Ryun Kim<sup>1</sup>, Hyongbum Kim<sup>2</sup>, Seung Chel Yang<sup>1</sup>, Mira Park<sup>1</sup>, Jung Ah Yoon<sup>3</sup>, Hyunjung J. Lim<sup>4</sup>, Seok-Ho Hong<sup>5</sup>, Francesco J. DeMayo<sup>6</sup>, John P. Lydon<sup>6</sup>, Youngsok Choi<sup>1</sup>, Dong Ryul Lee<sup>1,3</sup> & Haengseok Song<sup>1,3</sup>

DGCR8 is an RNA-binding protein that interacts with DROSHA to produce pre-microRNA in the nucleus, while DICER generates not only mature microRNA, but also endogenous small interfering RNAs in the cytoplasm. Here, we produced *Dgcr8* conditional knock-out mice using progesterone receptor (PR)-Cre (*Dgcr8<sup>d/d</sup>*) and demonstrated that canonical microRNAs dependent on the DROSHA-DGCR8 complex are required for uterine development as well as female fertility in mice. Adult *Dgcr8<sup>d/d</sup>* females neither underwent regular reproductive cycles nor produced pups, whereas administration of exogenous gonadotropins induced normal ovulation in these mice. Interestingly, immune cells associated with acute inflammation aberrantly infiltrated into reproductive organs of pregnant *Dgcr8<sup>d/d</sup>* mice. Regarding uterine development, multiple uterine abnormalities were noticeable at 4 weeks of age when PR is significantly increased, and the severity of these deformities increased over time. Gland formation and myometrial layers were significantly reduced, and the stromal cell compartment did not expand and became atrophic during uterine development in these mice. These results were consistent with aberrantly reduced stromal cell proliferation and completely failed decidualization. Collectively, we suggest that DGCR8-dependent canonical microRNAs are essential for uterine development and physiological processes such as proper immune modulation, reproductive cycle, and steroid hormone responsiveness in mice.

MicroRNAs are single-stranded non-coding RNAs that function as key elements of gene regulatory networks by directing the translational repression or degradation of complementary target mRNAs<sup>1,2</sup>. In general, canonical microRNAs are initially produced as the primary microRNA which is recognized by DGCR8, an RNA binding protein, forming the Microprocessor complex with DROSHA, an RNase III-containing enzyme. The Microprocessor complex cleaves the primary microRNA, resulting in stem-loop pre-microRNAs which are then exported from the nucleus to the cytoplasm by EXPORTIN-5 and subsequently processed to mature microRNAs by DICER. In addition to canonical microRNAs, mature microRNAs, such as mirtrons, can be produced via non-canonical pathways<sup>3,4</sup>, suggesting that some clusters of microRNAs could be distinctly regulated by the Microprocessor complex and DICER.

<sup>1</sup>Department of Biomedical Science, CHA University, Seongnam, Gyeonggi, 463-400 Korea. <sup>2</sup>Department of Pharmacology and Brain Korea 21 PLUS Project for Medical Science, Yonsei University College of Medicine, Seoul 120-752, Korea. <sup>3</sup>Fertility Center of CHA Gangnam Medical Center, CHA University, Seoul, Korea. <sup>4</sup>Department of Veterinary Medicine, College of Veterinary Medicine, Konkuk University, Seoul, 143-701 Korea. <sup>5</sup>Department of Internal Medicine, School of Medicine, Stem Cell Institute, Kangwon National University, Chuncheon, Kangwon 200-701, Korea. <sup>6</sup>Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, Texas, 77030 USA. Correspondence and requests for materials should be addressed to H.S. (email: hssong@cha.ac.kr)



**Figure 1. Conditional deletion of *Dgcr8* by PR-Cre in female reproductive tracts.** (a) Temporal expression profiling of *Dgcr8* and PR mRNAs in the uterus at various postnatal days (PND) by realtime RT-PCR. Experiments were performed with pooled RNA samples extracted from at least 3 mice for each stage. (b) Cell type-specific localization of PR during postnatal uterine development by immunohistochemistry. Arrowheads indicate PR positive cells. (c) Representative images of PCR results with genomic DNA of various tissues. Note that *Dgcr8* exon 3 is deleted in the uterus and oviduct but not in other tissues examined. Black and white arrowheads indicate PCR products for inclusion (1085 bp) and deletion (262 bp) of *Dgcr8* exon 3, respectively. (d) Western blotting for DGCR8 protein in the ovaries, oviducts, and uteri of 4-week-old *Dgcr8*<sup>d/d</sup> mice. Note that the genomic DNA PCR and Western blotting analyses both show incomplete *Dgcr8* deletion in the uterus and oviduct. Scale bar: 50  $\mu$ m. T, Tail; O, Ovary; Ov, Oviduct; U, Uterus; K; Kidney.


Mouse models with conditional deletions of microRNA processing factor(s) have provided evidence for their critical roles in various aspects of mammalian development and stem cell biology<sup>5–8</sup>. For instance, two independent studies with conditional deletions of *Dicer* and *Dgcr8* with *Zp3-Cre* (*Dicer*<sup>flox/flox</sup>; *Zp3-Cre* and *Dgcr8*<sup>flox/flox</sup>; *Zp3-Cre*) clearly demonstrated that microRNAs are globally suppressed in mouse oocytes<sup>6,7</sup>. Interestingly, meiotic abnormalities due to defective spindle formation occurred in *Dicer*<sup>flox/flox</sup>; *Zp3-Cre* oocytes but not in *Dgcr8*<sup>flox/flox</sup>; *Zp3-Cre* oocytes, suggesting that some phenotypes of *Dicer* deficient mice result from the dysregulation of endogenous small interfering RNAs (endo siRNAs), rather than microRNAs.

To examine the physiological function(s) of microRNAs in female reproductive tracts, the reproductive phenotypes of mice with conditional deletions of *Dicer* by anti-Müllerian hormone receptor 2 (*Amhr2*)-*Cre* (*Dicer*<sup>flox/flox</sup>; *Amhr2*<sup>cre/+</sup>) have been analyzed<sup>9–11</sup>. Although the spectrum of phenotypes is diverse, there are multiple shared abnormalities in female reproductive tracts, such as oviductal cysts, and a decreased weight and length of the uterus. *Dicer* conditional knockout mice by progesterone receptor (PR)-*Cre* (*Dicer*<sup>flox/flox</sup>; *PR-cre/+*) showed more severe reproductive phenotypes than those observed in *Dicer*<sup>flox/flox</sup>; *Amhr2*<sup>cre/+</sup> mice<sup>12</sup>. These results strongly suggest that spatiotemporal modes of CRE provide diverse reproductive phenotypes that could be affected by microRNAs. To precisely delineate the functions of microRNAs, especially canonical microRNAs, in female reproductive tracts, mouse models with conditional deletion(s) of not only *Dicer*, but also other gene(s) involved in microRNA biogenesis are unquestionably warranted. Here, we generated *Dgcr8* conditional knockout mice by PR-*Cre* and demonstrated that *Dgcr8*-dependent canonical microRNAs are critical for uterine morphogenesis and physiological actions of steroid hormones in female reproductive tracts suitable for embryo implantation in mice.

## Results

### *Dgcr8* is spatiotemporally deleted in female reproductive tracts of *Dgcr8*<sup>d/d</sup> mice in a PR-dependent manner.

To examine when and where *Dgcr8* is deleted in female reproductive tracts, particularly in the uterus of *Dgcr8*<sup>d/d</sup> mice, spatiotemporal expression profiles of *Dgcr8* and *Pgr* (PR) in the uterus were first examined (Fig. 1). Realtime RT-PCR results showed that expression levels of *Dgcr8* at postnatal day (PND) 3 were already comparable to those at PND 28 while PR expression is very low at PND 0 (birth) and 3 (Fig. 1a,b). The PR protein was merely localized in epithelial cells at PND 3 and its expression became stronger at PND 14. At PND 28, PR was exclusively localized not only in the epithelium, but also in the sub-epithelial stroma and myometrium<sup>13</sup>. These results were consistent with those of genotyping PCR for *Dgcr8*<sup>d/d</sup> mice at




**Figure 2. Acyclic infertility of  $Dgcr8^{d/d}$  ( $Dgcr8^{f/f}; PR^{cre/+}$ ) mice with normal ovarian function.** (a) Pregnancy outcomes in  $Dgcr8^{f/f}$  and  $Dgcr8^{d/d}$  mice. Numbers above bars indicate the number of mice with litters/total number of mice examined. (b) Examination of estrous cycle in  $Dgcr8^{d/d}$  mice. Graphs show % change in epithelial cells/total cells observed on slides by a vaginal smear method for a 2-week period ( $n = 3$  to 8 per each genotype). Note that a cyclic change of the epithelial cell percentage did not occur in  $Dgcr8^{d/d}$  mice. (c,d) Comparable ovulation, fertilization and embryo development in  $Dgcr8^{d/d}$  mice. Pregnancy of 8-week-old  $Dgcr8^{f/f}$  and  $Dgcr8^{d/d}$  mice ( $n = 4$  per each genotype) was induced by a superovulation regime with PMSG followed by hCG and mating with wild-type mature males with proven fertility. Microscopic images of 2-cell embryos flushed from oviducts at post-hCG 48 h (Day 2) and blastocysts cultured from the 2-cell stage for 3 days *in vitro* (Day 5). The percentage of 2-cell embryos that developed to blastocysts *in vitro* was not different from each other. The numbers above the bars indicate the number of blastocysts/total number of embryo. scale bar: 100  $\mu$ m. (e) Histological analyses for the ovaries of  $Dgcr8^{f/f}$  and  $Dgcr8^{d/d}$  mice collected at post-hCG 48 h. Note that the corpus luteum (CL) in  $Dgcr8^{d/d}$  mice was similar to that of control  $Dgcr8^{f/f}$  mice. \* indicates CL corpus luteum, scale bar: 200  $\mu$ m.

various stages (Fig. 1c). While we could barely detect PCR products for deleted allele(s) of  $Dgcr8$  prior to 3 weeks of age, they were prominent from 5 weeks of age. Western blotting supported the inference that uterine deletion of  $Dgcr8$  is PR-dependent (Fig. 1d).

**$Dgcr8^{d/d}$  mice are infertile with acyclic estrous cycles.** To examine whether  $Dgcr8^{d/d}$  mice are fertile, they were mated with mature males with proven fertility for 2 months (Fig. 2a).  $Dgcr8^{d/d}$  mice did not produce any litters although  $Dgcr8^{f/f}$  female mice produced 9 litters ( $6.9 \pm 0.3$  pups per litter). By monitoring estrous cycles with daily vaginal smears over a 2-week period, we observed that  $Dgcr8^{d/d}$  mice are anestrus whereas  $Dgcr8^{f/f}$  mice showed regular estrous cycles with respect to changes in the number of epithelial cells (Fig. 2b).

**$Dgcr8^{d/d}$  mice produce fertilizable oocytes with normal corpus lutea.** Previous reports have shown that PR is expressed in the pituitary as well as in female reproductive tracts<sup>13</sup>. Gross histology of the anterior pituitary in  $Dgcr8^{d/d}$  mice was not different from that of  $Dgcr8^{f/f}$  mice. RT-PCR for FSH $\beta$  and LH $\beta$  subunits showed production of gonadotropins in the pituitary of  $Dgcr8^{d/d}$  female mice (Supplementary Fig. 1). Furthermore, growing follicles at various stages were found in these mice (Data not shown). Thus, to examine whether aberrant control of gonadotropins in the pituitary may cause acyclic sterility in these mice, they were given a superovulation regime with exogenous gonadotropins and mated with fertile males. The total number of 2-cell embryos collected and the percentage of 2-cell embryos that developed to the blastocyst stage during *in vitro* culture were similar in  $Dgcr8^{f/f}$  and  $Dgcr8^{d/d}$  mice (Fig. 2c,d). In addition, histological analyses of ovaries at post-human chorionic gonadotropin (hCG) 48 h showed that the ovaries of  $Dgcr8^{d/d}$  mice had morphologically normal corpora lutea



**Figure 3. Aberrant acute immune responses in reproductive organs of pregnant  $Dgcr8^{d/d}$  mice.** (a) Gross morphology of female reproductive organs in  $Dgcr8^{f/f}$  and  $Dgcr8^{d/d}$  mice at post-hCG 24 h after mating with fertile males or no mating. Scale bar: 5 mm. (b) Histological analyses of female reproductive organs shown in (a). The bottom panel shows higher-magnification images of the boxed area in the middle panel. Note that acute inflammatory cells such as neutrophils and eosinophils are aberrantly abundant in female reproductive organs only in  $Dgcr8^{d/d}$  mice mated with fertile males. Scale bar: 100  $\mu\text{m}$ .


(Fig. 2e). Collectively, these results suggest that the acyclic sterility of  $Dgcr8^{d/d}$  mice is not related to intrinsic defects in  $Dgcr8$  deficient ovaries.

**Acute inflammatory infiltration of immune cells occurs in pregnant  $Dgcr8^{d/d}$  mice.** In addition to the acyclic infertility of  $Dgcr8^{d/d}$  mice, we observed acute inflammatory infiltration of immune cells into the ovaries, oviducts, and uteri of some  $Dgcr8^{d/d}$  mice that were given exogenous gonadotropins to induce pregnancy (Fig. 3a). To examine whether microRNA deficiency causes aberrant acute immune responses in these mice, their reproductive organs were histologically examined at post-hCG 24 h following pregnant mare's serum gonadotropin (PMSG) (Fig. 3b). Interestingly, while acute infiltration of neutrophils and eosinophils was clearly detected in these mice after they were mated with fertile males, gonadotropin treatments alone did not produce any obvious immune reactions. Furthermore, other insults such as oil did not produce any distinct immune reaction in these mice treated with gonadotropins (Supplementary Fig. 2). Thus, these results suggest that deficiency of canonical microRNAs leads to the recruitment of acute inflammatory cells that are reactive to semen containing numerous foreign antigens.

**Expression of microRNA processing factors is regulated by ovarian steroid hormones.** To understand whether microRNA processing is influenced in the uterus by ovarian steroid hormones, estrogen ( $E_2$ ) and progesterone ( $P_4$ ), we examined expression profiles of not only  $Dgcr8$  but also other factors involved in microRNA biogenesis in uteri of ovariectomized mice exposed to  $E_2$  and/or  $P_4$  at various time points. Although the expression of  $Dgcr8$  mRNA is not significantly altered by these hormones, mRNA expression of *Dicer*, *Exportin 5 (Xpo5)* and *Argonaute 2 (Argo2)* among factors examined in this study was temporally regulated by  $E_2$  (Supplementary Fig. 3). While  $P_4$  itself did not show any critical actions on expression profiles of these factors, it interfered with estrogen actions on expression patterns of these factors. Furthermore, pretreatment of ICI 182,780, a nuclear estrogen receptor antagonist, inhibited estrogenic actions on expression of these factors (Supplementary Fig. 4). These results collectively suggest that ovarian steroid hormones are involved in regulation of expression of microRNA processing factors with temporal manners in the uterus.

**Onset of multiple uterine abnormalities in  $Dgcr8^{d/d}$  mice coincides with cell- and time-specific expression of PR in the uterus.** To examine whether  $Dgcr8$  deficiency affects uterine development, the uteri of  $Dgcr8^{d/d}$  mice were grossly examined at various stages (Fig. 4a). At 3 weeks of age, the gross morphology, histology and uterine weight of  $Dgcr8^{d/d}$  mice were similar to those of  $Dgcr8^{f/f}$  mice. However, at 4 weeks of age when PR expression is significantly increased both in epithelial and stromal cell compartments (Fig. 1), some  $Dgcr8^{d/d}$  mice started to show multiple uterine defects, and the severity of these deformities increased over time. A comparison of uterine weight to body weight at the pre-pubertal and adult stages showed a progressive decrease in the uterine size of  $Dgcr8^{d/d}$  mice (Fig. 4b). To further investigate uterine abnormalities in  $Dgcr8^{d/d}$  mice, uterine sections of 3- to 8-week-old  $Dgcr8^{d/d}$  mice were examined histologically. Consistent with gross morphology, the uteri became severely abnormal at 5 weeks of age whereas it looked normal at 3 weeks of age (Fig. 4c). Furthermore, the number of glands was significantly reduced in the uteri of  $Dgcr8^{d/d}$  mice beginning at 5 weeks of age (Fig. 4c,d).

An immunohistochemical analysis for alpha smooth muscle actin ( $\alpha$ -SMA), a smooth muscle cell-specific marker, was performed to examine whether a loss of  $Dgcr8$  causes any abnormalities in the uterine myometrium




**Figure 4. Gross and histological analyses for uterine development of  $Dgcr8^{d/d}$  mice at various ages.** (a) Gross morphology of the uteri of 3- to 8-week-old  $Dgcr8^{d/d}$  mice. Note that while the uteri of 3-week-old  $Dgcr8^{d/d}$  mice look normal, they become severely abnormal beginning at 5 weeks of age. Scale bar: 5 mm. (b) Changes of uterine weight/total body weight of  $Dgcr8^{d/d}$  mice during uterine development. Consistent with the distinct morphologic abnormalities observed in mice beginning at 5 weeks of age, uterine weight was dramatically lower than that of control from 5-week-old stage ( $n = 4$  to 6 for each group). (c) Representative microscopic images of the uteri of 3- to 8-week-old  $Dgcr8^{ff}$  and  $Dgcr8^{d/d}$  mice. Scale bar: 100  $\mu$ m. (d) The number of uterine glands/section observed in  $Dgcr8^{ff}$  and  $Dgcr8^{d/d}$  mice ( $n = 4$  to 6 for each group). Unpaired Student's *t*-test, \* $p < 0.05$ , \*\* $p < 0.01$ . O, Ovary; Ov, Oviduct; U, Uterus; M, Myometrium; S, Stroma; GE, Gland epithelium; LE, Luminal epithelium. White arrowheads indicate glands.

(Fig. 5a,b). The myometrial thickness evaluated by  $\alpha$ -SMA was more or less similar to that of  $Dgcr8^{ff}$  mice at pre-pubertal stages, as expected from the PR expression patterns. However, it became significantly reduced at 8-week-old  $Dgcr8^{d/d}$  mice since the inner circular smooth muscle layer suddenly became narrow. Furthermore, the stromal cell compartment gradually dwindled in  $Dgcr8^{d/d}$  mice throughout uterine development whereas it became noticeably expanded in  $Dgcr8^{ff}$  mice between 4 and 5 weeks of age (Fig. 5c). Collectively, these results suggest that the development of all three major uterine cell compartments is severely impaired in  $Dgcr8^{d/d}$  mice.

**Disturbed hormone responses in stromal cells cause aberrantly reduced cell proliferation in  $Dgcr8^{d/d}$  mice.** To examine whether multiple uterine abnormalities could be due to aberrant response(s) to ovarian steroid hormones,  $E_2$  and  $P_4$ , BrdU incorporation experiments were performed with ovariectomized mice treated with vehicle,  $E_2$  or  $E_2 + P_4$  (Fig. 6)<sup>14,15</sup>.  $E_2$  induced epithelial cell proliferation in both  $Dgcr8^{ff}$  and  $Dgcr8^{d/d}$  mice. However, stromal cells did not undergo proliferation at all in  $Dgcr8^{d/d}$  mice under the influence of  $E_2 + P_4$ , whereas  $Dgcr8^{ff}$  mice showed typical stromal cell proliferation. Furthermore, Ki-67 immunofluorescence staining indicated that stromal cell proliferation is impaired in these mice throughout uterine development (Fig. 7a). Consistent with the impaired stromal cell proliferation (Figs 6 and 7), uterine stromal cells of  $Dgcr8^{d/d}$  mice pretreated with ovarian steroid hormones failed to show signs of decidualization when it was experimentally-induced by giving an oil injection into one of the uterine horns (Fig. 8).

Since aberrantly reduced PR expression could be a major cause of abnormal  $P_4$  responses in  $Dicer^{flx/flx}; PR^{cre/+}$  mice<sup>12</sup>, we examined whether PR is appropriately expressed in these mice during uterine development (Fig. 7b). In contrast to the significant reduction of uterine PR expression in  $Dicer^{flx/flx}; PR^{cre/+}$  mice<sup>12</sup>, PR was clearly detectable in all three major uterine cell types in  $Dgcr8^{d/d}$  mice, similar to its expression in  $Dgcr8^{ff}$  mice during uterine development. Collectively, these results suggest that  $Dgcr8$  deficient stromal cells are unable to properly respond to ovarian steroid hormones, leading to severe stromal atrophy. To investigate whether severe atrophy in stromal cells of  $Dgcr8^{d/d}$  mice is affected by facilitated apoptosis, TUNEL assays were performed in 3- and 5-week-old mice. Few TUNEL positive cells were detected in both  $Dgcr8^{ff}$  and  $Dgcr8^{d/d}$  mice (Supplementary Fig. 5). Furthermore, the expression levels of apoptosis-related genes such as *Bcl2l1*, *Aldh1a3* and *Fas* were not significantly different between them. These results suggest that severe atrophy in the stromal cell compartment of  $Dgcr8^{d/d}$  uteri is not associated with facilitated apoptosis.




**Figure 5. Stromal and myometrial abnormalities during uterine development of *Dgcr8*<sup>d/d</sup> mice.** (a) Immunohistochemical analyses for alpha smooth muscle actin ( $\alpha$ -SMA) in the uterine myometrial layers of *Dgcr8*<sup>d/d</sup> mice. Dark brown color indicates  $\alpha$ -SMA positive smooth muscle cells in the uterus. Scale bar: 100  $\mu$ m. (b) Myometrial thickness in the uteri of *Dgcr8*<sup>f/f</sup> and *Dgcr8*<sup>d/d</sup> mice (n = 4 to 6 for each group). Myometrial thickness was determined by the length of the area of  $\alpha$ -SMA positive cell layers. (c) Quantification of stromal area during uterine development in *Dgcr8*<sup>d/d</sup> mice. Stromal area was quantitatively measured by pixels of images for each uterine section of *Dgcr8*<sup>f/f</sup> and *Dgcr8*<sup>d/d</sup> mice. At least three independent sections were microscopically examined for each mouse (n = 4 to 6 mice for each group). Unpaired Student's *t*-test, \**p* < 0.05, \*\**p* < 0.01.

## Discussion


The endometrium is a complex tissue that undergoes cyclic proliferation and differentiation under the influence of ovarian steroid hormones,  $E_2$  and  $P_4$ . Imbalance between the actions and levels of the major regulators of endometrial function may cause several pathophysiological conditions such as endometriosis and endometrial cancer<sup>16,17</sup>. MicroRNAs have been proposed to regulate the stability of mRNAs that encode proteins involved in maintaining normal physiological processes such as cell proliferation and differentiation, apoptosis, angiogenesis, and inflammation<sup>18,19</sup>. Recent studies have demonstrated that dysregulated microRNA expression as well as hormonal discrepancy is also important for incidence of these conditions in the uterus<sup>20,21</sup>.

Previous reports have suggested that ovarian steroid hormones regulate expression profiles of factors participating in microRNA biogenesis as well as many microRNAs in the uterus<sup>22–24</sup>. It was shown that  $E_2$  and  $P_4$  significantly increase the levels of *Xpo5* mRNA, but only  $P_4$  increases *Dicer* expression<sup>22</sup>. However,  $E_2$  but not  $P_4$  influenced the expression patterns of *Xpo5* and *Dicer* at the mRNA level in this study (Supplementary Fig. 3). In addition, mRNA expression of *Argo2*, a member of the Argonaute family, showed similar patterns to those of *Dicer* and *Xpo5* after  $E_2$  treatment. While  $P_4$  did not have significant action on expression of all the genes examined in the uterus, it seemed to dampen and/or delay the action of  $E_2$  on the expression of *Dicer*, *Xpo5*, and *Argo2* (Supplementary Fig. 3). Although there are several incompatible results between a previous work<sup>22</sup> and this study, both demonstrated that action(s) of steroid hormones on the expression patterns of these genes are mediated via their nuclear receptors (Supplementary Fig. 4). To further understand exact actions of steroid hormones on the expression profiles of factors involved in microRNA biogenesis, cell type-specific localization of these factors in the uterus under the influence of ovarian steroid hormones needs to be examined.


Two Cre transgenic lines, *Amhr2*<sup>cre/+</sup> (*Amhr2*-Cre) and *PR*<sup>cre/+</sup> (*PR*-Cre), have been mainly utilized to delete genes of interest in female reproductive tracts<sup>9–12</sup>. *Amhr2*-Cre is expressed embryonically in the mesenchyme of the developing Mullerian ducts, and postnatally in ovarian granulosa cells and the stromal and myometrial layers of female reproductive tracts<sup>25–27</sup>. As expected from *Amhr2* expression profiles, *Dicer*<sup>fllox/fllox</sup>; *Amhr2*<sup>cre/+</sup> mice are infertile due to multiple defects in female reproductive tracts such as disorganized oviducts with cysts and shorter uterine horns, although these mice have normal mating behavior<sup>9–11</sup> that is not shown in *Dgcr8*<sup>d/d</sup> mice (Fig. 2).



**Figure 6.** Cell proliferation in the uteri of ovariectomized  $Dgcr8^{d/d}$  mice treated with ovarian steroid hormones. (a) BrdU incorporation experiments were performed to examine hormone-dependent endometrial cell proliferation in  $Dgcr8^{d/d}$  mice. Ovariectomized mice were sacrificed 24 h after injection with vehicle,  $E_2$  or  $E_2 + P_4$ . BrdU was given to these mice 3 h before sacrifice. Brown color indicates the nuclei of BrdU-incorporated cells. Scale bar: 25  $\mu\text{m}$ . (b,c) Graphs depicting the percentage of BrdU positive cells/total number of cells counted. Note that stromal cell proliferation was severely impaired in  $Dgcr8^{d/d}$  mice, whereas uterine epithelial cells normally responded to  $E_2$ . Unpaired Student's  $t$ -test, \* $p < 0.05$ , \*\* $p < 0.01$ .



**Figure 7.** Expression of Ki-67 and PR in uteri of  $Dgcr8^{d/d}$  mice at various stages. (a,b) Immunofluorescence of Ki-67 (a) and PR (b) on uterine sections from  $Dgcr8^{f/f}$  and  $Dgcr8^{d/d}$  mice at various ages. Ki-67 or PR was visualized as green and nuclei were stained with TO-PRO-3-Iodide (red). Yellow color indicates Ki-67 or PR positive cells. Scale bar: 50  $\mu\text{m}$ .




**Figure 8. Impaired decidualization of  $Dgcr8^{d/d}$  mice.** (a) Gross anatomy of  $Dgcr8^{d/d}$  uterus 4 days after decidualization was artificially induced by injecting oil into a uterine horn. Scale bar: 5 mm. (b) Decidual response was determined by uterine weight of stimulated horn/control horn. (n = 4 to 5 for each group). Unpaired Student's *t*-test, \**p* < 0.05. (c) Representative microscopic images of the uteri of  $Dgcr8^{f/f}$  and  $Dgcr8^{d/d}$  mice traumatized with oil. Scale bar: 100  $\mu$ m.

With respect to the severity and types of phenotypes in the uterus, conditional deletion of *Dicer* or *Dgcr8* by PR-Cre led to more severe abnormalities than those in  $Dicer^{flox/flox};Amhr2^{cre/+}$  mice. For example, decidualization was normally induced in  $Dicer^{flox/flox};Amhr2^{cre/+}$  mice<sup>11</sup>, but completely failed in  $Dgcr8^{d/d}$  mice (Fig. 8). In addition, the inner circular smooth muscle layer seems to be suddenly disintegrated in adult  $Dgcr8^{d/d}$  mice (Fig. 5). These differences could reflect the usage of different CRE systems in the uterus, considering that  $Dgcr8^{d/d}$  mice show various uterine defects that were shared with  $Dicer^{flox/flox};PR^{cre/+}$  mice. The reproductive abnormalities have many similarities, including shorter uterine horns, a reduced number of glands, and severe stromal atrophy with aberrant hormone responsiveness (Fig. 9), although the onset of reproductive phenotypes is somewhat different between  $Dicer^{flox/flox};PR^{cre/+}$  mice<sup>12</sup> and  $Dgcr8^{d/d}$  mice. These results indicate that canonical microRNAs are critical for postnatal uterine development, architecture, and physiological function. Moreover,  $Dicer^{flox/flox};PR^{cre/+}$  mice<sup>12</sup> and  $Dgcr8^{d/d}$  mice share stromal atrophy with aberrant progesterone signaling (Fig. 6), a unique phenotype that did not occur in  $Dicer^{flox/flox};Amhr2^{cre/+}$  mice.

A growing number of publications have provided evidence that specific microRNAs in many species, including humans and mice, are essential for the sequential events of embryo implantation from embryo-epithelial juxtacrine adhesion to decidualization by regulating critical factors for embryo implantation such as IGF1R, MUC1, LIF and COX-2<sup>28–32</sup>. While DGCR8 expression is not altered in the uterus during early pregnancy, DROSHA expression is increased during the decidualization of stromal cells *in vitro*<sup>24</sup>, implicating the importance of microRNAs during these events. In this respect, it is interesting that knockdown of *Dicer* dysregulates expression





**Figure 9.** A schematic cartoon to summarize major reproductive phenotypes of *Dicer* or *Dgcr8* conditional knockout mice by *Amhr2-Cre* or *PR-Cre*.

of only a few dozens of microRNAs and has only a minor effect on human endometrial stromal cells during *in vitro* decidualization<sup>33</sup>. Furthermore, *Dicer* deficiency causes a minor alteration in the microRNA signature in the oviducts (*Dicer<sup>fllox/fllox</sup>;Amhr2<sup>cre/+</sup>*) and uteri (*Dicer<sup>fllox/fllox</sup>;PR<sup>cre/+</sup>*) of *Dicer* conditional knockout mice<sup>9–12</sup>, suggesting that a large cohort of microRNAs could be produced in female reproductive tracts by non-canonical microRNA processing.

Unexpectedly, we observed acute immune cell infiltration in *Dgcr8<sup>d/d</sup>* mice mated with males for induction of pregnancy after a superovulation regime was given (Fig. 3). Administration of exogenous gonadotropins itself did not provoke any acute inflammatory responses unless *Dgcr8<sup>d/d</sup>* female mice were mated with fertile males. This suggests that microRNAs, especially those regulated by Microprocessor complex, could be a participant for immune modulation in these organs. MicroRNA-dependent signaling pathways may modulate cytokines to provoke immune cell infiltration, which can be caused by semen containing various foreign antigens. In addition, immune cells deficient of microRNAs may act aberrantly for themselves since immune cells such as T cells also express PR<sup>34,35</sup>. This phenotype is somewhat different from the chronic inflammation observed in immature non-pregnant *Dicer<sup>fllox/fllox</sup>;Amhr2<sup>cre/+</sup>* mice<sup>9</sup>. Further studies are warranted to delineate the regulatory action(s) of microRNAs on the aberrant immune responses in these organs.

Since DICER is involved in the production of not only canonical microRNAs but also endo siRNAs, non-canonical microRNAs, and other small RNAs in mammals<sup>4,36</sup>, *Dgcr8<sup>d/d</sup>* mice could be an alternative animal model to understand the precise action(s) of microRNAs in mammals. In fact, to clarify the distinct physiological roles of microRNAs and endo siRNAs, several studies have compared the phenotypes of *Dicer* and *Dgcr8* conditional knockout mice with tissue- or cell type-specific manners<sup>37–40</sup>. Our results in *Dgcr8<sup>d/d</sup>* mice reinforced the notion that reproductive abnormalities in the oviducts and uteri of *Dicer* conditional knockout mice with either *PR-Cre* or *Amhr2-Cre* are related to deficiencies in canonical microRNAs. However, several unique phenotypes including myometrial defect were observed in *Dgcr8<sup>d/d</sup>* mice (Fig. 5b). Moreover, *Dgcr8<sup>fllox/fllox</sup>;Amhr2<sup>cre/+</sup>* mice show subfertility with smaller litter size which has not been observed in any *Dicer* or *Dgcr8* conditional knockout mice to date (Kim *et al.* in preparation). These data suggest a possibility that the Microprocessor complex containing DGCR8 may have a distinct action other than canonical microRNA processing. In fact, there was an interesting report that the Microprocessor complex negatively regulates long interspersed element-1 and Alu retrotransposons in microRNA- and DICER-independent manners<sup>41</sup>. Furthermore, a recent study has provided evidence that DGCR8 itself acts for neural morphogenesis in a DROSHA-independent way in *Drosophila*<sup>42</sup>. Additional studies are definitely requested to determine whether DGCR8 has unique functions that are not associated with the sequential events of microRNA processing. In summary, here we provide a valuable animal model deficient of *Dgcr8* that have similar and distinct reproductive phenotypes compared with *Dicer* deficient mice. Our results clearly demonstrate that canonical microRNAs are critical for uterine morphogenesis and the physiological actions of steroid hormones in female reproductive tracts suitable for embryo implantation in mice.

## Methods

**Animals and genotyping of *Dgcr8<sup>d/d</sup>* mice.** All animal experiments were carried out in accordance with the protocols approved by the Institutional Animal Care and Use Committee of CHA University (IACUC),

No 140005). All animals used in this study were maintained and handled according to the policies approved by CHA University. Six to 8-week-old adult ICR mice were provided by Orient Bio (Gapyeong, Gyeonggi, Korea).

Dgcr8<sup>fllox/fllox</sup> mice were initially generated and provided by Dr. Elaine Fuchs' laboratory<sup>43</sup>. Genotyping PCR was performed using genomic DNA extracts from mouse tail biopsies. Dgcr8<sup>fllox/fllox</sup> mice were bred to PR<sup>cre/+</sup> (herein called PR-Cre) mice<sup>13</sup> to generate Dgcr8<sup>fllox/+</sup>;PR<sup>cre/+</sup> mice. These mice were then crossed to generate Dgcr8<sup>fllox/fllox</sup>;PR<sup>cre/+</sup> (designated as Dgcr8<sup>d/d</sup> mice hereafter) and Dgcr8<sup>fllox/fllox</sup>;PR<sup>+ /+</sup> mice (Dgcr8<sup>flf</sup> mice). To distinguish the Dgcr8 null allele (Dgcr8<sup>d/d</sup>) from Dgcr8<sup>flf</sup>, primers were designed to amplify 262 bp (deletion of exon 3) and 1085 bp (presence of exon 3) PCR products as shown in Fig. 1c.

**RNA extraction, reverse transcription-PCR (RT-PCR) and realtime RT-PCR.** Total RNA was extracted from mouse uterus by using Trizol Reagent (Invitrogen Life Technologies, San Diego, CA, USA) according to the manufacturer's protocols. The first-strand cDNA was synthesized from 2 µg of total RNA by using M-MLV reverse transcriptase (Promega, Madison, WI, USA) and RNasin Ribonuclease Inhibitor (Promega). The synthesized cDNA was utilized for PCR with specific primers at optimized cycles. For quantification of expression level, realtime RT-PCR was performed by using iQ<sup>TM</sup> SYBR Green Supermix (Bio-Rad, Hercules, CA, USA) on a BIO-RAD iCycler. To compare transcript levels between samples, a standard curve of cycle thresholds for several serial dilutions of a cDNA sample was established and then used to calculate the relative abundance of each gene. rPL7 was used as a reference gene in all the experiments performed in this study. All PCR reactions were performed in duplicate.

**Western blotting.** Tissues were lysed in lysis buffer including PRO-PREP (iNtRON, Seongnam, Korea) solution and 1X phosphatase inhibitor (Roche Applied Sciences, Indianapolis, IN, USA). The protein samples (10 µg/lane) were then separated by 8% SDS-PAGE, transferred onto nitrocellulose membrane (Bio-Rad) and blocked with 5% non-fat milk (Bio-Rad) in TBS (Bio-Rad) containing 0.1% Tween 20 (Sigma-Aldrich, St. Louis, MO, USA). Anti-DGCR8 (Proteintech, Chicago, IL, USA, 1:100) and anti-GAPDH (Cell Signaling, Danvers, MA, USA, 1:2000) were used for Western blotting analyses. The signals were developed using an ECL Western blotting substrate kit (Bio-Rad) and detected using a Chemidoc XRS+ (Bio-Rad) with Image Lab software.

**Vaginal smear assay and fertility analysis.** Vaginal smears from 6- to 7-week-old Dgcr8<sup>flf</sup> and Dgcr8<sup>d/d</sup> mice were collected daily over a 2-week period as previously described<sup>44</sup>. The stage of the estrous cycle (proestrus, estrus, metestrus, and diestrus) was determined based on the presence or absence of leukocytes, cornified epithelial cells, and nucleated epithelial cells. To evaluate reproductive performance, 8-week-old Dgcr8<sup>flf</sup> and Dgcr8<sup>d/d</sup> mice (n = 6 for each genotype) were individually bred to males with proven fertility. The numbers of litters and pups were recorded for a 2-month period.

**Superovulation and preimplantation embryo culture.** To induce ovulation, 8-week-old mice were given intraperitoneal (i.p) injections of 5 IU PMSG (Sigma-Aldrich) for 48 h, followed by i.p injections with 5 IU hCG (Sigma-Aldrich). Mice were then bred to wild-type males with known fertility and pregnancy was evaluated by the presence of vaginal plug next morning. The 2-cell embryos flushed from oviducts at post-hCG 48 h were cultured up to the blastocyst stage in 20 µl droplets of KSOM (Millipore, Danvers, MA, USA) covered with oil (SAGE *In-Vitro* Fertilization, Inc., Trumbull, CT, USA) in a petri dish.

**Tissue collection, H&E staining and immunostaining.** Female reproductive organs were dissected, and then fixed in 4% paraformaldehyde for histology or snap frozen for RNA and/or protein preparation. Fixed tissues were washed, dehydrated, and embedded in Paraplast (Merck KGaA, Darmstadt, Germany). Paraffin-embedded tissues were sectioned using a microtome, stained with hematoxylin and eosin (H&E) (Sigma-Aldrich), and observed by a light microscopy.

For immunostaining analyses, antibodies specific to PR (Thermo Scientific, Pierce Biotechnology, Rockford, IL, USA, 1:100), α-SMA (Abcam, Cambridge, UK, 1:100), and Ki-67 (Abcam, 1:100) were used in 5 µm thick paraffin-embedded sections. Blocking was carried out using protein block serum (Dako, Carpinteria, CA, USA) for 1 h, and then samples were incubated with an appropriate primary antibody at 4°C overnight. On the following day, sections were washed in PBS and incubated with HRP secondary antibody or FITC conjugated donkey anti-rabbit IgG (Jackson ImmunoResearch, West Grove, PA, USA) for 1 h at room temperature. Nuclear staining was performed using TO-PRO-3-iodide (Life Technologies, Carlsbad, CA, USA). For Immunohistochemistry, DAB reagent (Vector Laboratories, Inc., Burlingame, CA, USA) was applied to visualize signals. Slides were counterstained with hematoxylin, mounted with mounting solution and cover-slipped.

**Quantitative measurement of the number of uterine glands and the thickness of the uterine smooth muscle layer in Dgcr8<sup>d/d</sup> mice.** To quantitatively examine uterine gland formation in Dgcr8<sup>d/d</sup> mice, 4 independent uterine cross-sections for each 3- to 8-week-old mouse were H&E stained and the number of uterine glands was counted in a blind manner (n = 3 to 5 mice per genotype in each age group). Immunohistochemical staining for α-SMA, as a smooth muscle cell marker, was performed to evaluate the thickness of myometrial layers of Dgcr8<sup>d/d</sup> uteri. The captured microscopic images of α-SMA staining were applied to Image J software (National Institutes of Health, Bethesda, MD, USA) for measuring the thickness of myometrial layers of Dgcr8<sup>d/d</sup> mice at each age.

**BrdU incorporation and TUNEL assays.** Eight-week-old Dgcr8<sup>flf</sup> and Dgcr8<sup>d/d</sup> mice were ovariectomized, rested for 2 weeks, and treated subcutaneously with either 0.1 ml sesame oil as vehicle (Acros, NJ, USA), 200 ng E<sub>2</sub> (Sigma-Aldrich), or 200 ng E<sub>2</sub> + 2 mg P<sub>4</sub> (Sigma-Aldrich). Mice were sacrificed 24 h after

steroid hormone treatment(s). 5-Bromo-2'-Deoxyuridine (BrdU) (Invitrogen Life Technologies) was given to the mice 3 h before they were sacrificed. Uteri were dissected, fixed in 4% paraformaldehyde and then subjected to paraffin-embedded tissue processing. Uterine sections were utilized for BrdU immunostaining using a BrdU staining kit (Invitrogen Life Technologies), according to the manufacturer's instructions.

Uterine apoptosis of Dgcr8<sup>d/d</sup> mice was assessed using an *In Situ* Cell Death Detection Kit according to the manufacturer instructions (Roche, West Sussex, UK). Sections were deparaffinized and rehydrated in a graded alcohol series, and then processed for antigen retrieval. They were incubated with TUNEL reaction mixture for 1 h at 37 °C and then with DAPI for 10 min at room temperature, and observed under a fluorescence microscope.

**Artificial decidualization.** Briefly, 8-week-old female mice were ovariectomized and rested for 2 weeks, and then received daily injections of 100 ng E<sub>2</sub> for 3 days. After 2 days of resting, mice were then treated with daily injections of 1 mg P<sub>4</sub> and 6.7 ng E<sub>2</sub> for 3 days. At 6 h after the last injection, one uterine horn was traumatized by the injection of 50 µl of sesame oil. Mice were given daily injections of 1 mg P<sub>4</sub> and 6.7 ng E<sub>2</sub>/mouse following trauma. Mice were sacrificed at day 4 after the trauma.

**Statistics.** Realtime RT-PCR was performed in duplicate for at least three independent samples. All values represent the mean ± standard deviation. Unpaired Student's *t*-tests were used for statistical evaluation and *p* < 0.05 was considered statistically significant.

## References

- Dong, H. *et al.* MicroRNA: function, detection, and bioanalysis. *Chemical reviews* **113**, 6207–6233 (2013).
- Ha, M. & Kim, V. N. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol* **15**, 509–524 (2014).
- Havens, M. A., Reich, A. A., Duelli, D. M. & Hastings, M. L. Biogenesis of mammalian microRNAs by a non-canonical processing pathway. *Nucleic Acids Res* **40**, 4626–4640 (2012).
- Ha, M. & Kim, V. N. Regulation of microRNA biogenesis. *Nature reviews Molecular cell biology* **15**, 509–524 (2014).
- Gruber, A. J. *et al.* Embryonic stem cell-specific microRNAs contribute to pluripotency by inhibiting regulators of multiple differentiation pathways. *Nucleic acids research* **42**, 9313–9326 (2014).
- Ma, J. *et al.* MicroRNA activity is suppressed in mouse oocytes. *Current biology* **20**, 265–270 (2010).
- Suh, N. *et al.* MicroRNA function is globally suppressed in mouse oocytes and early embryos. *Current Biology* **20**, 271–277 (2010).
- Judson, R. L., Babiarz, J. E., Venero, M. & Belloch, R. Embryonic stem cell-specific microRNAs promote induced pluripotency. *Nature biotechnology* **27**, 459–461 (2009).
- Gonzalez, G. & Behringer, R. R. Dicer is required for female reproductive tract development and fertility in the mouse. *Mol Reprod Dev* **76**, 678–688 (2009).
- Hong, X., Luense, L. J., McGinnis, L. K., Nothnick, W. B. & Christenson, L. K. Dicer1 is essential for female fertility and normal development of the female reproductive system. *Endocrinology* **149**, 6207–6212 (2008).
- Nagaraja, A. K. *et al.* Deletion of Dicer in somatic cells of the female reproductive tract causes sterility. *Mol Endocrinol* **22**, 2336–2352 (2008).
- Hawkins, S. M. *et al.* Dysregulation of uterine signaling pathways in progesterone receptor-Cre knockout of dicer. *Mol Endocrinol* **26**, 1552–1566 (2012).
- Soyal, S. M. *et al.* Cre-mediated recombination in cell lineages that express the progesterone receptor. *Genesis* **41**, 58–66 (2005).
- Tong, W. & Pollard, J. W. Progesterone inhibits estrogen-induced cyclin D1 and cdk4 nuclear translocation, cyclin E-and cyclin A-cdk2 kinase activation, and cell proliferation in uterine epithelial cells in mice. *Molecular and cellular biology* **19**, 2251–2264 (1999).
- Franco, H. L. *et al.* Epithelial progesterone receptor exhibits pleiotropic roles in uterine development and function. *The FASEB Journal* **26**, 1218–1227 (2012).
- Brown, S. B. & Hankinson, S. E. Endogenous estrogens and the risk of breast, endometrial, and ovarian cancers. *Steroids* **99**, 8–10 (2015).
- Kim, J. J., Kurita, T. & Bulun, S. E. Progesterone action in endometrial cancer, endometriosis, uterine fibroids, and breast cancer. *Endocrine reviews* **34**, 130–162 (2013).
- Hwang, H. & Mendell, J. MicroRNAs in cell proliferation, cell death, and tumorigenesis. *British journal of cancer* **94**, 776–780 (2006).
- Lenkala, D. *et al.* The impact of microRNA expression on cellular proliferation. *Human genetics* **133**, 931–938 (2014).
- Pan, Q. & Chegini, N. MicroRNA Signature and Regulatory Functions in the Endometrium during Normal and Disease States. *Seminars in Reproductive Medicine* **26**, 479–493 (2008).
- Klinge, C. M. miRNAs and estrogen action. *Trends in Endocrinology & Metabolism* **23**, 223–233 (2012).
- Nothnick, W. B., Healy, C. & Hong, X. Steroidal regulation of uterine miRNAs is associated with modulation of the miRNA biogenesis components Exportin-5 and Dicer1. *Endocrine* **37**, 265–273 (2010).
- Fu, T.-Y., Lin, C.-T. & Tang, P.-C. Steroid hormone-regulated let-7b mediates cell proliferation and basigin expression in the mouse endometrium. *Journal of Reproduction and Development* **57**, 627–635 (2011).
- Zhang, C. *et al.* Expression of DROSHA in the Uterus of Mice in Early Pregnancy and Its Potential Significance During Embryo Implantation. *Reproductive Sciences*, doi: 10.1177/1933719115584444 (2015).
- Jamin, S. P., Arango, N. A., Mishina, Y., Hanks, M. C. & Behringer, R. R. Requirement of Bmpr1a for Müllerian duct regression during male sexual development. *Nature genetics* **32**, 408–410 (2002).
- Gifford, J. A. H., Hunzicker-Dunn, M. E. & Nilson, J. H. Conditional deletion of beta-catenin mediated by Amhr2cre in mice causes female infertility. *Biology of reproduction* **80**, 1282–1292 (2009).
- Wang, Y. *et al.* Loss of APC function in mesenchymal cells surrounding the Müllerian duct leads to myometrial defects in adult mice. *Molecular and cellular endocrinology* **341**, 48–54 (2011).
- Chakrabarty, A. *et al.* MicroRNA regulation of cyclooxygenase-2 during embryo implantation. *Proceedings of the National Academy of Sciences* **104**, 15144–15149 (2007).
- Chu, B. *et al.* miRNA-181 regulates embryo implantation in mice through targeting leukemia inhibitory factor. *Journal of molecular cell biology* **7**, 12–22 (2015).
- Inyawilert, W., Tzu-Yen, F., Chun-Ting, L. & Pin-Chi, T. Let-7-mediated suppression of mucin 1 expression in the mouse uterus during embryo implantation. *The Journal of reproduction and development* **61**, 138–144 (2015).
- Kang, Y.-J. *et al.* miR-145 suppresses embryo-epithelial juxtacrine communication at implantation by modulating maternal IGF1R. *Journal of cell science* **128**, 804–814 (2015).
- Li, Z., Gou, J., Jia, J. & Zhao, X. MicroRNA-429 functions as a regulator of epithelial-mesenchymal transition by targeting Pcdh8 during murine embryo implantation. *Human Reproduction* **30**, 507–518 (2015).

33. Estella, C. *et al.* miRNA signature and Dicer requirement during human endometrial stromal decidualization *in vitro*. *PLoS one* **7**, e41080 (2012).
34. Hughes, G. C., Clark, E. A. & Wong, A. H. The intracellular progesterone receptor regulates CD4+ T cells and T cell-dependent antibody responses. *Journal of leukocyte biology* **93**, 369–375 (2013).
35. Tan, I. J., Peeva, E. & Zandman-Goddard, G. Hormonal modulation of the immune system—A spotlight on the role of progestogens. *Autoimmunity reviews* **14**, 536–542 (2015).
36. Yang, J.-S. & Lai, E. C. Alternative miRNA biogenesis pathways and the interpretation of core miRNA pathway mutants. *Molecular cell* **43**, 892–903 (2011).
37. Kanellopoulou, C. Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes & Development* **19**, 489–501 (2005).
38. Modzelewski, A. J. *et al.* Dgcr8 and Dicer are essential for sex chromosome integrity during meiosis in males. *Journal of Cell Science* **128**, 2314–2327 (2015).
39. Wang, Y., Medvid, R., Melton, C., Jaenisch, R. & Blelloch, R. DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nature Genetics* **39**, 380–385 (2007).
40. White-Cooper, H. *et al.* Germ Cell-Specific Targeting of DICER or DGCR8 Reveals a Novel Role for Endo-siRNAs in the Progression of Mammalian Spermatogenesis and Male Fertility. *PLoS ONE* **9**, e107023 (2014).
41. Heras, S. R. *et al.* The Microprocessor controls the activity of mammalian retrotransposons. *Nature structural & molecular biology* **20**, 1173–1181 (2013).
42. Luhur, A., Chawla, G., Wu, Y.-C., Li, J. & Sokol, N. S. Drosha-independent DGCR8/Pasha pathway regulates neuronal morphogenesis. *Proceedings of the National Academy of Sciences* **111**, 1421–1426 (2014).
43. Yi, R. *et al.* DGCR8-dependent microRNA biogenesis is essential for skin development. *Proceedings of the National Academy of Sciences* **106**, 498–502 (2009).
44. Byers, S. L., Wiles, M. V., Dunn, S. L. & Taft, R. A. Mouse estrous cycle identification tool and images. *PLoS One* **7**, e35538 (2012).

### Acknowledgements

We thank Elaine Fuchs (Rockefeller University) for generously providing Dgcr8<sup>fl/fl</sup> mice. This work was supported in part by the National Research Foundation of Korea (NRF) grants funded by the Korea government (MEST) (NRF-2012R1A2A2A01011274 and NRF-2015R1A2A2A01006714), and by a grant from the National Institutes of Health (NIH) (R01 HD042311/NICHD to JPL).

### Author Contributions

Y.S.K. and H.S. conceived and designed the experiments. Y.S.K., H.K., H.K., S.C.Y., M.P. and J.A.Y. performed experiments. Y.S.K., H.K., H.J.L., S.H., F.J.D., J.P.L., Y.C., D.R.L. and H.S. analyzed the results. Y.S.K. and H.S. discussed the results and wrote the manuscript.

### Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Kim, Y. S. *et al.* Deficiency in DGCR8-dependent canonical microRNAs causes infertility due to multiple abnormalities during uterine development in mice. *Sci. Rep.* **6**, 20242; doi: 10.1038/srep20242 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>