

SCIENTIFIC REPORTS

OPEN

Influence of hypothermia and subsequent rewarming upon leukocyte-endothelial interactions and expression of Junctional-Adhesion-Molecules A and B

Received: 14 December 2015

Accepted: 04 February 2016

Published: 25 February 2016

Nicolai V. Bogert¹, Isabella Werner¹, Angela Kornberger¹, Patrick Meybohm², Anton Moritz¹, Till Keller³, Ulrich A. Stock¹ & Andres Beiras-Fernandez¹

Patients with risks of ischemic injury, e.g. during circulatory arrest in cardiac surgery, or after resuscitation are subjected to therapeutic hypothermia. For aortic surgery, the body is traditionally cooled down to 18 °C and then rewarmed to body temperature. The role of hypothermia and the subsequent rewarming process on leukocyte-endothelial interactions and expression of junctional-adhesion-molecules is not clarified yet. Thus, we investigated in an *in-vitro* model the influence of temperature modulation during activation and transendothelial migration of leukocytes through human endothelial cells. Additionally, we investigated the expression of JAMs in the rewarming phase. Exposure to low temperatures alone during transmigration scarcely affects leukocyte extravasation, whereas hypothermia during treatment and transendothelial migration improves leukocyte-endothelial interactions. Rewarming causes a significant up-regulation of transmigration with falling temperatures. JAM-A is significantly modulated during rewarming. Our data suggest that transendothelial migration of leukocytes is not only modulated by cell-activation itself. Activation temperatures and the rewarming process are essential. Continued hypothermia significantly inhibits transendothelial migration, whereas the rewarming process enhances transmigration strongly. The expression of JAMs, especially JAM-A, is strongly modulated during the rewarming process. Endothelial protection prior to warm reperfusion and mild hypothermic conditions reducing the difference between hypothermia and rewarming temperatures should be considered.

Therapeutic perioperative hypothermia during cardiopulmonary bypass (CPB) is standard of care in cardiac- and aortic-surgery since the first practical use in 1958¹. Reduction of temperature lowers endogenous metabolic rates and tissue oxygen consumption, which enables prolonged operations accompanied with a high risk of ischemia. However, cardiac surgical procedures supported by CPB generate a systemic inflammatory response syndrome (SIRS), which is associated with acute complications and relevant organ dysfunction and even organ failure². The activation of immune response during CPB is a complex process with many triggers. The exposure of circulating blood cells to the surface of an extracorporeal circuit with activation of the complement system³ leads to increased cytokine production⁴, such as interleukins (ILs), necrosis factors and bradykinin. Patients undergoing CPB may also show an endotoxemia⁵ with a strongly correlated risk of post-operative infections⁶. Different strategies have been tested to reduce this systemic inflammation, e.g. leukocyte filtration during CPB⁷ or prophylactic application of corticosteroids⁸ with limited success. With the application of therapeutic hypothermia during CPB, patients are not only exposed to the extracorporeal circuit, with the previous mentioned inflammatory response, but also to different temperature expositions during surgery. These temperature conditions in cardiac- and aortic surgery range from mild and moderate hypothermia (28–34 °C), to deep hypothermia conditions (16–18 °C).

¹Department of Thoracic and Cardiovascular Surgery, University Hospital Frankfurt, Goethe University, Frankfurt/Main, Germany. ²Department of Anaesthesiology, University Hospital Frankfurt, Goethe University, Frankfurt/Main, Germany. ³Department of Cardiology, University Hospital Frankfurt, Goethe University, Frankfurt/Main, Germany. Correspondence and requests for materials should be addressed to A.B.-F. (email: Andres.Beiras@kgu.de)

This range of temperatures should protect not only for ischemic events, but also decrease the systemic inflammatory response. Experimental studies investigated the influence of hypothermia on the production of chemokines and observed a decreasing production of IL-8^{9,10} and MCP-1⁹. Looking on the synthesis of pro-inflammatory cytokines, many studies observed a reduction of pro-inflammatory cytokines, e.g. TNF- α , IL-1 β ¹¹, IL-6¹² and IL-2¹³, whereas anti-inflammatory cytokines like IL-10^{11–13} increase under hypothermic conditions. The production of adhesion molecules is modified during hypothermia and results in a decreased expression of ICAM-1¹⁴, E-Selectin¹⁵, VCAM and ELAM¹⁶. Further clinical and experimental observations have shown that hypothermia is able to modulate the immune response towards an immunosuppressive effect. In contrast, a clinical analysis showed that patients undergoing post-hypothermic rewarming suffer from a higher risk of fever¹⁷, which could be associated with an over-activation of immune response. Experimental studies have demonstrated that post-hypothermic rewarming increases the production of pro-inflammatory cytokines like IL-6¹⁸ or IL-8¹⁹ and leads to an up-regulation of adhesion molecules, e.g. ICAM-1, E-Selectin and VCAM-1²⁰.

Therefore, a paradox effect upon inflammatory conditions may be considered. While hypothermia has a protective effect upon inflammation and endothelial modulation, post-hypothermic rewarming could move the immune response towards an activating effect. This pro-inflammatory effect is clinically relevant, as it triggers endothelial leakage and post-ischemic tissue infiltration through transendothelial migration (TEM) of leukocytes, which may result in disadvantages for patients subject to deep hypothermia and posterior rewarming. Endothelial leakage is mediated, among others, by Junctional-adhesion-molecules (JAM). These molecules are mainly located at endothelial or epithelial intercellular junctions and are associated with tight-junctions²¹, being partly responsible for modulation of TEM^{21,22}.

With an *in-vitro* model we simulated the previous mentioned acute systemic inflammatory response syndrome during CPB and investigated how temperature changes during cell activation and transendothelial migration affect leukocyte recruitment through endothelial cells. Temperature modulation was performed during cell activation, transmigration or during both periods. Furthermore we investigated the expression of JAM-A and JAM-B on endothelial cells after post-hypothermic rewarming to clarify their role in this process.

Results

Transmigration assay. *Hypothermic transmigration reduces transendothelial migration only partially.* In the first study-group, temperature modulation was performed as described in Fig. 1A. Results are summarized as transmigration-index (tmx) in Fig. 2A–C. Transmigration of activated PBL through non-activated HuMEC-1 under hypothermic conditions shows no significant changes with falling temperatures, whereas significant changes are noticeable compared to the control-group at 30°C ($p = <0.0001$) and 18°C ($p = 0.0207$). Hypothermic conditions during the transmigration of non-activated PBL through activated HuMEC-1 show significant changes in TEM with falling temperatures. Significant reductions of TEM are visible at 18°C compared to 37°C ($p = <0.0001$) and at 18°C compared to 30°C ($p = <0.0001$). Compared to the control-group, significant changes are observable at 37°C ($p = <0.0001$), 30°C ($p = <0.0001$) and 18°C ($p = 0.0049$). The application of hypothermia during transmigration of activated PBL through activated HuMEC-1 causes no significant changes in TEM. Significant changes are noticeable compared to the control-group at 37°C ($p = 0.0002$), 30°C ($p = <0.0001$) and 18°C ($p = 0.0055$).

Hypothermic treatment and migration reduces transendothelial migration significantly. In the second study-group, temperature modulation was performed as described in Fig. 1B. Results are summarized as transmigration-index (tmx) in Fig. 2D–F. Application of hypothermia during treatment and transmigration of activated PBL through non-activated HuMEC-1 causes no significant change in TEM within the temperature range, whereas significant changes are noticeable compared to the control-group at 30°C ($p = 0.0001$). Temperature modulation during cell treatment and transmigration of non-activated PBL through activated HuMEC-1 leads to a significant reduction of TEM at 18°C compared to 37°C ($p = 0.0003$) and at 18°C compared to 30°C ($p = 0.0003$). Compared to the control-group, significant changes are visible at 37°C ($p = <0.0001$) and 30°C ($p = <0.0001$). Transmigration of activated PBL through activated HuMEC-1 is affected under mild and severe hypothermic conditions, with significant diminished TEM at 18°C compared to 37°C ($p = 0.0003$) and at 18°C compared to 30°C ($p = 0.0472$). Significant changes are noticeable compared to the control-group at 37°C ($p = 0.0002$), 30°C ($p = <0.0001$) and 18°C ($p = 0.0241$).

Post-hypothermic rewarming strongly increases transendothelial migration. In the third study-group, temperature modulation was performed as described in Fig. 1C. Results are summarized as transmigration-index (tmx) in Fig. 2G–I. TEM of activated PBL through non-activated HuMEC-1 is not significantly affected after rewarming. Temperature modulation during treatment and following rewarming during transmigration of non-activated PBL through activated HuMEC-1 leads to a highly enhanced TEM with significant changes at 18°C compared to 37°C ($p = 0.0003$) and at 18°C compared to 30°C ($p = 0.0035$). Significant changes are noticeable compared to the control-group at 37°C ($p = <0.0001$), 30°C ($p = 0.0018$) and 18°C ($p = <0.0001$). Simultaneous activation of PBL and HuMEC-1 leads to the same enhanced TEM in accordance to a growing temperature difference. Significant increases of TEM can be observed at 18°C compared to 37°C ($p = <0.0001$) and 18°C compared to 30°C ($p = 0.0007$). Compared to the control-group, significant changes are visible at 37°C ($p = 0.0002$), 30°C ($p = 0.0015$) and 18°C ($p = <0.0001$).

Surface-expression-assay. *JAM-A-surface-expression is significantly modulated during post-hypothermic rewarming.* For surface-expression-assay, temperature modulation was performed as described in Fig. 1C and results of endothelial JAM-A-surface-expression are summarized as fluorescence-intensity (FI) in Fig. 3A–C.




Figure 1. Scheme of experimental temperature modulation. Cells were kept at 37°C initially. (A) HuMEC-1 and PBL were treated for 60 min at 37°C. TEM was performed for 120 min at 37°C, 30°C and 18°C. (B) HuMEC-1 and PBL were treated for 60 min at 37°C, 30°C and 18°C. TEM was performed for 120 min under the same temperature conditions as HuMEC-1 and PBL were treated before. (C) HuMEC-1 and PBL were treated for 60 min at 37°C, 30°C or 18°C with subsequent rewarming to 37°C during TEM for 120 min.




Figure 2. Influence of temperature modulation on transendothelial migration. (A–C) Influence on TEM when temperature modulation is only performed during transmigration in accordance to Fig. 1A. (A) Activated PBL show a significantly increased TEM with falling temperatures (37°C n = 11, 30°C n = 11, 18°C n = 18). (B) TEM of non-activated PBL through activated HuMEC-1 is significantly reduced with deeper temperatures (37°C n = 11, 30°C n = 12, 18°C n = 17), whereas (C) TEM of activated PBL through activated HuMEC-1 is not influenced with falling temperatures (37°C n = 12, 30°C n = 11, 18°C n = 17). (D–F) Influence on TEM when temperature modulation is applied during treatment and transmigration in accordance to Fig. 1B. (D) TEM of activated PBL through non-activated HuMEC-1 is not significantly modulated during cooling (37°C n = 11, 30°C n = 11, 18°C n = 12), whereas (E) TEM of non-activated PBL through activated HuMEC-1 is significantly reduced with falling temperatures (37°C n = 11, 30°C n = 12, 18°C n = 12). (F) Cooling shows a significant protective function with reduced TEM of activated PBL through activated HuMEC-1 (37°C n = 12, 30°C n = 11, 18°C n = 12). (G–I) Influence on TEM when temperature is modulated during treatment with following rewarming during transmigration in accordance to Fig. 1C. (G) Rewarming shows no influence on TEM of activated PBL through non-activated HuMEC-1 (37°C n = 11, 30°C n = 11, 18°C n = 11). (H) TEM of non-activated PBL through activated HuMEC-1 is significantly enhanced with falling temperatures (37°C n = 11, 30°C n = 12, 18°C n = 11). (I) Rewarming significantly enhances TEM of activated PBL through activated HuMEC-1 (37°C n = 12, 30°C n = 12, 18°C n = 11). *p < 0.05 vs. control, *p < 0.05.

Post-hypothermic rewarming of activated PBL in a co-culture with non-activated HuMEC-1 leads to significant decreased JAM-A-surface-expression at 18°C compared to 30°C (p = 0.0474) and a significant change at 18°C compared to the control-group (p = 0.0224). Activated HuMEC-1 in a co-culture with non-activated PBL show a significant decreased JAM-A-surface-expression at 30°C compared to 37°C (p = 0.0431) and at 18°C compared to 37°C (p = 0.0027). The control-group shows increased JAM-A-surface-expression compared to 30°C (p = <0.0001) and to 18°C (p = 0.0001). Co-culture of simultaneously activated HuMEC-1 and PBL leads to significant decreased JAM-A-surface-expression at 18°C compared to 37°C (p = 0.0039). Compared to the control-group, significant changes are visible at 37°C (p = 0.0217), 30°C (p = 0.0198) and 18°C (p = 0.0001).

JAM-B-surface-expression is not significantly modulated during post-hypothermic rewarming. For surface-expression assay, temperature modulation was performed as described in Fig. 1C and results of endothelial JAM-B-surface-expression are summarized as fluorescence-intensity (FI) in Fig. 3. JAM-B-surface-expression is not significantly modulated within the applied temperature-range and significant changes are only noticeable in comparison to the control-group. Activated PBL in a co-culture with non-activated HuMEC-1 leads to significant decreased JAM-B-surface-expression at 18°C compared to the control-group (p = 0.0012). Activated HuMEC-1 in a co-culture with non-activated PBL show a significant decreased JAM-B-surface-expression compared to control at 37°C (p = 0.0088), 30°C (p = 0.0082) and 18°C (p = 0.0108). Co-culture of simultaneously activated HuMEC-1 and PBL leads to significant decreased JAM-B-surface-expression compared to control at 37°C (p = 0.0009), 30°C (p = 0.0002) and 18°C (p = 0.0003).




Figure 3. Modulation of JAM-A- and JAM-B-surface-expression during rewarming. (A) Co-culture of activated PBL and non-activated HuMEC-1 significantly decreases expression of JAM-A (37°C n = 7, 30°C n = 6, 18°C n = 6) with falling temperatures. JAM-B (37°C n = 8, 30°C n = 6, 18°C n = 5) shows no significant change during temperature modulation. (B) Rewarming in a co-culture of non-activated PBL and activated HuMEC-1 shows significantly decreased expression of JAM-A (37°C n = 8, 30°C n = 6, 18°C n = 6) with growing temperature differences, whereas JAM-B is not affected (37°C n = 8, 30°C n = 6, 18°C n = 6). (C) Co-culture with activated PBL and activated HuMEC-1 leads to a significant down-regulation of JAM-A expression (37°C n = 8, 30°C n = 6, 18°C n = 6) with falling temperatures, whereas JAM-B expression is not affected (37°C n = 8, 30°C n = 6, 18°C n = 6). #p < 0.05 vs. control, *p < 0.05.

Total-expression-assay. *JAM-total-protein-expression is not significantly modulated during rewarming.* For total-expression-assay, temperature modulation was performed as described in Fig. 1C and results of endothelial JAM-total-expression are summarized as relative amount of protein in Fig. 4. Neither JAM-A-total-expression nor JAM-B-total-expression is significantly modulated during post-hypothermic rewarming. As well no significant changes are noticeable compared to the control-group.

Analysis of IL-6 and IL-10. *Rewarming leads to significant modulation of IL-6 expression.* For analysis of IL-6 temperature modulation was performed as described in Fig. 1C and results are summarized in Fig. 5A–C. Activation of PBL in co-culture with non-activated HuMEC-1 leads to a significant up-regulation of IL-6 at 30°C compared to 37°C (p = 0.0431). Significant changes are visible at 30°C (p = 0.002) and 18°C (p = 0.0483) compared to control. Rewarming in a co-culture of non-activated PBL and activated HuMEC-1 causes a significant increase of IL-6 at 30°C compared to 37°C (p = <0.0001) and a significant decrease of IL-6 at 18°C compared to 30°C (p = 0.0008). Compared to the control-group significant changes are visible at 37°C (p = 0.0003), 30°C (p = <0.0001) and 18°C (p = <0.0001). Co-culture of simultaneously activated HuMEC-1 and PBL leads to a significantly enhanced IL-6 expression at 30°C (p = <0.0001) and at 18°C (p = 0.0128) compared to 37°C. Temperature modulation from 30°C to 18°C causes a significant decrease of IL-6 (p = 0.0031). Compared to the control-group, significant changes are visible at 37°C (p = 0.0005), 30°C (p = <0.0001) and 18°C (p = <0.0001).

Post-hypothermic rewarming causes no significant changes of IL-10. For analysis of IL-10 temperature modulation was performed as described in Fig. 1C and results are summarized in Fig. 5D–F. Rewarming causes no significant changes of IL-10 with falling temperature conditions.

Post-hypothermic rewarming significantly raises the IL-6/IL-10 ratio. For analysis of IL-6/IL-10 ratio temperature modulation was performed as described in Fig. 1C and results are summarized in Fig. 5G–I. Activation of PBL in co-culture with non-activated HuMEC-1 leads to a significant change of IL-6/IL-10 ratio at 37°C compared to control (p = 0.0002). Post-hypothermic rewarming in a co-culture of non-activated PBL and activated HuMEC-1 causes a significant increase of IL-6/IL-10 ratio compared to the control-group at 37°C (p = <0.0001), 30°C (p = 0.0249) and 18°C (p = 0.0327). Co-culture of simultaneously activated HuMEC-1 and PBL leads to a significantly enhanced IL-6/IL-10 ratio at 18°C compared to 37°C (p = 0.0250). Compared to the control-group, significant changes are visible at 37°C (p = 0.0028), 30°C (p = 0.0023) and 18°C (p = 0.0199).

Discussion

Several studies have analyzed the single molecular effects of hypothermia or post-hypothermic rewarming during inflammatory processes, but the final common effects on leukocyte recruitment remain unclear. The current study investigates TEM of PBL through an endothelial monolayer in an *in-vitro* inflammatory model, using typical clinical temperature conditions. Additionally, we investigated the expression of JAM-A and JAM-B after post-hypothermic rewarming. Cooling of normothermic cells during cell activation and transmigration or the




Figure 4. Modulation of JAM-A- and JAM-B-total-expression during rewarming. (A) Co-culture of activated PBL and non-activated HuMEC-1 shows no significant changes on expression of JAM-A (37°C n = 2, 30°C n = 2, 18°C n = 2) and JAM-B (37°C n = 2, 30°C n = 2, 18°C n = 2) with falling temperatures. (B) Rewarming in a co-culture of non-activated PBL and activated HuMEC-1 shows no significant influence on expression of JAM-A (37°C n = 2, 30°C n = 3, 18°C n = 2) and JAM-B (37°C n = 2, 30°C n = 2, 18°C n = 2). (C) Co-culture with activated PBL and activated HuMEC-1 leads to no significant regulation of JAM-A (37°C n = 2, 30°C n = 2, 18°C n = 2) and JAM-B (37°C n = 2, 30°C n = 2, 18°C n = 2) expression. *p < 0.05 vs. control, *p < 0.05. Representative western-blot are presented for every experimental group and illustrate the JAM-A, JAM-B and β-actin-expression. Full-length blots are presented in Supplementary Figure 1.

single application of hypothermia during transmigration reduces TEM and promotes an anti-inflammatory effect. Hypothermic cell activation and subsequent rewarming during transmigration leads to enhanced TEM of PBL through an endothelial monolayer and evokes a pro-inflammatory effect. Post-hypothermic rewarming modulates endothelial JAM-expression. Especially the down regulation of JAM-A expression could be associated with the enhanced TEM of PBL during the rewarming process.

It is well established that inflammatory processes induce several intracellular cascades and signaling pathways. One of the acting pathways during inflammation is the activation of NF-κB, which is associated with the expression of cytokines, chemokines or adhesion molecules^{23,24}. Several studies have shown that NF-κB is modulated during hypothermic conditions^{9,25,26}. Furthermore, some of them show that hypothermia leads to an activation of NF-κB^{9,25}, whereas other studies show an inhibition of NF-κB²⁶. The same variable effects of hypothermia on cellular modulation are controversial discussed in studies which analyzed cytokine expression^{9,26,27}. Furthermore, the cells used in our study were stimulated concomitantly to hypothermia with LPS to mimic an inflammatory process, as would happen in a real clinical situation. Recently, Thiel *et al.*²⁸ used a similar strategy to reproduce a pro-inflammatory status in a murine model of lupus nephritis. Hypothermia is able to modulate cytokine expression, which could lead to an anti-inflammatory⁹ or a pro-inflammatory effect²⁶. Adhesion molecules play an essential role in initiating transendothelial migration. Typical adhesion molecules such as ICAM-1¹⁴, E-Selectin¹⁵, VCAM or ELAM¹⁶ show a decreased expression during hypothermic conditions. Down regulation of adhesion molecules could be associated with decreased TEM during hypothermic conditions. Seo *et al.*²⁷ showed that long-term hypothermic conditions over 6 h are able to prevent microglial migration. Our study confirms this observation and illustrates the final common pathway of short-term hypothermia, which leads to decreased TEM of PBL through an endothelial monolayer. Short-term hypothermia applied during transmigration or during activation and transmigration may simultaneously lead to reduced leukocyte infiltration. This process depends on the depth of applied hypothermic conditions. Deeper temperature conditions lead to a higher inhibitory effect on TEM.

Clinical application of hypothermia requires as well rewarming processes to regain normothermic conditions. Several studies show that post-hypothermic rewarming modulates inflammatory responses. Not only hypothermic conditions are able to modulate NF-κB, but also post-hypothermic rewarming is able to modulate this essential inflammatory pathway^{20,29,30}. Diestel *et al.*²⁹ describe a diminished degradation of the inhibitor of regulation of NF-κB (IκB-α) in the first 30 min during rewarming, followed of a delayed and higher degradation of IκB-α after 50 min during the rewarming process compared to normothermic conditions. Other studies describe an




Figure 5. IL-6 and IL-10 during rewarming. (A–C) Influence of rewarming on IL-6. (A) Co-culture of activated PBL and non-activated HuMEC-1 shows significant influence on IL-6 (37°C n = 5, 30°C n = 5, 18°C n = 3). (B) Rewarming in a co-culture of non-activated PBL and activated HuMEC-1 leads to a significant modulation of IL-6 with falling temperatures (37°C n = 5, 30°C n = 5, 18°C n = 6). (C) This effect is also visible in a co-culture of activated PBL and activated HuMEC-1 (37°C n = 6, 30°C n = 5, 18°C n = 5). (D–F) Influence of rewarming on IL-10. Temperature modulation shows no significant influence on IL-10 expression. (D) Co-culture of activated PBL and non-activated HuMEC-1 (37°C n = 5, 30°C n = 5, 18°C n = 3). (E) Co-culture of non-activated PBL and activated HuMEC-1 (37°C n = 5, 30°C n = 5, 18°C n = 6). (F) Co-culture of activated PBL and activated HuMEC-1 (37°C n = 6, 30°C n = 5, 18°C n = 5). (G–I) Influence of rewarming on IL-6/IL-10 ratio. (G) Co-culture of activated PBL and non-activated HMEC-1 shows no significant changes in IL-6/IL-10 ratio (37°C n = 5, 30°C n = 5, 18°C n = 3). (H) Co-culture of non-activated PBL and activated HuMEC-1 show no significant influence on IL-6/IL-10 ratio with falling temperatures (37°C n = 5, 30°C n = 5, 18°C n = 6). (I) Co-culture with activated PBL and activated HuMEC-1 leads to significant up regulation of IL-6/IL-10 ratio (37°C n = 6, 30°C n = 5, 18°C n = 5). *p < 0.05 vs. control, #p < 0.05.

augmentation of NF- κ B during rewarming²⁰ and show that NF- κ B is up regulated on the first postoperative day after deep hypothermic conditions followed by fast rewarming compared to slow rewarming³⁰. The effect of post-hypothermic rewarming on cytokines is also controversially discussed, which could lead to a pro-inflammatory cytokine expression^{19,31} or to an anti-inflammatory cytokine expression^{32,33}. Awad *et al.*²⁰ show that posthypothermic rewarming causes an up regulation of typical adhesion molecules such as ICAM-1, E-Selectin and VCAM-1. Up regulation of adhesion molecules could be associated with higher transmigration potency during posthypothermic rewarming. Our study shows an increased TEM of PBL through an endothelial monolayer during post-hypothermic rewarming. This enhancement of TEM seems to be temperature dependent. As the temperature differences between initial-point, hypothermic conditions and rewarming grows, the leukocyte recruitment gets more essentially influenced.

We hypothesized that this increasing influence of rewarming is caused by regulation of JAMs. JAMs are as well modulated during inflammatory immune response and could be expressed through the activation of the typical inflammatory NF- κ B pathway. Azari *et al.*³⁴ and Ohkuni *et al.*³⁵ show that inhibition of NF- κ B resulted in a decreased JAM-A expression. In another study, JAM-A expression was decreased during induction of a pro-inflammatory response through an TLR3 ligand, whereas TNF- α and IL-8 expression increases. Both effects are mediated through NF- κ B³⁵. Comparing this report to the previous described changes, post-hypothermic

rewarming could lead to a pro-inflammatory response with enhanced cytokine production and reduced JAM-A expression. Laukoetter *et al.*³⁶ observed that JAM-A deficient animals suffer from a breakdown of intestinal barrier function, associated with higher leukocyte recruitment and increased permeability. Our data suggest that decreased JAM-A expression leads to higher leukocyte recruitment and show that posthypothermic rewarming leads to a decreased JAM-A expression in a temperature dependent manner. JAM-B expression is less modulated during temperature changes and shows no correlation with TEM. Temperature shift during rewarming after short-term hypothermia causes a breakdown of endothelial barrier function and leads to higher TEM of PBL.

The *in vitro* re-enactment of clinical practice has shown a significant influence of temperature modulation on TEM. Our data show that TEM of PBL through endothelial cells is not only modulated by cell-activation itself - the activation temperature and the rewarming processes are essential. Prolonged hypothermia without rewarming is able to inhibit TEM, whereas hypothermia with post-hypothermic rewarming enhances TEM strongly, mainly depending on the difference between hypothermic temperature and rewarming temperature, with increased inflammatory activation in higher temperature differences. The IL-6/IL-10 ratio clearly shows an increased pro-inflammatory status depending on the delta between the target hypothermia temperature and 37 °C. The modifications in JAMs, especially JAM-A, provide a potential mechanistic explanation for the observed changes in TEM during the rewarming process. Endothelial protection prior to warm reperfusion and mild hypothermic conditions should be considered in clinical practice.

Methods

Ethics Statement. This study was approved by the ethics committee of the University of Frankfurt, Germany (GN.:227/13). All blood samples were obtained under informed consent and according to the declaration of Helsinki. All participants provided their written informed consent to participate in this study.

Cell culture. Human microvascular endothelial cells-1 (HuMEC-1) were kindly provided by Dr. V. Mirakaj (University Tübingen, Department of Anesthesiology and Intensive Care Medicine) and cultured in MCDB-131 (Life Technologies, Darmstadt, Germany) supplemented with 10% fetal calf serum (Gibco, Karlsruhe, Germany), 1% glutamine (Gibco, Karlsruhe, Germany), 1% pen/strep solution (Sigma Chemical Co. St. Louis, USA), 10 ng/ml Epidermal Growth Factor (Sigma Chemical Co. St. Louis, USA) and 1 µg/ml hydrocortisone (Sigma Chemical Co. St. Louis, USA) at 37 °C and 5% CO₂ atmosphere.

Isolation of peripheral blood leukocytes (PBL). Blood was drawn into EDTA tubes from healthy male and female volunteers (n = 5). For density gradient centrifugation, the fresh blood samples were underlaid with 5 ml Polymorphprep[®] (Axis-Shield, Oslo, Norway) and centrifuged at 450 g for 35 min. The layers of peripheral blood leukocytes (PBL), containing peripheral blood mononuclear cells (PBMC) and polymorph nuclear cells (PMNC)³⁷, were aspirated and washed with PBS (Gibco, Karlsruhe, Germany). PBL were suspended in 37 °C pre-warmed RPMI 1640 (Life Technologies, Darmstadt, Germany).

Influence of temperature modulation on transendothelial migration. *Cultivation of an endothelial monolayer on cell culture inserts.* Transmigration assays were performed with the use of 3-µm pore size FluoroBlok inserts (BD, Heidelberg, Germany) in triplicate. BrdU-labeled non-proliferating HuMEC-1 were placed on fibronectin-coated inserts and were allowed to form an endothelial monolayer for approximately 24 h at 37 °C, 5% CO₂.

Treatment of cultivated HuMEC-1. As previously described³⁷, HuMEC-1 were partially activated with 100 µg/ml LPS (Sigma Chemical Co. St. Louis, USA) in MCDB-131 (Life Technologies, Darmstadt, Germany), whereas the other fraction of HuMEC-1 remained non-activated. The treatment of every study group was performed for 60 min under different temperature conditions (Fig. 1A–C). After treatment, medium was removed and transmigration-assay was started.

Treatment of PBL. PBL were activated with 10 µg/ml LPS in RPMI 1640 with 10% fetal bovine serum (Gibco, Karlsruhe, Germany), whereas the other fraction of PBL remained non-activated. Treatment of every study group was performed for 60 min under different temperature conditions (Fig. 1A–C). Cells were washed after incubation with PBS.

Study groups. Following subassembly-groups were used within the main-groups: A) non-activated PBL and non-activated HuMEC-1 (control); B) activated PBL and non-activated HuMEC-1; C) non-activated PBL and activated HuMEC-1; D) activated PBL and activated HuMEC-1. Temperature modulation (Fig. 1) was performed to every subassembly group in the same manner, whether if they were activated or remained non-activated. In the first main-group (Fig. 1A), temperature modulation (37 °C, 30 °C and 18 °C) was only performed during transmigration. In the second main-group (Fig. 1B), temperature modulation was performed during treatment and during transmigration. In the third main-group (Fig. 1C), temperature modulation was only performed during treatment following normothermic transmigration, which mimics the posthypothermic rewarming process. The control group consisted in all cases of non-activated PBL and non-activated HuMEC-1 undergoing the same temperature modulation strategies as the study groups.

Transmigration-assay. PBL stained with Calcein-AM (ABD-Bioquest, Biomol, Hamburg, Germany) were suspended in assay-medium (RPMI-1640 + 0.5% BSA) and applied on top of the endothelial monolayer. The transmigration assay was performed for 120 min under normothermic and hypothermic conditions (37 °C, 30 °C, 18 °C). Transmigration conditions of all three main-groups are shown in Fig. 1A–C. The fluorescence signal

of transmigrated PBL to the bottom surface of the membrane insert was measured with a microplate reader (Berthold Technologies, Bad Wildbad, Germany) at 485/535 nm. Transmigrated cells are expressed as percent of control (transmigration index, tmx).

JAM-A- and JAM-B-surface-expression during rewarming. *Cultivation of HuMEC-1.* Surface-expression-assay was performed with the use of black 96-Well-Plates with clear-bottom (BD, Heidelberg, Germany). BrdU-labeled non-proliferating HuMEC-1 were placed on fibronectin-coated wells and were allowed to form an endothelial monolayer for approximately 24 h at 37 °C, 5% CO.

Treatment of cells and Co-culture conditions. HuMEC-1 and PBL were treated as previously described. Normothermic cells were treated at different temperature conditions (37 °C, 30 °C, 18 °C) for 60 min (Fig. 1C). Activated or non-activated HuMEC-1 and activated or non-activated PBL were incubated together in a co-culture with transmigration-medium (RPMI-1640 + 0.5% BSA). Co-culture was performed at 37 °C for 120 min to simulate the rewarming process (Fig. 1C). After incubation, PBL were removed and HuMEC-1 were washed with PBS.

Immunofluorescence analysis. HuMEC-1 were fixed with 4% PFA (AppliChem, Darmstadt, Germany) at RT and were blocked with PBS supplemented with 10% donkey serum (Jackson ImmunoResearch Europe Ltd., Suffolk, UK). HuMEC-1 were primarily stained with rabbit anti-JAM-A-IgG [1:200] (Santa Cruz Biotechnology, Heidelberg, Germany) and goat anti-JAM-B-IgG [1:100] (Santa Cruz Biotechnology, Heidelberg, Germany). Secondary antibody incubation was performed with donkey anti-rabbit-IgG-PE [1:100] (Santa Cruz Biotechnology, Heidelberg, Germany) or donkey anti-goat-IgG-FITC [1:100] (Santa Cruz Biotechnology, Heidelberg, Germany) in PBS containing 1% donkey serum. JAM-surface-protein expression was measured with a microplate reader (Berthold Technologies, Bad Wildbad, Germany) at 485/535 nm and 530/590 nm. Fluorescence intensity (FI) was calculated in relation to control.

JAM-A- and JAM-B-total-expression during rewarming. *Cultivation of HuMEC-1.* Total-expression-assay was performed with the use of T25 flasks (Sarstedt, Nümbrecht, Germany) and HuMEC-1 were allowed to form an endothelial monolayer for approximately 24 h at 37 °C, 5% CO.

Treatment of cells and Co-culture conditions. HuMEC-1 and PBL were treated in the same manner than previously described in immunofluorescence-analysis of surface-expression.

Protein isolation and Western Blot Analysis. Cells were harvested with cell scrapers and proteins were isolated from cell pellet under constant agitation with cell lysis buffer for 10 min at 4 °C. Cell lysates were applied to a 4–20% Mini-Protein TGX gel (BioRad, München, Germany) and electrophoresis was applied at 200 V for 20 min. Proteins were transferred to PVDF-membranes (BioRad, München, Germany) at 2,5 mA for 5 min. Membranes were blocked for 60 min with blocking-buffer containing 5% serum (Jackson ImmunoResearch Europe Ltd., Suffolk, UK) of the species where the second antibody was raised in. Primary antibody incubation of rabbit anti-JAM-A-IgG [1:200] (Santa Cruz Biotechnology, Heidelberg, Germany) and goat anti-JAM-B-IgG [1:200] (Santa Cruz Biotechnology, Heidelberg, Germany) was performed over night at 4 °C. Secondary antibody incubation of donkey anti-goat-IgG-HRP [1:5000] (Santa Cruz Biotechnology, Heidelberg, Germany) and goat anti-rabbit-IgG-HRP [1:5000] (Merck, Darmstadt, Germany) was performed for 60 min at RT. Membranes were incubated with HRP substrate solution (Merck, Darmstadt, Germany) and proteins were visualized with Fusion FX7 system (Pierce, Erlangen, Germany). β -Actin was used as an internal control for protein loading. The visualized protein bands were analyzed using ImageJ. The relative amount of protein was calculated in relation to β -actin expression and to control-group. All gels were ran under the same experimental conditions. Cropped western-blots are presented in Fig. 4 and full-length blots are presented in supplementary Figure 1.

Analysis of IL-6 and IL-10 during rewarming. *Collection of Co-culture supernatants and ELISA procedure.* Co-culture supernatants from western blot analysis were collected before cells were harvested. Supernatants were centrifuged to remove the PBL cell pellet. IL-6 (R&D, Abingdon, Great Britain) and IL-10 (R&D, Abingdon, Great Britain) ELISA were performed in accordance to the manufacturers guide. IL-6 and IL-10 are expressed as percent of control (IL-6 Index, IL-10 Index) or were evaluated in relation to each other to calculate an IL-6/IL-10-Ratio.

Statistical analysis. All data represent mean \pm standard error of the mean (SEM). Statistical analysis was performed with Prism 6 software (Graph Pad, La Jolla, USA) using Analysis of Variance (ANOVA) and *post-hoc* Tukey's-multiple-comparisons-test. Differences with $p < 0.05$ were considered statistically significant.

References

1. Sealy, W. C., Young, G. Jr., Brown, I. W. Jr., Smith, W. W. & Lesage, A. M. Profound hypothermia combined with extracorporeal circulation for open heart surgery. *Surgery* **48**, 432–438 (1960).
2. Asimakopoulos, G. Systemic inflammation and cardiac surgery: an update. *Perfusion* **16**, 353–360 (2001).
3. Chenoweth, D. E. *et al.* Complement activation during cardiopulmonary bypass: evidence for generation of C3a and C5a anaphylatoxins. *The New England journal of medicine* **304**, 497–503, doi: 10.1056/NEJM198102263040901 (1981).
4. Takabayashi, T. *et al.* A new biologic role for C3a and C3a desArg: regulation of TNF-alpha and IL-1 beta synthesis. *Journal of immunology* **156**, 3455–3460 (1996).
5. Andersen, L. W., Landow, L., Baek, L., Jansen, E. & Baker, S. Association between gastric intramucosal pH and splanchnic endotoxin, antibody to endotoxin, and tumor necrosis factor-alpha concentrations in patients undergoing cardiopulmonary bypass. *Critical care medicine* **21**, 210–217 (1993).

6. Klein, D. J., Briet, F., Nisenbaum, R., Romaschin, A. D. & Mazer, C. D. Endotoxemia related to cardiopulmonary bypass is associated with increased risk of infection after cardiac surgery: a prospective observational study. *Critical care* **15**, R69, doi: 10.1186/cc10051 (2011).
7. de Amorim, C. G. *et al.* Leukocyte Depletion During CPB: Effects on Inflammation and Lung Function. *Inflammation*, doi: 10.1007/s10753-013-9730-z (2013).
8. Dieleman, J. M. *et al.* Prophylactic corticosteroids for cardiopulmonary bypass in adults. *The Cochrane database of systematic reviews*, CD005566, doi: 10.1002/14651858.CD005566.pub3 (2011).
9. Diestel, A., Roessler, J., Berger, F. & Schmitt, K. R. Hypothermia downregulates inflammation but enhances IL-6 secretion by stimulated endothelial cells. *Cryobiology* **57**, 216–222, doi: 10.1016/j.cryobiol.2008.08.005 (2008).
10. Sutcliffe, I. T., Smith, H. A., Stanimirovic, D. & Hutchison, J. S. Effects of moderate hypothermia on IL-1 beta-induced leukocyte rolling and adhesion in pial microcirculation of mice and on proinflammatory gene expression in human cerebral endothelial cells. *Journal of cerebral blood flow and metabolism: official journal of the International Society of Cerebral Blood Flow and Metabolism* **21**, 1310–1319, doi: 10.1097/00004647-200111000-00007 (2001).
11. Lim, C. M. *et al.* Hypothermia protects against endotoxin-induced acute lung injury in rats. *Intensive care medicine* **29**, 453–459, doi: 10.1007/s00134-002-1529-6 (2003).
12. Fujimoto, K. *et al.* Early induction of moderate hypothermia suppresses systemic inflammatory cytokines and intracellular adhesion molecule-1 in rats with caerulein-induced pancreatitis and endotoxemia. *Pancreas* **37**, 176–181, doi: 10.1097/MPA.0b013e318162cb26 (2008).
13. Lee, S. L., Battistella, F. D. & Go, K. Hypothermia induces T-cell production of immunosuppressive cytokines. *The Journal of surgical research* **100**, 150–153, doi: 10.1006/jsre.2001.6230 (2001).
14. Meybohm, P. *et al.* Mild hypothermia alone or in combination with anesthetic post-conditioning reduces expression of inflammatory cytokines in the cerebral cortex of pigs after cardiopulmonary resuscitation. *Critical care* **14**, R21, doi: 10.1186/cc8879 (2010).
15. Johnson, M., Haddix, T., Pohlman, T. & Verrier, E. D. Hypothermia reversibly inhibits endothelial cell expression of E-selectin and tissue factor. *Journal of cardiac surgery* **10**, 428–435 (1995).
16. Hanusch, C. *et al.* Hypothermic preservation of lung allograft inhibits cytokine-induced chemoattractant-1, endothelial leucocyte adhesion molecule, vascular cell adhesion molecule-1 and intracellular adhesion molecule-1 expression. *Clinical and experimental immunology* **149**, 364–371, doi: 10.1111/j.1365-2249.2007.03417.x (2007).
17. Cocchi, M. N. *et al.* Fever After Rewarming: Incidence of Pyrexia in Postcardiac Arrest Patients Who Have Undergone Mild Therapeutic Hypothermia. *Journal of intensive care medicine*, doi: 10.1177/0885066613491932 (2013).
18. Bisschops, L. L., Hoedemaekers, C. W., Mollnes, T. E. & van der Hoeven, J. G. Rewarming after hypothermia after cardiac arrest shifts the inflammatory balance. *Critical care medicine* **40**, 1136–1142, doi: 10.1097/CCM.0b013e3182377050 (2012).
19. Inoue, K. *et al.* Effects of rewarming on nuclear factor-kappaB and interleukin 8 expression in cold-preserved alveolar epithelial cells. *Transplantation* **76**, 409–415, doi: 10.1097/01.TP.0000076095.51697.5E (2003).
20. Awad, E. M. *et al.* Cold induces reactive oxygen species production and activation of the NF-kappa B response in endothelial cells and inflammation *in vivo*. *Journal of thrombosis and haemostasis: JTH* **11**, 1716–1726, doi: 10.1111/jth.12357 (2013).
21. Martin-Padura, I. *et al.* Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *The Journal of cell biology* **142**, 117–127 (1998).
22. Ostermann, G., Weber, K. S., Zerneck, A., Schroder, A. & Weber, C. JAM-1 is a ligand of the beta(2) integrin LFA-1 involved in transendothelial migration of leukocytes. *Nature immunology* **3**, 151–158, doi: 10.1038/ni755 (2002).
23. Kempe, S., Kestler, H., Lasar, A. & Wirth, T. NF-kappaB controls the global pro-inflammatory response in endothelial cells: evidence for the regulation of a pro-atherogenic program. *Nucleic acids research* **33**, 5308–5319, doi: 10.1093/nar/gki836 (2005).
24. Chen, C. C., Rosenbloom, C. L., Anderson, D. C. & Manning, A. M. Selective inhibition of E-selectin, vascular cell adhesion molecule-1, and intercellular adhesion molecule-1 expression by inhibitors of I kappa B-alpha phosphorylation. *Journal of immunology* **155**, 3538–3545 (1995).
25. Han, H. S., Karabiyikoglu, M., Kelly, S., Sobel, R. A. & Yenari, M. A. Mild hypothermia inhibits nuclear factor-kappaB translocation in experimental stroke. *Journal of cerebral blood flow and metabolism: official journal of the International Society of Cerebral Blood Flow and Metabolism* **23**, 589–598, doi: 10.1097/01.WCB.0000059566.39780.8D (2003).
26. Arai, T. *et al.* Hypothermia augments NF-kappaB activity and the production of IL-12 and IFN-gamma. *Allergology international: official journal of the Japanese Society of Allergology* **57**, 331–338, doi: 10.2332/allergolint.O-08-532 (2008).
27. Seo, J. W. *et al.* Time-dependent effects of hypothermia on microglial activation and migration. *Journal of neuroinflammation* **9**, 164, doi: 10.1186/1742-2094-9-164 (2012).
28. Thiel, A. *et al.* Human embryonic stem cell-derived mesenchymal cells preserve kidney function and extend lifespan in NZB/W F1 mouse model of lupus nephritis. *Scientific reports* **5**, 17685, doi: 10.1038/srep17685 (2015).
29. Diestel, A. *et al.* Mechanisms of hypothermia-induced cell protection mediated by microglial cells *in vitro*. *The European journal of neuroscience* **31**, 779–787, doi: 10.1111/j.1460-9568.2010.07128.x (2010).
30. Gordan, M. L. *et al.* Fast rewarming after deep hypothermic circulatory arrest in rats impairs histologic outcome and increases NFkappaB expression in the brain. *Perfusion* **25**, 349–354, doi: 10.1177/0267659110377946 (2010).
31. Aibiki, M., Maekawa, S., Nishiyama, T., Seki, K. & Yokono, S. Activated cytokine production in patients with accidental hypothermia. *Resuscitation* **41**, 263–268 (1999).
32. Mizushima, Y., Wang, P., Cioffi, W. G., Bland, K. I. & Chaudry, I. H. Restoration of body temperature to normothermia during resuscitation following trauma-hemorrhage improves the depressed cardiovascular and hepatocellular functions. *Archives of surgery* **135**, 175–181 (2000).
33. McInerney, J. J., Breakell, A., Madira, W., Davies, T. G. & Evans, P. A. Accidental hypothermia and active rewarming: the metabolic and inflammatory changes observed above and below 32 degrees C. *Emergency medicine journal: EMJ* **19**, 219–223 (2002).
34. Azari, B. M. *et al.* Transcription and translation of human F11R gene are required for an initial step of atherogenesis induced by inflammatory cytokines. *Journal of translational medicine* **9**, 98, doi: 10.1186/1479-5876-9-98 (2011).
35. Ohkuni, T. *et al.* Poly(I:C) reduces expression of JAM-A and induces secretion of IL-8 and TNF-alpha via distinct NF-kappaB pathways in human nasal epithelial cells. *Toxicology and applied pharmacology* **250**, 29–38, doi: 10.1016/j.taap.2010.09.023 (2011).
36. Laukoetter, M. G. *et al.* JAM-A regulates permeability and inflammation in the intestine *in vivo*. *The Journal of experimental medicine* **204**, 3067–3076, doi: 10.1084/jem.20071416 (2007).
37. Werner, I. *et al.* Methylene blue modulates transendothelial migration of peripheral blood cells. *PloS one* **8**, e82214, doi: 10.1371/journal.pone.0082214 (2013).

Acknowledgements

We thank Maryam Tabib and Falko Seyffarth for their excellent technical support.

Author Contributions

N.V.B. & I.W. PhD: Conducted experiments; Analyzed Data, Prepared Draft. A.K. MD: Analyzed Data; Reviewed Manuscript. P.M. MD: Analyzed Data; Reviewed Manuscript. A.M. MD: Analyzed Data; Reviewed Manuscript. T.K. MD: Analyzed Data; Reviewed Manuscript. U.A.S. MD: Analyzed Data; Reviewed Manuscript. A.B.-F. MD, PhD: Designed Experiments; Analyzed Data; Prepared Draft, Reviewed Manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Bogert, N. V. *et al.* Influence of hypothermia and subsequent rewarming upon leukocyte-endothelial interactions and expression of Junctional-Adhesion-Molecules A and B. *Sci. Rep.* **6**, 21996; doi: 10.1038/srep21996 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>