SCIENTIFIC **REPORTS**

There are amendments to this paper

OPEN

Received: 25 November 2015 Accepted: 12 February 2016 Published: 02 March 2016

Hydrogen Sulfide Recruits Macrophage Migration by Integrin β1-Src-FAK/Pyk2-Rac Pathway in Myocardial Infarction

Lei Miao^{1,*}, Xiaoming Xin^{1,*}, Hong Xin¹, Xiaoyan Shen¹ & Yi-Zhun Zhu^{1,2}

Myocardial infarction (MI) triggers an inflammatory reaction, in which macrophages are of key importance for tissue repairing. Infiltration and/or migration of macrophages into the infarct area early after MI is critical for infarct healing, vascularization, and cardiac function. Hydrogen sulfide (H₂S) has been demonstrated to possess cardioprotective effects post MI and during the progress of cardiac remodeling. However, the specific molecular and cellular mechanisms involved in macrophage recruitment by H₂S remain to be identified. In this study, the NaHS (exogenous sources of H₂S) treatment exerted an increased infiltration of macrophages into the infarcted myocardium at early stage of MI cardiac tissues in both wild type (WT) and cystathionine- γ -lyase-knockout (CSE-KO) mice. And NaHS accelerated the migration of macrophage cells *in vitro*. While, the inhibitors not only significantly diminished the migratory ability in response to NaHS, but also blocked the activation of phospho-Src, -Pyk2, -FAK³⁹⁷, and -FAK⁹²⁵. Furthermore, NaHS induced the internalization of integrin β 1 on macrophage surface, but, integrin β 1 silencing inhibited macrophage migration and Src signaling activation. These results indicate that H₂S may have the potential as an anti-infarct of MI by governing macrophage migration, which was achieved by accelerating internalization of integrin β 1 and activating downstream Src-FAK/Pyk2-Rac pathway.

Myocardial infarction (MI) occurs, resulting in an inadequate substrate or oxygen supply of the downstream myocardium, and further leading to cardiomyocytes deterioration¹. These processes are mediated by a wide array of inflammatory reactions or factors². Monocytes/macrophages, the major source of the inflammatory factors, are of central importance for healing after MI^{3,4}. Macrophages reside in both healthy and injured heart, and increase in number during disease^{5,6}. And macrophage is a primary responder cell type that involved in the regulation of post-MI wound healing at multiple levels^{2,4}.

Induced by MI, macrophages migrate into the infarct zone, initiate intracellular signaling, which localizes the inflammatory response to clean the debris and subsequently induce scar formation⁷. The infiltration of macrophages into the infarct area early after MI, is critical for infarct healing, vascularization, and cardiac function⁸. Early depletion of infiltrating macrophages impaire wound healing, provoke adverse left ventricular (LV) remodeling, and increase mortality after MI⁹. Injection of human activated macrophage suspension early after rat MI, promotes recruitment of resident macrophages and accelerates vascularization, tissue repair, and improves cardiac remodeling and function¹⁰. Therefore, manipulating macrophage migration and function could be a promising therapeutic strategy in optimizing the process of infarct repair.

Hydrogen sulfide (H₂S), as a gasotransmitter, has been demonstrated to possess cardioprotective function in various models of cardiac injury^{11,12}. CSE (cystathionine- γ -lyase) is the predominant H₂S-generating enzyme in the cardiovascular system, and its deficiency significantly attenuates endogenous H₂S and results in exacerbating myocardial ischemia/reperfusion injury^{13,14}. Our previous studies, together with others, have demonstrated that H₂S plays diverse roles in protecting against cardiovascular diseases such as atherosclerosis, myocardial ischemia and heart failure^{15–19}. Some mechanisms are considered to contribute to the cardioprotection of H₂S, such as

¹Department of Pharmacology, School of Pharmacy and Institutes of Biomedical Sciences, Fudan University, Shanghai, China. ²Department of Pharmacology, School of Pharmacy, Macau University of Science & Technology, Macau, China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to X.S. (email: shxiaoy@fudan.edu.cn) orY.-Z.Z. (email: zhuyz@fudan.edu.cn) protecting cells against oxidative stress by increasing glutathione, promoting the translocation of the nuclear transcription factor Nrf2 to induce the activation of numerous detoxifying genes²⁰. However, the influence of H_2S in macrophage migration and the contribution to infarct repair have not been clarified. Thus, the objective of this study was to elucidate the impact of H_2S on macrophage infiltration and/or migration after MI and investigate the involved mechanisms, to offer a new dimension to our understanding of macrophage recruitment post MI.

Methods

Reagents and antibodies. NaHS was administered instead of H_2S . NaHS and the Pyk2 inhibitor (PF431396) were obtained from Sigma-Aldrich; The FAK specific inhibitor (PF573228), the Src inhibitor (PP2), and the Rac1 inhibitor (NSC23766) were purchased from Selleck Chemicals. CES-siRNA (sc-142618) was obtained from Santa Cruz. Rhodamine was obtain from Cytoskeleton, and DAPI from Beyotime; Following antibodies were used: anti-Rac, anti-Fak, anti-p-FAK³⁹⁷, anti-p-FAK⁹²⁵, anti-Src, anti-p-Src, anti- β -actin, anti-Integrin β 1, anti-Integrin β 3, anti-Cavenolin-1,anti-p-Pyk2 (Cell Signaling); anti-CSE (Santa Cruz); Anti-Integrin β 1-FITC, anti-Galectin-3 (eBbioscience); anti-CD68 (Biolegend); Lactate dehydrogenase (LDH) assay from BeyotimeInstitute of Biotechnology.

Animals. Mice devoid of CSE (KO mice) were normal in growth rate and reproduction, but had markedly reduced endogenous H_2S production. WT and KO mice used in this study were littermates obtained via heterozygous breeding. The animal procedures were performed in accordance with the Animal Management Rules of the local authorities and were approved by the ethics committee of Experimental Research, Shanghai Medical College, Fudan University.

MI Models. The left coronary artery was ligated permanently to induce myocardial infarction model as previously reported¹⁷.

Immunohistochemical analysis. The hearts excised from sacrificed animals were fixed in 4% paraformaldehyde, sectioned at 5 µm following embedded in paraffin. The immunohistochemical staining was performed with an EnVision Kit (Dako, Carpinteria, CA). Antibody specific for macrophages (anti-CD68, Biolegend and Galectin-3, eBioscience) was used to selectively detect macrophages.

Cell line culture. The murine macrophage cell line, RAW264.7, was purchased from American Type Culture Collection and maintained in RPMI 1640 medium supplemented with 10% FBS (HyClone, Logan, UT), at 37 °C in 5% CO_2 . The cells were treated with various concentrations of NaHS diluted in the RPMI 1640 medium (50, 100 and 200 mM) for different time points. Pharmacological inhibitors (5 μ M of PF431396 or PF573228, 10 μ M of PP2 and 25 μ M of NSC23766) were treated for 1 hour before NaHS was added.

Isolation of macrophages from bone marrow. Bone marrow-derived macrophages (BMMs) were isolated using standard protocols^{21,22}. After differentiation for 7 days in RPMI-1640 containing 10 ng/ml recombinant murine M-CSF, cells were either untreated or incubated with NaSH for Co-culturing.

Neonatal Mouse Primary Cardiomyocyte isolation. Primary cardiomyocytes were obtained from the ventricles of One- to two-day-old neonatal mice according to the method described²³. The isolated primary cardiomyocytes were seeded into a 24-well plateat the density of $1*10^6$ /ml then subjected to hypoxia for 4 h, in accordance with the technique described²⁴.

Co-culturing of BMMs and Cardiomyocyte. For transwell co-culturing, the 0.4-mm-pore size transwell inserts containing $2*10^5$ BMMs were placed into the 24-well plate with cardiomyocytes that were hypoxia initially. Co-culture system was incubated for 12h.

Lactate dehydrogenaseassays. Lactate dehydrogenase (LDH) was detected to evaluate the severity of cardiomyocyte injury. LDH released in the culture medium was measured using commercial kits, according to the manufacturer's instruction.

Real-time PCR. For CSE gene expression analysis, total RNA was extracted from RAW264.7 cells using TRIzol (Takara) method. cDNAs were synthesized with the RevertAidtm First Strand cDNA Synthesis Kit #1622 (Fermentas). The primer sequences for the CSE were previously described¹³.

Immunofluorescence and confocal microscopy. RAW264.7 cells plated onto glass coverslips were treated with various concentrations of NaHS for indicated time. The cells were fixed by 4% paraformaldehyde and blocked with 5% BSA (Amresco), following permeabilization by 0.5% Triton X-100. The cells were incubated with Rhodamine-conjugated phalloidin (red) for F-actin staining and DAPI (blue) for nucleus staining, or with primary antibody against integrin β 1 followed by the incubation of appropriately labeled secondary antibodies. Confocal laser scanning was carried out with Zeiss710 confocal imaging system.

Migration. The migration of RAW264.7 cells exposed to NaHS was determined by Transwell assays using polycarbonate transwell filters (Corning, 8 μ m). Cells incubated in the presence or absence of NaHS, or pharma-cological inhibitors were seeded into the upper compartment (containing 1% FBS, while the lower compartment contains 10% FBS only). The cells were allowed to migrate for 6 h before they were fixed in cold methanol. The non-migratory cells in the upper compartment were removed with a cotton swab, and the migrated cells were stained with 0.4% crystal violet (Sigma). For each experiment, the number of transmigrated cells in five random fields on the underside of the filter was counted and photographed, and three independent filters were analyzed.



Figure 1. Myocardial immunostaining of CD68 after 3, 5, or 8 days of post-MI treatment with NaHS in both WT and KO mice. Scale bars, 200 µm and 500 µm.

RNA interference. The siRNA to CSE (sc-142618) was obtained from Santa Cruz Biotechnology. And the siRNA to mouse integrin β 1 were chemically synthesised by Shanghai GenePharma Co., Ltd. The siRNA sequences for integrin β 1 were designed as follows: 5'-CAG AGA CAUUACUCAGAUdTdT-3' (forward) and 5'-AUC UGA GUA AUG UCU UCU G dTdT-3' (re-verse). A scrambled small RNA sequence was used as a negative control. The siRNAs were transfect into RAW264.7 cells in OPTI-MEM I Reduced SerumMedium (Gibco) for 24 or 48 h. The medium could be changed 6–8 h after transfection.

Protein isolation. Total proteins or cell surface proteins of cells were extracted by M-PER Mammalian Protein Extraction Reagent (Pierce) and Pierce Cell Surface Protein Isolation Kit (Thermo Scientific) respectively according to the manufacturer's instructions. Membrane proteins were extracted by Membrane and Cytosol Protein Extraction Kit (Beyotime). The protein concentration was quantified using a BCA protein assay kit (Beyotime).

Western blot. The extracted proteins were separated and transferred to nitrocellulose membranes. The membranes were blocked, following immunoblotted with appropriate antibodies. Then, the blots were developed by ImmobilonTM Western Chemiluminescent HRP Substrate (Millipore). And the immunoreactivity was visualized with a chemoluminescence reagent.

Statistical analyses. Quantitative data were presented as mean \pm SEM. The calculations were performed with GraphPad Prism 6.0 software. Differences between two groups were analyzed by two-tailed Student's t-tests, or assessed by one way analysis of variance with Tukey's post-hoc test when more than two groups were compared. The differences were considered significant with P < 0.05.

Results

NaHS increased the infiltration of macrophages early after MI. Macrophage infiltrate into the myocardium is required for proper healing of infracted myocardium⁸. To identify the role of NaHS in cardiac remodeling, the recruitment of macrophages into the heart following MI was investigated by immunohistochemical analysis of CD68 and galectin-3, the macrophage markers. As shown in Fig. 1, the density of CD68 immunostaining in KO (CSE knock out)-MI mice was apparently less than that in WT-MI mice. In both WT and KO mice, NaHS treatment significantly increased the density of CD68 staining. Interestingly, the expression of the macrophage marker was increased by NaHS treatment in WT infarct LV at day 3, peaked at day 5, and weakened at day 8 post-MI. Moreover, in the KO mice, the NaHS treatment rescued the decrease of infiltrating macrophages caused by CES depletion, and exhibited a similar time course for CD68 staining. Similarly, the immunostaining tendency of galectin-3 was consistent with CD68 (Supplemental Fig. 1). These findings indicate that H₂S promotes macrophage infiltration in the early, but not later stage of MI.

To test the potent cardioprotective efficacy of NaHS, the Lactate dehydrogenase (LDH) level in BMMs cell supernatant was determined in NaHS treatment assay. First, the isolated neonatal mouse primary cardiomyocytes were impaired by hypoxia, and LDH in supernatant was significantly increased, indicating a severity of cardiomyocyte injury (Fig. 2a). Then, BMMs, isolated from bone marrow of WT or KO mice, were co-cultured with the cardiomyocytes in the NaHS presence or not. As shown in Fig. 2b, although the LDH level was not apparently diminished in WT-BMMs and KO-BMMs groups, NaHS treated BMMs abrogated the LDH level largely compared to control group, demonstrating that macrophage infiltration triggered by NaHS antagonizes the damage of cardiomyocytes induced by hypoxia.

NaHS enhanced the migratory ability of macrophages. Cellular infiltration is suggested to be the migration of cells from their original sources. Given that NaHS preserved the cardiac function following MI through the increased infiltration of macrophages, it is still unclear whether and how NaHS stimulates



Figure 2. NaHS antagonized hypoxia-induced cardiomyocytesdamage. (a) The impair model of neonatal mouse primary cardiomyocytewas was developed, and the LDH in the cardiomyocyte supernatant was analyzed. (b) The BMMs were co-cultured with the cardiomyocytes in the NaHS presence or not, then the LDH in the cardiomyocyte supernatant was analyzed.

macrophages to move from the border zone into the core of an infarct. To address these issues, the effect of NaHS on actin cytoskeleton was analyzed by immunofluorescent microscopy in the murine macrophage cells, RAW264.7. As shown in Fig. 3a, NaHS treatment resulted in significant rearrangement of the cellular actin cytoskeleton with the elongated lamellipodia, indicating a higher migration capability. The transwell assay also confirmed that NaHS dose-dependently increased the number of migratory RAW264.7 cells (Fig. 3b,c).

It is well known that Src/FAK axis played a pivotal role in macrophage mobilization, and Pyk2 and Rac activation were involved in the destabilization of endothelial cell contacts²⁵. The Src-mediated FAK phosphorylation on several tyrosine residues plays an essential role in macrophage locomotion²⁶. Thus, to ascertain the involvement of this signaling pathway in H₂S-promoted macrophage migration, the phosphorylated status of Src, Pyk2 and FAK and activation of Rac in RAW264.7 cells were examined. As expected, significant increases in phospho-Src, -Pyk2, -FAK³⁹⁷, and -FAK⁹²⁵ were observed in RAW264.7 cells exposed to NaHS for 0.5, 1 or 2h. However, total Src and FAK were unaltered. Additionally, membrane bound Rac, the active form of Rac in the down-stream of Src/FAK signaling, was also increased under NaHS treatment for 3 h, while the loading control Caveolin-1 in cell membrane was not altered (Fig. 3d). Thus, those results indicate that H₂S indeed accelerates the migration of macrophages, and the Src-FAK/Pyk2-Rac signaling is involved in the process.

NaHS triggered macrophage migration through Src-FAK/Pyk2-Rac axis. To substantiate the Src-FAK/Pyk2-Rac axis is responsible for H_2S triggered migration, the RAW264.7 cells were pre-incubated with the Pyk2 inhibitor PF431396, the FAK specific inhibitor PF573228, the Src inhibitor PP2, or the Rac inhibitor NSC23766, and then NaHS was added and incubated for 6 h. The migratory ability of the treated cells was determined by transwell assay. As shown in Fig. 4a,b, NaHS accelerated the migration of RAW264.7 cells, which were diminished by all the four inhibitors. Western blot results were also consistent with the results of transwell assay, revealed that PF431396 suppressed the NaHS-induced activation of Pyk2. Additionally, PF573228 significantly decreased the NaHS triggered phosphorylation of Fak, but had a weak effect on phosphorylated Src. PP2 completely blocked the phosphorylation of Src, Pyk2 and FAK induced by NaHS (Fig. 4c). These results confirm that the Src-FAK/Pyk2-Rac signaling is critical in H_2 S-induced macrophages migration, and also indicate that the direct target of H_2 S is in the up-stream of this signaling pathway.

NaHS promoted the internalization of macrophage surface integrin β **1**. To pinpoint the direct target of H₂S in triggering macrophage migration, the levels of integrin β 1 and β 3 in the surface of RAW264.7 cells with or without NaHS treatment were examined. The total level of integrin β 1 or β 3 did not change within the time course of NaHS treatment (Supplemental Fig. 2). While, the internalization of surface integrin β 1 dramatically increased under NaHS treatment. And the surface integrin β 1 decreased much more quickly in NaHS treated cells than in control cells. However, only a weak effect was observed on the internalization of integrin β 3 (Fig. 5a). In addition, flow cytometric analysis also revealed that the surface integrin β 1 was decreased much more quickly following NaHS treatment (Fig. 5b). To further verify the effect of NaHS on the internalization of integrin β 1, confocal microscopic analyses were performed. As shown in Fig. 5c, within the indicated time course of treatment, the labeled integrin β 1 was gradually internalized in the NaHS groups, but mainly presented on the plasma membrane in the untreated control. Collectively, these findings, on one side, suggest a function of H₂S in





indicated concentrations of NaHS for 6 h, then subjected to immunofluorescence analysis using Rhodamineconjugated phalloidin (red) for F-actin, DAPI (blue) for nucleus. Scale bar, 5 μ m. (**b**) After treatment with indicated concentrations of NaHS for 6 h, the migratory ability of RAW264.7 cells was analyzed by transwell assay. Scale bar, 50 μ m. (**c**) The number of migrated cells per field was statistically analyzed. Values are the mean \pm SEM from three independent experiments. *P < 0.05, **P < 0.01 versus control. (**d**) Western blot images of proteins from NaHS treated RAW264.7 cells, the probed antibodies as indicated (I, total cell lysates; II, membrane extracts; p, phosphorylated; t, total).

regulating the internalization of macrophage surface integrin β 1; on the other side, indicate that H₂S may activate the Src-FAK/Pyk2-Rac signaling *via* an integrin β 1-dependent pathway.

Integrin β 1 silencing inhibited macrophage migration and Src-FAK/Pyk2 signaling activation induced by NaHS. To ascertain that integrin β 1 is the target of H₂S for promoting macrophages migration through activation of Src-FAK/Pyk2-Rac signaling, the integrin β 1-specific-siRNA was introduced into RAW264.7 cells. The transfection efficiency was confirmed by western blot, and the data showed that integrin β 1 expression in the siRNA- β 1 group was reduced significantly (Supplemental Fig. 3). Next, migratory capacity of the siRNA- β 1 transfected cells in response to NaHS was evaluated by transwell assay. As shown in Fig. 5d, the increased migratory capacity of H₂S treated macrophages was significantly impaired by integrin β 1-siRNA. Furthermore, the silence of integrin β 1 also inhibited the activation of Src-FAK/Pyk2 signaling induced by NaHS (Fig. 5e). Thus, these results confirm that integrin β 1-Src-FAK/Pyk2-Rac signaling played a pivotal role in H₂S-induced macrophage migration.

Endogenous CSE knockdown diminished macrophage migration, integrin β 1 internalization and Src-FAK/Pyk2 signaling activation. To further explore the influence of H₂S on macrophages migration, endogenous CSE was knocked down by specific siRNA in RAW264.7 cells before the NaHS treatment.



Figure 4. The Src-FAK/Pyk2-Rac pathway responded to macrophage migration triggered by NaHS. (a) RAW264.7 cells were incubated for 1 h in the presence or absence of PF431396 (5 μ M), PF573228 (5 μ M), PP2 (10 μ M) or NSC23766 (25 μ M) respectively, followed by incubation with additional 100 μ M NaHS for 6 h. The migratory ability of RAW264.7 cells was determined by transwell assay. Scale bar, 50 μ m. (b) The numbers of the migrated cells per field were shown as the mean ± SEM from three independent experiments. *P < 0.05 versus Control, **P < 0.01 versus NaHS treatment. (c) RAW264.7 cells were incubated with PF431396 (5 μ M), PF573228 (5 μ M) or PP2 (10 μ M) respectively for 1 h before 100 μ M NaHS treatment. The expressions of indicated proteins were assessed by western blot (I, total cell lysates; II, membrane extracts; p, phosphorylated; t, total).

Compared to siRNA-NC, the siRNA-CSE group significantly reduced the mRNA level of CSE (Fig. 6a). When the endogenous CSE was down-regulated, the NaHS induced migration of macrophages was abolished (Fig. 6b,c). Additionally, CSE silencing also abrogated the activation of phospho-Src, -Pyk2, and -FAK induced by NaHS. Further, although endogenous CSE knockdown did not alter the total amount of integrin β 1, increased the cell surface integrin β 1 (Fig. 6d). Taken together, these findings suggest that H₂S indeed promotes macrophage migration *via* accelerating internalization of integrin β 1 and activating the down-stream Src-FAK/Pyk2-Rac signaling.

Discussion

Up to date, therapeutic effects of clinically applicable drugs for MI are still not satisfactory^{1,27}. H_2S as a novel gaseous signaling molecule has been reported a potent cardioprotective effect^{16,17}. Our previous study showed that increasing endogenous H_2S by a water-soluble modulator S-Propargyl-Cysteine ameliorated ischemic conditions through angiogenesis promotion¹⁹. In our recent study, we also proved that decrease of H_2S by CSE KO aggravated cardiac dysfunction and increased mortality post-MI; while, an improved cardiac remodeling and function accompanied with decreased mortality was observed in both WT mice and CSE KO-MI mice treated with different concentrations of NaHS, suggesting that a significant efficacy of H_2S for MI treatment (unpublished data).

Macrophages digest debris and dominate inflammation resolution following MI^{28} . At the early stage after MI, depletion of infiltrating macrophage aggravates pathological infarct healing, resulting in an increased left ventricular dilatation and wall thinning⁹. Acute MI alters the macrophage phenotype and supply chain²⁹. While, increase in macrophage recruitment accelerates infarct repair, and improves cardiac remodeling and function¹⁰. In the present study, by immunostaining with the antibodies against CD68 and galectin-3, the markers of macrophage^{30,31}, we found that H₂S could *in vivo* increase the infiltration of macrophages early after MI, suggesting a protective role of H₂S in early post-MI inflammation and the subsequent healing process³².

Accentuation, prolongation, or expansion of the post infarction inflammatory response leads to a worse remodeling or dysfunction in MI^{33–35}. Within the initial stage, MI, on one side, results in the migration of



Figure 5. The influence of NaHS in the internalization of surface integrin β 1 and Src-FAK/Pyk2 signaling. (a) Surface proteins of RAW264.7 cells were biotinylated and separated by streptavidin beads. Integrin β 1 and integrin β 3 levels were detected by western blot (I: internalized proteins from cell surface; II: total cell lysates; III: cell surface proteins). (b) RAW264.7 cells were incubated with NaHS (100 µM) for the indicated time and surface levels of integrin β 1 were labeled with the FITC-conjugated secondary antibody, followed by Flow cytometric analysis. (c) RAW264.7 cells were incubated with NaHS (100 µM) for the indicated time. Surface Integrin β 1 was stained with FITC-conjugated antibody (green); F-actin bundles were detected with Rhodamine (Red); Nuclei were counterstained with DAPI (blue). Scale bar, 5 µm. All the data are representative of three independent experiments. (d) After siRNA-integrin β 1 or siRNA-NC transfection for 24 h, RAW264.7 cells were incubated by transwell assay. Values are the mean \pm SD from three independent experiments. (e) RAW264.7 cells transfected with NaHS (100 µM) for 6 h and the phosphor-Src, -Pyk2, -FAK³⁹⁷, -FAK⁹²⁵, total Src and FAK were analyzed by western blot (NC, negative control; p, phosphorylated; t, total).

macrophages into the infarcted myocardium, initiating intracellular signaling; on the other side, macrophages responding to signals, could also move from the border zone into the core of an infarct². The phase and rate of macrophage infiltration is orchestrated by a wide range of factors. And yet, the specific signals and mechanism responsible for macrophage migration remain unclear. In our present study, increased infiltrating macrophages raised an important question of how H_2S triggers the migration of macrophages in the cardiac wound.

Previous report by Frangogiannis showed that macrophage infiltration and migration into the myocardium post-MI involved upregulation of both integrins and adhesion molecules³⁶. Integrins, especially β 1 and β 3, are highly expressed isoforms in macrophages, and have been reported to play important roles in receiving ECM signals and regulating cell migration^{37–39}. Integrin trafficking, including internalization, is the crucial step in cell communication³⁸. Downregulation of integrin β 1 at the plasma membrane is a potential mechanism for





enhancing cell motility⁴⁰. Thus, through controlling integrin $\beta 1$ internalization and recycling, junctional adhesion molecule could promote neutrophil chemotaxis⁴¹. Here, by biotin-based experiments, we found that H₂S dramatically induced the internalization of integrin $\beta 1$, but not integrin $\beta 3$; and silence of integrin $\beta 1$ inhibited the migration of macrophages, indicating that integrin $\beta 1$ might be the possible target of H₂S in regulating macrophage migration.

The internalization of integrin β1 can activate down-stream FAK/Src signaling, which is an important pathway involving in cell migration. The expression or activation of Src, a cellular nonreceptor tyrosine kinase in the downstream of integrin signaling, could reflect the migratory ability of macrophage⁴². As a substrate of Src, FAK serves as an adaptor that connects integrins with actin cytoskeleton⁴³. Upon integrin activation, FAK is phosphorylated at tyrosine residues Tyr³⁹⁷ or/and Tyr⁹²⁵, which in turn activates the binding site Src⁴²⁻⁴⁴. Recently, Liu at al proved that integrins, phospho-FAK and phospho-Src participate in MI microenvironment improvement for myocardial repair⁴⁵. By analogy to the related FAK kinase, Pyk2 is important in macrophage activation⁴⁶. Additionally, evidence also shows that through controlling phosphorylation of Pyk2 and paxillin, Pyk2 regulates the polarization, migration, and spreading of macrophage⁴⁷. Rac, a member of Rho GTPase family, its activation is also related to the cell polarization and migration in macrophages⁴⁸. In this study, the phospho-Src, -Pyk2, -FAK³⁹⁷, and -FAK⁹²⁵ were significantly upregulated upon H₂S treatment, and the downstream Rac was activated relatively later. Inhibitors for FAK, Src, Pyk2, and Rac could effectively diminish the migratory ability of macrophages stimulated by H₂S and block the activation of related signaling molecules in response to H₂S. Those results hint that H₂S triggers macrophage migration *via* the Src-FAK/Pyk2-Rac axis.

In conclusion, our results revealed that H₂S treatment exerted an increased infiltration of macrophages into the infarcted myocardium at early stage of MI in both WT mice and CSE-KO mice, and the increased macrophage

infiltrating antagonized the damage of cardiomyocytes induced by hypoxia. The favorable effects of H₂S on the infarcted myocardium can be attributed to the promotion of infiltrating macrophages. Further, we present evidence for a novel mechanism that macrophage migration triggered by H₂S is mediated by integrin β 1-Src-FAK/ Pyk2-Rac pathway. Thus, activation of macrophages by H₂S supplement can be a promising therapeutic strategy for MI treatment.

References

- 1. Downey, J. M. & Cohen, M. V. Why do we still not have cardioprotective drugs? Circ. J. 73, 1171-1177 (2009).
- 2. Lambert, J. M., Lopez, E. F. & Lindsey, M. L. Macrophage roles following myocardial infarction. Int. J. Cardiol. 130, 147-158 (2008).
- 3. Frantz, S. *et al.* Monocytes/macrophages prevent healing defects and left ventricular thrombus formation after myocardial infarction. *FASEB J.* 27, 871–881 (2013).
- Weinberger, T. & Schulz, C. Myocardial infarction: a critical role of macrophages in cardiac remodeling. *Front. Physiol.* 6, 107 (2015).
 Nahrendorf, M. *et al.* The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions. *J. Exp. Med.* 204, 3037–3047 (2007).
- 6. Schultze, J. L., Schmieder, A. & Goerdt, S. Macrophage activation in human diseases. Semin. Immunol. 27, 249-56 (2015).
- 7. Sutton, M. G. & Sharpe, N. Left ventricular remodeling after myocardial infarction: pathophysiology and therapy. *Circulation* 101, 2981–2988 (2000).
- Nahrendorf, M. & Swirski, F. K. Monocyte and macrophage heterogeneity in the heart. *Circ. Res.* 112, 1624–1633 (2013).
 van Amerongen, M. J., Harmsen, M. C., van Rooijen, N., Petersen, A. H. & van Luyn, M. J. Macrophage depletion impairs wound
- healing and increases left ventricular remodeling after myocardial injury in mice. *Am. J. Pathol.* **170**, 818–829 (2007).
- 0. Leor, J. et al. Ex vivo activated human macrophages improve healing, remodeling, and function of the infarcted heart. *Circulation* **114**, 194–100 (2006).
- 11. Qipshidze, N., Metreveli, N., Mishra, P. K., Lominadze, D. & Tyagi, S. C. Hydrogen sulfide mitigates cardiac remodeling during myocardial infarction via improvement of angiogenesis. *Int. J. Biol. Sci.* **8**, 430–441 (2012).
- Shen, Y., Shen, Z., Luo, S., Guo, W. & Zhu, Y. Z. The Cardioprotective Effects of Hydrogen Sulfide in Heart Diseases: From Molecular Mechanisms to Therapeutic Potential. Oxid. Med. Cell Longev. 2015, 925167 (2015).
- 13. Pan, L. L., Liu, X. H., Gong, Q. H., Yang, H. B. & Zhu, T. Z. Role of cystathionine gamma-lyase/hydrogen sulfide pathway in cardiovascular disease: a novel therapeutic strategy? *Antioxid. Redox Signal.* **17**, 106–118 (2012).
- 14. Calvert, J. W., Coetzee, W. A. & Lefer, D. J. Novel insights into hydrogen sulfide-mediated cytoprotection. *Antioxid. Redox Signal.* 12, 1203–1217 (2010).
- Wang, Y. et al. Role of hydrogen sulfide in the development of atherosclerotic lesions in apolipoprotein E knockout mice. Arterioscler. Thromb. Vasc. Biol. 29, 173–179 (2009).
- Chuah, S. C., Moore, P. K. & Zhu, Y. Z. S-allylcysteine mediates cardioprotection in an acute myocardial infarction rat model via a hydrogen sulfide-mediated pathway. Am. J. Physiol. Heart Circ. Physiol. 293, H2693–2701 (2007).
- 17. Zhu, Y. Z. et al. Hydrogen sulfide and its possible roles in myocardial ischemia in experimental rats. J. Appl. Physiol. 102, 261–268 (2007).
- Huang, C. et al. Cardioprotective effects of a novel hydrogen sulfide agent-controlled release formulation of S-propargyl-cysteine on heart failure rats and molecular mechanisms. PloS one 8, e69205 (2013).
- Kan, J. et al. S-propargyl-cysteine, a novel water-soluble modulator of endogenous hydrogen sulfide, promotes angiogenesis through activation of signal transducer and activator of transcription 3. Antioxid. Redox Signal. 20, 2303–2316 (2014).
- Polhemus, D. J., Calvert, J. W., Butler, J. & Lefer, D. J. The cardioprotective actions of hydrogen sulfide in acute myocardial infarction and heart failure. Scientifica (Cairo) 2014, 768607 (2014).
- 21. Zhang, X., Goncalves, R. & Mosser, D. M. The isolation and characterization of murine macrophages. *Curr. Protoc. Immunol.* Chapter 14, Unit 14 (2008).
- 22. Weischenfeldt, J. & Porse, B. Bone Marrow-Derived Macrophages (BMM): Isolation and Applications. CSH Protoc. 2008, pdb prot5080 (2008).
- Wang, G. W., Schuschke, D. A. & Kang, Y. J. Metallothionein-overexpressing neonatal mouse cardiomyocytes are resistant to H₂O₂ toxicity. Am. J. Physiol. 276, H167–175 (1999).
- Rakhit, R. D. et al. Nitric oxide-induced cardioprotection in cultured rat ventricular myocytes. Am. J. Physiol. Heart Circ. Physiol. 278, H1211–1217 (2000).
- van Buul, J. D., Anthony, E. C., Fernandez-Borja, M., Burridge, K. & Hordijk, P. L. Proline-rich tyrosine kinase 2 (Pyk2) mediates vascular endothelial-cadherin-based cell-cell adhesion by regulating beta-catenin tyrosine phosphorylation. *J. Biol. Chem.* 280, 21129–21136 (2005).
- 26. Owen, K. A. *et al.* Regulation of lamellipodial persistence, adhesion turnover, and motility in macrophages by focal adhesion kinase. *J. Cell Biol.* **179**, 1275–1287 (2007).
- Rossello, X., Pocock, S. J. & Julian, D. G. Long-Term Use of Cardiovascular Drugs: Challenges for Research and for Patient Care. J. Am. Coll. Cardiol. 66, 1273–1285 (2015).
- Kain, V. & Halade, G. V. Big eater macrophages dominate inflammation resolution following myocardial infarction. J. Mol. Cell. Cardiol. 87, 225–227 (2015).
- 29. Nahrendorf, M. & Swirski, F. K. Innate immune cells in ischaemic heart disease: does myocardial infarction beget myocardial infarction? *Eur. Heart J.* [Epub ahead of print] (2015).
- Li, X. et al. Myocardial injury after ischemia-reperfusion in mice deficient in Akt2 is associated with increased cardiac macrophage density. Am. J. Physiol. Heart Circ. Physiol. 301, H1932–1940 (2011).
- 31. Cha, S. J. et al. CD68 acts as a major gateway for malaria sporozoite liver infection. J. Exp. Med. 212, 1391–1403 (2015).
- 32. Nagai, T. *et al.* Decreased myocardial dendritic cells is associated with impaired reparative fibrosis and development of cardiac rupture after myocardial infarction in humans. *J. Am. Heart Assoc.* **3**, e000839 (2014).
- 33. Frangogiannis, N. G. Regulation of the inflammatory response in cardiac repair. Circ. Res. 110, 159-173 (2012).
- 34. Mann, D. L. The emerging role of innate immunity in the heart and vascular system: for whom the cell tolls. *Circ. Res.* 108, 1133–1145 (2011).
- Liehn, E. A., Postea, O., Curaj, A. & Marx, N. Repair after myocardial infarction, between fantasy and reality: the role of chemokines. J. Am. Coll. Cardiol. 58, 2357–2362 (2011).
- 36. Frangogiannis, N. G. The mechanistic basis of infarct healing. Antioxid. Redox Signal. 8, 1907–1939 (2006).
- 37. Caswell, P. T. & Norman, J. C. Integrin trafficking and the control of cell migration. Traffic 7, 14-21 (2006).
- Lee, H. D., Kim, Y. H. & Kim, D. S. Exosomes derived from human macrophages suppress endothelial cell migration by controlling integrin trafficking. *Eur. J. Immunol.* 44, 1156–1169 (2014).
- Tanimizu, N., Kikkawa, Y., Mitaka, T. & Miyajima, A. alpha1- and alpha5-containing laminins regulate the development of bile ducts via beta1 integrin signals. J. Biol. Chem. 287, 28586–28597 (2012).
- Panicker, A. K., Buhusi, M., Erickson, A. & Maness, P. F. Endocytosis of beta1 integrins is an early event in migration promoted by the cell adhesion molecule L1. *Exp. Cell Res.* 312, 299–307 (2006).

- 41. Cera, M. R. et al. JAM-A promotes neutrophil chemotaxis by controlling integrin internalization and recycling. J. Cell Sci. 122, 268–277 (2009).
- Maa, M. C. et al. Requirement of inducible nitric-oxide synthase in lipopolysaccharide-mediated Src induction and macrophage migration. J. Biol. Chem. 283, 31408–31416 (2008).
- Thapa, B., Koo, B. H., Kim, Y. H., Kwon, H. J. & Kim, D. S. Plasminogen activator inhibitor-1 regulates infiltration of macrophages into melanoma via phosphorylation of FAK-Tyr(9)(2)(5). Biochem. Biophys. Res. Commun. 450, 1696–1701 (2014).
- 44. Mitra, S. K., Hanson, D. A. & Schlaepfer, D. D. Focal adhesion kinase: in command and control of cell motility. *Nat. Rev. Mol. Cell Biol.* **6**, 56–68 (2005).
- 45. Liu, Z. *et al.* The influence of chitosan hydrogel on stem cell engraftment, survival and homing in the ischemic myocardial microenvironment. *Biomaterials* **33**, 3093–3106 (2012).
- 46. Kamen, L. A., Schlessinger, J. & Lowell, C. A. Pyk2 is required for neutrophil degranulation and host defense responses to bacterial infection. J. Immunol. 186, 1656–1665 (2011).
- Rhee, I., Davidson, D., Souza, C. M., Vacher, J. & Veillette, A. Macrophage fusion is controlled by the cytoplasmic protein tyrosine phosphatase PTP-PEST/PTPN12. *Mol. Cell. Biol.* 33, 2458–2469 (2013).
- Aflaki, E. et al. Impaired Rho GTPase activation abrogates cell polarization and migration in macrophages with defective lipolysis. Cell. Mol. Life Sci. 68, 3933–3947 (2011).

Acknowledgements

This study was supported by the Key Program of National Nature Science Foundation of China (No. 81330080) and the Key Program of Shanghai Committee of Science and Technology in China (No. 14JC1401100).

Author Contributions

L.M., X.S. and Y.-Z.Z. designed the research; L.M., X.X. and H.X. performed the experiments. L.M. wrote the manuscript. X.X. and H.X. performed array analysis. Y.-Z.Z. and X.S. supervised the work and helped write the manuscript.

Additional Information

Supplementary information accompanies this paper at http://www.nature.com/srep

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Miao, L. *et al*. Hydrogen Sulfide Recruits Macrophage Migration by Integrin β1-Src-FAK/Pyk2-Rac Pathway in Myocardial Infarction. *Sci. Rep.* **6**, 22363; doi: 10.1038/srep22363 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/