

SCIENTIFIC REPORTS



OPEN

Dust Rains Deliver Diverse Assemblages of Microorganisms to the Eastern Mediterranean

Ghida Nouhad Itani & Colin Andrew Smith

Received: 15 November 2015

Accepted: 18 February 2016

Published: 04 March 2016

Dust rains may be particularly effective at delivering microorganisms, yet their biodiversities have been seldom examined. During 2011 and 2012 in Beirut, Lebanon, 16 of 21 collected rainfalls appeared dusty. Trajectory modelling of air mass origins was consistent with North African sources and at least one Southwest Asian source. As much as ~4 g particulate matter, ~20 µg DNA, and 50 million colony forming units were found deposited per square meter during rainfalls each lasting less than one day. Sequencing of 93 bacteria and 25 fungi cultured from rain samples revealed diverse bacterial phyla, both Gram positive and negative, and Ascomycota fungi. Denaturing Gradient Gel Electrophoresis of amplified 16S rDNA of 13 rains revealed distinct and diverse assemblages of bacteria. Dust rain 16S libraries yielded 131 sequences matching, in decreasing order of abundance, Betaproteobacteria, Alphaproteobacteria, Firmicutes, Actinobacteria, Bacteroidetes, Cyanobacteria, Epsilonproteobacteria, Gammaproteobacteria, and Deltaproteobacteria. Clean rain 16S libraries yielded 33 sequences matching only Betaproteobacteria family Oxalobacteraceae. Microbial composition varied between dust rains, and more diverse and different microbes were found in dust rains than clean rains. These results show that dust rains deliver diverse communities of microorganisms that may be complex products of revived desert soil species and fertilized cloud species.

Deserts supply prodigious amounts of dust¹ that affect many human concerns, including weather², climate³, and health⁴, and dust is transported large distances⁵. Satellites allow comprehensive monitoring and supply dramatic imagery⁶, and with trajectory modelling⁷, help reveal origins and dispersal of desert dust. The largest sources of dust are in North Africa, and the Arabian Peninsula is a significant source⁸.

The mineral and microbial composition of desert dusts and their effects on downwind ecologies and human health has attracted much interest^{9–13}. Aeolian dust supplies important mineral nutrients to aquatic ecosystems^{3,14}. Particulate matter⁴ and pathogens^{9,15} are of greatest health concerns. In addition to dry deposition of desert dust, wet deposition occurs in the Mediterranean region and is known as dust rain, red rain, bloody rain, coloured rain, and muddy rain. Dust rains have been noted¹⁶ and natural origins posited¹⁷ since antiquity. In the modern era, the biogeochemical and ecological roles of dust rains attract scientific interest^{18–20}.

Wet deposition of dust by rain is much less studied than dry deposition, and very few reports of microbial composition are available^{21,22}. Aware of the widespread transport of microorganisms in aeolian dust¹⁰ and clouds²³, having observed occasionally dramatic dust rains in Lebanon (see Supplementary Fig. S1), and considering that rain clouds could provide additional protection against desiccation and solar radiation and efficiently wash microorganisms into recipient soils, we wondered as to the diversity of microorganisms delivered to the eastern Mediterranean by dust rains.

Estimates suggest that each gram of arid soil contains as many as 10⁹ prokaryotes²⁴ that may be aerosolized with soil particles. Estimates of bacteria in clouds²⁵ exceed 10⁵ ml⁻¹. Modern molecular biology offers tools to identify species in environmental samples by isolation of DNA, amplification of small subunit ribosomal DNA genes, and DNA sequencing²⁶. Metagenomic approaches have revolutionized environmental microbiology²⁷ and allow identification of species without culturing^{28,29}, although biases may still exist³⁰.

Here, we examined the microbial composition of dust rains falling in Beirut, Lebanon using total DNA extraction, solid media cultures, amplification of 16S and 18S rDNA, and sequencing.

Department of Biology, American University of Beirut, Beirut, Lebanon. Correspondence and requests for materials should be addressed to C.A.S. (email: colin.smith@aub.edu.lb)

Rain date ^a	Trajectory ^b	Observations ^c	Sources ^d	Exposure ^e , day	Rainfall, mm	PM ^f , mg/m ²	DNA, ng/m ²	CFU, 10 ³ /m ²
20110923	2011092310	muddy, dusty skies	SE, M, Naf	0.75	12	120	900	600
20110924	2011092400	muddy	SE, M	0.75	0.3	20	300	700
<i>20111001</i>	<i>2011093023</i>	<i>clean rain</i>	WAs, M	<i>0.75</i>	<i>20</i>	<i>100</i>	<i>NP^g</i>	<i>7,000</i>
20111025	2011102518	muddy, dusty skies	uncertain	0.65	20	300	5,000	3,000
20111026	2011102610	muddy	uncertain	0.15	5	30	3,000	300
<i>20111103</i>	<i>2011110312</i>	<i>clean rain</i>	SE, WAs, M	<i>0.35</i>	<i>30</i>	<i>80</i>	<i>10,000</i>	<i>3,000</i>
20111114	2011111408	very muddy, dusty skies	uncertain	0.5	0.4	30	10,000	200
20111115	2011111518	clean rain and hail	WAs, M, Naf	1	100	200	16,000	50,000
20111224	2011122400	muddy, dusty skies	SE, M, Naf	0.75	7	300	3000	4,000
20120111	2012011112	muddy	E, M, Naf	1	17	80	NP ^g	6,000
<i>20120113</i>	<i>2012011300</i>	<i>clean rain</i>	<i>E, M, Naf</i>	<i>0.75</i>	<i>30</i>	<i>20</i>	<i>200</i>	<i>1,000</i>
20120208	2012020812	very muddy	E, M, Naf	1	6	500	3,000	4,000
20120215	2012021512	very muddy	Naf	1	5	300	11,000	700
20120314	2012031400	very muddy	SE, M, Naf	0.75	5	100	20,000	30,000
20120402	2012040200	very muddy	M, Naf, SWAs	0.75	1.0	300	9,000	500
20120420	2012042000	very muddy	E, Naf, M	0.75	0.3	130	300	1,000
20120430	2012043018	very muddy	SWAs	0.75	0.04	500	6,000	3,000
20120502	2012050215	very muddy	Naf, M, SWAs	0.75	0.5	4,000	12,000	1,600
20120529	2012052906	muddy, dusty skies	Naf, M	0.15	0.6	80	150	200
20121220	2012122000	muddy and hail	M, Naf	1	30	600	7,000	15,000
<i>20121221</i>	<i>2012122100</i>	<i>clean, smell of snow</i>	SE, WAs, M	<i>0.75</i>	<i>70</i>	<i>170</i>	<i>1,700</i>	<i>8,000</i>

Table 1. Characterisation of rains. ^aRain date: the nominal date assigned to the rainfall, usually the calendar date of the morning after precipitation. Clean rain entries are italicized, and rains selected for more intense study are bold. ^bTrajectory: the specific date and time used for the backward trajectory (format: YYYYMMDDHH). ^cObservations: the general appearance of the rain water and weather when notable. ^dSources: wind backward trajectories passed over Europe (E), southern Europe (SE), the Mediterranean Sea (M), North Africa (Naf), western Asia (WAs), and southwestern Asia (SWAs). Most backward trajectories passed over multiple regions. Some did not yield coherent predictions and are labelled “uncertain.” ^eNo exposure time exceeded one day, and rain deposition was frequently of much shorter duration than exposure. ^fPM: particulate matter. ^gNP: Not processed; DNA was not extracted from this sample.

Results

Amount of particulate matter, DNA, and CFU found in rains varied widely. During 2011 and 2012 in Beirut, Lebanon, 21 rainfalls were collected, of which 16 appeared to contain dust (appeared muddy) before filtration. Rainfalls were filtered and the residues analysed. The mass of particulate matter in each rain was measured and found to vary widely, ranging between 20 and 4,000 mg m⁻² (Table 1). The characterisation of rains as dusty was subjective and did not relate directly to mass of particulate matter deposited in the collector, rather, the appearance of dustiness (tan or ochre and turbid) usually correlated to a high ratio of particulate matter to rain.

Total DNA was extracted from 19 rainfalls by bead-beating of residues. DNA masses ranged from 150 to ~20,000 ng m⁻² (Table 1). The mass of particulate matter did not correlate well to that of DNA ($n = 18$, Spearman's correlation = 0.33; $p = 0.19$). Colony forming units (CFU) were determined by plating dilutions on solid media and growth for 3 days at room temperature. Culturable microbes ranged widely from 200 thousand to 50 million CFU m⁻² (Table 1). No strong relationships were observed between CFU and DNA mass ($n = 18$, Spearman's correlation = 0.36; $p = 0.14$), and between CFU and mass of particulate matter ($n = 18$, Spearman's correlation = 0.48; $p = 0.04$).

Most dust rain backward trajectories passed over arid regions of North Africa. The Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPLIT) was used to trace air volumes backwards in time to predict plausible origins of solids and microorganisms in rains⁷. Most rains had trajectories consistent with acquisition of dust or soils from arid regions of North Africa, although one traced only from Southwest Asia, and several had mixed or uncertain trajectories (Fig. 1, see Supplementary Fig. S2).

Dust rains contain diverse culturable microorganisms. Reasoner's 2A agar was chosen for its ability to support growth of bacteria found in water and its relatively lower nutrient composition in the expectation that growth-rate differences between slow- and fast-growing microbes would be reduced. Rain cultures on solid media revealed live, diverse, and often pigmented colonies (see Supplementary Fig. S3). Cultures of two clean rains, 3 November 2011 and 15 November 2011, were notable for high CFU, less apparent diversity, and fewer pigmented colonies. Avoiding duplication of colonies of similar appearance, a few sample colonies were picked randomly from each rainfall for identification by amplification and sequencing of bacterial 16S rDNA and fungal 18S rDNA. Ninety-three sequences matched 16S database entries, comprising 5 phyla, including 3




Figure 1. Wind Backward Trajectories of Four Selected Rains. (a) 25 October 2011. (b) 26 October 2011. (c) 15 November 2011. (d) 24 December 2011. Trajectories are for 72 hours ending at the described date and time. Red traces with triangles follow the air volume at 500 m; blue traces are for 1000 m; green traces are for 2000 m. Symbols mark 3-hour intervals with large symbols at 24-hour intervals. The images were obtained using the online HYSPLIT model from the US National Oceanic and Atmospheric Administration using Global Data Assimilation System meteorological data (http://ready.arl.noaa.gov/HYSPLIT_traj.php). All trajectories are found in Supplementary Fig. S2.

classes of phylum Proteobacteria, in decreasing order of relative abundance: Actinobacteria (33/93), Firmicutes (19/93), Betaproteobacteria (16/93), Bacteroidetes (9/93), Gammaproteobacteria (9/93), Alphaproteobacteria (6/93), and Deinococcus-Thermus (1/93) (Table 2, see Supplementary Table S1, S2). No Bacteroidetes or Deinococcus-Thermus were cultured from clean rains. Twenty-five sequences matched 18S database entries, all from the phylum Ascomycota (Table 3). Ascomycota classes included Dothideo (17/25), Eurotiomycetes (6/25), and Sordariomycetes (2/25). Only Dothideomycetes were found in clean rain cultures, although only 3 cultures were sequenced (Table 3, see Supplementary Table S3).

DGGE analysis of 16S rDNA shows diversity of bacteria in dust rains. In order to display the diversity of bacteria in dust rains including unculturable species, Denaturing Gradient Gel Electrophoresis (DGGE) was used to separate and visualize a 233 bp amplicon of the V3 variable region of 16S rDNA in 13 rainfalls (Fig. 2). Each rain presented a pattern of bands consistent with different and diverse assemblages of bacteria. Several samples were dominated by one or two prominent bands, and 16 bands were excised for sequencing, of which 9 yielded readable sequences. Bands 3, 6, and 7 yielded readable sequences closely matching Betaproteobacteria class Burkholderiales genera *Pelomonas*, *Ralstonia*, and *Herbaspirillum*, respectively (see Supplementary Table S4). Bands 5 and 9 were short, but matched Burkholderiales entries. Although the sequence of band 9 of clean rain of 15 November 2011 was short and of poorer quality, it aligned precisely with band 7 of clean rain of 3 November 2011 and matched *Herbaspirillum* as well as any other entry, suggesting it represents the same species as band 7 (Fig. 2). Other than the alignment of bands 7 and 9, no other shared bands were identified with confidence.

Rain date	Matches to phyla in cultured bacteria ^a	Matches to phyla in uncultured bacteria ^b
20110923	3 Actino, 1 Beta, 1 Gamma	1 Alpha, 1 Cyano
20110924	2 Actino, 1 Deino	1 Alpha, 1 Beta, 1 Firmi
20111001	3 Actino, 1 Firmi	NP ^c
20111025	ND ^d	5 Alpha, 23 Beta
20111026	1 Actino, 1 Firmi, 1 Gamma	11 Alpha, 28 Beta
20111103	2 Actino, 4 Beta, 1 Gamma	5 Beta
20111114	2 Actino, 1 Bacter, 1 Beta, 1 Firmi	1 Actino, 3 Cyano
20111115	1 Actino, 1 Alpha, 2 Beta, 1 Gamma	28 Beta
20111224	1 Actino, 1 Bacter, 2 Firmi	4 Actino, 13 Alpha, 2 Bacter, 2 Beta, 1 Delta, 3 Epsilon, 6 Firmi, 1 Gamma
20120111	1 Actino, 1 Alpha, 1 Beta, 2 Firmi	NP
20120113	2 Actino	0 ^e
20120208	1 Actino, 3 Bacter, 2 Firmi	1 Actino, 1 Alpha, 1 Bacter, 2 Beta, 3 Firmi
20120215	2 Bacter, 3 Beta,	3 Bacter
20120314	3 Actino, 1 Bacter, 1 Gamma,	2 Beta, 1 Gamma
20120402	1 Actino, 3 Beta, 2 Gamma,	1 Actino, 5 Beta, 3 Firmi
20120420	1 Actino, 1 Alpha, 1 Firmi, 1 Gamma	NA ^f
20120430	3 Actino, 1 Firmi	NA ^f
20120502	1 Actino, 1 Beta, 3 Firmi,	NA ^f
20120529	1 Actino, 1 Alpha, 1 Firmi	NA ^f
20121220	3 Actino, 1 Alpha, 1 Bacter, 1 Gamma	0 ^e
20121221	1 Actino, 1 Alpha, 4 Firmi	0 ^e

Table 2. Numbers of bacterial sequences obtained from cultures and uncultured total DNA. ^aRain residues were cultured on Reasoner's 2A agar and sample colonies identified by sequencing of amplified 16S rDNA. Identification abbreviations are Actino: Actinobacteria, Alpha: Alphaproteobacteria, Bacter: Bacteroidetes, Beta: Betaproteobacteria, Cyano: Cyanobacteria, Deino: Deinococcus-Thermus, Delta: Deltaproteobacteria, Epsilon: Epsilonproteobacteria, Firmi: Firmicutes, Gamma: Gammaproteobacteria. ^bDNA was extracted from rain residues, 16S rDNA amplified, cloned, and sequenced. Identification abbreviations as in *a*. ^cNP: Not processed. ^dND: Not determined. ^e0: no readable sequences recovered. ^fNA: Not amplifiable; PCR amplification was unsuccessful.

Sequencing dust rain DNA reveals diverse compositions. Because the majority of bacteria would not have been cultured^{28,29}, we made plasmid libraries from amplified 16S rDNA extracted from rain residues. Individual clones were sequenced from each rain to yield a total of 164 sequences (Table 2, see Supplementary Table S2). To understand how the microbiota of dust rains might be distinct from clean rains and dry depositions of desert dust, we selected 4 distinct rains for more extensive sequencing from their 16S libraries: one appearing to be clean rain and hail, 15 November 2011, one having a canonical North Africa backward trajectory, 24 December 2011, one of uncertain origin and high CFU, 25 October 2011, and one of uncertain origin and low CFU, 26 October 2011 (Fig. 1). Across 5 dates, 33 clean rain sequences displayed little diversity, matching only class Burkholderiales of Betaproteobacteria. In strong contrast, across 10 dates, dust rains displayed great diversity, with 101 of 131 having matches within 5 classes of Proteobacteria, and the remainder having matches to 7 Actinobacteria, 6 Bacteroidetes, 4 Cyanobacteria, and 13 Firmicutes. In accord with the DGGE banding, each rain presented a different collection of bacterial sequences, and 3 sequenced DGGE bands from clean and dust rains matched the same genera as sequences found in their cloned 16S libraries (see Supplementary Table S4).

Fungal 18S rDNA was amplified from rain residues and yielded 35 sequences matching Ascomycota and Basidiomycota (Table 3, see Supplementary Table S3). In clean rains were found only Basidiomycota classes Agaricomycetes and Tremellomycetes and matches to uncultured eukaryotes. Interestingly, fungal sequences from the clean rain of 15 November 2011 matched only uncultured eukaryotes. Many dust rain 18S sequences matched Basidiomycota classes Agaricomycetes and Tremellomycetes, and unlike clean rain sequences, many Ascomycota class Dothideomycetes. A few sequences matched other classes of Ascomycota and Basidiomycota, and two matched uncultured eukaryotes.

All sequences were categorized by genus, and rains were analyzed for specific and shared genera in different groupings (Fig. 3). Each rain selected for more 16S rDNA sequencing included genera specific to that rain (Fig. 3a). Interestingly, the rain of 24 December 2011, a dust rain with a backward trajectory passing over North Africa, had the greatest diversity, Chao1 index = 47.17 and Shannon index = 2.75, and the clean rain of 15 November 2011 had the least diversity, Chao1 index = 2 and Shannon index = 0.67 (Table 4). Analysis of separately combined clean and combined dust rain microbial composition with regard to sequences found in cultured or uncultured bacteria indicated both specific and shared genera. Dust rain contained more specific genera, and its diversity of uncultured genera was higher, Chao1 index = 88.1 and Shannon index = 3.38, than uncultured genera found in clean rain, Chao1 index = 4 and Shannon index = 1.07 (Table 4). Relatively few sequences were shared in cultured and uncultured groupings. Although many fewer fungal sequences were obtained, similar results were observed (Fig. 3c; Table 4). In order to determine whether microbial compositions

Rain date	Matches to classes in cultured fungi ^a	Matches to classes in uncultured fungi ^b
20110923	1 Dothideo, 1 Sordario	1 Dothideo
20110924	1 Dothideo	1 Agaricomycetes, 1 Basidiomycota
20111001	1 Dothideo	NP ^c
20111025	ND ^d	1 Dothideo, 1 Basidiomycota, 1 uncultured
20111026	1 Dothideo	2 Basidiomycota
20111103	ND ^d	1 Agaricomycetes, 2 uncultured
20111114	1 Dothideo	1 Dothideo
20111115	ND ^d	4 uncultured
20111224	1 Dothideo, 1 Eurotio	2 Dothideo, 1 Tremellomycetes
20120111	1 Eurotio, 1 Sordario	NP ^c
20120113	1 Dothideo	2 Agaricomycetes, 1 Dothideo, 2 Tremellomycetes
20120208	1 Dothideo	No clones found
20120215	ND ^d	NA ^e
20120314	ND ^d	NA ^e
20120402	1 Eurotio	2 Tremellomycetes
20120420	2 Dothideo	1 Agaricomycetes, 2 Dothideo, 1 Eurotio, 1 Malasseziomycetes, 2 Tremellomycetes, 1 uncultured
20120430	1 Dothideo	NA ^e
20120502	1 Dothideo, 1 Eurotio	NA ^e
20120529	2 Dothideo, 1 Eurotio	NA ^e
20121220	2 Dothideo, 1 Eurotio	0 ^f
20121221	1 Dothideo	2 Agaricomycetes

Table 3. Numbers of fungal sequences obtained from cultures and uncultured total DNA. ^aRain residues were cultured on Reasoner's 2A agar and sample colonies identified by sequencing of amplified 18S rDNA. Identification abbreviations are Dothideo: Dothideomycetes, Eurotio: Eurotiomycetes, Sordario: Sordariomycetes. ^bDNA was extracted from rain residues, 18S rDNA amplified, cloned, and sequenced. Identification abbreviations as in *a*. ^cNP: Not processed. ^dND: Not determined. ^eNA: Not amplifiable. PCR amplification was unsuccessful. ^f0: no clones found.

correlated with backward trajectories, cultured and uncultured bacteria genera were categorized North African and dusty, Mediterranean and dusty, other origins and dusty, or clean of all origins (Fig. 3d). The North African and Mediterranean genera were probably shared, mostly because their backward-trajectories were shared, yet many genera found in dust rains of other and uncertain origins were specific.

Discussion

Although the biogeochemistry of dust rains is an active area of research^{18,19}, the microbiota of dust rains is much less studied. To our knowledge, this is the first study of the microbiota of dust rains in the eastern Mediterranean. Thus, our results may help address important gaps in the literature concerning airborne microbial communities, their origins, and possible ecological roles. Interestingly, we found each dust rain to display a different set of characteristics, that the composition of dust rain was significantly more diverse than that of clean rain, and that each rain appears to deliver a different assemblage of microorganisms. Dust rains with backward trajectories passing over North Africa or the Mediterranean shared relatively few bacterial genera with dust rains of other origins. The heaviest deposition within one day, of 4 g m⁻², occurred in a light rain of 0.5 mm. This compares to yearly desert dust depositions estimated as 30–60 g m⁻² yr⁻¹ in the eastern Mediterranean³¹. Although the phenomenon of dust rain is often dramatic, heavy rainfalls depositing significant masses of particulate matter, especially at night, may be so diluted and washed that they pass without remark²⁰. The distinction between clean and dust rains, stark as it sometimes appears, is too subjective to predict microbial composition and survival, and some clean rains also had significant particulate matter. Nonetheless, backward trajectories were consistent with most apparently dust rains acquiring their particulate matter from arid regions of North Africa and moisture from the Mediterranean Sea with subsequent orographic precipitation upon encountering Mount Lebanon³², similar to what has been observed in Anatolia³³. Backward trajectories also suggest other mechanisms occur, including complex air mixtures and solids originating from arid regions in Southwest Asia, consistent with known weather patterns and dust sources in the region^{11,31,33,34}. Importantly, clean rain backward trajectories avoided or skirted arid regions. Solid media cultures and DGGE analysis revealed most rains to have distinct microbial compositions, sometimes with prominent species. This is consistent with each rain acquiring microorganisms from different and possibly multiple sources integrated along complex paths that likely include non-arid terrestrial and marine surfaces³².

Most culture-based approaches likely miss a large proportion of species^{28,29}, and our use of one type of solid media, R2A, limits ascertainable compositions. Thus, the CFU and diverse compositions observed in solid media cultures served primarily to indicate that many microorganisms had not been sterilized by desiccation or solar radiation (see Supplementary Fig. S3). Comparison of our observed CFU m⁻² with published values of aerosolized desert dust³⁵ CFU ranging as high as 15,000 m⁻³ and cloud water²⁵ CFU approximately 400 ml⁻¹ is




Figure 2. Denaturing Gradient Gel Electrophoresis of Bacterial 16S of Selected Rain Residues. Lanes are labelled by the date of the rain (YYYY.MM.DD). Clean rain headings are italicized, and headings of rains selected for greater sequencing are bold. Lane MI is a mix of *Escherichia coli*, *Bacillus cereus*, *Proteus vulgaris* and *Salmonella enterica*; lane MII is *E. coli* and *B. cereus*; Lane MIII is *P. vulgaris* and *S. enterica*. Prominent bands that were excised for sequencing are numbered directly above. The image was generated by Quantity One (version 4.6.3, Bio-Rad Laboratories) with adjustments in Photoshop CS6 (Adobe Systems).




Figure 3. Comparisons of Rains. Venn diagrams showing number of specific and shared genera identified. (a) all identified bacterial genera of the four rains of more intense study (25 October 2011, 26 October 2011, 15 November 2011, and 24 December 2011). (b) Bacterial genera identified in combined clean rains and combined dust rains separated by culture-dependent and culture-independent isolation. (c) Fungal genera identified in combined clean rains and combined dust rains separated by culture-dependent and culture-independent isolation. (d) All identified bacteria categorized according to Table 1 as being dust rains with back-trajectories including North Africa, the Mediterranean, neither North Africa or the Mediterranean, or clean rains of all origins. Images were generated using Venny 2.1 (<http://bioinfogp.cnb.csic.es/tools/venny/index.html>).

Grouped sets of sequences ^a	Sequences ^b	Genera ^c	Chao1 ^d	Shannon ^e
20111025 uncultured bacteria	28	10	22.5	1.99
20111026 uncultured bacteria	39	9	17	1.68
20111115 uncultured bacteria	28	2	2	0.67
20111224 uncultured bacteria	32	19	47.17	2.75
clean uncultured bacteria	33	4	4	1.07
dust uncultured bacteria	131	44	88.1	3.38
clean uncultured fungi	13	6	6.13	1.71
dust uncultured fungi	22	14	39	2.46
dust NAF origins uncultured bacteria	57	27	60	3.10
dust Med origins uncultured bacteria	57	30	75.13	3.15
dust other origins uncultured bacteria	71	18	30.25	2.47

Table 4. Number of genera found in uncultured microbes and diversity indices. ^aSequences of uncultured bacteria and fungi were grouped by date of rain, rain classification as clean or dusty as in Table 1, and having back-trajectories passing over North Africa or the Mediterranean as in Table 1. ^bThe number of readable sequences in each category. ^cThe number of genera in each category. ^dChao1 index calculated with respect to genera. ^eShannon index calculated with respect to genera.

difficult, but it suggests rains may scavenge large air volumes, or as discussed below, promote the growth of microorganisms resident in dust and clouds.

DGGE profiles, plate cultures, and sequence data all are consistent with each dust rain delivering a different composition of organisms. The selection imposed by culturing, especially on only one medium, may obscure distinctions between rains. Conversely, cultures revealed many Gram-positive bacteria that would otherwise not have been noted without extensive sequencing. Culture-independent analysis was attractive based on the significant masses of DNA extracted and difficulties of culturing most microorganisms^{28,29}. Although it is difficult to rule out bias resulting from DNA extractions, amplification, and sequencing, most DNA samples were amplifiable and yielded readable sequences matching far more Alphaproteobacteria and Betaproteobacteria than did cultures. Interestingly, we were unable to amplify bacterial DNA from some light rains, consistent with DNA damage incurred from solar radiation or desiccation of dust before acquisition by clouds. Desert dust in the Mediterranean may sometimes be lofted and precipitated under continuous cloud cover³⁶, and sometimes lofted before cloud formation³⁷, the extremes of which may yield very different survival rates for many species.

Our observations are consistent with observed microbial compositions and variability of air^{38,39}, rains and clouds^{23,40–43}, desert dust^{9,10,15,35,44–47}, and dust rains in the Alps^{21,22}. Our data support the suggestion of Chuvochina *et al.*²¹ that delivered bacterial assemblages are affected more by the peculiarities of the transport and precipitation event more than by the source of particulate matter. Thus, each rain delivers a different, complex assemblage of microorganisms, and the microbial composition of dust rains needs further study. Our results encourage future studies to employ more sophisticated collection schemes, record meteorological data, conduct mineralogical analysis, use next-generation sequencing, and apply advanced statistical analysis.

In contrast to dust rains, the clean rain of 15 November 2011 appears to have a very low diversity (Table 4) and a high proportion of plant-associated members of family Oxalobacteraceae, consistent with its European backward trajectory and what is observed in clouds^{23,40,41,43}. Interestingly, this torrential rain included hail and had the highest observed CFU, approximately 50,000 m⁻² (~500 CFU ml⁻¹). Hailstones have been observed to contain diverse bacteria, including Burkholderiales, which are often plant-associated and capable of nucleating ice, and can have CFU exceeding 5,000 ml⁻¹^{43,48}. The few library sequences of another clean rain on 3 November 2011 matched only members of order Burkholderiales, mostly *Ralstonia*, a common environmental genus⁴⁹. Particularly interesting is the dominance of Oxalobacteraceae, a family that has been observed in the upper troposphere⁴¹. The sequences of cultured bacteria, DGGE bands, and 16S library sequences of clean rains on 3 November 2011 and 15 November 2011 all corroborate the dominant representation of *Ralstonia* and *Herbaspirillum* genera of order Burkholderiales in these rains.

In strong contrast to the clean rains, the dust rain of 24 December 2011, which had a canonical backward trajectory consistent with North African origins to its particulate matter, had sequences matching diverse bacterial phyla, including Gram-positive Actinobacteria and Firmicutes, as well as Bacteroidetes and several classes of Proteobacteria. Its diversity was high (Table 4), and it contained many specific bacterial genera (Fig. 3a). These observations, in particular the presence of Firmicutes, Actinobacteria, Firmicutes, Bacteroidetes, Alphaproteobacteria, and Burkholderiales (Betaproteobacteria), are similar to those observed in dust events^{9,10} and are consistent with soils from arid regions of North Africa^{15,35,44}, including dust collected in the central and eastern Mediterranean^{45–47}. Library and colony fungal sequences from this dust rain matched *Aspergillus*, *Cladosporidium*, and *Cryptococcus* fungal species, which have been found in dry dust depositions^{9,35,45}. Intriguingly, many sequences match Gram-negative bacteria less typical of desert dust, such as aquatic Rhodobacteriaceae species, *Roseovarius crassostreae* and *Loktanella hongkongensis*, suggesting dust rains deliver species representing both desert dusts and rain.

In the uncertain-origin dust rains of 25 October 2011 and 26 October 2011, only sequences matching Alphaproteobacteria and Betaproteobacteria were found. Interestingly, they shared few bacterial genera with the dust rain of 24 December 2011. The lack of Gram-positive representation in 16S sequences in these two dust rains

despite finding matches to Gram-positive bacteria in cultures emphasizes the utility of culturing when interests are directed to specific, culturable species. The representation of Burkholderiales and Sphingomonas sequences suggests a large proportion of non-dust species. Their different sequence compositions within Proteobacteria highlight the variability between events closely spaced in time.

Few reports of the microbial composition of dust rains are available^{21,22}. In a study of three rains with and three without Saharan dust influence, Peter *et al.*²² find rains of Atlantic and continental origin dominated by Betaproteobacteria of genera *Massilia* and Sphingobacteria and rains with Saharan dust dominated by Gammaproteobacteria. Alphaproteobacteria and Gammaproteobacteria were found in both non-dust and dust influenced rains. *Massilia* dominated two of three dust influenced rains, and Bacilli were found only in dust rains. In contrast, we find few Gammaproteobacteria in dust rains, *Massilia* and Sphingobacteria only in dust rains, and clean rains dominated by Betaproteobacteria of Burkholderiaceae and Oxalobacteraceae. Our dust rains have similar Chao1 and Shannon values to the dust rains of Peter *et al.*²², yet our clean rains are much less diverse than our dust rains. A metagenomic study of Saharan dust in Alpine snow²¹, which included one dust rain in Grenoble, France, and a Saharan sand sample, also found great differences between samples. Actinobacteria (*Blastococcus*), Alphaproteobacteria (*Sphingomonas* and *Rubellimicrobium*), Bacteroidetes, Cyanobacteria, and Deinococcus-Thermus were particularly common. Compared to Chuvochina *et al.*²¹, we find more Betaproteobacteria and Firmicutes, yet fewer Deinococcus-Thermus. Our intensely studied dust rain of 24 December 2011 has Chao1 and Shannon values within the ranges they report for their libraries. Our results are not inconsistent with their findings, especially their observation that individual rains can deliver very different assemblages.

The data herein suggest that dust rains are an effective means by which microorganisms from desert dust and clouds are transported and delivered to surface habitats. More broadly, the culture and culture-independent sequencing, DGGE analysis, and solid-culture observations herein support the conclusion that each dust rain in the eastern Mediterranean likely delivers a unique collection of microorganisms, and that an understanding of their ecological significance, especially in expectation of natural and anthropogenic desertification and climate change, will require comprehensive examination. Extensive sampling of potential sources^{50,51}, regular surveys, and standardization of methods will be needed.

One could imagine that transport in a raincloud would confer protection against desiccation and solar radiation and promote the survival of species lofted from arid soils. Less clear is whether dust rains offer a qualitatively different means of microbial transport beyond the efficiency of wet terrestrial deposition. Is dust rain microbiology a simple combination of that of rainclouds and desert dust? Fahlgren *et al.*³⁸ speculate bacteria can grow in the atmosphere. Dust microorganisms can rapidly colonize aquatic habitats⁵² and dust provides nutrients to oligotrophic aquatic habitats^{53–56}. Peter *et al.*²² find an approximately 100-fold increase in cell counts upon wet deposition of desert dust in Alpine lakes. Thus, dust rains may sustain and deliver microbiota that are complex products of revived desert species and fertilized cloud species. One might further speculate that the mixing of desert dust and clouds can create persistent habitats with water, soil, air, and light that propagate airborne microbial ecologies¹².

Methods

Rain collection and processing. For each collection, a clean, new, polyethylene tarpaulin in a 2.1 square meter frame was placed on the southwest corner of the American University of Beirut biology building roof (33° 54' 8" N, 35° 28' 45" E, approximately 20 m above sea level) before expected rains. Each rain was collected 4–18 hour after falling and filtered through a 0.22 µm cellulose acetate membrane filter. No collection exceeded 24 h. One-sixteenth of each filter with residue was reserved for a glycerol stock, and the remainder divided into two portions, one portion stored in 1 ml of DNA extraction buffer EB (20 mM Tris-HCl pH 7.8, 50 mM ethylenediamine tetraacetic acid [EDTA], and 20 mM NaCl) in screw-cap, 2 ml, polypropylene microcentrifuge tubes for later extraction. The other portion was further divided in two, one portion separated from the filter, washed, and dried 24 h at 65 °C for mass measurement on an analytical balance, and the other stored in 0.5 ml EB.

HYSPLIT backward trajectories. For each collected rainfall, wind backward trajectories were determined using the online Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPLIT) model⁷ from the US National Oceanic and Atmospheric Administration using Global Data Assimilation System meteorological data (http://ready.arl.noaa.gov/HYSPLIT_traj.php). Latitude (33.902° N) and longitude (35.479° E), total run time (at least 72 hrs), and altitudes (500 m, 1000 m, 2000 m) were entered for each rainfall date. The output was a map with the trajectory of the wind starting from the day of rainfall (Fig. 1, see Supplementary Fig. 2).

Rain culturing. For each rainfall, the filter residue was stored in glycerol (20% final concentration) at –70 °C. Dilutions of each sample were plated on Reasoner's 2A (R2A) agar using glass beads and cultured at room temperature. A small, non-random variety of colonies were restreaked for amplification and sequencing of bacterial 16S rDNA and fungal 18S rDNA as described below.

DNA extraction and quantification. Total DNA was extracted from filters of rain residues using bead beating. To each 2 ml, screw-cap microcentrifuge tube containing filter and residue was added sodium dodecyl sulfate and RNaseA to final concentrations of 3% and 10 µl/ml, respectively. Spherical glass beads of 1 mm diameter were added to fill remaining volume. The samples were beaten at maximum speed on a Mini-Beadbeater-1 (Biospec Products, Bartlesville, OK, USA) for 2 min followed by placement on ice for 1 min. The beating was repeated twice, with an additional beating of 1 min. The tubes were centrifuged, supernatant collected, 500 µl of extraction buffer EB added, beaten for 2 min, and centrifuged. First and second supernatants of each sample were combined and extracted twice with phenol-chloroform-isoamyl alcohol (25:24:1) and once with chloroform.

5. Engelstaedter, S., Tegen, I. & Washington, R. North African dust emissions and transport. *Earth-Sci. Rev.* **79**, 73–100 (2006).
6. Legrand, M., Plana-Fattori, A. & N'doume, C. Satellite detection of dust using the IR imagery of Meteosat 1. Infrared difference dust index. *J. Geophys. Res.-Atmos.* **106**, 18251–18274 (2001).
7. Draxler, R. R. & Hess, G. D. An overview of the HYSPLIT₄ modelling system for trajectories. *Austr. Meteorol. Mag.* **47**, 295–308 (1998).
8. Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E. & Gill, T. E. Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product. *Rev. Geophys.* **40**, 2–1–2–31 (2002).
9. Griffin, D. W. Atmospheric movement of microorganisms in clouds of desert dust and implications for human health. *Clin. Microbiol. Rev.* **20**, 459–477 (2007).
10. Kellogg, C. A. & Griffin, D. W. Aerobiology and the global transport of desert dust. *Trends Ecol. Evol.* **21**, 638–644 (2006).
11. Varga, G., Újvári, G. & Kovács, J. Spatiotemporal patterns of Saharan dust outbreaks in the Mediterranean Basin. *Aeolian Research* **15**, 151–160 (2014).
12. Womack, A. M., Bohannan, B. J. & Green, J. L. Biodiversity and biogeography of the atmosphere. *Philos. T. Roy. Soc. B.* **365**, 3645–3653 (2010).
13. Yamaguchi, N., Ichijo, T., Sakotani, A., Baba, T. & Nasu, M. Global dispersion of bacterial cells on Asian dust. *Sci. Rep.* **2**, 525 (2015).
14. Psenner, R. Living in a dusty world: airborne dust as a key factor for alpine lakes. *Water Air Soil Poll.* **112**, 217–227 (1999).
15. Giongo, A. *et al.* Microbial hitchhikers on intercontinental dust: high-throughput sequencing to catalogue microbes in small sand samples. *Aerobiologia* **29**, 71–84 (2013).
16. Homer. *Iliad*, 11.54; 16.459 (1250 BCE).
17. Cicero, M. T. *De divinatione* ii. 27 (44 BCE).
18. Guerzoni, S. *et al.* The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea. *Prog. Oceanogr.* **44**, 147–190 (1999).
19. Avila, A., Alarcon, M. & Queralt, I. The chemical composition of dust transported in red rains—its contribution to the biogeochemical cycle of a Holm Oak forest in Catalonia (Spain). *Atmos. Environ.* **32**, 179–191 (1998).
20. Sala, J. Q., Cantos, J. O. & Chiva, E. M. Red dust rain within the Spanish Mediterranean area. *Climatic Change* **32**, 215–228 (1996).
21. Chuvochina, M. S. *et al.* Community variability of bacteria in alpine snow (Mont Blanc) containing Saharan dust deposition and their snow colonisation potential. *Microbes Environ.* **26**, 237–247 (2011).
22. Peter, H., Hörtnagl, P., Reche, I. & Sommaruga, R. Bacterial diversity and composition during rain events with and without Saharan dust influence reaching a high mountain lake in the Alps. *Environ. Microbiol. Reports* **6**, 618–624 (2014).
23. Delort, A. M. *et al.* A short overview of the microbial population in clouds: Potential roles in atmospheric chemistry and nucleation processes. *Atmos. Res.* **98**, 249–260 (2010).
24. Whitman, W. B., Coleman, D. C. & Wiebe, W. J. Prokaryotes: the unseen majority. *Proc. Natl. Acad. Sci. USA* **95**, 6578–6583 (1998).
25. Amato, P. *et al.* An important oceanic source of micro-organisms for cloud water at the Puy de Dôme (France). *Atmos. Environ.* **41**, 8253–8263 (2007).
26. Gilbert, J. A. & Dupont, C. L. Microbial metagenomics: beyond the genome. *Ann. Rev. Mar. Sci.* **3**, 347–371 (2011).
27. Pace, N. R. A molecular view of microbial diversity and the biosphere. *Science* **276**, 734–740 (1997).
28. Amann, R. L., Ludwig, W. & Schleifer, K. H. Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. *Microbiol. Rev.* **59**, 143–169 (1995).
29. Ward, D. M., Weller, R. & Bateson, M. M. 16S rRNA sequences reveal numerous uncultured microorganisms in a natural community. *Nature* **345**, 63–65 (1990).
30. Forney, L. J., Zhou, X. & Brown, C. J. Molecular microbial ecology: land of the one-eyed king. *Curr. Opin. Microbiol.* **7**, 210–220 (2004).
31. Ganor, E. & Foner, H. A. (1996). The mineralogical and chemical properties and the behaviour of aeolian Saharan dust over Israel In *The Impact Of Desert Dust Across The Mediterranean* (eds Guerzoni, S. & Chester, R.) 163–172 (Kluwer Academic Publishers, 1996).
32. Gat, J. R. & Carmi, I. Effect of climate changes on the precipitation patterns and isotopic composition of water in a climate transition zone: case of the Eastern Mediterranean Sea area. *IAHS Spec. Publ.* **168**, 513–523 (1987).
33. Kubilay, N., Nickovic, S., Moulin, C. & Dulac, F. An illustration of the transport and deposition of mineral dust onto the eastern Mediterranean. *Atmos. Environ.* **34**, 1293–1303 (2000).
34. Saliba, N. A., El Jam, F., El Tayar, G., Obeid, W. & Roumie, M. Origin and variability of particulate matter (PM10 and PM2.5) mass concentrations over an Eastern Mediterranean city. *Atmos. Res.* **97**, 106–114 (2010).
35. Kellogg, C. A. *et al.* Characterization of aerosolized bacteria and fungi from desert dust events in Mali, West Africa. *Aerobiologia* **20**, 99–110 (2004).
36. Tilev-Tanriover, Ş. & Kahraman, A. Saharan dust transport by Mediterranean cyclones causing mud rain in Istanbul. *Weather* **70**, 145–150 (2015).
37. Levin, Z., Ganor, E. & Gladstein, V. The effects of desert particles coated with sulfate on rain formation in the eastern Mediterranean. *J. Appl. Meteorol.* **35**, 1511–1523 (1996).
38. Fahlgren, C., Hagström, Å., Nilsson, D. & Zweifel, U. L. Annual variations in the diversity, viability, and origin of airborne bacteria. *Appl. Environ. Microbiol.* **76**, 3015–3025 (2010).
39. Fierer, N. *et al.* Short-term temporal variability in airborne bacterial and fungal populations. *Appl. Environ. Microbiol.* **74**, 200–207 (2008).
40. Cho, B. C. & Jang, G. I. Active and diverse rainwater bacteria collected at an inland site in spring and summer 2011. *Atmos. Environ.* **94**, 409–416 (2014).
41. DeLeon-Rodriguez, N. *et al.* Microbiome of the upper troposphere: Species composition and prevalence, effects of tropical storms, and atmospheric implications. *Proc. Natl. Acad. Sci. USA* **110**, 2575–2580 (2013).
42. Kourtev, P. S., Hill, K. A., Shepson, P. B. & Konopka, A. Atmospheric cloud water contains a diverse bacterial community. *Atmos. Environ.* **45**, 5399–5405 (2011).
43. Šantl-Temkiv, T. *et al.* Hailstones: a window into the microbial and chemical inventory of a storm cloud. *PLoS One* **8**, e53550 (2013).
44. Chanal, A. *et al.* The desert of Tataouine: an extreme environment that hosts a wide diversity of microorganisms and radiotolerant bacteria. *Environ. Microbiol.* **8**, 514–525 (2006).
45. Katra, I. *et al.* Richness and diversity in dust stormborne biomes at the Southeast Mediterranean. *Sci. Rep.* **4**, 5265 (2014).
46. Rosselli, R. *et al.* Microbial immigration across the Mediterranean via airborne dust. *Sci. Rep.* **5**, 16306 (2015).
47. Polymenakou, P. N., Mandalakis, M., Stephanou, E. G. & Tselepidis, A. Particle size distribution of airborne microorganisms and pathogens during an Intense African dust event in the eastern Mediterranean. *Environ Health Perspect.* **116**, 292–296 (2008).
48. Bowers, R. M. *et al.* Characterization of airborne microbial communities at a high-elevation site and their potential to act as atmospheric ice nuclei. *Appl. Environ. Microbiol.* **75**, 5121–5130 (2009).
49. Ryan, M. P., Pembroke, J. T. & Adley, C. C. Genotypic and phenotypic diversity of *Ralstonia pickettii* and *Ralstonia insidiosa* isolates from clinical and environmental sources including High-purity Water. Diversity in *Ralstonia pickettii*. *BMC Microbiol.* **11**, 194 (2011).
50. Martiny, J. B. H. *et al.* Microbial biogeography: putting microorganisms on the map. *Nat. Rev. Microbiol.* **4**, 102–112 (2006).

51. Barberán, A. *et al.* Continental-scale distributions of dust-associated bacteria and fungi. *Proc. Natl. Acad. Sci. USA* **112**, 5756–5761 (2015).
52. Hervás, A., Camarero, L., Reche, I. & Casamayor, E. O. Viability and potential for immigration of airborne bacteria from Africa that reach high mountain lakes in Europe. *Environ. Microbiol.* **11**, 1612–1623 (2009).
53. Herut, B. *et al.* Response of East Mediterranean surface water to Saharan dust: On-board microcosm experiment and field observations. *Deep-Sea Res. Pt. II* **52**, 3024–3040 (2005).
54. Morales-Baquero, R., Pulido-Villena, E. & Reche, I. Atmospheric inputs of phosphorus and nitrogen to the southwest Mediterranean region: Biogeochemical responses of high mountain lakes. *Limnol. Oceanogr.* **51**, 830–837 (2006).
55. Pulido-Villena, E., Reche, I. & Morales-Baquero, R. Evidence of an atmospheric forcing on bacterioplankton and phytoplankton dynamics in a high mountain lake. *Aquat. Sci.* **70**, 1–9 (2008).
56. Reche, I. *et al.* Effect of Saharan dust inputs on bacterial activity and community composition in Mediterranean lakes and reservoirs. *Limnol. Oceanogr.* **54**, 869–879 (2009).
57. Muyzer, G., De Waal, E. C. & Uitterlinden, A. G. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* **59**, 695–700 (1993).
58. Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. 16S ribosomal DNA amplification for phylogenetic study. *J. Bact.* **173**, 697–703 (1991).
59. Griffin, D. W., Kellogg, C. A., Peak, K. K. & Shinn, E. A. A rapid and efficient assay for extracting DNA from fungi. *Letts. Appl. Microbiol.* **34**, 210–214 (2002).
60. Sambrook, J., Fritsch, E. F. & Maniatis, T. *Molecular Cloning* 2nd edn, Vol. 2, 1.43–1.84. (Cold Spring Harbor Laboratory Press 1989).
61. Oliveros, J. C. *Venny. An interactive tool for comparing lists with Venn's diagrams.* (2007–2015) Available at: <http://bioinfo.cnb.csic.es/tools/venny/index.html>. (Accessed 02 January 2016).
62. Hughes, J. B., Hellmann, J. J., Ricketts, T. H. & Bohannan, B. J. M. Counting the uncountable: Statistical approaches to estimating microbial diversity. *Appl. Environ. Microbiol.* **67**, 4399–4406 (2001).

Acknowledgements

This work was supported by the Lebanese National Council for Scientific Research. We thank Nicole Raad for assistance with statistics. The authors acknowledge the US National Oceanic and Atmospheric Administration (NOAA) Air Resources Laboratory (ARL) for the provision of the HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) model via NOAA ARL READY website (<http://ready.arl.noaa.gov>) used in this publication.

Author Contributions

C.A.S. obtained funding and supervised the project. G.N.I. and C.A.S. designed experiments and collected rainfalls. G.N.I. conducted experiments, analysed sequences, and drafted the manuscript. C.A.S. prepared figures and edited the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Itani, G. N. and Smith, C. A. Dust Rains Deliver Diverse Assemblages of Microorganisms to the Eastern Mediterranean. *Sci. Rep.* **6**, 22657; doi: 10.1038/srep22657 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>