

# SCIENTIFIC REPORTS



OPEN

## Soybean cyclophilin GmCYP1 interacts with an isoflavonoid regulator GmMYB176

Hemanta Raj Mainali<sup>1</sup>, Arun Kumaran Anguraj Vadivel<sup>1,2</sup>, Xuyan Li<sup>2,3</sup>, Mark Gijzen<sup>1,2</sup> & Sangeeta Dhaubhadel<sup>1,2</sup>

Received: 02 September 2016

Accepted: 23 November 2016

Published: 11 January 2017

Cyclophilins (CYPs) belong to the immunophilin superfamily with peptidyl-prolyl *cis-trans* isomerase (PPIase) activity. They catalyze the interconversion of the *cis*- and *trans*-rotamers of the peptidyl-prolyl amide bond of peptides. A yeast-two-hybrid screening using the isoflavonoid regulator GmMYB176 as bait identified GmCYP1 as one of the interacting proteins in soybean embryos. GmCYP1 localizes both in the nucleus and cytoplasm, and interacts *in planta* with GmMYB176, in the nucleus, and with SGF141 (a soybean 14-3-3 protein) in the nucleus and the cytoplasm. GmCYP1 contains a single cyclophilin-like domain and displays a high sequence identity with other plant CYPs that are known to have stress-specific function. Tissue-specific expression of *GmCYP1* revealed higher expression in developing seeds compared to other vegetative tissues, suggesting their seed-specific role. Furthermore, *GmCYP1* transcript level was reduced in response to stress. Since isoflavonoids are involved in plant stress resistance against biotic and abiotic factors, the interaction of GmCYP1 with the isoflavonoid regulators GmMYB176 and 14-3-3 protein suggests its role in defense in soybean.

Cyclophilins (CYPs) are a group of proteins that possess peptidyl prolyl *cis/trans* isomerase (PPIase) activity. They are involved in protein folding by interconverting the *cis*- and *trans*-rotamers of the peptidyl prolyl amide bond of peptides, and are broadly classified into three major classes: parvulins, FK506 binding proteins (FKBP) and cyclophilins<sup>1</sup>. FKBP and CYPs are collectively called immunophilins as they were originally identified as receptors for immunosuppressive drugs, FK506 and cyclosporine A, respectively<sup>2-4</sup>. CYPs are present in a wide range of organisms, from archaea, bacteria to plants and animals<sup>5-7</sup>. Genome-wide analyses of *CYP* genes in various organisms revealed disparity in the number of genes, ranging from 8 to 16 in organisms such as *Drosophila*<sup>8</sup>, *Caenorhabditis elegans*<sup>9</sup>, *Saccharomyces cerevisiae*<sup>10</sup>, and human<sup>8</sup>. A large number of studies focussed on human CYP, hCYP, have shown its crucial role in protein folding, signal transduction, cell signaling, regulation of gene expression, immune response, and disease conditions<sup>11-13</sup>.

The first plant CYPs were identified concomitantly in 1990, from tomato (*Lycopersicon esculentum*), maize (*Zea mays*), and oilseed rape (*Brassica napus*)<sup>14</sup>. With the advancement in genome sequencing technology, and availability of plant genome sequence data in public domain, the identification and characterization of plant CYPs has progressed significantly in the recent years. Compared to other organisms, photosynthetic organisms contain significantly higher number of CYPs; 62 in soybean<sup>15</sup>, 35 in *Arabidopsis*<sup>16,17</sup>, 28 in rice<sup>17,18</sup>, and 26 in *Chlamydomonas*<sup>19</sup>.

In order to combat biotic and abiotic stress, sessile organisms like plants have developed several sophisticated mechanisms at the cellular and molecular levels<sup>20,21</sup>. One of the consequences of abiotic stress is the denaturation and aggregation of cellular proteins leading to cell death. The chaperone-like activity of CYPs and their role in the rate-limiting step of protein folding by peptidyl prolyl bond isomerization<sup>22</sup> is associated with their involvement in stress responses. Expression of many plant CYPs is induced in response to stress suggesting their possible function in stress tolerance. For example, expression of the *Arabidopsis CYP*, *ROTAMASE CYCLOPHILIN 1 (ROC1)*, increases upon wounding<sup>23</sup>. Similarly, maize and bean CYP gene expression increases in response to heat stress, wounding, high salinity, or low temperature<sup>24</sup>. *Solanum commersonii CYP* gene expression is also up-regulated by low temperature, abscisic acid, drought, or wounding<sup>25</sup>. Pepper CYPs are differentially regulated during abiotic

<sup>1</sup>Department of Biology, University of Western Ontario, 1151 Richmond St, London, ON, N6A 5B7, Canada. <sup>2</sup>London Research and Development Centre, Agriculture and Agri-Food Canada, 1391 Sandford Street, London, ON, N5V 4T3, Canada. <sup>3</sup>College of Plant Science, Jilin University, Changchun, 130062, China. Correspondence and requests for materials should be addressed to S.D. (email: sangeeta.dhaubhadel@canada.ca)

stress and pathogen infection<sup>26</sup>. Ectopic expression of *Thellungiella halophila* CYP, *ThCYP1*, in fission yeast and tobacco cells increased salt tolerance<sup>27</sup>. Transgenic *Arabidopsis* plants overexpressing pigeon pea CYP (*CcCYP1*) showed enhanced PPIase activity under stressed conditions, which correlated with their increased tolerance against drought, salinity and high temperature<sup>28</sup>. Similarly, overexpression of cotton CYP (*GhCYP1*) in tobacco plants conferred tolerance against salt stress and fire-blight disease<sup>29</sup>. Together, these findings clearly demonstrate a role for plant CYPs in stress tolerance.

Soybean (*Glycine max*) is a grain legume belonging to the family *Fabaceae*. Soybean seeds provide a major supply of oil, protein and beneficial plant natural compounds such as isoflavonoids and saponins. The soybean genome contains 88,647 predicted transcripts and 56,044 protein coding loci located on 20 different chromosomes<sup>30</sup>. Previously, we performed a genome-wide analysis of soybean CYPs and identified 62 CYP genes<sup>15</sup>. Among these, GmCYP1 has been shown to act as a “helper” to *Phytophthora sojae* RXLR effector Avr3b by activating its hydrolase activity in plant cells<sup>31</sup>. The protein-protein interaction between GmCYP1 and Avr3b was shown to be isoform-specific since GmCYP1 paralogs failed to interact with Avr3b. Here we present a molecular characterization of *GmCYP1* covering its sequence analysis, phylogeny, temporal and spatial expression, subcellular localization, and provide the evidence for its possible role in isoflavonoid biosynthesis and stress response in soybean.

## Results and Discussion

**Isolation, sequence analysis and phylogeny of GmCYP1.** GmCYP1 was identified in our Y2H screening as a protein that demonstrated protein-protein interaction with the isoflavonoid regulator GmMYB176. The Y2H assay was performed to identify GmMYB176-interacting proteins using GmMYB176 as the bait protein and proteins from soybean embryos (50–60 days after pollination) as prey. Of the several hundred yeast colonies screened, 6.5% contained a sequence corresponding to *GmCYP1* (accession #AF456323, locus Glyma.11G098700). *GmCYP1* is predicted to contain only one exon (519 bp), and is located on the long arm (q arm) of chromosome 11, approximately 16 Mb from the centromere. It encodes a single domain protein of 172 amino acid residues with a calculated molecular mass of 18.22 kDa and a pI of 8.69. The cyclophilin-like domain in GmCYP1 is predicted in between the amino acid residues 7 and 169.


In order to find sequences closely related to GmCYP1, a protein-protein BLAST (BLASTP) was performed using GmCYP1 as a query against the NCBI non-redundant protein database. A list of 12 high-scoring and previously characterized CYPs is shown in Table S1. Alignment of the deduced sequence of GmCYP1 with previously characterized CYPs from several different plant species, human, yeast, and two *Arabidopsis* multi-domain CYPs (AtCYP40 and AtCYP63) revealed two general features (Fig. 1). First, three amino acid residues that critically affect PPIase activity (R<sub>55</sub>, F<sub>60</sub> and H<sub>126</sub>)<sup>32</sup> are conserved in all CYPs aligned. Second, the tryptophan residue (W<sub>121</sub>) implicated in substrate cyclosporinA binding<sup>32,33</sup> is present in all of the CYPs studied except in the multi domain CYPs.

A phylogenetic analysis of GmCYP1 and other functionally characterized plant CYPs clustered GmCYP1 close to *GhCYP1*, *CcCYP1* and *PvCYP* from cotton, pigeon pea and common bean, respectively. Both *GhCYP1* and *CcCYP1* are known to have stress-specific function (Fig. 2). Overexpression of *GhCYP1* in tobacco conferred increased tolerance to biotic and abiotic stress<sup>29</sup>. Similarly, *Arabidopsis* plants overexpressing *CcCYP1* showed higher PPIase activity during stress and increased tolerance against multiple abiotic stresses as compared to control<sup>28</sup>. Differential accumulation of *PvCYP* transcripts in response to various external stimuli suggested that it may possess a stress-related function<sup>24</sup>. The high amino acid sequence identity of GmCYP1 with *GhCYP1* (96%), *CcCYP1* (81%) and *PvCYP* (96%), suggests similar possible functions of GmCYP1 in stress response.

**GmCYP1 localizes in the nucleus and the cytoplasm.** To further study the subcellular localization of GmCYP1, a translation fusion of GmCYP1 with YFP was created under the control of 35S promoter, and transiently expressed in *N. benthamiana* leaves. Although there was no predicted nuclear localization sequence in GmCYP1, confocal imaging of the GmCYP1-YFP-infiltrated tobacco leaves showed both nuclear and cytoplasmic localization (Fig. 3a). Nuclear localization of GmCYP1 was confirmed by co-expression of GmCYP1-YFP and NLS-CFP.

Molecules of size smaller than 20–40 kDa, such as ions, water, and small proteins, can pass through the nuclear pore complex by diffusion<sup>34</sup>, whereas movement of larger molecules (70 kDa or higher) entails an active transport system<sup>35</sup>, mediated by transport receptors and signal peptides<sup>36</sup>. The size of GmCYP1-YFP (45.22 kDa) is not considerably larger than the size of molecules that can pass through the nuclear pore complex by diffusion. It is also possible that heterologous protein GmCYP1-YFP, expressed in *N. benthamiana* is cleaved by the endogenous host proteases and only the cleaved YFP fragments are localized to the nucleus. The confirmation of the YFP signal in the nucleus arising from the intact GmCYP1-YFP and not from the cleaved product of a fusion protein was performed by Western blot analysis (Fig. 3b). Therefore, it is not clear whether GmCYP1-YFP localization in the nucleus was due to passive diffusion or to active transport. Regardless, its nuclear localization indicates a possible role in the regulation of gene expression.

**GmCYP1 interacts with GmMYB176 in planta.** GmCYP1 was identified as one of the interacting proteins of GmMYB176 in the Y2H assay. Protein chaperones often bind with misfolded bait proteins when a protein is overexpressed or heterologously expressed. To confirm that GmCYP1 is a true GmMYB176-interacting protein, we performed a targeted Y2H assay using GmCYP1 and GmMYB176. However, our targeted Y2H assay failed to verify the interaction between GmCYP1 with GmMYB176. This result led us to hypothesize that there may be an indirect interaction between GmCYP1 and GmMYB176 via involvement of protein (s) that may not be conserved between the species. Therefore, a BiFC assay<sup>37</sup> was carried to further investigate *in planta* interaction between GmCYP1 and GmMYB176. Translational fusions of GmMYB176 and GmCYP1 were created with N-terminal




**Figure 1. Multiple sequence alignment of deduced amino acid sequence of GmCYP1 with CYPs from other species.** Amino acid sequences of ROC3, BnCYP, GhCYP1, CcCYP1, ROC6, ROC1, ROC5, Cpr1, hCYP-D, hCYP-A, AtCYP40, AtCYP63 and GmCYP1 (refer to Table S1 for accession numbers) were aligned by ClustalW, and imported into BOXSHADE 3.21 for shading. Identical amino acids are shown in the dark box and similar amino acids are indicated by the grey box. Amino acid residues involved in PPIase activity (R<sub>55</sub>, F<sub>60</sub> and H<sub>126</sub>) (Zydowsky *et al.*<sup>32</sup>) and CsA binding (W<sub>121</sub>) (Liu *et al.*<sup>33</sup>; Zydowsky *et al.*<sup>32</sup>) are highlighted with green and red, respectively. Secondary structure is shown below the alignment. The relative positions of amino acids indicated for PPIase activity and CsA binding sites, and the secondary structure features are based on hCYP-A (Kallen *et al.*, 1991).

(YN) or C-terminal (YC) halves of YFP<sup>38</sup> and co-expressed in *N. benthamiana* leaf epidermal cells in the following combinations: (A) GmCYP1-YN and GmMYB176-YC, (B) GmMYB176-YN and GmCYP1-YC. The negative controls used for the experiment were co-expression of GmCYP1-YN and -YC only or GmCYP1-YC and -YN only. As shown in Fig. 4a, GmCYP1 interacts with GmMYB176 *in planta*, and the interaction between GmCYP1 and GmMYB176 was strong in the nucleus. Similar results were obtained for the reciprocal combinations. No YFP signal was detected during co-expression of control constructs -YC or -YN with GmCYP1-YN or GmCYP1-YC, respectively, confirming that YFP signal was due to the protein-protein interaction *in planta*. Furthermore, we measured the strength of interaction by using FRET approach. It is a powerful tool for non-invasive monitoring of protein-protein interactions that involves transfer of energy between two closely positioned fluorophores<sup>39</sup>, and has been widely used to determine the interactions between proteins in living plant cells<sup>40–42</sup>. Translational fusions of GmMYB176 with YFP and GmCYP1 with CFP were co-expressed in *N. benthamiana* leaf epidermal cells and analysed for FRET efficiency between the reporters *in vivo*. The results revealed a FRET efficiency of 23.8%, indicating a close proximity of GmMYB176 and GmCYP1 and their co-existence in a complex (Fig. 4b). Empty vectors containing YFP only and CFP only were used as negative control which showed a FRET efficiency of 0.5%. A FRET value of 8–10% was considered as the background level in previous studies<sup>43,44</sup>.

When *A. tumefaciens* GV3101 strains carrying BiFC plasmids containing GmCYP1-YN or GmCYP1-YC were co-infiltrated into *N. benthamiana* leaves and visualized by confocal microscope, a strong yellow fluorescence was observed in the nucleus and relatively weaker fluorescence in the cytoplasm, suggesting that GmCYP1 forms a homodimer in both cellular compartments *in planta* (Fig. 4a). Even though recombinant hCYP-A has been reported to form monomers, dimers, and trimers when expressed in *E. coli*<sup>45</sup>, there is no published literature on homo-dimerization of plant CYPs. Search for predicted motifs in GmCYP1 identified a putative phosphorylation (pST binding) site within the GmCYP1 sequence (97-ENFVKKHTGPGILSM-112), where T<sub>105</sub> is potentially phosphorylated. The pST binding motifs are binding sites for 14-3-3 family of proteins. 14-3-3 protein functions as a dimer to bind with its client proteins where each monomer in the dimer is capable of interacting with a






**Figure 3. Subcellular localization of GmCYP1.** *A. tumefaciens* GV3101 carrying the plasmids with GmCYP1-YFP and nuclear localizing CFP (NLS-CFP) constructs were co-infiltrated into *N. benthamiana* leaves and visualized by confocal microscopy. Expression of (a) GmCYP1-YFP, (b) NLS-CFP, (c) images of A and B merged to confirm nuclear localization of GmCYP1. Scale bars indicate 50  $\mu$ m. (d) Western blot analysis of translational fusion of GmCYP1-YFP proteins. GmCYP1-YFP was transiently expressed in *N. benthamiana* leaves for 1, 2 or 3 days, and protein accumulation was measured. Proteins (30  $\mu$ g) were separated on a SDS-PAGE and transferred to PVDF membrane by electroblotting. GmCYP1-YFP protein was detected by sequential incubation of the blot with anti-GFP antibody and anti-mouse IgG conjugated with HRP, followed by chemiluminescent reaction. eGFP is shown as a positive control.


activity was found in the hypocotyl. This study revealed additional information on tissue-specific expression of *GmCYP1*. For example, in leaves, GUS staining was more pronounced in the vascular tissues. Similarly, *GmCYP1* promoter was active in flower buds, stamen, stigma and silique walls but not in flower stem, style, petal or seeds. Failure to observe GUS activity in seeds could be due to weaker activity of the *GmCYP1* promoter in the seed coat, and is supported by *GmCYP1* expression in seed coat as shown in Fig. 5a. Similar results were observed for soybean *chalcone synthase* (*CHS*) gene promoters. Both *CHS7* and *CHS8* promoter driven GUS activities were absent in *Arabidopsis* seeds, despite the fact that *CHS7* and *CHS8* transcripts were present in the seed coat in soybean<sup>48</sup>. It is possible that the observed differences of gene expression in soybean and *Arabidopsis* may be caused by the presence or absence of the required regulatory factors in the specific tissue or developmental stage. Like *GmCYP1*, the *Arabidopsis* ortholog of *GmCYP1*, *ROC1* (AGI:At4g38740), also exhibited higher transcript accumulation in seeds than in other tissues (Fig. 5c). The normalized mean expression data of *ROC1* was compiled from AtGenExpress Visualization Tool (<http://jsp.weigelworld.org/expviz/expviz.jsp>). The expression of *ROC1* increased gradually, and approx. 7-fold throughout seed development, from a relatively low value (0.55) at the mid globular stage (stage 3), to values of 3.2–3.75 during the later developmental stages (early curled cotyledon embryos, stage 8) to green cotyledon embryos (stage 10) (Fig. 5d). The similar expression pattern during embryo development of *GmCYP1* and its *Arabidopsis* ortholog *ROC1* suggests a conserved role for *GmCYP1* and *ROC1* in seed development.

To identify putative *cis*-elements that regulate the expression of *GmCYP1* gene in soybean, we performed *in silico* motif analysis of 1148 bp upstream of translational start site using PlantPan2.0 database (<http://plantpan2.itsp.ncku.edu.tw/promoter.php>). The transcription factors specific to soybean was selected during the analysis. Transcription factors that are known for their role in stress and hormonal pathways are shown in Fig. 6. The analysis identified several sequence motifs that are recognized by a number of key factors such as bZIP, MYB, WRKY, bHLH, AP2, NAC factors have been identified. Besides their role in normal plant growth and development, the stress-specific roles of these factors have been well documented in crop plant and model species<sup>49–51</sup>.


***GmCYP1* expression is reduced in response to stress.** *P. sojae* effector Avr3b contains Nudix hydrolase activity *in planta* that is required for the virulence of the pathogen in soybean<sup>52</sup>. Recently, it has been shown



**Figure 4. GmCYP1 interacts with GmMYB176 and SGF14 in planta.** *N. benthamiana* leaves were co-transformed with *A. tumefaciens* carrying (a) GmCYP1-YN and SGF141-YC or their reciprocal combination, GmCYP1-YN and GmMYB176-YC or their reciprocal combination, GmCYP1-YN and GmCYP1-YC with vector only control (-YN or -YC only), GmCYP1-YN and GmCYP1-YC, and observed by confocal microscopy. Protein-protein interactions were visualized by a strong yellow fluorescence. Scale bars indicate 50  $\mu$ m. (b) FRET analysis demonstrating protein-protein interactions between GmMYB176 and GmCYP1. The CFP and YFP channels were excited with 458 nm and 514 nm lasers respectively, and FRET efficiencies were calculated in multiple samples ( $n > 15$ ). The empty pEG101 (YFP) and pEG102 (CFP) vector pairs were used as a FRET signal control.



**Figure 5. Expression analysis of *GmCYP1* in soybean.** (a) Total RNA extracted from soybean root, stem, leaf, flower bud, flower, embryo (30, 40, 50, 60, and 70 days after pollination), seed coat and pod wall (30 and 40 days after pollination) were used for quantitative RT-PCR analysis of *GmCYP1*. Two biological replicates and three technical replicates for each biological replicate were carried out. The standard error of the mean is represented by an error bar. The data were normalized against *SUB1-3* gene. (b) Histochemical analysis of *GmCYP1* promoter-GUS activity in vegetative and reproductive tissues during various stages of development in *Arabidopsis*. Construct containing a *GmCYP1* promoter driven *GUS* gene was transformed into *Arabidopsis* and selected T2 transgenic plants were used for analysis. (c) The mean normalized expression values of *ROC1* in different *Arabidopsis* tissues and (d) stages of seed development were obtained from AtGenExpress Visualization Tool (<http://jsp.weigelworld.org/expviz/expviz.jsp>). Error bars indicate the standard deviation of the mean. The stages of seed developments are: stage 3, mid globular to early heart embryos; stage 4, early to late heart embryos; stage 5, late heart to mid torpedo embryos; stage 6, mid to late torpedo embryos; stage 7, late torpedo to early walking-stick embryos; stage 8, walking-stick to early curled cotyledons embryos; stage 9, curled cotyledons to early green cotyledon embryos; and stage 10, green cotyledon embryos.



**Figure 6. Promoter analysis of *GmCYP1*.** A 1148 bp upstream of translation start site of *GmCYP1* was used for analysis using PlantPan2.0 database (<http://plantpan2.itps.ncku.edu.tw/promoter.php>). The transcription factors that are known for their role in stress and hormonal pathways are indicated.


that *GmCYP1* acts as a ‘helper’ by directly interacting with *Avr3b*, and modulating the hydrolase activity of it in soybean<sup>31</sup>. This *GmCYP1*-*Avr3b* interaction is required for the virulence and avirulence functions of *Avr3b* in soybean. Treatment of soybean hypocotyls with  $\text{AgNO}_3$  is known to induce defense response and phytoalexin accumulation<sup>53</sup>. Here we measured the accumulation of *GmCYP1* transcripts in response to stress by treating etiolated soybean hypocotyls with  $\text{AgNO}_3$  and monitoring *GmCYP1* expression 24, 48 or 72 h post-treatment. The results revealed that  $\text{AgNO}_3$  treated soybean hypocotyls accumulate reduced level of *GmCYP1* transcripts compared to the control hypocotyls at all the time points under the study (Fig. 7a). The difference in the level of *GmCYP1* transcript accumulation between  $\text{AgNO}_3$  treated and control hypocotyls was more pronounced at 24 h compared to 48 or 72 h post-treatment.

Isoflavonoid phytoalexins are host-produced antimicrobial compounds that are massively induced by pathogen attack or any other stress<sup>54,55</sup>. *GmMYB176* regulates isoflavonoid biosynthesis by regulating *CHS8* gene expression<sup>56</sup>. Since *GmCYP1* interacts with *GmMYB176*, and its expression is down-regulated upon stress, whereas phytoalexin biosynthesis is induced upon stress, we measured the expression levels of isoflavonoid biosynthetic genes *GmCHS8*, *isoflavone synthase* (*GmIFS2*) and a isoflavonoid-specific *prenyltransferase* (*GmPT*) in response to  $\text{AgNO}_3$  treatment. Our results revealed that expression of all these genes were induced upon  $\text{AgNO}_3$  treatment albeit at different levels (Fig. 7b). *GmCHS8* is the first enzyme in the flavonoid pathway, and as compared to control, its transcripts were accumulated at 6, 12 and 5.8 fold higher at 24, 48 and 72 hours, respectively after  $\text{AgNO}_3$  treatment. *GmIFS2* is a key legume-specific enzyme that introduces the isoflavonoid branch in the flavonoid pathway in legumes. Transcript levels of *GmIFS2* were 11.6, 17 and 3.7 fold higher after 24, 48 and 72 hours, respectively in  $\text{AgNO}_3$  treated samples compared to control. To confirm if downstream phytoalexin biosynthetic genes are induced upon stress, we measured the transcript levels of *GmPT*. Our results demonstrated that *GmPT* transcripts accumulated at 121, 64 and 3.5 fold greater than control at 24, 48 and 72 hours, respectively after  $\text{AgNO}_3$  treatment (Fig. 7b). A significantly higher difference in the expression of isoflavonoid genes were observed only when there was a significant reduction of *GmCYP1* gene expression (Fig. 7).

Several studies have shown that pathogen effectors interact with plant helper proteins for their activation and proper function<sup>57,58</sup>. Activated effector proteins bind with plant targets to suppress plant defenses and otherwise enable pathogen growth. For effectors encoded by *Avr* genes, activation can also result in detection by an immune receptor encoded by a resistance (*R*) gene. Thus, activation of *Avr3b* by *GmCYP1* triggers immunity in soybean cultivars containing *Rps3b*, whereas activation of *Avr3b* in soybean cultivars lacking *Rps3b* enables pathogen growth<sup>31</sup>. Unlike many other plant CYPs that function in protecting plants during biotic and abiotic stress<sup>29,59</sup>, *GmCYP1* acts as a susceptibility factor in soybean, at least in instances when *Avr3b* does not trigger immunity<sup>31</sup>. The expression of susceptibility proteins is generally induced during infection in susceptible plants<sup>60</sup>. However, our results show that *GmCYP1* transcripts are reduced in stressed plants compared to controls. This finding, together with the results that show *GmCYP1* interacts with *GmMYB176*, suggests a role for *GmCYP1* as a negative regulator of plant defense and isoflavonoid biosynthesis in soybean. The high expression of *GmCYP1* in the late stage developing embryos cannot be explained by this hypothesis because isoflavonoid biosynthesis is active in seed tissues, albeit this is seed isoflavonoid and not phytoalexin isoflavonoid biosynthesis.

Overall, this study presents a detailed analysis of *GmCYP1*. The presence of predicted cyclophilin domain, subcellular localization and its sequence homology with other identified CYPs from other organisms provides insights into its putative function. The interaction of *GmCYP1* and *GmMYB176* is particularly intriguing because of the conditional functionality of *GmCYP1* in effector-triggered immunity or susceptibility to the pathogen *P. sojae*. It seems more than coincidence that the biosynthesis of isoflavonoid phytoalexins, being necessary for resistance to *P. sojae*, is also connected to *GmCYP1*. Since isoflavonoids are involved in plant stress resistance against biotic and abiotic factors<sup>61–63</sup>, the interaction of *GmCYP1* with isoflavonoid regulators and its potential role as a suppressor of plant defense merits further investigation.





**Figure 7. Expression of *GmCYP1* and isoflavonoid biosynthetic genes in response to stress.** Etiolated soybean hypocotyls were treated with either 1 mM AgNO<sub>3</sub> or water (control) for 24, 48 or 72 h, and tissues were used to evaluate (a) *GmCYP1* (b) isoflavonoid biosynthetic gene (*GmCHS8*, *GmCHI1B1*, *GmIFS2*, *GmPT*) transcript accumulation using quantitative RT-PCR. Error bars indicate SEM of two biological replicates and three technical replicates for each biological replicate. The data were normalized against the *SUB1-3* gene for *GmCYP1* and *CON4* gene for isoflavonoid genes. Asterisk (\*) indicate significant difference between the samples using Student's *t*-test.

## Methods

**Plant growth conditions.** Soybean (*Glycine max* [L.] Merr) cultivar Harosoy 63 was grown in AAFC-London field plots during 2011 and 2012 for tissue collection. *Nicotiana benthamiana* plants were grown in pots under 16 h light at 25 °C and 8 h dark 20 °C cycle with 70–80% relative humidity and 100–150 μmol m<sup>2</sup>/s light intensity.

**RNA extraction and quantitative RT-PCR analysis.** Total RNA was extracted from soybean tissues according to Wang and Vodkin<sup>64</sup>. The RNA samples were quantified using a NanoDrop spectrophotometer (Thermo Scientific, USA), and their integrity was checked. Total RNA (1 μg) from each sample was used for cDNA synthesis using the QuantiTect® Reverse Transcription Kit (Qiagen, USA). For quantitative RT-PCR, SsoFast™ EvaGreen® Supermix (Bio-Rad, USA) was used with the CFX96 real-time PCR detection system (Bio-Rad, USA). Quantitative analysis of *GmCYP1* and isoflavonoid biosynthetic gene expression was carried out

using the primers listed in Table S2. The amplicons were cloned into a pGEM-T Easy vector (Promega, USA), and its sequence verified. *SOYBEAN UBIQUITIN-3 (SUBI-3)* or *CON4* was used as a reference gene for data normalization and to calculate the relative mRNA levels. The data were analyzed using CFX manager (Bio-Rad, USA).

**Plasmid constructions.** Full length *GmCYP1* was amplified from soybean cDNA constructed from mature embryo (60 and 70 DAP) using the primers GmCYP1-Gate-F: 5'-GGGGAC AAGTTTGTACAAAAAGCAGGCTTCATGCCTAACCCCTAAGTCTTCTTC-3' and GmCYP1-Gate-R: 5'-GGGGACCACTTTGTACAAGAAAGCTGGGTCAGAGGGTTGACCGCAGTTG-3'. The PCR product was recombined into pDONR-Zeo (Invitrogen, USA) using the BP clonase reaction mix (Invitrogen, USA), transformed into *Escherichia coli* DH5 $\alpha$ , and grown on LB media supplemented with zeocin (50  $\mu$ g/mL). The *E. coli* colonies containing recombinant plasmids were screened by colony PCR using gene-specific primers to identify pDONR-Zeo-GmCYP1. For subcellular localization study, the pDONR-Zeo-GmCYP1 was recombined with the destination vector pEarlyGate101<sup>65</sup> using the LR clonase reaction mix (Invitrogen, USA). The LR reaction was transformed into *E. coli* DH5 $\alpha$ , PCR screened, then transformed into *Agrobacterium tumefaciens* GV3101 for plant transformation.

For *in planta* protein-protein interaction study, pDONR-Zeo-GmCYP1 was recombined separately with pEarlyGate201-YN and pEarlyGate202-YC to obtain pEG201-GmCYP1-YN and pEG202-GmCYP1-YC, respectively. The recombinant plasmids were transformed into *E. coli* DH5 $\alpha$ , PCR screened, and then transformed into *A. tumefaciens* GV3101. For FRET analysis, pDONR221-GmMYB176 and pDONR-Zeo-GmCYP1 were recombined into the pEG101 or pEG102 upstream of the YFP or CFP sequence under the control of the CaMV 35 S promoter. The recombinant plasmids were transformed into *E. coli* DH5 $\alpha$ , PCR screened, and then transformed into *A. tumefaciens* GV3101.

The promoter fragment of *GmCYP1* (1148 bp) was amplified using the primers GmCYP1-P-F: 5'-GGGGACAAGTTTGTACAAAAAGCAGGCTCATCGTACTGCGATTTGAACGCAAGACTTC-3', and GmCYP1-P-R: 5'-GGGGACCACTTTGTACAAGAAAGCTGGGTAGTTGCTGCAGAGAAGAGAAGCGTA AATGAC-3', cloned into pDONR-Zeo (Invitrogen, USA), as described previously, followed by the recombination in the destination vector pMDC162 to obtain pGmCYP1pro-GUS. The pGmCYP1pro-GUS was transformed into *A. tumefaciens* GV3101 by electroporation, and then into wild-type *Arabidopsis* Col-0 by floral dip method<sup>66</sup>.

**Subcellular localization and bimolecular fluorescent complementation assay.** The subcellular localization of GmCYP1 was studied by infiltrating *A. tumefaciens* GV3101 carrying pEG101-GmCYP1 into *N. benthamiana* leaves, as described by Sparkes *et al.*<sup>67</sup>. For co-expression, equal volumes of two construct-bearing strains, suspended in Gamborg's solution, were mixed together and then infiltrated into *N. benthamiana* leaf epidermal cells. The protein expression was visualized by confocal microscopy using a Leica TCS SP2 inverted confocal microscope. An excitation wavelength of 514 nm was used for YFP imaging, and 525–545 nm emissions were collected. For visualization of CFP, an excitation wavelength of 458 nm was used, and emissions were collected between 465–495 nm.

**Fluorescence resonance energy transfer (FRET) assay.** Equal volumes of two construct-bearing strains containing YFP and CFP fusions (in Gamborg's solution), were mixed together and then infiltrated into *N. benthamiana* leaf epidermal cells. The protein expression was visualized by confocal microscopy using a Leica TCS SP2 inverted confocal microscope. An excitation wavelength of 458 nm and 514 nm were used for CFP and YFP imaging respectively. FRET acceptor bleaching, with CFP as donor and YFP as acceptor, was carried out by following Leica confocal application manual. The average of the FRET efficiency was calculated from multiple samples ( $n > 15$ ).

**Histochemical GUS assay.** For histochemical GUS staining, T<sub>2</sub> transgenic *Arabidopsis* tissues were used. Tissues were incubated in dark in a solution containing 100 mM sodium phosphate buffer pH 7.0, 10 mM EDTA, 0.05% Triton X-100, 1 mM potassium ferricyanide, 1 mM potassium ferrocyanide and 1 mM X-Gluc (5-Bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronide) for 16 h at 37 °C with gentle shaking. De-staining was carried out with 95% ethanol for 4 to 6 times. A LeicaM2 FLIII<sup>TM</sup> microscope with a QImaging Retiga 2000 R camera was used to take the photographs. For staining of silique and seeds, improved clearing method was used<sup>68</sup>.

**Yeast two-hybrid assay.** Yeast two-hybrid assay (Y2H) was performed using the Matchmaker<sup>®</sup> Gold Two-Hybrid System (Clontech Laboratories, Inc., USA). Briefly, GmMYB176 was cloned into the vector pGBKT7 as the bait. The cDNA library as prey was generated by SMART<sup>TM</sup> cDNA Synthesis technology (Clontech Laboratories, Inc., USA) from soybean embryos (50–60 days after pollination) and fused to GAL4 activation domain. Screening was performed by co-transformation of bait and prey using Mate & Plate<sup>TM</sup> library system (Clontech Laboratories, Inc., USA). After transformation, yeast cells were spread on SD/Ade/-His/-Leu/-Trp agar plates, and incubated at 30 °C for 5 days.

## References

- Dilworth, D., Gudavicius, G., Leung, A. & Nelson, C. J. The roles of peptidyl-proline isomerases in gene regulation. *Biochem. Cell Biol.* **90**, 55–69, doi: 10.1139/o11-045 (2012).
- Harding, M. W., Galat, A., Uehling, D. E. & Schreiber, S. L. A receptor for the immunosuppressant FK506 is a cis-trans peptidyl-prolyl isomerase. *Nature* **341**, 758–760, doi: 10.1038/341758a0 (1989).
- Siekierka, J. J., Hung, S. H., Poe, M., Lin, C. S. & Sigal, N. H. A cytosolic binding protein for the immunosuppressant FK506 has peptidyl-prolyl isomerase activity but is distinct from cyclophilin. *Nature* **341**, 755–757, doi: 10.1038/341755a0 (1989).
- Takahashi, N., Hayano, T. & Suzuki, M. Peptidyl-prolyl cis-trans isomerase is the cyclosporin A-binding protein cyclophilin. *Nature* **337**, 473–475, doi: 10.1038/337473a0 (1989).

5. Galat, A. Variations of sequences and amino acid compositions of proteins that sustain their biological functions: An analysis of the cyclophilin family of proteins. *Arch. Biochem. Biophys.* **371**, 149–162, doi: 10.1006/abbi.1999.1434 (1999).
6. Maruyama, T., Suzuki, R. & Furutani, M. Archaeal peptidyl prolyl cis-trans isomerases (PPIases) update 2004. *Front. Biosci.* **9**, 1680–1720 (2004).
7. Kumari, S., Roy, S., Singh, P., Singla-Pareek, S. & Pareek, A. Cyclophilins: Proteins in search of function. *Plant Signaling & Behavior* **8**, e22734, doi: 10.4161/psb.22734 (2013).
8. Galat, A. Peptidylprolyl cis/trans isomerases (immunophilins): biological diversity-targets-functions. *Curr Top Med Chem* **3**, 1315–1347 (2003).
9. Page, A. P., MacNiven, K. & Hengartner, M. O. Cloning and biochemical characterization of the cyclophilin homologues from the free-living nematode *Caenorhabditis elegans*. *Biochem. J.* **317**, 179–185 (1996).
10. Arevalo-Rodriguez, M., Wu, X., Hanes, S. D. & Heitman, J. Prolyl isomerases in yeast. *Front. Biosci.* **9**, 2420–2446 (2004).
11. Braaten, D. & Luban, J. Cyclophilin A regulates HIV-1 infectivity, as demonstrated by gene targeting in human T cells. *EMBO J.* **20**, 1300–1309, doi: 10.1093/emboj/20.6.1300 (2001).
12. Lu, K. P., Finn, G., Lee, T. H. & Nicholson, L. K. Prolyl cis-trans isomerization as a molecular timer. *Nat. Chem. Biol.* **3**, 619–629, doi: 10.1038/nchembio.2007.35 (2007).
13. Nigro, P., Pompilio, G. & Capogrossi, M. C. Cyclophilin A: A key player for human disease. *Cell Death Dis.* **4**, doi: 10.1038/cddis.2013.410 (2013).
14. Gasser, C. S., Gunning, D. A., Budelier, K. A. & Brown, S. M. Structure and expression of cytosolic cyclophilin/peptidyl-prolyl cis-trans isomerase of higher plants and production of active tomato cyclophilin in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **87**, 9519–9523 (1990).
15. Mainali, H., Chapman, P. & Dhaubhadel, S. Genome-wide analysis of *Cyclophilin* gene family in soybean (*Glycine max*). *BMC Plant Biol.* **14**, 282, doi: 10.1186/s12870-014-0282-7 (2014).
16. Romano, P. G., Horton, P. & Gray, J. E. The Arabidopsis cyclophilin gene family. *Plant Physiol.* **134**, 1268–1282, doi: 10.1104/pp.103.022160 (2004).
17. Trivedi, D., Yadav, S., Vaid, N. & Tuteja, N. Genome wide analysis of Cyclophilin gene family from rice and Arabidopsis and its comparison with yeast. *Plant Signaling & Behavior* **7**, 1653–1666 (2012).
18. Ahn, J. C. *et al.* Classification of rice (*Oryza sativa* L. Japonica nipponbare) immunophilins (FKBPs, CYPs) and expression patterns under water stress. *BMC Plant Biol.* **10**, 253, doi: 10.1186/1471-2229-10-253 (2010).
19. Vallon, O. Chlamydomonas Immunophilins and Parvulins: Survey and Critical Assessment of Gene Models. *Eukaryotic Cell* **4**, 230–241, doi: 10.1128/ec.4.2.230-241.2005 (2005).
20. Fujita, M. *et al.* Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Curr. Opin. Plant Biol.* **9**, 436–442 (2006).
21. Atkinson, N. J. & Urwin, P. E. The interaction of plant biotic and abiotic stresses: from genes to the field. *J. Exp. Bot.* **63**, 3523–3543 (2012).
22. Brandts, J. F., Halvorson, H. R. & Brennan, M. Consideration of the Possibility that the slow step in protein denaturation reactions is due to cis-trans isomerism of proline residues. *Biochemistry* **14**, 4953–4963 (1975).
23. Chou, I. T. & Gasser, C. S. Characterization of the cyclophilin gene family of Arabidopsis thaliana and phylogenetic analysis of known cyclophilin proteins. *Plant Mol. Biol.* **35**, 873–892 (1997).
24. Marivet, J., Margis-Pinheiro, M., Frendo, P. & Burkard, G. Bean cyclophilin gene expression during plant development and stress conditions. *Plant Mol. Biol.* **26**, 1181–1189, doi: 10.1007/bf00040698 (1994).
25. Meza-Zepeda, L. A., Baudo, M. M., Palva, E. T. & Heino, P. Isolation and characterization of a cDNA corresponding to a stress-activated cyclophilin gene in *Solanum commersonii*. *J. Exp. Bot.* **49**, 1451–1452, doi: 10.1093/jxb/49.325.1451 (1998).
26. Kong, H. Y., Lee, S. C. & Hwang, B. K. Expression of pepper cyclophilin gene is differentially regulated during the pathogen infection and abiotic stress conditions. *Physiol. Mol. Plant Pathol.* **59**, 189–199, doi: 10.1006/pmpp.2001.0356 (2001).
27. Chen, A. P. *et al.* Ectopic expression of ThCYP1, a stress-responsive cyclophilin gene from *Thellungiella halophila*, confers salt tolerance in fission yeast and tobacco cells. *Plant Cell Rep.* **26**, 237–245, doi: 10.1007/s00299-006-0238-y (2007).
28. Sekhar, K., Priyanka, B., Reddy, V. D. & Rao, K. V. Isolation and characterization of a pigeonpea cyclophilin (CcCYP) gene, and its over-expression in Arabidopsis confers multiple abiotic stress tolerance. *Plant, Cell Environ.* **33**, 1324–1338, doi: 10.1111/j.1365-3040.2010.02151.x (2010).
29. Zhu, C. *et al.* Overexpression of a cotton cyclophilin gene (GhCyp1) in transgenic tobacco plants confers dual tolerance to salt stress and *Pseudomonas syringae* pv. tabaci infection. *Plant Physiol. Biochem.* **49**, 1264–1271, doi: 10.1016/j.plaphy.2011.09.001 (2011).
30. Schmutz, J. *et al.* Genome sequence of the palaeopolyploid soybean. *Nature* **463**, 178–183, doi: 10.1038/nature08670 (2010).
31. Kong, G. *et al.* The Activation of *Phytophthora* Effector Avr3b by Plant Cyclophilin is Required for the Nudix Hydrolase Activity of Avr3b. *PLoS Pathog.* **11**, e1005139, doi: 10.1371/journal.ppat.1005139 (2015).
32. Zydowsky, L. D. *et al.* Active site mutants of human cyclophilin A separate peptidyl-prolyl isomerase activity from cyclosporin A binding and calcineurin inhibition. *Protein Sci.* **1**, 1092–1099, doi: 10.1002/pro.5560010903 (1992).
33. Liu, J., Chen, C. M. & Walsh, C. T. Human and *Escherichia coli* cyclophilins: sensitivity to inhibition by the immunosuppressant cyclosporin A correlates with a specific tryptophan residue. *Biochemistry* **30**, 2306–2310 (1991).
34. Fried, H. & Kutay, U. Nucleocytoplasmic transport: taking an inventory. *Cellular and Molecular Life Sciences CMLS* **60**, 1659–1688 (2003).
35. D'Angelo, M. A., Raices, M., Panowski, S. H. & Hetzer, M. W. Age-dependent deterioration of nuclear pore complexes causes a loss of nuclear integrity in postmitotic cells. *Cell* **136**, 284–295 (2009).
36. Harel, A. & Forbes, D. J. Importin beta: conducting a much larger cellular symphony. *Mol. Cell* **16**, 319–330, doi: 10.1016/j.molcel.2004.10.026 (2004).
37. Ohad, N., Shichrur, K. & Yalovsky, S. The analysis of protein-protein interactions in plants by bimolecular fluorescence complementation. *Plant Physiol.* **145**, 1090–1099 (2007).
38. Li, X., Chen, L. & Dhaubhadel, S. 14-3-3 proteins regulate the intracellular localization of the transcriptional activator GmMYB176 and affect isoflavonoid synthesis in soybean. *Plant J.* **71**, 239–250 (2012).
39. Day, R. N., Periasamy, A. & Schaufele, F. Fluorescence Resonance Energy Transfer Microscopy of Localized Protein Interactions in the Living Cell Nucleus. *Methods* **25**, 4–18, doi: http://dx.doi.org/10.1006/meth.2001.1211 (2001).
40. Immink, R. G. H., Gadella, T. W. J., Ferrario, S., Busscher, M. & Angenent, G. C. Analysis of MAD5 box protein-protein interactions in living plant cells. *Proc. Natl. Acad. Sci. USA* **99**, 2416–2421, doi: 10.1073/pnas.042677699 (2002).
41. Kato, N., Pontier, D. & Lam, E. Spectral Profiling for the Simultaneous Observation of Four Distinct Fluorescent Proteins and Detection of Protein-Protein Interaction via Fluorescence Resonance Energy Transfer in Tobacco Leaf Nuclei. *Plant Physiol.* **129**, 931–942, doi: 10.1104/pp.005496 (2002).
42. Achnine, L., Blancaflor, E. B., Rasmussen, S. & Dixon, R. A. Colocalization of L-phenylalanine ammonia-lyase and cinnamate 4-hydroxylase for metabolic channeling in phenylpropanoid biosynthesis. *Plant Cell* **16**, 3098–3109, doi: 10.1105/tpc.104.024406 (2004).
43. Seidel, T. *et al.* Colocalization and FRET-analysis of subunits c and a of the vacuolar H<sup>+</sup>-ATPase in living plant cells. *J. Biotechnol.* **112**, 165–175, doi: http://dx.doi.org/10.1016/j.jbiotec.2004.04.027 (2004).
44. Seidel, T., Gollack, D. & Dietz, K.-J. Mapping of C-termini of V-ATPase subunits by *in vivo*-FRET measurements. *FEBS Lett.* **579**, 4374–4382, doi: http://dx.doi.org/10.1016/j.febslet.2005.06.077 (2005).

45. Zhang, X. C., Wang, W. D., Wang, J. S., Pan, J. C. & Zou, G. L. Evidences of monomer, dimer and trimer of recombinant human cyclophilin A. *Protein Pept Lett* **18**, 1188–1193 (2011).
46. Tzivion, G., Shen, Y. H. & Zhu, J. 14-3-3 proteins; bringing new definitions to scaffolding. *Oncogene* **20**, 6331–6338 (2001).
47. Gijzen, M., Kuflu, K., Qutob, D. & Chernys, J. T. A class I chitinase from soybean seed coat. *J. Exp. Bot.* **52**, 2283–2289, doi: 10.1093/jxbbot/52.365.2283 (2001).
48. Yi, J., Derynck, M. R., Chen, L. & Dhaubhadel, S. Differential expression of *CHS7* and *CHS8* genes in soybean. *Planta* **231**, 741–753, doi: 10.1007/s00425-009-1079-z (2010).
49. Puranik, S., Sahu, P. P., Srivastava, P. S. & Prasad, M. NAC proteins: regulation and role in stress tolerance. *Trends Plant Sci.* **17**, 369–381, doi: <http://dx.doi.org/10.1016/j.tplants.2012.02.004> (2012).
50. Gutterson, N. & Reuber, T. L. Regulation of disease resistance pathways by AP2/ERF transcription factors. *Curr. Opin. Plant Biol.* **7**, 465–471, doi: <http://dx.doi.org/10.1016/j.pbi.2004.04.007> (2004).
51. Dong, J., Chen, C. & Chen, Z. Expression profiles of the Arabidopsis WRKY gene superfamily during plant defense response. *Plant Mol. Biol.* **51**, 21–37, doi: 10.1023/a:1020780022549 (2003).
52. Dong, S. *et al.* *Phytophthora sojae* Avirulence Effector Avr3b is a Secreted NADH and ADP-ribose Pyrophosphorylase that Modulates Plant Immunity. *PLoS Pathog.* **7**, e1002353, doi: 10.1371/journal.ppat.1002353 (2011).
53. Bhattacharyya, M. K. & Ward, E. W. B. Resistance, susceptibility and accumulation of glyceollins I–III in soybean organs inoculated with *Phytophthora megasperma* f. sp. *glycinea*. *Physiol. Mol. Plant Pathol.* **29**, 227–237, doi: [http://dx.doi.org/10.1016/S0048-4059\(86\)80023-6](http://dx.doi.org/10.1016/S0048-4059(86)80023-6) (1986).
54. Phillips, D. A. & Kapulnik, Y. Plant isoflavonoids, pathogens and symbionts. *Trends Microbiol.* **3**, 58–64, doi: 10.1016/s0966-842x(00)88876-9 (1995).
55. Subramanian, S., Graham, M. A., Yu, O. & Graham, T. L. RNA interference of soybean isoflavone synthase genes leads to silencing in tissues distal to the transformation site and to enhanced susceptibility to *Phytophthora sojae*. *Plant Physiol.* **137**, 1345–1353 (2005).
56. Yi, J. *et al.* A single-repeat MYB transcription factor, GmMYB176, regulates CHS8 gene expression and affects isoflavonoid biosynthesis in soybean. *Plant J.* **62**, 1019–1034, doi: 10.1111/j.1365-313X.2010.04214.x (2010).
57. Win, J. *et al.* Effector Biology of Plant-Associated Organisms: Concepts and Perspectives. *Cold Spring Harbor Symp. Quant. Biol.* **77**, 235–247, doi: 10.1101/sqb.2012.77.015933 (2012).
58. Lapin, D. & Van den Ackerveken, G. Susceptibility to plant disease: more than a failure of host immunity. *Trends Plant Sci.* **18**, 546–554, doi: <http://dx.doi.org/10.1016/j.tplants.2013.05.005> (2013).
59. Coaker, G., Falick, A. & Staskawicz, B. Activation of a Phytopathogenic Bacterial Effector Protein by a Eukaryotic Cyclophilin. *Science* **308**, 548–550, doi: 10.1126/science.1108633 (2005).
60. Huibers, R. P., de Jong, M., Dekter, R. W. & Van den Ackerveken, G. Disease-Specific Expression of Host Genes During Downy Mildew Infection of Arabidopsis. *Mol. Plant-Microbe Interact.* **22**, 1104–1115, doi: 10.1094/MPMI-22-9-1104 (2009).
61. Graham, T. L., Graham, M. Y., Subramanian, S. & Yu, O. RNAi silencing of genes for elicitation or biosynthesis of 5-deoxyisoflavonoids suppresses race-specific resistance and hypersensitive cell death in *phytophthora sojae* infected tissues. *Plant Physiol.* **144**, 728–740 (2007).
62. Gutierrez-Gonzalez, J. J. *et al.* Differential Expression of Isoflavone Biosynthetic Genes in Soybean During Water Deficits. *Plant Cell Physiol.* **51**, 936–948, doi: 10.1093/pcp/pcq065 (2010).
63. Chennupati, P., Seguin, P., Chamoun, R. & Jabaji, S. Effects of High-Temperature Stress on Soybean Isoflavone Concentration and Expression of Key Genes Involved in Isoflavone Synthesis. *J. Agric. Food Chem.* **60**, 12421–12427, doi: 10.1021/jf3036319 (2012).
64. Wang, C. S. & Vodkin, L. O. Extraction of RNA from tissues containing high levels of procyranidins that bind RNA. *Plant Mol. Biol. Rep.* **12**, 132–145, doi: 10.1007/BF02668374 (1994).
65. Earley, K. W. *et al.* Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J.* **45**, 616–629, doi: 10.1111/j.1365-313X.2005.02617.x (2006).
66. Zhang, X., Henriques, R., Lin, S. S., Niu, Q. W. & Chua, N. H. Agrobacterium-mediated transformation of *Arabidopsis thaliana* using the floral dip method. *Nature Prot.* **1**, 641–646, doi: 10.1038/nprot.2006.97 (2006).
67. Sparkes, I. A., Runions, J., Kearns, A. & Hawes, C. Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. *Nature Prot.* **1**, 2019–2025 (2006).
68. Stangeland, B. & Salehian, Z. An improved clearing method for GUS assay in *Arabidopsis* endosperm and seeds. *Plant Mol. Biol. Rep.* **20**, 107–114 (2002).

## Acknowledgements

The authors thank Alex Molnar for graphic designs, and Ling Chen, Arjun Sukumaran and Dorothy Drew for technical assistance.

## Author Contributions

H.R.M. performed experiments, analysed data and prepared draft manuscript. A.K.A.V. and X.L. performed experiments. M.G. contributed to data interpretation and manuscript preparation. S.D. conceived and designed experiments, prepared manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Mainali, H. R. *et al.* Soybean cyclophilin GmCYP1 interacts with an isoflavonoid regulator GmMYB176. *Sci. Rep.* **7**, 39550; doi: 10.1038/srep39550 (2017).

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017