

SCIENTIFIC REPORTS



OPEN

Memory and synaptic plasticity are impaired by dysregulated hippocampal O-GlcNAcylation

Yong Ryoul Yang^{1,2,*}, Seungju Song^{3,4,*}, Hongik Hwang³, Jung Hoon Jung⁵, Su-Jeong Kim⁵, Sora Yoon¹, Jin-Hoe Hur¹, Jae-Il Park⁶, Cheol Lee¹, Dougu Nam¹, Young-Kyo Seo¹, Jung-Hun Kim⁵, Hyewhon Rhim^{3,4} & Pann-Ghill Suh¹

Received: 21 July 2016
Accepted: 14 February 2017
Published: 03 April 2017

O-GlcNAcylated proteins are abundant in the brain and are associated with neuronal functions and neurodegenerative diseases. Although several studies have reported the effects of aberrant regulation of O-GlcNAcylation on brain function, the roles of O-GlcNAcylation in synaptic function remain unclear. To understand the effect of aberrant O-GlcNAcylation on the brain, we used *Oga*^{+/-} mice which have an increased level of O-GlcNAcylation, and found that *Oga*^{+/-} mice exhibited impaired spatial learning and memory. Consistent with this result, *Oga*^{+/-} mice showed a defect in hippocampal synaptic plasticity. *Oga* heterozygosity causes impairment of both long-term potentiation and long-term depression due to dysregulation of AMPA receptor phosphorylation. These results demonstrate a role for hyper-O-GlcNAcylation in learning and memory.

O-GlcNAcylation is a posttranslational modification involving attachment of beta-*N*-acetylglucosamine (GlcNAc) to Ser/Thr residues. The addition and removal of O-GlcNAc to target proteins is regulated by O-GlcNAc transferase (OGT) and O-GlcNAcase (OGA), respectively¹, and both OGT and OGA are abundantly expressed in the brain^{2,3}. The roles of O-GlcNAcylation have been extensively investigated in aging-associated neurodegenerative diseases, such as Alzheimer's disease and Parkinson's disease^{4,5}. Several aggregation-prone proteins involved in neurodegeneration are O-GlcNAcylated, including Tau^{6,7} and α -synuclein^{8,9}. Elevated O-GlcNAcylation prevents protein aggregation and slows neurodegeneration^{4,5}. β -amyloid precursor protein (APP) involved in amyloid plaque formation is also O-GlcNAcylated¹⁰. Increased total O-GlcNAcylation attenuates the production of oligomeric A β , the main component of senile plaques, by lowering the activity of γ -secretase¹¹. Oligomeric forms of amyloid- β (A β) are known to acutely increase excitatory synaptic transmission and interfere with synaptic plasticity^{12,13}. Previous evidence suggests that aberrant regulation of O-GlcNAcylation may contribute to the impaired synaptic plasticity in neurodegeneration.

O-GlcNAcylation is particularly enriched in neuronal synapses^{14,15}, and proteomic studies have identified many postsynaptic density proteins modified by O-GlcNAc^{8,16}. In addition, the activity of the brain OGT is ten-fold higher than that of peripheral tissues². Various neuronal proteins are O-GlcNAcylated and involved in synaptic functions^{1,17}. GluA2, a subunit of glutamatergic α -amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA) receptors, interacts with OGT, and is O-GlcNAcylated. Although phosphorylation of GluA2 is not affected by increased O-GlcNAcylation, O-GlcNAcylation of GluA2 is required for synaptic plasticity¹⁸. Furthermore, synapsin I, a synaptic vesicle-associated protein, is O-GlcNAcylated, with suggested roles in the localization and function of synapsin I^{19,20}. O-GlcNAcylation of synapsin I has been implicated in the modulation of synaptic plasticity. In addition, O-GlcNAcylation is present on proteins important for neuronal signaling. Calcium/calmodulin-dependent kinase II (CaMKII), CaMKIV, and the transcription factor cyclic adenosine monophosphate (AMP)-response element binding protein (CREB) are O-GlcNAcylated, which

¹School of Life Sciences, Ulsan National Institute of Science and Technology, Ulsan 44919, Republic of Korea. ²Aging Research Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Daejeon 34141, Republic of Korea. ³Center for Neuroscience, Brain Science Institute, Korea Institute of Science and Technology (KIST), Seoul 136-791, South Korea. ⁴Department of Neuroscience, University of Science and Technology (UST), Daejeon 305-333, South Korea. ⁵Department of Life Science, Pohang University of Science and Technology, Pohang, Gyungbuk 790-784, Republic of Korea. ⁶Korea Basic Science Institute, Gwangju 61186, Republic of Korea. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to H.R. (email: hrhim@kist.re.kr) or P-G.S. (email: pgsuh@unist.ac.kr)

influences synaptic plasticity in the hippocampus^{21–23}. Although it is clear that O-GlcNAcylation is abundant in synapses and that O-GlcNAcylation affects synaptic plasticity and learning and memory in the hippocampus, past studies have used different methods to modulate O-GlcNAcylation levels, resulting in conflicting results. Decreased O-GlcNAc levels by alloxan treatment (OGT inhibitor) impairs high-frequency stimulation (HFS)-induced long-term potentiation (LTP) in the Schaffer Collateral (SC)-CA1 Pathway²⁴. In contrast, the elevation of O-GlcNAcylation induced by Thiamet-G (OGA inhibitor) inhibits HFS-LTP and impairs hippocampal learning¹⁸.

Here we assessed how chronic elevations of O-GlcNAcylation in the hippocampus affect synaptic function, behavioral traits, and spatial learning and memory, using *Oga*^{+/-} mice with constitutively increased O-GlcNAc levels.

Results

***Oga*^{+/-} brains have normal morphology and dendritic spine density.** Consistent with previous studies showing enriched expression of O-GlcNAc cycling enzymes, OGT and OGA, we detected high levels of O-GlcNAcase expression in the hippocampus, which was visualized by beta galactosidase (LacZ) staining of an *Oga*^{+/-} brain section (Fig. 1a). To assess the effect of increased O-GlcNAcylation on hippocampus-dependent function, we used *Oga*^{+/-} mice with chronically elevated O-GlcNAcylation. The hippocampal lysates prepared from *Oga*^{+/-} mice showed an increase in the overall O-GlcNAcylation levels (Fig. 1b). To verify the elevation of O-GlcNAcylation in the hippocampus, we used immunohistochemistry with an anti-O-GlcNAc antibody. As expected, increased immunoreactivity was observed throughout all regions of the hippocampus in *Oga*^{+/-} mice compared to WT. (Fig. 1c). Next, we tested whether *Oga* heterozygosity leads to morphological changes in the brain. Morphological analysis of neurons in the hippocampus by Nissl staining revealed that the *Oga*^{+/-} hippocampus shows no morphological changes in hippocampal CA1, CA3, or dentate gyrus (DG) (Fig. 1d). In addition, we found that there were no differences in the numbers of cells immunostained for the neuronal marker neuronal nuclei (NeuN) and for the astrocyte marker glial fibrillary acidic protein (GFAP) in the hippocampal CA1, CA3, or DG (Fig. 1e). The average brain weight also was not affected by *Oga* heterozygosity (Fig. 1f and g). Lastly, dendritic spine density was not altered in the *Oga*^{+/-} hippocampal CA1 pyramidal neurons (Fig. 1h and i). These results together suggest that synaptic development and hippocampal structure are not affected by the elevation of O-GlcNAcylation.

***Oga*^{+/-} mice display impaired spatial learning and memory.** Various neuronal proteins involved in synaptic function and learning and memory are known to be O-GlcNAcylated^{8,9,15}. To assess whether hyper-O-GlcNAcylation affects hippocampal-dependent spatial learning and memory, we employed the Barnes circular maze test. In this test, mice were trained to escape a brightly lighted circular field by discovering the escape hole at its periphery. Compared with wild-type (WT) mice, *Oga*^{+/-} mice showed impaired learning performance during four days of training (Fig. 2a–c). To assess memory formation, we performed probe trials on days 5 and 12. WT and *Oga*^{+/-} mice performed similarly during the probe trials when total distance was measured (Fig. 2d and g). *Oga*^{+/-} mice exhibited increased latency to the target region during probe trials (Fig. 2f and i). However, no significant difference were observed in the time spent in the target region between WT and *Oga*^{+/-} mice (Fig. 2e and h). To further verify the impairment in spatial learning and memory of *Oga*^{+/-} mice, we performed a context fear conditioning test. Compared with WT mice, *Oga*^{+/-} mice failed to retain fear memory 24 h after fear conditioning (Fig. 2j and k). Our data indicate that proper removal of O-GlcNAc modification by OGA is required for hippocampal-dependent spatial learning and memory.

To examine motor coordination in these mice, we tested motor performance using the rotarod task. *Oga*^{+/-} mice did not display a defect in motor function during the rotarod test (Fig. S1A). During the open field test, *Oga*^{+/-} mice showed normal locomotor activities (Fig. S1B and C). Anxiety-related behaviors were also tested using the elevated plus maze. Compared with WT mice, *Oga*^{+/-} mice exhibited no significant differences in the number of entries to the open arms and amount of time spent in the open arms (Fig. S1D–G). Considered collectively, these data indicate that *Oga*^{+/-} mice show normal locomotor activity and anxiety levels.

Glutamatergic and GABAergic synaptic transmission in the hippocampus is normal in *Oga*^{+/-} mice.

We next explored the effect of heterozygous loss of *Oga* on intrinsic neuronal excitability and excitatory synaptic transmission in hippocampal CA1 pyramidal neurons. Excitability was tested by injecting step depolarizing currents, and we found that intrinsic excitability of hippocampal CA1 pyramidal neurons remains unchanged in *Oga*^{+/-} mice (Fig. 3a and b). The frequency and amplitude of miniature excitatory and inhibitory postsynaptic currents (mEPSCs and mIPSCs, respectively) in hippocampal CA1 pyramidal neurons were also comparable in WT and *Oga*^{+/-} pyramidal neurons (Fig. 3c and d), indicating that basal synaptic responses are not affected in *Oga*^{+/-} mice. In addition, we measured the ratio of AMPA to N-methyl-D-aspartate (NMDA) receptor-mediated synaptic currents in SC–CA1 synapses, and the AMPA/NMDA ratio was similar between WT and *Oga*^{+/-} synapses (Fig. 3e). These results suggest that *Oga* heterozygosity does not affect basal SC–CA1 synaptic transmission or short-term plasticity.

Impaired NMDA receptor (NMDAR)-dependent synaptic plasticity in *Oga*^{+/-} mice. Previous studies investigating the effects of increased O-GlcNAcylation on synaptic plasticity have generated conflicting results^{18,24}. Therefore, we next assessed whether increased O-GlcNAcylation resulting from *Oga* haploinsufficiency alters synaptic plasticity. The slope of the field excitatory postsynaptic potential (fEPSP) to fiber volley amplitudes (input-output curves) was not changed in *Oga*^{+/-} mice (Fig. 4a). Presynaptic release probability, as measured by paired-pulse facilitation (PPF), also remained unaffected in *Oga*^{+/-} mice (Fig. 4b). When we measured NMDAR-mediated LTP in *Oga*^{+/-} mice, the magnitude of LTP induced by high-frequency stimulation




Figure 1. Normal morphological features and dendritic spine density in *Oga*^{+/-} brains (a) Beta galactosidase (LacZ) staining of the *Oga*^{+/-} adult brain, confirming the expression pattern of OGA in the hippocampus. Scale bar, 200 µm. (b) Immunoblot analysis showing elevated O-GlcNAcylation in *Oga*^{+/-} hippocampal lysates compared with that in the WT hippocampal lysates. (c) Representative images of hippocampal neurons from WT and *Oga*^{+/-} mice immunolabeled for O-GlcNAc (green). Scale bar, 200 µm. (d) Nissl staining of the hippocampus from coronal brain sections. Scale bar, 100 µm. (e) Immunostaining for a neuronal cell marker (neuronal nuclei; NeuN), glial marker (glial fibrillary acidic protein; GFAP), and nuclei (4',6-diamidino-2-phenylindole; DAPI) in the hippocampus of WT and *Oga*^{+/-} mice at eight weeks of age. Three sections were obtained from three mice. Scale bar, 100 µm. (f) Representative pictures of brain tissues. (g) Whole-brain weight (without skull) isolated from the WT and *Oga*^{+/-} mice at eight weeks of age (n = 5). (h) Representative images of dendritic segments of Golgi-stained CA1 pyramidal neurons from WT and *Oga*^{+/-} mice (scale bar = 10 µm) (i) Spine number was quantified along a 10 µm segment from the primary apical dendritic branch origin of Golgi-impregnated CA1 pyramidal neurons (n = 15–20 dendrites from three mice). Error bars represent ± standard error of the mean (SEM). NS: not significant (unpaired *t*-test). Full-length blots/gels are presented in Supplementary Figure S4.

(HFS) at the SC–CA1 pathway was reduced compared to WT mice (Fig. 4c). Moreover, in the same hippocampal pathway, low-frequency stimulation (LFS)-induced long-term depression (LTD) was impaired in *Oga*^{+/-} mice in comparison to WT mice (Fig. 4d). These results suggest that the removal of O-GlcNAcylation mediated by OGA is required for NMDAR-dependent LTP and LTD at SC–CA1 synapses.

Impaired modulation of AMPA receptor during LTP/LTD in *Oga*^{+/-} mice. Regulation of AMPA receptor trafficking is crucial for controlling the strength of synaptic transmission during LTP/LTD. In particular, phosphorylation of GluA1 AMPAR subunit at S845 and S831 play key roles in AMPA receptor trafficking and synaptic plasticity²⁵. We thus decided to examine whether GluA1 phosphorylation is altered in response to chemically induced LTP and LTD in *Oga*^{+/-} hippocampus. We briefly stimulated hippocampal slices from WT




Figure 2. Spatial learning memory deficits in *Oga*^{+/-} mice (a) Latency in attaining the goal and (b) number of errors committed before reaching the goal in the Barnes maze task. (WT, n = 9; *Oga*^{+/-}, n = 9, one-way ANOVA followed by the Tukey's test to unpaired t-test) (c) Representative traces of WT and *Oga*^{+/-} mice in the Barnes maze task. (d) Total distance, (e) time spent in the target area, and (f) time of latency to attaining the target hole during a probe trial on day 5. (g) Total distance, (h) time spent in the target area, and (i) time of latency to attain the target hole during a probe trial on day 12. (WT, n = 9; *Oga*^{+/-}, n = 9) (j) Schematic diagram of the context-dependent fear conditioning procedure (k) Percentage time freezing during the 3 min of the context test 24 h after fear conditioning. (WT, n = 9; *Oga*^{+/-}, n = 8). Error bars represent \pm standard error of the mean (SEM). NS: not significant, *** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$ (unpaired *t*-test).

and *Oga*^{+/-} mice with glycine for LTP or NMDA for LTD^{26,27}. WT hippocampal slices showed elevated phosphorylation of the S845 and S831 GluA1 in chemical LTP, and the phosphorylation of the S845 GluA1 was decreased in chemical LTD. However, in *Oga*^{+/-} hippocampal slices, phosphorylation of the S845 and S831 GluA1 were not properly regulated following chemical LTP or LTD (Fig. 5a–c). These results indicate that *Oga* heterozygosity impairs the proper regulation of AMPA receptor phosphorylation during synaptic plasticity.

Diverse neuronal proteins and signaling mediators are O-GlcNAcylated and involved in synaptic functions. For example, AMPA receptor subunit GluA2 is O-GlcNAcylated by OGT¹⁸, and CaMKII, CaMKIV, and CREB whose alternation of activity affects synaptic plasticity^{28,29}, are also O-GlcNAcylated^{21–23}. *Oga* heterozygosity did not alter the basal phosphorylation or total protein levels AMPA receptor subunits (GluA1, GluA2) and NMDA receptor subunits (GluN1, GluN2A, GluN2B) (Fig. S2A). In addition, no difference was observed in phosphorylation of CaMKII, CaMKIV, or CREB between the hippocampus of WT and *Oga*^{+/-} mice (Fig. S2B).

Discussion

Although increasing evidence has been generated by various studies regarding the significance of O-GlcNAcylation in regulating synaptic functions, the different experimental designs used have resulted in conflicting conclusions on the impact of changing the levels of O-GlcNAcylation. Here we used mice with a heterozygous loss-of-function mutation in OGA which have elevated O-GlcNAc levels. We found that OGA is highly expressed in the hippocampus, suggesting that O-GlcNAc modification of neuronal proteins is closely related to hippocampus-dependent functions. *Oga*^{+/-} mice exhibited impaired synaptic plasticity in the hippocampus at SC-CA1 synapses, dysregulated phosphorylation of AMPA receptor subunit GluA1 in chemically induced LTP




Figure 3. Normal synaptic transmission in *Oga*^{+/-} mice. (a) Representative traces of action potentials triggered by -100 – 200 pA current injection in the hippocampal CA1 region of WT and *Oga*^{+/-} mice. (b) Number of action potentials triggered by the injection of current at different levels (WT, $n = 11$; *Oga*^{+/-}, $n = 9$; unpaired t -test, not significant) (c) Representative mEPSC traces from WT and *Oga*^{+/-} hippocampal CA1 pyramidal neurons (upper). Average values for mEPSC amplitude (lower left) and frequency (lower right) (WT, $n = 9$; *Oga*^{+/-}, $n = 10$; unpaired t -test, NS: not significant) (d) Representative mIPSC traces from WT and *Oga*^{+/-} hippocampal CA1 pyramidal neurons (upper). Average values for mIPSC amplitude (lower left) and frequency (lower right). (WT, $n = 10$; *Oga*^{+/-}, $n = 8$; unpaired t -test, NS: not significant) (e) AMPA/NMDA current ratio (WT, $n = 20$; *Oga*^{+/-}, $n = 17$; Mann-Whitney U test, NS: not significant).

and LTD, and deficits in hippocampus-dependent learning and memory. This result together demonstrates that increased levels of O-GlcNAcylation lead to altered synaptic plasticity in the hippocampus, which may underlie the impairment of learning and memory observed in *Oga*^{+/-} mice.

Several studies have shown that synaptic plasticity is variably affected by O-GlcNAcylation. Tallent *et al.* showed that the elevation of O-GlcNAcylation induced by OGA inhibitor (9d) decreases PPF and increases LTP induction, and suggested that the elevation of O-GlcNAcylation facilitated LTP by modulating the interplay between phosphorylation and O-GlcNAcylation of signaling molecules, such as synapsin I/II, ERK, and CaMKII²⁴. In the same study, the authors also reported that reduced O-GlcNAcylation with OGT inhibitor (Alloxan) prevents LTP induction²⁴. However, contrary to this result, Kanno *et al.* found that Alloxan enhances hippocampal SC-CA1 LTP by regulating AMPA receptor trafficking³⁰. Taylor *et al.* also showed that acutely elevated O-GlcNAcylation by OGA inhibitor (Thiamet-G) or glucosamine induces LTD, but impairs LTP at CA3-CA1 synapses, which also led to a deficit in novel object recognition¹⁸. Each study mentioned above used different methods to change the levels of O-GlcNAcylation. Alloxan is known as a weak OGT inhibitor and thus likely to have off-target effects^{31,32}. Furthermore, as we previously reported, the OGA inhibitor (Thiamet-G) increases the levels of OGA expression^{33,34}. Glucosamine also affects




Figure 4. Impaired long-term potentiation (LTP) and long-term depression (LTD) in *Oga*^{+/-} mice.

(a) Input-output curves for basal synaptic transmission in area CA1 of the hippocampus. Representative traces are shown for the input (fiber volley) and the output (field excitatory postsynaptic potential; fEPSP). (WT, $n = 9$; *Oga*^{+/-}, $n = 10$; unpaired *t*-test, not significant) (b) Paired-pulse facilitation (PPF) in WT and *Oga*^{+/-} hippocampal CA1 pyramidal neurons. Representative traces from WT and *Oga*^{+/-} at 50 ms interstimulus interval are shown. (WT, $n = 8$; *Oga*^{+/-}, $n = 10$; unpaired *t*-test, not significant) (c) High frequency stimulation (HFS)-induced LTP (WT, $n = 10$; *Oga*^{+/-}, $n = 9$). Traces show averaged fEPSP indicated with 1 and 2. A bar graph is depicted 50 min after LTP. (d) LFS-induced LTD (WT, $n = 7$; *Oga*^{+/-}, $n = 8$). Traces show averaged fEPSP indicated with 1 and 2. A bar graph is depicted 50 min after LTD. Error bars represent \pm standard error of the mean (SEM). *** $P < 0.001$ (unpaired *t*-test).

various intracellular signaling pathways^{35–37}. Each experiment was performed in acutely elevated O-GlcNAcylation by pretreatment of OGA inhibitors or glucosamine. The different experimental designs might have resulted in conflicting results. Previously, the discrepancy in the effect of dysregulated O-GlcNAcylation was also observed in other intracellular signaling pathways and physiological functions^{33,34,38}. Despite this discrepancy, earlier studies suggest that dysregulated O-GlcNAcylation can affect synaptic plasticity. Here, we used OGA heterozygous mice that have elevated O-GlcNAcylation levels. *Oga*^{+/-} hippocampus displayed impaired regulation of AMPAR GluA1 phosphorylation which plays an important role in mediating AMPAR trafficking during synaptic plasticity. Although Taylor *et al.* showed that GluA1 is not O-GlcNAcylation¹⁸, the phosphorylation of GluA1 can be indirectly regulated by activation of upstream signaling molecules, including protein kinase C (PKC), CaMKII, and protein kinase A (PKA)^{39–41}. Importantly, both PKC and CaMKII are modified by O-GlcNAcylation^{21,42}, and the dynamic interplay between O-GlcNAcylation and phosphorylation in neurons was shown to be involved in hippocampal synaptic plasticity²⁴. Activation of PKC or PKA reduces global O-GlcNAc levels in cytoskeletal fraction of cultured cerebellar neurons⁴³. We speculate that GluA1 phosphorylation might be affected by altered O-GlcNAcylation levels. However, we cannot rule out the possibility that *Oga* heterozygosity affects multiple signaling pathways involved in hippocampal LTP and LTD.

ROS play an important role in synaptic plasticity⁴⁴ by regulating synaptic plasticity-related signaling molecules, receptors, and channels^{45–47}. Importantly, O-GlcNAcylation have been shown to affect ROS generation⁴⁸. In addition, forkhead box O1 (FoxO1), a regulator of the transcription of the oxidative stress responsive enzymes catalase and MnSOD (SOD2), is O-GlcNAcylation⁴⁹. We therefore examined whether ROS levels are affected in *Oga*^{+/-} hippocampus compared to WT hippocampus. Despite *Oga* heterozygosity, the ROS levels were not altered in *Oga*^{+/-} hippocampus (Fig. S3).

Aging is associated with impairments in cognitive and synaptic function⁵⁰. Dysfunction of the aging brain is not caused by neuronal loss⁵¹ but by specific alterations in neuronal morphology, cell-cell interactions, and gene expression⁵⁰. The hippocampus appears to be particularly vulnerable to the effects of aging on cognitive function and synaptic plasticity. O-GlcNAcylation and its regulatory enzymes are highly detected in the hippocampus⁵². O-GlcNAcylation modulates neuronal cell signaling processes and gene expression, which is critical for




Figure 5. Deficit in chemically induced LTP and LTD in *Oga*^{+/-} mice. (a) Representative immunoblots showing effects of chemical LTP and LTD induction on AMPAR subunits GluA1 S845, S831 phosphorylation. Acute hippocampal slices were stimulated with either glycine (200 μ M) for chemical LTP or NMDA (20 μ M) for chemical LTD. Levels of phospho-S845 and phospho-S831 of GluA1, and levels of total O-GlcNAc-modified proteins were analyzed by immunoblotting. (b) Chemical LTP and LTD significantly increases and decreases the levels of GluA1 S845 phosphorylation, respectively in WT ($n = 4$, normalized to control). However, acute hippocampal slices from *Oga*^{+/-} mice failed to exhibit a significant change in GluA1 S845 phosphorylation following chemical LTP and LTD induction ($n = 3$, normalized to control). One-way ANOVA followed by the Tukey's test was used. (c) Chemical LTP significantly increases the levels of GluA1 S831 phosphorylation in WT ($n = 3$, normalized to control), but not in *Oga*^{+/-} mice ($n = 3$, normalized to control). Error bars represent \pm standard error of the mean (SEM). NS: not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; one-way ANOVA followed by the Tukey's test. Full-length blots/gels are presented in Supplementary Figure S4.

proper neuronal function^{1,53}. Interestingly, we previously reported that the brains of older mice show significantly increased levels of O-GlcNAcylation compared with those in younger mice³. However, the mechanism underlying the chronic elevations in O-GlcNAcylation on brain aging remains unknown. Based on our observations, in the normally aged brain, we speculate that chronically elevated O-GlcNAcylation contributes to impairment of synaptic plasticity and learning and memory.

Methods

Mice. *Oga*^{+/-} mice (C57BL/6J) were generated as described previously³. All mice were housed under a 12-hour light/dark cycle and given *ad libitum* access to food and water. All experimental protocols were approved by Institutional Animal Care and Use Committee of the Ulsan National Institute of Science and Technology (UNIST-IACUC-14-018) and all methods were performed in accordance with the relevant guidelines and regulations.

Barnes maze. The paradigm of Barnes circular maze consists of white circular platform (92 cm diameter), with 20 evenly spaced holes (5 cm diameter) located 7.5 cm from the perimeter and is elevated 100 cm above the floor. Several spatial cues with distinct shapes were placed near the walls of the testing room. A black target box (20 \times 10 \times 10 cm) was placed under one hole. The mice were encouraged to find this box by aversive noise (85 dB) on the platform. Barnes maze were run for 4 consecutive days, and 3 trials were carried out each day with 20 min inter-trial intervals. The mouse was allowed to search for the target box for 3 min. Distance, latency, and numbers of errors to reach the target hole were recorded during training trials by video tracking software. On day 5, a probe test was performed without the escape box. Mice were allowed to freely find the target box for 3 min. Time spent around each hole, total distance travelled, and the latency to find target hole were recorded.

Golgi staining. Brains from 8-week-old mice were processed with the FD Rapid GolgiStain™ Kit (NeuroTechnologies) according to the instructions of the manufacturer. Images of dendritic spines (apical dendrites of CA1 pyramidal neurons) were acquired using an Olympus Cell^TIRF Xcellence microscope in UNIST-Olympus Biomed Imaging Center (UOBC).

Chemical LTP and LTD induction. Acute hippocampal slices (300- μ m thick) from WT or *Oga*^{+/-} mice (8–10 weeks) were prepared in a sucrose-cutting buffer containing (in mM) 234 sucrose, 2.5 KCl, 1.25 NaH₂PO₄,

24 NaHCO₃, 11 glucose, 10 MgSO₄, 0.5 CaCl₂ bubbled with 95% O₂ and 5% CO₂. The slices were recovered at 35 °C for one hour in a recovery buffer containing (in mM) 124 NaCl, 3 KCl, 1.25 NaH₂PO₄, 26 NaHCO₃, 10 glucose, 6.5 MgSO₄, 1 CaCl₂ bubbled with 95% O₂ and 5% CO₂. Following the recovery, the slices were further incubated at 37 °C for one hour in an extracellular fluid containing (in mM) 125 NaCl, 2.5 KCl, 1 MgCl₂, 2 CaCl₂, 33 glucose, 25 HEPES, and then treated with 20 μM D-AP5 and 0.5 μM TTX for 20 min. The slices were subsequently treated with 3 μM strychnine, 20 μM bicuculline and 200 μM glycine for 10 min to induce chemical LTP, or with 20 μM NMDA for 3 min to induce chemical LTD in a Mg-free extracellular fluid, and transferred back to a regular extracellular fluid for 30 min prior to sample collection.

Statistical analysis. The Student's unpaired T-test or non-parametric Mann-Whitney U-test was used to compare two independent groups. For multiple comparisons, a one-way repeated measures ANOVA with Tukey's *post hoc* test was utilized, as specified in the Figure legends. All data are expressed as the mean ± SEM and significance indicated by *P < 0.05, **P < 0.01, and ***P < 0.001.

References

- Hart, G. W., Slawson, C., Ramirez-Correa, G. & Lagerlof, O. Cross talk between O-GlcNAcylation and phosphorylation: roles in signaling, transcription, and chronic disease. *Annu Rev Biochem* **80**, 825–858, doi: 10.1146/annurev-biochem-060608-102511 (2011).
- Okuyama, R. & Marshall, S. UDP-N-acetylglucosaminyl transferase (OGT) in brain tissue: temperature sensitivity and subcellular distribution of cytosolic and nuclear enzyme. *J Neurochem* **86**, 1271–1280 (2003).
- Yang, Y. R. *et al.* O-GlcNAcase is essential for embryonic development and maintenance of genomic stability. *Aging Cell* **11**, 439–448, doi: 10.1111/j.1474-9726.2012.00801.x (2012).
- Yuzwa, S. A. *et al.* Increasing O-GlcNAc slows neurodegeneration and stabilizes tau against aggregation. *Nat Chem Biol* **8**, 393–399, doi: 10.1038/nchembio.797 (2012).
- Marotta, N. P. *et al.* O-GlcNAc modification blocks the aggregation and toxicity of the protein alpha-synuclein associated with Parkinson's disease. *Nat Chem* **7**, 913–920, doi: 10.1038/nchem.2361 (2015).
- Arnold, C. S. *et al.* The microtubule-associated protein tau is extensively modified with O-linked N-acetylglucosamine. *J Biol Chem* **271**, 28741–28744 (1996).
- Liu, F., Iqbal, K., Grundke-Iqbal, I., Hart, G. W. & Gong, C. X. O-GlcNAcylation regulates phosphorylation of tau: a mechanism involved in Alzheimer's disease. *Proc Natl Acad Sci USA* **101**, 10804–10809, doi: 10.1073/pnas.0400348101 (2004).
- Alfaro, J. F. *et al.* Tandem mass spectrometry identifies many mouse brain O-GlcNAcylated proteins including EGF domain-specific O-GlcNAc transferase targets. *Proc Natl Acad Sci USA* **109**, 7280–7285, doi: 10.1073/pnas.1200425109 (2012).
- Wang, Z. *et al.* Enrichment and site mapping of O-linked N-acetylglucosamine by a combination of chemical/enzymatic tagging, photochemical cleavage, and electron transfer dissociation mass spectrometry. *Mol Cell Proteomics* **9**, 153–160, doi: 10.1074/mcp.M900268-MCP200 (2010).
- Griffith, L. S., Mathes, M. & Schmitz, B. Beta-amyloid precursor protein is modified with O-linked N-acetylglucosamine. *J Neurosci Res* **41**, 270–278, doi: 10.1002/jnr.490410214 (1995).
- Kim, C. *et al.* O-linked beta-N-acetylglucosaminidase inhibitor attenuates beta-amyloid plaque and rescues memory impairment. *Neurobiol Aging* **34**, 275–285, doi: 10.1016/j.neurobiolaging.2012.03.001 (2013).
- Whitcomb, D. J. *et al.* Intracellular oligomeric amyloid-beta rapidly regulates GluA1 subunit of AMPA receptor in the hippocampus. *Sci Rep* **5**, 10934, doi: 10.1038/srep10934 (2015).
- Jo, J. *et al.* Abeta(1-42) inhibition of LTP is mediated by a signaling pathway involving caspase-3, Akt1 and GSK-3beta. *Nat Neurosci* **14**, 545–547, doi: 10.1038/nn.2785 (2011).
- Akimoto, Y. *et al.* Localization of the O-GlcNAc transferase and O-GlcNAc-modified proteins in rat cerebellar cortex. *Brain Res* **966**, 194–205 (2003).
- Cole, R. N. & Hart, G. W. Cytosolic O-glycosylation is abundant in nerve terminals. *Journal of Neurochemistry* **79**, 1080–1089, doi: DOI 10.1046/j.1471-4159.2001.00655.x (2001).
- Vosseller, K. *et al.* O-linked N-acetylglucosamine proteomics of postsynaptic density preparations using lectin weak affinity chromatography and mass spectrometry. *Mol Cell Proteomics* **5**, 923–934, doi: 10.1074/mcp.T500040-MCP200 (2006).
- Ogawa, M., Sawaguchi, S., Kamemura, K. & Okajima, T. Intracellular and extracellular O-linked N-acetylglucosamine in the nervous system. *Exp Neurol* **274**, 166–174, doi: 10.1016/j.expneurol.2015.08.009 (2015).
- Taylor, E. W. *et al.* O-GlcNAcylation of AMPA Receptor GluA2 Is Associated with a Novel Form of Long-Term Depression at Hippocampal Synapses. *J Neurosci* **34**, 10–21, doi: 10.1523/Jneurosci.4761-12.2014 (2014).
- Cole, R. N. & Hart, G. W. Glycosylation sites flank phosphorylation sites on synapsin I: O-linked N-acetylglucosamine residues are localized within domains mediating synapsin I interactions. *Journal of Neurochemistry* **73**, 418–428, doi: DOI 10.1046/j.1471-4159.1999.0730418.x (1999).
- Skorobogatko, Y. *et al.* O-linked beta-N-acetylglucosamine (O-GlcNAc) site thr-87 regulates synapsin I localization to synapses and size of the reserve pool of synaptic vesicles. *J Biol Chem* **289**, 3602–3612, doi: 10.1074/jbc.M113.512814 (2014).
- Erickson, J. R. *et al.* Diabetic hyperglycaemia activates CaMKII and arrhythmias by O-linked glycosylation. *Nature* **502**, 372–376, doi: 10.1038/nature12537 (2013).
- Rexach, J. E. *et al.* Dynamic O-GlcNAc modification regulates CREB-mediated gene expression and memory formation. *Nat Chem Biol* **8**, 253–261, doi: 10.1038/nchembio.770 (2012).
- Dias, W. B., Cheung, W. D., Wang, Z. & Hart, G. W. Regulation of calcium/calmodulin-dependent kinase IV by O-GlcNAc modification. *J Biol Chem* **284**, 21327–21337, doi: 10.1074/jbc.M109.007310 (2009).
- Tallent, M. K. *et al.* *In vivo* modulation of O-GlcNAc levels regulates hippocampal synaptic plasticity through interplay with phosphorylation. *J Biol Chem* **284**, 174–181, doi: 10.1074/jbc.M807431200 (2009).
- Huganir, R. L. & Nicoll, R. A. AMPARs and synaptic plasticity: the last 25 years. *Neuron* **80**, 704–717, doi: 10.1016/j.neuron.2013.10.025 (2013).
- Lee, H. K., Kameyama, K., Huganir, R. L. & Bear, M. F. NMDA induces long-term synaptic depression and dephosphorylation of the GluR1 subunit of AMPA receptors in hippocampus. *Neuron* **21**, 1151–1162 (1998).
- Hosokawa, T., Mitsushima, D., Kaneko, R. & Hayashi, Y. Stoichiometry and phosphoisotypes of hippocampal AMPA-type glutamate receptor phosphorylation. *Neuron* **85**, 60–67, doi: 10.1016/j.neuron.2014.11.026 (2015).
- Lisman, J., Schulman, H. & Cline, H. The molecular basis of CaMKII function in synaptic and behavioural memory. *Nat Rev Neurosci* **3**, 175–190, doi: 10.1038/nrn753 (2002).
- Benito, E. & Barco, A. CREB's control of intrinsic and synaptic plasticity: implications for CREB-dependent memory models. *Trends Neurosci* **33**, 230–240, doi: 10.1016/j.tins.2010.02.001 (2010).
- Kanno, T., Yaguchi, T., Nagata, T., Mukasa, T. & Nishizaki, T. Regulation of AMPA receptor trafficking by O-glycosylation. *Neurochem Res* **35**, 782–788, doi: 10.1007/s11064-010-0135-1 (2010).

31. Konrad, R. J. *et al.* Alloxan is an inhibitor of the enzyme O-linked N-acetylglucosamine transferase. *Biochem Biophys Res Commun* **293**, 207–212, doi: Pii S0006-291x(02)00200-0 10.1016/S0006-291x(02)00200-0 (2002).
32. Tiedge, M., Richter, T. & Lenzen, S. Importance of cysteine residues for the stability and catalytic activity of human pancreatic beta cell glucokinase. *Arch Biochem Biophys* **375**, 251–260, doi: 10.1006/abbi.1999.1666 (2000).
33. Yang, Y. R. *et al.* Elevated O-GlcNAcylation promotes colonic inflammation and tumorigenesis by modulating NF-kappaB signaling. *Oncotarget* **6**, 12529–12542, doi: 10.18632/oncotarget.3725 (2015).
34. Yang, Y. R. *et al.* OGA heterozygosity suppresses intestinal tumorigenesis in Apc(min/+) mice. *Oncogenesis* **3**, e109, doi: 10.1038/oncsis.2014.24 (2014).
35. d'Abusco, A. S. *et al.* Glucosamine affects intracellular signalling through inhibition of mitogen-activated protein kinase phosphorylation in human chondrocytes. *Arthritis Res Ther* **9**, doi: ARTN R104 10.1186/ar2307 (2007).
36. Goldberg, H. J., Scholey, J. & Fantus, I. G. Glucosamine activates the plasminogen activator inhibitor 1 gene promoter through Sp1 DNA binding sites in glomerular mesangial cells. *Diabetes* **49**, 863–871 (2000).
37. Liu, B. Q. *et al.* Glucosamine induces cell death via proteasome inhibition in human ALVA41 prostate cancer cell. *Exp Mol Med* **43**, 487–493, doi: 10.3858/emmm.2011.43.9.055 (2011).
38. Yang, Y. R. *et al.* Obesity resistance and increased energy expenditure by white adipose tissue browning in Oga(+/-) mice. *Diabetologia* **58**, 2867–2876, doi: 10.1007/s00125-015-3736-z (2015).
39. Mammen, A. L., Kameyama, K., Roche, K. W. & Haganir, R. L. Phosphorylation of the alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor GluR1 subunit by calcium/calmodulin-dependent kinase II. *J Biol Chem* **272**, 32528–32533 (1997).
40. Roche, K. W., O'Brien, R. J., Mammen, A. L., Bernhardt, J. & Haganir, R. L. Characterization of multiple phosphorylation sites on the AMPA receptor GluR1 subunit. *Neuron* **16**, 1179–1188 (1996).
41. Boehm, J. *et al.* Synaptic incorporation of AMPA receptors during LTP is controlled by a PKC phosphorylation site on GluR1. *Neuron* **51**, 213–225, doi: 10.1016/j.neuron.2006.06.013 (2006).
42. Robles-Flores, M. *et al.* Posttranslational modifications on protein kinase C isozymes. Effects of epinephrine and phorbol esters. *Biochim Biophys Acta* **1783**, 695–712, doi: 10.1016/j.bbamcr.2007.07.011 (2008).
43. Griffith, L. S. & Schmitz, B. O-linked N-acetylglucosamine levels in cerebellar neurons respond reciprocally to perturbations of phosphorylation. *Eur J Biochem* **262**, 824–831 (1999).
44. Knapp, L. T. & Klann, E. Role of reactive oxygen species in hippocampal long-term potentiation: contributory or inhibitory? *J Neurosci Res* **70**, 1–7, doi: 10.1002/jnr.10371 (2002).
45. Betzen, C. *et al.* Oxidative stress upregulates the NMDA receptor on cerebrovascular endothelium. *Free Radic Biol Med* **47**, 1212–1220, doi: 10.1016/j.freeradbiomed.2009.07.034 (2009).
46. Huddleston, A. T., Tang, W., Takeshima, H., Hamilton, S. L. & Klann, E. Superoxide-induced potentiation in the hippocampus requires activation of ryanodine receptor type 3 and ERK. *J Neurophysiol* **99**, 1565–1571, doi: 10.1152/jn.00659.2007 (2008).
47. Gong, L., Gao, T. M., Huang, H. & Tong, Z. Redox modulation of large conductance calcium-activated potassium channels in CA1 pyramidal neurons from adult rat hippocampus. *Neurosci Lett* **286**, 191–194 (2000).
48. Ngho, G. A., Watson, L. J., Facundo, H. T. & Jones, S. P. Augmented O-GlcNAc signaling attenuates oxidative stress and calcium overload in cardiomyocytes. *Amino Acids* **40**, 895–911, doi: 10.1007/s00726-010-0728-7 (2011).
49. Housley, M. P. *et al.* A PGC-1alpha-O-GlcNAc transferase complex regulates FoxO transcription factor activity in response to glucose. *J Biol Chem* **284**, 5148–5157, doi: 10.1074/jbc.M808890200 (2009).
50. Burke, S. N. & Barnes, C. A. Neural plasticity in the ageing brain. *Nat Rev Neurosci* **7**, 30–40, doi: 10.1038/nrn1809 (2006).
51. Morrison, J. H. & Hof, P. R. Life and death of neurons in the aging brain. *Science* **278**, 412–419 (1997).
52. Liu, K. *et al.* Accumulation of protein O-GlcNAc modification inhibits proteasomes in the brain and coincides with neuronal apoptosis in brain areas with high O-GlcNAc metabolism. *J Neurochem* **89**, 1044–1055, doi: 10.1111/j.1471-4159.2004.02389.x (2004).
53. Lazarus, B. D., Love, D. C. & Hanover, J. A. O-GlcNAc cycling: implications for neurodegenerative disorders. *Int J Biochem Cell Biol* **41**, 2134–2146, doi: 10.1016/j.biocel.2009.03.008 (2009).

Acknowledgements

This research was supported by the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (NRF-2013R1A1A2064434 and NRF-2016M3C7A1913845), the Korean Government (MSIP) (2010-0028684), and an intramural funding from Korea Institute of Science and Technology (2E26820).

Author Contributions

Y.R.Y. and S.S. contributed equally in designing and conducting the experiments and performing data analysis. J.H.J., S.K. and H.H. conducted the electrophysiological, biochemical experiments and behavioral tests. J.H.H. assisted with microscopic image analysis. S.Y., D.N., J.P., C.L., Y.K.S., and J.H.K. discussed the hypothesis and interpreted the data. P.G.S. and H.R. directed the project and advised during the designing of the figures and writing of the manuscript. All the authors have read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing Interests: The authors declare no competing financial interests.

How to cite this article: Yang, Y. R. *et al.* Memory and synaptic plasticity are impaired by dysregulated hippocampal O-GlcNAcylation. *Sci. Rep.* **7**, 44921; doi: 10.1038/srep44921 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017