

# SCIENTIFIC REPORTS



OPEN

## ATHB17 enhances stress tolerance by coordinating photosynthesis associated nuclear gene and *ATSIG5* expression in response to abiotic stress

Received: 20 October 2016  
Accepted: 28 February 2017  
Published: 30 March 2017

Ping Zhao\*, Rong Cui\*, Ping Xu, Jie Wu, Jie-Li Mao, Yu Chen, Cong-Zhao Zhou, Lin-Hui Yu & Cheng-Bin Xiang

Photosynthesis is sensitive to environmental stress and must be efficiently modulated in response to abiotic stress. However, the underlying mechanisms are not well understood. Here we report that *ARABIDOPSIS THALIANA* HOMEBOX 17 (*ATHB17*), an *Arabidopsis* HD-Zip transcription factor, regulated the expression of a number of photosynthesis associated nuclear genes (PhANGs) involved in the light reaction and *ATSIG5* in response to abiotic stress. *ATHB17* was responsive to ABA and multiple stress treatments. *ATHB17*-overexpressing plants displayed enhanced stress tolerance, whereas its knockout mutant was more sensitive compared to the wild type. Through RNA-seq and quantitative real-time reverse transcription PCR (qRT-PCR) analysis, we found that *ATHB17* did not affect the expression of many known stress-responsive marker genes. Interestingly, we found that *ATHB17* down-regulated many PhANGs and could directly modulate the expression of several PhANGs by binding to their promoters. Moreover, we identified *ATSIG5*, encoding a plastid sigma factor, as one of the target genes of *ATHB17*. Loss of *ATSIG5* reduced salt tolerance while overexpression of *ATSIG5* enhanced salt tolerance, similar to that of *ATHB17*. *ATHB17* can positively modulate the expression of many plastid encoded genes (PEGs) through regulation of *ATSIG5*. Taken together, our results suggest that *ATHB17* may play an important role in protecting plants by adjusting expression of PhANGs and PEGs in response to abiotic stresses.

Abiotic stresses such as salinity, drought, high light and unfavorable temperatures, adversely affect the growth and development of plants. Photosynthesis in chloroplasts is one of the primary processes to be affected by abiotic stress<sup>1</sup>. The effects can be direct, such as decreased CO<sub>2</sub> diffusion caused by stomata closure<sup>2</sup> or affecting ribulose biphosphate carboxylase/oxygenase (Rubisco) activity<sup>3</sup>. More importantly, abiotic stress reduces the threshold intensity for the onset of photo-inhibition, and results in over excitation of the photosystems (PSs), dramatically increasing reactive oxygen species (ROS) production<sup>4,5</sup>. Therefore, rapid response of photosynthetic machinery and metabolism is key for plants to cope with the fluctuating environment<sup>6</sup>.

Chloroplasts are genetically semi-autonomous organelles that evolutionarily retain an eubacteria-type of circular genome DNA. In higher plants, the chloroplast 120–150 kb genome encodes only about 120 genes<sup>7</sup>. More required proteins are encoded by the nuclear genome and imported to play roles in the chloroplasts after translation in the cytosol<sup>8</sup>.

Chloroplast gene transcription in higher plants is performed by at least two types of RNA polymerases, plastid encoded RNA polymerase (PEP) and nuclear encoded RNA polymerase (NEP). PEP holoenzyme is a complex composed by five subunits:  $\alpha$ ,  $\beta$ ,  $\beta'$ ,  $\omega$  (omega),  $\sigma$  (sigma), in which the  $\alpha\beta\beta'\omega$  constitutes the catalytic core, while the nuclear-encoded  $\sigma$  subunit recognizes the specific promoter region and initiates transcription of the core complex<sup>9</sup>.

School of Life Sciences, University of Science and Technology of China, Hefei, Anhui Province, 230027, PR China.  
\*These authors contributed equally to this work. Correspondence and requests for materials should be addressed to L.-H.Y. (email: yulh@ustc.edu.cn) or C.-B.X. (email: xiangcb@ustc.edu.cn)

The transcription of photosynthesis related genes in chloroplasts is mainly dependent on PEP, and the nuclear-encoded sigma factors play special roles in regulating the chloroplast transcription<sup>10,11</sup>. Since the first chloroplast sigma factor gene was isolated from red algae nuclear genome<sup>12,13</sup>, more chloroplast sigma factors (e.g. *ATSIG1-6*) had been identified in different plant species<sup>14,15</sup>. Most plastid encoded genes appear to be regulated by several sigma factors with overlapping functions. However, within a particular time frame during plant development, plastid genes are likely to be coordinated by a distinct sigma factor, for example, *psaA* and *rbcl* by *ATSIG1*<sup>16</sup>, *psaJ* by *ATSIG2*<sup>17</sup>, *psbN* by *ATSIG3*<sup>18</sup>, *ndhF* by *ATSIG4*<sup>19</sup>, *psbBT*, *psbD*, *psbC*, *psbZ*, *psbA* and *psaAB* by *ATSIG5*<sup>17,20–22</sup>.

Among the six sigma factors in *Arabidopsis*, only *ATSIG5* expression is stress inducible and phylogenetically specific<sup>23–25</sup>. It is induced by high light, low temperature, high salt and osmotic stress<sup>17</sup>, as well as blue light<sup>23</sup>. Beside these stresses, *MpSIG5* of liverwort *Marchantia polymorpha* is significantly induced by reactive oxygen species (ROS) stress<sup>26</sup>. *ATSIG5* regulates the repair capacity from injury to the PS II reaction center by salt stress. It does this by determining the promoter recognition specificity of PEP in plastid gene expression that activates *psbD* from the blue-light responsive promoter (BLRP)<sup>9,27</sup>. In addition, *ATSIG5* regulates chloroplast *psbD* and *psbA* coding for the PS II core proteins D1 and D2 in response to light quality and intensity, and combines extrinsic and intrinsic signals important for adjusting nuclear and plastid gene transcription in light acclimation processes<sup>28</sup>.

Coordinating the transcription of photosynthesis associated nuclear genes (PhANGs) and plastid encoded genes (PEGs) to maintain the proper stoichiometry of nuclear encoded proteins, plastid proteins, carotenoids and chlorophylls, is critical for correct assembly of functional photoprotective and photosynthetic complexes within chloroplasts under stress conditions<sup>29</sup>. Although many abiotic stress-responsive transcription factors (TFs) have been studied, very few are known to modulate the expression of photosynthesis-related genes<sup>5</sup>.

The homeodomain leucine-zipper (HD-Zip) TFs are the most abundant group of homeobox genes in plants, which have diverse functions during plant development and stress adaptation<sup>30,31</sup>. According to their distinctive features such as gene structures, DNA-binding specificities, additional common motifs and physiological functions, HD-Zip TFs can be classified into four subfamilies<sup>30</sup>. There are 10 HD-Zip II genes in the *Arabidopsis* genome, which play important roles ranging from auxin response to shade avoidance. Five HD-Zip class II genes, including *HOMEODOMAIN ARABIDOPSIS THALIANA 1 (HAT1)*, *HAT2*, *HAT3*, *ATHB2*, *ATHB4*, are known to respond to light quality changes<sup>32</sup>. Auxin response analyses strongly suggest that *HAT1*, *HAT3* and *ATHB4* are under the control of the phytochrome system as is *ATHB2*<sup>33</sup>. Of the remaining class II HD-Zip members, little is known about their functions, excepting *ATHB17*.

*ATHB17* localizes to both the cytoplasm and nuclei; the distribution is determined by its unique N-terminus. Overexpression of *ATHB17* in *Arabidopsis* enhances chlorophyll content in the leaves, while expression of a truncated *ATHB17* protein in maize increases ear weight at silking<sup>34,35</sup>. *ATHB17*-overexpressing *Arabidopsis* plants are sensitive to ABA and NaCl, whereas *ATHB17* knockout mutants are insensitive to ABA and NaCl at post germination stage. However, these phenotypes are weak and the expression of ABA-responsive genes is not significantly altered in the *ATHB17*-overexpressing plants compared with wild-type (WT) plants<sup>36</sup>. At this stage it remains unclear if the phenotypes result from modulating ABA signaling or other mechanisms.


In contrast, in this study, we found that *ATHB17* regulated the expression of PhANGs and *ATSIG5* to cope with environmental stresses. *ATHB17* responded to multiple abiotic stresses. Overexpression of *ATHB17* enhanced plant tolerance to salt, drought and oxidative stresses, and *ATHB17* knockout resulted in the opposite phenotypes. By RNA-seq analysis, we find *ATHB17* repressed the expression of many PhANGs while it activated the transcription of *ATSIG5*. Further analysis revealed that *ATHB17* directly bound to the promoter of several PhANGs and most likely regulated their expression. Moreover, *ATHB17* directly bound to the *ATSIG5* promoter and activated *ATSIG5* expression to regulate some PEG expression to deal with environmental stress. Our study revealed another stress response pathway that modulates the photosynthesis light reaction by down regulating PhANGs and up regulating PEGs in response to multiple abiotic stresses, therefore protecting plants from photosynthesis-derived damage.

## Results

***ATHB17* is preferentially expressed in roots and responsive to multiple stress signals.** To reveal the expression pattern of *ATHB17*, we analyzed transgenic plants harboring the *ATHB17* promoter-GUS reporter construct (*pATHB17::GUS*). Strong GUS activity was detected in roots of seedlings at different ages (Fig. 1A-a to A-e and A-g). *ATHB17* was also expressed in rosette leaves with much higher expression levels in the leaf veins (Fig. 1A-f). However, at mature stage, *ATHB17* was mainly expressed in roots (Fig. 1A-g), in agreement with the results of Park *et al.*<sup>36</sup>. There was only weak expression in other organs, such as rosette leaf (Fig. 1A-h), cauline leaf (Fig. 1A-i), flower and young silique (Fig. 1A-j), and mature silique (Fig. 1A-l). These results were confirmed by quantitative real-time reverse transcription PCR (qRT-PCR) analysis shown in Fig. 1B, implying a potential function of *ATHB17* in roots.

Moreover, we found that *ATHB17* could be significantly induced by ABA (abscisic acid), paraquat, drought, and NaCl treatments (Fig. 1C). Consistent with this result, GUS staining of the *pATHB17::GUS* reporter line treated with NaCl, ABA, PQ, and mannitol also showed strongly induced expression of *ATHB17*, especially in the leaves (Fig. 1D).

To study the localization of *ATHB17* protein in plant cell, we obtained transgenic plants expressing green fluorescent protein (GFP) and *ATHB17* fusion protein under the control of its own promoter (*pATHB17::ATHB17::GFP*). By examining GFP fluorescence in root of the transgenic plant, we found that *ATHB17* was predominantly localized in the nucleus (Fig. 1E), which is consistent with the previous report<sup>36</sup>.




**Figure 1. Expression pattern of *ATHB17* and response to different stress signals.** (A) The expression pattern of *ATHB17* was revealed by GUS staining of *pATHB17::GUS* transgenic plants. GUS activity was observed in 2-day-old (a), 4-day-old (b), 7-day-old (c), 9-day-old (d), or 14-day-old (e) seedlings, leaves of 9-day-old plant (f), roots of mature plants (g), rosette leaf (h), cauline leaf (i), flower and young silique (j), old silique (k). (B) Analysis of the *ATHB17* expression patterns by qRT-PCR. UBQ5 was used as an internal control. Values are mean  $\pm$  SD of three replica experiments. (C) Expression levels of *ATHB17* after different treatments. Induction levels of *ATHB17* in 10-day-old plants by ABA (100  $\mu$ M, 4 h), paraquat (5  $\mu$ M, 4 h), drought (4 h), NaCl (200 mM, 4 h) were determined by qRT-PCR. Values are mean  $\pm$  SD of three replica experiments (Student's t-test \* $P < 0.05$ , \*\* $P < 0.01$ ). (D) Effects of different stress treatments on *pATHB17::GUS* expression, 7-day-old or 14-day-old *pATHB17::GUS* transgenic seedlings grown on MS medium were transferred to MS liquid medium or MS liquid medium containing NaCl (200 mM), ABA (10  $\mu$ M), paraquat (5  $\mu$ M), mannitol (200 mM) for 1–6 h and then the seedlings were harvested for GUS staining for 3 h. (E) *ATHB17* protein localization in the root cells of transgenic plants expressing *ATHB17*-GFP under the control of the *ATHB17* native promoter.

***ATHB17* is a positive regulator of tolerance to multiple abiotic stresses.** In order to uncover the functions of *ATHB17*, we generated 35S:: *ATHB17* overexpression (OX) transgenic *Arabidopsis* plants, and obtained an *ATHB17* knockout (KO) mutant from Arabidopsis Biological Resource Center (Supplementary Fig. S2). To test whether *ATHB17* was involved in salt tolerance, we determined the NaCl sensitivity of the *ATHB17* OX and KO lines. Firstly, the NaCl sensitivity at germination and seedling establishment stage was assayed. Seeds were germinated and grown on MS (Murashige and Skoog) medium containing different concentration of NaCl. The results showed that the root growth of *ATHB17* OX lines was more resistant to salt stress compared with the WT plants. In contrast, the *ATHB17* KO plants were more sensitive to NaCl stress with significantly reduced root growth under stress. However, there was no difference in germination or root elongation on MS medium without NaCl (Fig. 2A,B). In another salt tolerance assay, 7-day-old *ATHB17* OX, KO and WT plants were transferred to MS medium or MS medium containing NaCl. After another 7 days growth, the *ATHB17* OX plants showed higher survival ratio whereas the *ATHB17* KO plants showed opposite phenotypes compared with the WT (Fig. 2C,D). Moreover, salt tolerance assay with older plants grown in soil showed that *ATHB17* positively regulated plant survival to salt stress (Supplementary Fig. S2).

To confirm this, we conducted complementation analysis by expressing *ATHB17* in the *ATHB17* KO plants, which restored the salt tolerance of KO plants to WT level (Supplementary Fig. S3). In addition, *ATHB17* was able to confer salt tolerance in tobacco at different developmental stages when overexpressed (Supplementary Fig. S4).

Since *ATHB17* was responsive to multiple stresses as demonstrated above, we also analyzed the role of *ATHB17* in drought stress and oxidative stress. The *ATHB17* OX *Arabidopsis* plants showed enhanced drought tolerance



**Figure 2.** *ATHB17* is a positive regulator for salt stress resistance. (A,B) Salt tolerance assay of *ATHB17* OX and KO Phenotypes of seedlings (A) and root length (B) of *ATHB17* OX, KO and WT plants grown on medium containing indicated concentrations of NaCl. Seeds were germinated and grown vertically on MS medium or MS medium containing the indicated concentration of NaCl for 10 days. Values are mean  $\pm$  SD ( $n = 60$ , \* $P < 0.05$ , \*\* $P < 0.01$ ). Bar = 1 cm. (C) Phenotype of *ATHB17* OX, KO and WT plants on MS medium with different concentrations of NaCl. Seven-day-old plants on MS medium were transferred to MS medium or MS medium containing the indicated concentrations of NaCl for 7 days. Bar = 4 cm. (D) Survival ratio of the *ATHB17* OX, KO and WT plants on MS medium or MS medium with indicated concentrations of NaCl. Values are mean  $\pm$  SD of three biological replicates each containing 14–16 plants per genotype (\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ).

while the *ATHB17* KO plants were drought sensitive compared with WT plants (Supplementary Fig. S5A,B), which is consistent with the result of Park *et al.*<sup>36</sup>. In addition, overexpression of *ATHB17* also conferred improved tolerance to paraquat (Supplementary Fig. S5C,D). These data indicate *ATHB17* is a positive regulator of multiple abiotic stresses.

**The expression of stress-responsive marker genes was not significantly affected by *ATHB17*.** To investigate the mechanism of *ATHB17* modulated stress tolerance, we compared the transcriptomes of *ATHB17* OX, KO and WT plants grown on NaCl-free MS medium and MS medium containing 200 mM NaCl using RNA-seq method. Surprisingly, we found the expression of classic stress-induced marker genes, such as *responsive to desiccation 29A* (*RD29A*), *RD29B*, *RD20*, *RD22*, *cold-regulated 47* (*COR47*), *COR45B*, *CRT/DRE binding factor 1* (*CBF1*), *salt overly sensitive 2* (*SOS2*) and *SOS3*, were not significantly influenced by *ATHB17* under both normal and salt treated conditions. Expression levels of ABA signaling and synthesis pathway genes were also found not significantly changed in the *ATHB17* OX and KO plants (Table 1). To validate the results of RNA-seq profiling analysis, eight stress-responsive genes were selected for qRT-PCR analysis. The results showed that the response of these genes was not significantly affected by the expression level of *ATHB17* (Fig. 3). These results implicate that *ATHB17*-conferred stress tolerance, probably arose not through the classical stress response pathway, but through a new pathway.

***ATHB17* negatively regulates some PhANGs.** Based on Gene Ontology (GO) term enrichment analysis of the RNA-seq profiling data, we found that the PhANGs were significantly enriched among the different expressed genes between the *ATHB17* OX and *ATHB17* KO plants both under normal and salt stress conditions (Supplementary Data 1 and 2). The expression of 26 of the 63 PhANGs in the *Arabidopsis* genome were down-regulated (>2 fold) in the *ATHB17* OX compared with *ATHB17* KO plants under normal or NaCl treated conditions. More specifically, under normal condition, 20 of the 26 PhANGs had reduced expression (>1.5 fold)

Gene ID	Annotation	Normal (TPM*)			NaCl (TPM*)		
		WT	OX	KO	WT	OX	KO
AT5G52310	RD29A (responsive to desiccation 29A)	0.99	0.38	0.01	71.54	73.85	52.09
AT5G52300	RD29B (responsive to desiccation 29B)	0.01	0.01	0.28	16.32	14.61	12.21
AT5G25610	RD22 (responsive to desiccation 22)	36.67	11.53	44.81	182.8	280.38	373.55
AT2G33380	RD20 (responsive to desiccation 20)	16.52	20.36	10.65	116.37	146.30	133.47
AT1G20440	COR47 (cold regulated 47)	943.44	770.26	1405.98	3953.93	4602.60	5296.48
AT2G42530	COR15B (cold regulated 15B)	0.01	0.01	0.01	4.12	3.80	6.10
AT4G25490	CBF1 (C-repeat/DRE binding factor 1)	0.01	0.01	0.01	15.49	20.61	19.94
AT4G26080	ABI1 (abscisic acid insensitive 1)	36.01	46.48	44.95	122.14	122.88	186.78
AT5G57050	ABI2 (abscisic acid insensitive 2)	0.66	5.57	2.63	74.01	64.04	61.45
AT2G36270	ABI5 (abscisic acid insensitive 3)	3.96	3.07	4.29	17.47	23.62	16.68
AT5G67030	ABA1 (abscisic acid deficient 1)	49.55	44.95	75.24	287.96	282.38	264.50
AT1G16540	ABA3 (abscisic acid deficient 3)	8.26	17.86	8.71	24.23	35.42	17.09
AT3G14440	NCED3 (9-cis-epoxycarotenoid dioxygenase 3)	0.99	0.38	0.01	71.54	73.85	52.09
AT5G35410	SOS2 (salt overly sensitive 2)	9.58	10.95	6.64	7.75	10.21	7.32
AT5G24270	SOS3 (salt overly sensitive 3)	1.32	4.03	1.38	0.82	0.40	0.01

**Table 1. Relative expression levels of stress-responsive genes identified by RNA-seq profiling.** \*TPM: number of transcripts per million tags.


in the *ATHB17* OX plants compared with *ATHB17* KO plants, with 16 of these 20 PhANGs up-regulated in the *ATHB17* KO plants while down-regulated in the *ATHB17* OX plants compared with WT plants (Table 2).

Under salt treatment, 21 of the 26 PhANGs were down-regulated in the *ATHB17* OX plants. In contrast, 16 PhANGs had higher expression in *ATHB17* KO plants while lower expression in *ATHB17* OX plants compared with WT plants (Table 2). These genes mainly encode proteins in PS I complexes (*PSAF*, *PSAH2*, *PSAH1*, *PSAD1*, *PSAG*), light-harvesting complexes (*LHCA1*, *LHCA2*, *LHCA3*, *LHCA4*, *LHCA5*, *LHCB3*, *LHCB5*, *LHCB6*, *LHCB7*, *LHB1B1*, *LHB1B2*), chlorophyll a/b-binding (*CAB1*, *CAB3*) proteins, and PS II oxygen evolving complex (*PSBO1*, *PSBO2*, *PPL1*, *PPL2*).

qRT-PCR analysis was carried out to validate the results of expression profiling analysis by RNA-seq. The data in Fig. 4A showed that 11 of the 13 tested PhANGs had higher transcript levels in *ATHB17* KO plants and lower expression levels in *ATHB17* OX plants compared with WT plants. This result is consistent with the data of RNA-seq profiling analysis. Together, these data indicate that *ATHB17* may act as a repressor of PhANGs, which agrees with its function as a repressor reported by Rice *et al.*<sup>35</sup>.

**ATHB17 can directly bind to the promoters of several PhANGs.** HD-ZIP transcription factors show a binding preference for variant HD-binding sequences<sup>30,37</sup>. According to our bioinformatic analysis, combined with the information from AGRIS (*Arabidopsis* Gene Regulatory Information Server), we selected six types of 8–9 bp sequences as potential HD binding sites. We firstly tested the binding affinities of *ATHB17* to these sequences by yeast one-hybrid (Y1H) assay. We found full length *ATHB17* protein had strong self-activation activity. It has been reported that *ATHB17* protein lacking the first 113 amino acids (*ATHB17*Δ113) can still homo-dimerize and specifically recognize the sequence requirements for DNA binding<sup>35</sup>. Thus we used *ATHB17* protein lacking the first 107 amino acids (*ATHB17*Δ107) for Y1H and electrophoretic mobility shift assay (EMSA). As results shown in Supplementary Fig. S6A, *ATHB17*Δ107 had no self-activation activity. Y1H assay revealed *ATHB17* had strong binding affinities to these HD binding *cis*-elements: aaattagt, ttaatt and taaatga (Supplementary Fig. S6B). Then we performed EMSA to confirm the binding affinities of *ATHB17*Δ107 to the potential *ATHB17* binding *cis*-elements *in vitro*. Supplementary Fig. S6C shows that *ATHB17*Δ107 could directly bind to the sequences aaattagt and ttaatt, but failed to bind to taaatga *in vitro*. The results of EMSA indicate that *ATHB17*Δ107 protein was able to directly bind to the ttaatt motif *in vitro*, and the binding was specific as demonstrated by competition assay using unlabelled (competitor) and non-specific probes (non-competitor) (Supplementary Fig. S6D).

Based on the above analysis, we chose aaattagt and ttaatt as the *ATHB17* binding *cis*-elements. Promoter sequence analyses of 19 *ATHB17* down-regulated PhANGs revealed that 10 of the 19 genes had at least one *ATHB17* binding *cis*-element (Fig. 5A). To analyze whether *ATHB17* could directly regulate transcription of all these 10 genes by binding to the *ATHB17* *cis*-elements in their promoters, we performed a Y1H assay. As shown in Fig. 5B, *ATHB17*Δ107 could only bind to the promoters of 5 PhANGs (*FDA6*, *LHCA2*, *LHB1B1*, *LHB1B2*, *PSBO1*) with different binding affinities. Subsequently, chromatin immunoprecipitation (ChIP) assays using the transgenic plants expressing 35S-haemagglutinin (HA)-*ATHB17* plants were conducted to validate the binding affinities *in vivo*. ChIP-qPCR showed that *ATHB17* could directly bind to *ATHB17* binding *cis*-element motifs in the promoters of the 5 genes screened out by Y1H. However, *ATHB17* failed to bind the h fragment in *PSBO1* promoter, which also had no binding affinity to *ATHB17*Δ107 in yeast (Fig. 5C). These results are consistent with the data of Y1H and the expression pattern of these genes in the *ATHB17* OX, KO and WT plants (Table 2), indicating that *ATHB17* can directly bind to the promoter of a number of PhANGs to regulate their expression.



**Figure 3. qRT-PCR validation of the data of stress-responsive marker genes in the RNA-seq profiling.**

About 0.1 g of 12-day old seedlings of the *ATHB17* OX, KO lines and WT plants were treated with liquid MS medium containing 0 or 200 mM NaCl for 5 h. Total RNA were extracted and reverse-transcribed as templates for qRT-PCR. UBQ5 was used as an internal control. Values are mean  $\pm$  SD of three independent experiments (\* $P < 0.05$ ).

#### **ATHB17 binds to the HD binding *cis*-elements in the *ATSIG5* promoter.**

Through RNA-seq profiling analysis, we found a nuclear encoded sigma factor, *ATSIG5*, had higher expression in *ATHB17* OX plants than in WT and *ATHB17* KO plants both under normal and salt stress conditions (Fig. 6A). qRT-PCR validation showed that *ATHB17* OX plants had increased *ATSIG5* transcription compared with WT and *ATHB17* KO plants under normal condition. After salt treatment, *ATSIG5* transcript was significantly up-regulated in *ATHB17* OX plants, while down-regulated in *ATHB17* KO plants compared with WT plants (Fig. 6B). Promoter sequence analysis revealed two potential ATHB17 binding *cis*-elements in the *ATSIG5* promoter: *cis1* ttaattt and *cis2* aaattagt located at 1148 bp and 1054 bp upstream of start codon, respectively (Fig. 6C). These data suggest that *ATSIG5* may be a candidate target of ATHB17.

ChIP and Y1H assay was conducted to determine whether ATHB17 could directly bind to the ATHB17 binding *cis*-elements in the promoter of *ATSIG5*. The results of ChIP-qPCR showed in Fig. 6D revealed that ATHB17 was able to bind to the *ATSIG5* promoter DNA fragments I and II which contained the ATHB17 binding *cis*-element. However, the DNA fragment III containing no ATHB17 binding *cis*-element was not enriched, suggesting that ATHB17 specifically bound to the ATHB17 binding *cis*-element motifs in the *ATSIG5* promoter. Binding of ATHB17 to the *ATSIG5* promoter was further tested by Y1H assay. Consistent with the ChIP assay results, ATHB17 could directly bind to both of the promoter fragments containing ATHB17 binding *cis*-element *cis1* and *cis2* in yeast cells, with much stronger binding affinities to the region containing *cis1*, but could not bind to the negative control promoter fragment without HD binding *cis*-element (Fig. 6E). These data indicate that *ATSIG5* was a direct target of ATHB17.

#### **ATHB17 regulates salt tolerance partly by modulating *ATSIG5*.**

*ATSIG5* was reported to respond to multiple stress signals, including salt stress. The *sig5-1* mutant is hypersensitive to NaCl treatment<sup>17</sup>. To study the functions of *ATSIG5*, we generated 35S::*ATSIG5* (OX) lines and obtained *ATSIG5* knockout mutants *sig5-1* and *sig5-4* (Supplementary Fig. S7). After germination and growth vertically on MS or MS medium containing NaCl for 10 days, the *ATSIG5* OX showed salt tolerant phenotypes with better root growth while *sig5-1* and *sig5-4* showed salt sensitive phenotypes compared with the WT (Fig. 7). These data agree with the results reported by Nagashima *et al.*<sup>17</sup>.

To investigate further whether *ATHB17* modulated salt stress through regulating *ATSIG5* expression, we introduced *ATHB17* OX into *sig5-1* background by crossing. The *ATHB17* OX *sig5-1* offspring were tested for salt tolerance. Although *ATHB17* gene was overexpressed in the hybrid offspring, the *ATHB17* OX *sig5-1* seedlings did not show salt stress resistant phenotypes as *ATHB17*OX plants did but an intermediate phenotype between the two parents (Fig. 8), implying that the ATHB17-conferred stress tolerance was partially dependent on *ATSIG5*.

*ATSIG5* was found to play an important role in regulating many PEGs which respond to environmental signals<sup>17,20,21</sup>. In order to study whether ATHB17 could modulate PEGs transcription through regulating *ATSIG5*, we investigated the expression of several downstream PEGs of *ATSIG5* which encoded components of PS I and II in *ATHB17* OX and KO plants by qRT-PCR. As shown in Fig. 9A, under normal condition, all the 6 PEGs tested were down-regulated in *ATHB17* KO plants, while 4 were up-regulated in *ATHB17* OX plants, compared with WT plants. However, after NaCl stress treatment, the expression of all these genes was reduced in *ATHB17* KO plants, and increased in *ATHB17* OX plants (Fig. 9B). These results indicated that ATHB17 might positively regulate PEG expression through *ATSIG5*.


Gene ID	Description	Normal (TPM*)			NaCl (TPM*)		
		KO	WT	OX	KO	WT	OX
AT1g29930	CAB1 (chlorophyll a/b binding protein 1)	2659.41	1921.89	172.11	5531.28	4153.04	1530.00
AT1g31330	PSAF (PS I subunit F)	1117.89	912.05	284.09	1296.45	946.94	291.59
AT2g34420	LHB1B2 (light-harvesting chlorophyll protein complex II subunit B2)	2781.39	2899.35	712.06	4826.90	4362.05	1421.93
AT3g55330	PPL1 (PsbP-like protein 1)	37.48	14.53	5.57	29.30	17.14	9.41
AT3g61470	LHCA2 (PS I light harvesting complex gene 2)	2269.95	2062.94	705.34	4063.92	3235.44	1422.93
AT4g10340	LHCB5 (light harvesting complex of PS II subunit 5)	836.73	592.29	86.05	1080.78	707.61	357.03
AT4g30950	FAD6 (fatty acid desaturase 6)	182.28	154.60	67.61	260.43	212.79	127.28
AT5g66570	PSBO1 (PS II oxygen-evolving complex 1)	2749.86	2251.56	485.78	4219.77	3937.28	1953.87
AT1g52230	PSAH2 (PS I subunit H2)	69.84	92.16	26.70	72.84	60.00	30.22
AT2g34430	LHB1B1 (light-harvesting chlorophyll protein complex II subunit B1)	1449.82	1655.64	306.57	4226.69	3826.35	2079.15
AT3g16140	PSAH1 (PS I subunit H1)	143.97	158.89	57.24	196.95	166.81	48.03
AT1g15820	LHCB6 (light harvesting complex of PS II subunit 6)	1560.05	1324.97	614.10	1194.31	1474.56	735.88
AT1g29910	CAB3 (chlorophyll a/b binding protein 3)	821.65	702.62	110.45	1632.16	1800.76	510.13
AT2g39470	PPL2 (PsbP-like protein 2)	46.05	21.47	15.17	63.48	79.12	26.82
AT3g47470	LHCA4 (light-harvesting chlorophyll protein complex I subunit A4)	1025.23	863.49	428.16	1665.12	2171.46	1119.13
AT3g50820	PSBO2 (PS II subunit O-2)	146.05	87.21	46.29	316.18	454.27	462.50
AT3g54110	PUMP1 (plant uncoupling mitochondrial protein 1)	87.13	71.68	40.53	43.95	49.28	64.84
AT4g02770	PSAD1 (PS I subunit D-1)	2057.38	1748.79	900.69	1283.02	1341.55	856.36
AT5g54270	LHCB3 (light-harvesting chlorophyll B-binding protein 3)	294.44	231.23	70.69	320.65	348.94	127.28
AT1g61520	LHCA3 (PS I light harvesting complex gene 3)	2178.95	1902.07	1325.00	3721.29	2438.82	1668.09
AT1g45474	LHCA5 (PS I light harvesting complex gene 5)	55.87	67.72	31.50	46.39	35.77	20.21
AT1g55670	PSAG (PS I subunit G)	37.48	37.33	23.63	38.66	24.56	13.81
AT1g67740	PSBY (PS II BY)	89.90	63.75	142.91	135.10	154.12	64.44
AT1g76570	LHCB7 (light-harvesting complex B7)	8.44	7.93	9.03	24.82	11.21	10.81
AT3g54890	LHCA1 (PS I light harvesting complex gene 1)	53.38	122.55	222.82	103.76	94.78	50.03
AT4g28660	PSB28 (PS II reaction centre PSB28 protein)	33.33	10.90	20.75	13.43	10.71	4.00

**Table 2.** Expression levels of PhANGs in *ATHB17* OX, KO and WT plants under normal or NaCl treated conditions identified by RNA-seq profiling. \*TPM: transcripts per million tags.


## Discussion

Previous studies on salt stress tolerance have mainly focused on the genes related to ion homeostasis, metabolites or osmo-protectants, antioxidant, hormones, ABA synthesis and signaling, and stress-responsive TFs<sup>38–40</sup>. Very few researches focus on the coordination of photosynthetic genes to improve plant salt tolerance. In this study, we did not find any significant difference in the expression of classic stress-induced marker genes between *ATHB17* OX and KO plants, including many stress-responsive genes, ABA synthesis and signaling pathway genes, as well as SOS genes (Table 1 and Fig. 3). These results partially agree with the results by Park *et al.*<sup>36</sup>, who reported that *ATHB17* overexpression did not affect the expression of a number of ABA-responsive genes. These findings implicate that *ATHB17*-conferred stress tolerance may arise through a pathway different from the traditional ones.

Environmental stresses present great challenges to the normal development and growth of plants. In addition to developmental changes, chloroplasts constantly experience changing environments, thus a tight coordination between the nucleus and chloroplast is crucial to the survival of plants. These genome-coordinating mechanisms are achieved through both anterograde (nucleus to organelle) and retrograde (organelle to nucleus) signals<sup>41,42</sup>. Most of chloroplast proteins are nuclear-encoded, and the concentrations of these proteins are efficiently regulated by nuclear transcription<sup>43</sup>. TFs play important roles in the nuclear-chloroplast communication. Nuclear-encoded sigma factors regulate PEP activity to modulate the expression of different sets of genes responding to the external environmental signals<sup>44</sup>. Besides sigma factors, very few TFs have been isolated that regulate transcripts of nuclear photosynthetic genes and chloroplast genes to date. GATA-type TFs GNC and CGA1 were reported to modulate the expression of chloroplast protein genes *genomes uncoupled 4* (*GUN4*) and *HEMA1*<sup>45</sup>. Abscisic acid insensitive 4 (*ABI4*) represses the expression of photosynthetic nuclear genes, potentially acting as a master switch required for the modulation of nuclear genes in response to environmental signals and developmental cues<sup>46</sup>. Higher yield rice (*HYR*) is a master regulator in rice, responding to environmental



**Figure 4.** qRT-PCR analysis of the expression level of PhANGs and PEGs. About 0.1 g 12-day-old plants grown under normal condition were used for RNA extraction. First-strand cDNA was synthesized from 1  $\mu$ g total RNA with Oligo dT primer for qRT-PCR analysis. Relative transcription levels of the genes in *ATHB17* OX and KO are normalized to levels in WT control (WT = 0). Values are mean  $\pm$  SD of three independent experiments.




**Figure 5.** *ATHB17* binds to the promoters of several PhANGs. (A) Location of *ATHB17* binding cis-elements in the promoter of several PhANGs. The *ATHB17* binding cis-elements are indicated with filled rectangles, above/below which the sites of the last base of the cis-elements relative to the start code are shown. (B) Y1H assay for *ATHB17* binding to the 25 bp fragment containing *ATHB17* binding cis-element from the promoter of five PhANGs, respectively. (C) ChIP assay. About 70–200 bp promoter fragments containing *ATHB17* binding cis-element were enriched by anti-HA antibodies in ChIP-qPCR analysis. Values are mean  $\pm$  SD of three independent experiments (\* $P$  < 0.05, \*\* $P$  < 0.01, \*\*\* $P$  < 0.001).

stress, directly activating several photosynthesis genes by binding to their promoters<sup>47</sup>. In addition, Golden2-like 1 (GLK1) and GLK2 TFs coordinate the expression of the photosynthetic apparatus genes in *Arabidopsis*<sup>48</sup>.

On the other hand, the chloroplast can act as an environmental sensor. Metabolite and protein signals, such as the recently discovered dihydroxyacetone phosphate (DHAP), heme, methylerythritolcyclodiphosphate (MEcPP), 3'-phosphoadenosine 5'-phosphate (PAP), GUN1, PTM, A-type heat-shock transcription factors (HSEA1D, HSEA2 and HSEA3), and Heat Shock 90 (HSP90) produced in chloroplasts act as novel types of retrograde signals to regulate PhANGs in the nucleus<sup>42,49–53</sup>.






**Figure 6.** *ATHB17* directly regulates the transcription of *ATSIG5*. (A) Expression levels of *ATSIG5* in the RNA-seq profiling data. (B) qRT-PCR validation of the expression levels of *ATSIG5* in RNA-seq profiling. About 0.1 g 12-day-old plants grown on MS medium were used transferred to liquid MS medium containing 0 or 200 mM NaCl for 5 h. The plants were harvested for qRT-PCR analysis. Values are mean  $\pm$  SD of three independent experiments and asterisks denote Student's *t*-test significance compared with KO (\* $P$  < 0.05, \*\* $P$  < 0.01, \*\*\* $P$  < 0.001). (C) The schematic illustration of the locations of *ATHB17* binding *cis*-element in the promoters of *ATSIG5* and the fragments (short lines) used in ChIP-qPCR assay. *cis1* is ttaatt located 1148 bp upstream of start codon, *cis2* is aaattagt located 1054 bp upstream of start codon, control is a random 8 bp sequence. (D) qPCR data from ChIP assay with antibody against HA. A fragment without *ATHB17* binding *cis*-element (fragment III) was used as negative control. Values are mean  $\pm$  SD of three replica experiments (\* $P$  < 0.05, \*\*\* $P$  < 0.001); (E) Y1H assay for *ATHB17* binding to the 25 bp fragment containing *ATHB17* binding *cis*-element from the promoter of *ATSIG5*. Fragment containing no *ATHB17* binding *cis*-element was used as a negative control.

In this study, we found that *ATHB17*, a HD-ZIP TF, played important roles in regulating the expression of PhANGs in response to salt stress. By genetic analysis with *ATHB17* knockout mutants and *ATHB17*-overexpressing lines, we have demonstrated that *ATHB17* is a positive regulator in response to salt stress (Fig. 2). RNA-seq profiling analysis revealed that *ATHB17* acted as a repressor of PhANGs. *ATHB17* was reported as a transcriptional repressor containing EAR (ERF-associated amphiphilic repression)-like motif<sup>35</sup>. Consistent with this result, we found that *ATHB17* repressed the transcription of 26 PhANGs under normal or salt stress conditions (Table 2 and Fig. 4), with 5 of them (*FDA6*, *LHCA2*, *LHB1B1*, *LHB1B2* and *PSBO1*) directly regulated by *ATHB17* (Fig. 5). *ATHB17* repressed the expression of genes related to light-harvesting complexes (*LHCA1*, *LHCA2*, *LHCA3*, *LHCA4*, *LHCA5*, *LHCB3*, *LHCB5*, *LHCB6*, *LHCB7*, *LHB1B1*, *LHB1B2*), chlorophyll a/b-binding proteins (*CAB1*, *CAB3*), PS II oxygen evolving complex (*PSBO1*, *PSBO2*, *PPL1*, *PPL2*), thus reducing light harvesting to alleviate photo-oxidative damage. The light-harvesting protein complex together with chlorophyll captures light energy and delivers it to the PSs. Under stress conditions, light harvesting must be reduced in order to avoid over-excitation and damage of PSs. Therefore, we can speculate that decreasing light capture by *ATHB17* is one of the ways to protect plants from photo-damage under stress conditions.


Furthermore, we also found that *ATHB17* could directly activate the transcript of *ATSIG5* (Fig. 6), a multiple-stress responsive TF which can be induced by various stresses, such as high light, salt, osmotic stress and




**Figure 7.** *ATSIG5*-overexpressing transgenic plants were more tolerant while its knockout mutants were more sensitive to salt stress. (A) Salt tolerance assay of *ATSIG5*-overexpressing and knockout plants. Seeds were sowed on MS medium containing 0, 100, 120, 150 mM NaCl and grown vertically for 10 days. Bar = 1 cm; (B,C) Root length of the 10-day-old plants grown on MS medium containing indicated concentrations of NaCl. Values are mean  $\pm$  SD (n = 45, \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001).

low temperature<sup>17</sup>. Light induction of *ATSIG5* expression was strongly dependent on cryptochrome1 (CRY1), and long hypocotyl 5 (HY5) which could directly bind to *ATSIG5* promoter to regulate *ATSIG5* expression in a light responsive manner<sup>28</sup>. However, stress and light responsiveness of *ATSIG5* seemed to be independent<sup>17</sup>. It remains unclear what factors are involved in modulating *ATSIG5* expression in response to other stresses. Here, we found HB17 is one of the upstream regulators of *ATSIG5* in response to salt stress. We found that, similar with *ATSIG5*, *ATHB17* is also a multiple stress-responsive TF. NaCl, mannitol, paraquat, and ABA could highly induce its expression in leaves (Fig. 1). Knockout of *ATHB17* impaired the induction of *ATSIG5* by salt stress (Fig. 6B). Through Y1H and ChIP assay, we found *ATHB17* could directly bind to the *ATSIG5* promoter to regulate its transcription (Fig. 6). Overexpression of *ATSIG5* in *Arabidopsis* enhanced its salt tolerance, while *sig5* knockout plants became salt sensitive (Fig. 7). However, when *ATHB17* was overexpressed in the *ATSIG5* knockout background, salt tolerance of the plants was partially impaired (Fig. 8). These results indicated that *ATHB17* is one of the upstream regulators of *ATSIG5* in response to salt, and *ATHB17*-conferred salt resistance partly depends on *ATSIG5*.

The activity of PS II can be inactivated by a variety of environmental stresses, which inhibit the repair of PS II rather than directly attacking it<sup>54,55</sup>. D1 and D2, which are encoded by chloroplast gene *psbA* and *psbD*, bind all the redox-active components related to electron transfer of PS II and create oxidative power to break water molecules. Therefore, D1 and D2 are the main targets of oxidative damage<sup>56,57</sup>. After degradation of the photo-damaged D1 protein, a new functional copy is inserted into the core complex following *de novo* synthesis, to repair of PS II in response to stresses<sup>58</sup>. *ATSIG5* was not only involved in the response of plants to blue light,



**Figure 8.** *ATSIG5* acts downstream of *ATHB17*. (A) The phenotypes of *ATHB17 OX*, *ATHB17 OX/sig5-1*, *sig5-1* and WT seedlings grown on MS medium containing indicated concentrations of NaCl. Seeds were sowed on MS medium containing 0, 100, 125, 150 and 175 mM NaCl and grown vertically for 10 days. Bar = 1 cm. (B) root length of the 10-day-old plants grown on MS medium containing different concentrations of NaCl. Values are mean  $\pm$  SD (n = 45, \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001).



**Figure 9.** The expression levels of several PEGs in *ATHB17 OX* and KO plants. Twelve-day-old plants grown under normal conditions were transferred to liquid MS medium containing 0 or 200 mM NaCl for 6 h. The plants were harvested for RNA extraction. First-strand cDNA was synthesized from 1  $\mu$ g total RNA with random primer for qRT-PCR analysis. Relative transcription levels of the genes in *ATHB17 OX* and KO were normalized to levels in WT control (WT = 0). Values are mean  $\pm$  SD of three independent experiments. (A) PEGs expression levels in *ATHB17 OX*, KO and WT plants treated with NaCl-free liquid MS. (B) PEG expression levels in *ATHB17 OX*, KO and WT plants treated with liquid MS containing 200 mM NaCl.

but also involved in protection of chloroplasts under various stress conditions through enhancing repair of the PS II reaction center<sup>17</sup>. *ATSIG5* binds to the *psbD* promoter, which drives the expression of *psbD-psbC-psbZ* operon encoding core-proteins of PS II, in response to blue light and various abiotic stress<sup>17,59</sup>. *ATSIG5* also activates expression of the *psbA* in chloroplasts<sup>20</sup>, and acts as a key mediator of circadian regulation of expression of several chloroplast genes<sup>21</sup>. Therefore, *ATSIG5* is likely to combine extrinsic and intrinsic signals, which are important in adjusting plastid and nuclear gene expression, to light and environmental stress to successfully adapt plants to changing environments<sup>20</sup>. Under stress conditions, *ATHB17* may coordinate the expression of PEGs through regulating *ATSIG5* (Fig. 9), thus enhancing PS II reaction center repair to survive better.

Moreover, *ATHB17* may be involved in balancing the stoichiometry of PS I to PS II under salt stress condition. It is well-known that PS II is the main target of oxidative damage under stress conditions, therefore, the subunit proteins of PS II are maintained with a rapid turnover rate to facilitate the repair cycle<sup>56,57</sup>. Recent studies found the PS II photo-inhibition-repair cycle can also protect PS I from irreversible damage<sup>60</sup>. Although PS II in chloroplasts undergoes a frequent repair cycle, the functional PS II is decreased, and becomes rate-limiting to photosynthesis under stress conditions. During acclimation, when either PS becomes rate-limiting to photosynthesis, expression of genes for its reaction center proteins are up-regulated, while genes for reaction center proteins of the other PS, which has surplus photochemical capacity, are simultaneously decreased<sup>61</sup>. Thus, in order to balance the stoichiometry of PS I to PS II under salt stress condition, expression of genes encoding subunits of PS II should be induced and genes encoding subunits of PS I should be simultaneously repressed. By RNA-seq profiling and qRT-PCR analyses, we found that *ATHB17* repressed the expression of several nuclear genes which encoded PS I reaction center subunits (*psaF*, *psaH1*, *psaH2*, *psaD1*, *psaG*) (Table 2, Fig. 4), and activated the expression *ATSIG5*, which may up-regulate several chloroplast encoded PS II subunit genes (Fig. 9), thus benefiting the stoichiometry of PS I to PS II.

More interestingly, *ATHB17* is a multiple stress responsive gene. Under normal conditions, we found that *ATHB17* was primarily expressed in the root, with weak expression in the shoot, consistent with the previous report<sup>36</sup>. However, Hymus *et al.*<sup>34</sup> only detected very weak *ATHB17* signal in specific cells in the embryonic, QC area of primary and lateral roots, through *in situ* hybridization analysis. These differences may result from low sensitivity of *in situ* hybridization. Under stress conditions, *ATHB17* was significantly induced, especially in the leaves (Fig. 1C and D). As discussed above, the induction of *ATHB17* expression in the leaves may alleviate the damage to chloroplast, thus enhance plant tolerance to environmental stresses. Consistent with this hypothesis, overexpression of *ATHB17* could also enhance plant tolerance to drought and oxidative stresses (Supplementary Fig. S5), indicating a mechanism similar to that of salt stress may exist in responding to these stresses. Taken together, our results imply that *ATHB17* may act as an important TF regulating both PhANGs and PEGs to decrease light harvesting, enhance PS repair, and balance PS stoichiometry under stress conditions, thus improving the tolerance of plants to multiple stresses.

In conclusion, our study demonstrated that *ATHB17* was an important TF in coordinating the expression of PhANGs and PEGs to cope with multiple stresses. Under stress conditions, the expression of *ATHB17* was induced, especially in the photosynthetic leaves. As a TF, *ATHB17* repressed the transcription of many PhANGs indirectly or directly by binding to their promoters. *ATHB17* also directly activated transcription of *ATSIG5*, whose protein is then translocated to chloroplast as an important regulator of many chloroplast genes, such as *psbA*, *psbB*, *psbC*, *psbD* and *psBT*. However, it may modulate several other chloroplast genes through other unknown pathways. Overall, it eventually reduces light harvest under stress, enhances PS repair cycle and balances the stoichiometry of PS I to PS II in the chloroplast, therefore alleviating the damage to chloroplast under stress conditions and improving plant stress tolerance.

## Material and Methods

**Plant materials and growth conditions.** *Arabidopsis thaliana* (Col-0) and tobacco (*Nicotiana tabacum*, NC89) were used for transformation. Seeds were sterilized in 10% (w/v) bleach for 10 min, and washed five to six times with sterile water. For *Arabidopsis*, the seeds were first treated at 4 °C for 3 days vernalization, then sowed on MS medium. Seven-day-old *Arabidopsis* seedlings were transferred to soil. For tobacco, the washed seeds were directly germinated on MS medium. The growth house was controlled at 22 ± 1 °C with a 16 h a photoperiod.

**Constructs and preparation of transgenic plants.** To generate 35S::*ATHB17* and 35S::*ATSIG5* over-expression binary vectors, the *ATHB17* or *ATSIG5* cDNA was isolated by RT-PCR with *ATHB17*-attb-LP and *ATHB17*-attb-RP or *ATSIG5*-attb-LP and *ATSIG5*-attb-RP and cloned into pCB2004<sup>62</sup> via the GATEWAY cloning system. To analyze the *ATHB17* expression pattern, a 3.0 kbp promoter fragment was amplified with the primers *pATHB17*-LP and *pATHB17*-RP and then shuttled into the vector pCB308R<sup>62</sup>. To get the native promoter-gene fusion construct, a DNA fragment containing *ATHB17* promoter and coding region amplified by genomic PCR with primers *ATHB17*-attb-LP2 and *ATHB17*-attb-RP2 was cloned into pMDC110 to fuse with GFP<sup>63</sup>. For ChIP assays, *ATHB17* full-length coding sequence amplified by RT-PCR using specific primers *ATHB17*-HA-attb-LP and *ATHB17*-attb-RP was inserted into pCB2004 to get *pCB2004::ATHB17-HA* by the GATEWAY cloning system. All the constructs were electroporated into *Agrobacterium tumefaciens* C58C1, which were used to transform wild type *Arabidopsis* plants as described<sup>64,65</sup>. All the primers used are listed in Supplementary Table 1.

**Identification of *ATHB17* and *ATSIG5* knockout mutants and overexpression plants.** The T-DNA insertion site of the SALK\_095524 mutant in *ATHB17* was confirmed by genomic PCR using three specific primers: SALK\_095524-LP, SALK\_095524-RP, LBb1.3. Similarly, the *sig5-1* (SALK\_049021) mutant was identified by genomic PCR with three primers: *sig5-1* LP, *sig5-1* RP and LBb1.3. Mutant of *sig5-4* (SALK\_101921) were identified by genomic PCR with three primers: *sig5-4* LP, *sig5-4* RP and LBb1.3. For expression analysis of the knockout mutants and overexpression transgenic plants, Quantitative RT-PCR was used with primers for the full-length coding sequence. All the primers used are listed in Supplementary Table 1.

**GUS activity staining assay.** GUS staining of *pATHB17::GUS* transgenic plants was performed as described<sup>66</sup>. *Arabidopsis* seedlings were soaked in staining buffer containing 1 mM 5-bromo-4-chloro-3-indoryl-β-D-glucuronide (X-Gluc, Rose Scientific Inc., Somerset, NJ, U.S.A.) overnight, then decolorized using an ethanol series. Individual representative seedlings were photographed.

**Hydroponic culture.** 20-day-old seedlings germinated on MS agar medium were used for hydroponic culture as previously described<sup>67</sup>. The root of each tobacco seedling was wrapped in a sponge strip and inserted into a hole made in thick polystyrene foam board. The foam boards were floated on MS hydroponic solution with or without NaCl. Plants were cultured at 22 °C under 16-h photoperiod. Nutrient solution was changed every 7 days.

**RNA-seq.** Two-week-old seedlings of *ATHB17* OX, KO, and the WT control were treated with or without 200 mM NaCl in liquid MS medium, 60 rpm shaking in the air for 6 h, then total RNA isolated with TRIzol reagent (Invitrogen Inc.). RNA from 3 independent replicates were mixed by equal volume. RNA-seq was performed and analyzed by BGI (Beijing Genome Institute, Shenzhen) corporation following the protocol provided by the manufacturer. The Illumina HiSeq™ 2000 platform was used for sequencing. For annotation, all clean tags were mapped to the reference sequence and allowed no more than one nucleotide mismatch. Clean tags mapped to reference sequences from multiple genes were filtered. For gene expression analysis, the number of clean tags for each gene was calculated and then normalized to the number of transcripts per million clean tags (TPM)<sup>68</sup>.

**qRT-PCR.** Total RNA was extracted using TRIzol reagent (Invitrogen Inc.), and first-strand cDNA synthesized from 1 µg total RNA in a 20 µl reaction mixture with Prime Script RT reagent kit (TAKARA BIOTECHNOLOGY CO., LTD). For nuclear encoded gene expression analysis, oligo dT primer was used for cDNA synthesis. For PEGs expression analysis, random primers was used for cDNA synthesis. The transcript levels of classic stress-related genes and other genes were examined with specific primers. The PCR was performed on ABI step-one instrument with the amplification conditions of 40 cycles of 95 °C for 1 min, 62 °C for 20 seconds and 72 °C for 30 seconds. *UBQ5* was used as the internal control. The relative expression levels were calculated by the  $2^{-\Delta\Delta CT}$  method. All the primers used are listed in Supplementary Table 1.

**Yeast one-hybrid assay (Y1H).** A cDNA fragment encoding *ATHB17*Δ107 was amplified by the primers: the forward with restriction endonuclease BamH I, 5'-CGGGATCCGACCAGCTA AGGCTAGACATGAA-3' and the reverse with restriction endonuclease Xba I, 5'-GCTCTAGATCAACGATCACGCTCTTGCG-3', and inserted into plasmid pAD-GAL4-2.1 (pAD) to get AD/*ATHB17*Δ107.

To get the report vectors containing HD-binding sequences, three copies of the HD-binding sequence, containing Sac I and Mlu I adaptors, were annealed and inserted into the Sac I and Mlu I sites of pHIS2. To get the report vectors containing promoter sequence of different genes, 25 bp promoter segment containing HD-binding sequence with Sac I and Mlu I adaptors was annealed and inserted into pHIS2, respectively. The constructs were confirmed by sequencing. The pAD and pHIS2 empty vector were used for negative control. AD/*ATHB17*Δ107 and the reporter pHIS2 containing different DNA sequence were co-transfected into yeast cells Y187 respectively. The yeast was first grown on SD-Trp-Leu medium for 3 days at 30 °C, and then transferred to SD-Trp-Leu-His medium with 10 mM or 20 mM 3-aminotriazole (3-AT, sigma) at different dilutions. The yeasts were incubated at 30 °C for 5 days and the extent of yeast growth was determined.

**ChIP-PCR assay.** Leaves of 10-day-old *T<sub>3</sub>* homozygote of *35S::ATHB17-HA* transgenic plants and anti-HA tag antibody (Cali-Bio, CB100005M) were used for pulling down the chromatin, as previously described<sup>69</sup>. After degrading the associated proteins with proteinase K, the chromatin DNA samples were treated using phenol/chloroform, then precipitated and finally eluted in 30 µl TE buffer. ChIP-qPCR was then used to verify the promoter segment of related genes using the primers listed in Supplementary Table 1.

**EMSA assay.** *ATHB17*Δ107 protein was expressed in *Escherichia coli* with pET28a (+) protein expression system (Novagen) and purified by Ni<sup>2+</sup> chromatography. Three copies of complementary single-stranded HD-binding sequence were synthesized and annealed to form double-stranded DNA fragment. The DNA fragments were marked with α-<sup>32</sup>P-dCTP and gel purified as probes. Purified *ATHB17*Δ107 protein (200 ng) was incubated with probes for 30 min on ice. For the competition test, non-labeled probe and non-specific probes were added into the binding reaction. Each reaction was loaded on a 4.5% (w/v) native polyacrylamide gel with 0.5 × TBE buffer. The gel then exposed to X-ray film.

**Statistical analysis.** Statistically significant differences ( $P < 0.05$  or  $P < 0.01$  or  $P < 0.001$ ) were computed based on the Student's *t*-tests.

## References

- Munns, R., James, R. A. & Lauchli, A. Approaches to increasing the salt tolerance of wheat and other cereals. *Journal of Experimental Botany* **57**, 1025–1043 (2006).
- Flexas, J., Bota, J., Loreto, F., Cornic, G. & Sharkey, T. D. Diffusive and metabolic limitations to photosynthesis under drought and salinity in C(3) plants. *Plant Biology* **6**, 269–279 (2004).
- Lawlor, D. W. & Cornic, G. Photosynthetic carbon assimilation and associated metabolism in relation to water deficits in higher plants. *Plant Cell Environ* **25**, 275–294 (2002).
- Vasilikiotis, C. & Melis, A. Photosystem-II Reaction-Center Damage and Repair Cycle - Chloroplast Acclimation Strategy to Irradiance Stress. *P Natl Acad Sci USA* **91**, 7222–7226 (1994).
- Saibo, N. J. M., Lourenco, T. & Oliveira, M. M. Transcription factors and regulation of photosynthetic and related metabolism under environmental stresses. *Ann Bot-London* **103**, 609–623 (2009).
- Nouri, M. Z., Moumeni, A. & Komatsu, S. Abiotic Stresses: Insight into Gene Regulation and Protein Expression in Photosynthetic Pathways of Plants. *Int J Mol Sci* **16**, 20392–20416 (2015).
- Sato, S., Nakamura, Y., Kaneko, T., Asamizu, E. & Tabata, S. Complete structure of the chloroplast genome of *Arabidopsis thaliana*. *DNA research* **6**, 283–290 (1999).
- Pfannschmidt, T., Schutze, K., Fey, V., Sherameti, I. & Oelmuller, R. Chloroplast redox control of nuclear gene expression—a new class of plastid signals in interorganellar communication. *Antioxidants & redox signaling* **5**, 95–101 (2003).

9. Kanamaru, K. & Tanaka, K. Roles of chloroplast RNA polymerase sigma factors in chloroplast development and stress response in higher plants. *Bioscience, biotechnology, and biochemistry* **68**, 2215–2223 (2004).
10. Legen, J. *et al.* Comparative analysis of plastid transcription profiles of entire plastid chromosomes from tobacco attributed to wild-type and PEP-deficient transcription machineries. *The Plant Journal* **31**, 171–188 (2002).
11. Krause, K., Maier, R. M., Kofer, W., Krupinska, K. & Herrmann, R. G. Disruption of plastid-encoded RNA polymerase genes in tobacco: expression of only a distinct set of genes is not based on selective transcription of the plastid chromosome. *Molecular & general genetics: MGG* **263**, 1022–1030 (2000).
12. Liu, B. & Troxler, R. F. Molecular characterization of a positively photoregulated nuclear gene for a chloroplast RNA polymerase sigma factor in *Cyanidium caldarium*. *P Natl Acad Sci USA* **93**, 3313–3318 (1996).
13. Tanaka, K. *et al.* Nuclear encoding of a chloroplast RNA polymerase sigma subunit in a red alga. *Science* **272**, 1932–1935 (1996).
14. Lysenko, E. A. Analysis of the evolution of the family of the sig genes encoding plant sigma factors. *Russ J Plant Physiol+* **53**, 605–614 (2006).
15. Yao, J., Roy-Chowdhury, S. & Allison, L. A. AtSig5 is an essential nucleus-encoded Arabidopsis sigma-like factor. *Plant physiology* **132**, 739–747 (2003).
16. Schweer, J., Turkeri, H., Kolpack, A. & Link, G. Role and regulation of plastid sigma factors and their functional interactors during chloroplast transcription - Recent lessons from *Arabidopsis thaliana*. *European Journal of Cell Biology* **89**, 940–946 (2010).
17. Nagashima, A. *et al.* The multiple-stress responsive plastid sigma factor, SIG5, directs activation of the psbD blue light-responsive promoter (BLRP) in *Arabidopsis thaliana*. *Plant Cell Physiol* **45**, 357–368 (2004).
18. Zghidi, W., Merendino, L., Cottet, A., Mache, R. & Lerbs-Mache, S. Nucleus-encoded plastid sigma factor SIG3 transcribes specifically the psbN gene in plastids. *Nucleic Acids Res* **35**, 455–464 (2007).
19. Favory, J. J. *et al.* Specific function of a plastid sigma factor for ndhF gene transcription. *Nucleic Acids Res* **33**, 5991–5999 (2005).
20. Tsunoyama, Y. *et al.* Blue light-induced transcription of plastid-encoded psbD gene is mediated by a nuclear-encoded transcription initiation factor, AtSig5. *P Natl Acad Sci USA* **101**, 3304–3309 (2004).
21. Noordally, Z. B. *et al.* Circadian Control of Chloroplast Transcription by a Nuclear-Encoded Timing Signal. *Science* **339**, 1316–1319 (2013).
22. Yamburenko, M. V., Zubo, Y. O. & Borner, T. Abscisic acid affects transcription of chloroplast genes via protein phosphatase 2C-dependent activation of nuclear genes: repression by guanosine-3'-5'-bisdiphosphate and activation by sigma factor 5. *Plant J* **82**, 1030–1041 (2015).
23. Tsunoyama, Y., Morikawa, K., Shiina, T. & Toyoshima, Y. Blue light specific and differential expression of a plastid sigma factor, Sig5 in *Arabidopsis thaliana*. *FEBS Lett* **516**, 225–228 (2002).
24. Lerbs-Mache, S. Function of plastid sigma factors in higher plants: regulation of gene expression or just preservation of constitutive transcription? *Plant molecular biology* **76**, 235–249 (2011).
25. Lysenko, E. A. Plant sigma factors and their role in plastid transcription. *Plant cell reports* **26**, 845–859 (2007).
26. Kanazawa, T., Ishizaki, K., Kohchi, T., Hanaoka, M. & Tanaka, K. Characterization of four nuclear-encoded plastid RNA polymerase sigma factor genes in the liverwort *Marchantia polymorpha*: blue-light- and multiple stress-responsive SIG5 was acquired early in the emergence of terrestrial plants. *Plant Cell Physiol* **54**, 1736–1748 (2013).
27. Thum, K. E., Kim, M., Christopher, D. A. & Mullet, J. E. Cryptochrome 1, cryptochrome 2, and phytochrome co-activate the chloroplast psbD blue light-responsive promoter. *The Plant Cell* **13**, 2747–2760 (2001).
28. Mellenthin, M., Ellersiek, U., Börger, A. & Baier, M. Expression of the Arabidopsis Sigma Factor SIG5 Is Photoreceptor and Photosynthesis Controlled. *Plants* **3**, 359–391 (2014).
29. Oh, S. & Montgomery, B. L. Phytochrome-dependent coordinate control of distinct aspects of nuclear and plastid gene expression during anterograde signaling and photomorphogenesis. *Front Plant Sci* **5**, 171 (2014).
30. Ariel, F. D., Manavella, P. A., Dezar, C. A. & Chan, R. L. The true story of the HD-Zip family. *Trends in Plant Science* **12**, 419–426 (2007).
31. Turchi, L., Baima, S., Morelli, G. & Ruberti, I. Interplay of HD-Zip II and III transcription factors in auxin-regulated plant development. *Journal of Experimental Botany* **66**, 5043–5053 (2015).
32. Ciarbelli, A. R. *et al.* The Arabidopsis homeodomain-leucine zipper II gene family: diversity and redundancy. *Plant Mol Biol* **68**, 465–478 (2008).
33. Kunihiro, A. *et al.* Phytochrome-interacting factor 4 and 5 (PIF4 and PIF5) activate the homeobox ATHB2 and auxin-inducible IAA29 genes in the coincidence mechanism underlying photoperiodic control of plant growth of *Arabidopsis thaliana*. *Plant & cell physiology* **52**, 1315–1329 (2011).
34. Hymus, G. J. *et al.* Application of HB17, an Arabidopsis class II homeodomain-leucine zipper transcription factor, to regulate chloroplast number and photosynthetic capacity. *J Exp Bot* **64**, 4479–4490 (2013).
35. Rice, E. A. *et al.* Expression of a Truncated ATHB17 Protein in Maize Increases Ear Weight at Silking. *Plos One* **9**(4): e94238. doi 10.1371/journal.pone.0094238.
36. Park, M. Y., Kim, S. A., Lee, S. J. & Kim, S. Y. ATHB17 is a positive regulator of abscisic acid response during early seedling growth. *Mol Cells* **35**, 125–133 (2013).
37. Tominaga-Wada, R. *et al.* The GLABRA2 homeodomain protein directly regulates CESA5 and XTH17 gene expression in Arabidopsis roots. *Plant J* **60**, 564–574 (2009).
38. Gupta, B. & Huang, B. R. Mechanism of Salinity Tolerance in Plants: Physiological, Biochemical, and Molecular Characterization. *Int J Genomics* (2014).
39. Zhu, J. K. Salt and drought stress signal transduction in plants. *Annual Review of Plant Biology* **53**, 247–273 (2002).
40. Liang, C. *et al.* GhABF2, a bZIP transcription factor, confers drought and salinity tolerance in cotton (*Gossypium hirsutum* L.). *Sci Rep* **6**, 35040 doi: 10.1038/srep35040 (2016).
41. Woodson, J. D. & Chory, J. Coordination of gene expression between organellar and nuclear genomes. *Nature Reviews Genetics* **9**, 383–395 (2008).
42. Chan, K. X., Phua, S. Y., Crisp, P., McQuinn, R. & Pogson, B. J. Learning the Languages of the Chloroplast: Retrograde Signaling and Beyond. *Annu Rev Plant Biol* **67**, 25–53 (2016).
43. Kleffmann, T. *et al.* The Arabidopsis thaliana chloroplast proteome reveals pathway abundance and novel protein functions. *Curr Biol* **14**, 354–362 (2004).
44. Kanamaru, K. & Tanaka, K. Roles of chloroplast RNA polymerase sigma factors in chloroplast development and stress response in higher plants. *Biosci Biotech Bioch* **68**, 2215–2223 (2004).
45. Hudson, D. *et al.* GNC and CGA1 Modulate Chlorophyll Biosynthesis and Glutamate Synthase (GLU1/Fd-GOGAT) Expression in Arabidopsis. *Plos One* **6** (11): e26765. doi 10.1371/journal.pone.0026765.
46. Leon, P., Gregorio, J. & Cordoba, E. ABI4 and its role in chloroplast retrograde communication. *Frontiers in Plant Science* **3** (2013).
47. Ambavaram, M. M. R. *et al.* Coordinated regulation of photosynthesis in rice increases yield and tolerance to environmental stress. *Nat Commun* **5**: 5302 doi: 10.1038/ncomms6302 (2014).
48. Waters, M. T. *et al.* GLK Transcription Factors Coordinate Expression of the Photosynthetic Apparatus in Arabidopsis. *Plant Cell* **21**, 1109–1128 (2009).
49. Estavillo, G. M. *et al.* Evidence for a SAL1-PAP chloroplast retrograde pathway that functions in drought and high light signaling in Arabidopsis. *Plant Cell* **23**, 3992–4012 (2011).

50. Woodson, J. D., Perez-Ruiz, J. M. & Chory, J. Heme synthesis by plastid ferrochelatase I regulates nuclear gene expression in plants. *Curr Biol* **21**, 897–903 (2011).
51. Xiao, Y. M. *et al.* Retrograde Signaling by the Plastidial Metabolite MEcPP Regulates Expression of Nuclear Stress-Response Genes. *Cell* **149**, 1525–1535 (2012).
52. Chi, W., Feng, P., Ma, J. & Zhang, L. Metabolites and chloroplast retrograde signaling. *Curr Opin Plant Biol* **25**, 32–38 (2015).
53. Jung, H. S. *et al.* Subset of heat-shock transcription factors required for the early response of Arabidopsis to excess light. *Proc Natl Acad Sci USA* **110**, 14474–14479 (2013).
54. Murata, N., Takahashi, S., Nishiyama, Y. & Allakhverdiev, S. I. Photoinhibition of photosystem II under environmental stress. *Bba-Bioenergetics* **1767**, 414–421 (2007).
55. Allakhverdiev, S. I. *et al.* Salt stress inhibits the repair of photodamaged photosystem II by suppressing the transcription and translation of psbA genes in Synechocystis. *Plant Physiology* **130**, 1443–1453 (2002).
56. Kettunen, R., Tyystjarvi, E. & Aro, E. M. Degradation pattern of photosystem II reaction center protein D1 in intact leaves. The major photoinhibition-induced cleavage site in D1 polypeptide is located amino terminally of the DE loop. *Plant Physiol* **111**, 1183–1190 (1996).
57. Schuster, G., Timberg, R. & Ohad, I. Turnover of thylakoid photosystem II proteins during photoinhibition of *Chlamydomonas reinhardtii*. *Eur J Biochem* **177**, 403–410 (1988).
58. Theis, J. & Schroda, M. Revisiting the photosystem II repair cycle. *Plant Signal Behav* **11**, e1218587 (2016).
59. Hoffer, P. H. & Christopher, D. A. Structure and blue-light-responsive transcription of a chloroplast psbD promoter from *Arabidopsis thaliana*. *Plant Physiol* **115**, 213–222 (1997).
60. Tikkanen, M., Mekala, N. R. & Aro, E. M. Photosystem II photoinhibition-repair cycle protects Photosystem I from irreversible damage. *Biochim Biophys Acta* **1837**, 210–215 (2014).
61. Rochaix, J. D. Posttranscriptional control of chloroplast gene expression. From RNA to photosynthetic complex. *Plant Physiol* **125**, 142–144 (2001).
62. Lei, Z. Y. *et al.* High-throughput binary vectors for plant gene function analysis. *Journal of Integrative Plant Biology* **49**, 556–567 (2007).
63. Curtis, M. D. & Grossniklaus, U. A gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiology* **133**, 462–469 (2003).
64. Horsch, R. B. *et al.* Analysis of *Agrobacterium tumefaciens* virulence mutants in leaf discs. *Proc Natl Acad Sci USA* **83**, 2571–2575 (1986).
65. Clough, S. J. & Bent, A. F. Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**, 735–743 (1998).
66. Jefferson, R. A., Kavanagh, T. A. & Bevan, M. W. Gus Fusions - Beta-Glucuronidase as a Sensitive and Versatile Gene Fusion Marker in Higher-Plants. *Embo J* **6**, 3901–3907 (1987).
67. Yu, L. H. *et al.* Overexpression of Arabidopsis NLP7 improves plant growth under both nitrogen-limiting and -sufficient conditions by enhancing nitrogen and carbon assimilation. *Sci. Rep.* **6**, 27795; doi: 10.1038/srep27795 (2016).
68. t Hoen, P. A. C. *et al.* Deep sequencing-based expression analysis shows major advances in robustness, resolution and inter-lab portability over five microarray platforms. *Nucleic Acids Res* **36** (2008).
69. Lee, J. *et al.* Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell* **19**, 731–749 (2007).

## Acknowledgements

This work was supported by Ministry of Science and Technology of China (Grant # 2012CB114304 to C.-B.X), NNSFC (Grant # 30700051 to C.-B.X) and the Chinese Academy of Sciences (Grant # KSCX3-EW-N-007 to C.-B. X). We thank ABRC for offering T-DNA insertion lines.

## Author Contributions

C.X. and L.Y. designed the experiments. P.Z., R.C., and L.Y. performed most of the experiments and data analyses. P.X., J.W., J.-L.M., Y.C., and C.-Z.Z. assisted in experiments and data analyses. P.Z. and L.Y. wrote the manuscript. C.X. supervised the project and revised the manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing Interests:** The authors declare no competing financial interests.

**How to cite this article:** Zhao, P. *et al.* ATHB17 enhances stress tolerance by coordinating photosynthesis associated nuclear gene and *ATSIG5* expression in response to abiotic stress. *Sci. Rep.* **7**, 45492; doi: 10.1038/srep45492 (2017).

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017