

SCIENTIFIC REPORTS



OPEN

IgE binds asymmetrically to its B cell receptor CD23

Balvinder Dhaliwal¹, Marie O. Y. Pang^{2,3}, Anthony H. Keeble^{2,3,†}, Louisa K. James^{2,3,‡}, Hannah J. Gould^{2,3}, James M. McDonnell^{2,3}, Brian J. Sutton^{2,3} & Andrew J. Beavil^{2,3}

Received: 10 November 2016

Accepted: 21 February 2017

Published: 31 March 2017

The antibody IgE plays a central role in allergic disease mechanisms. Its effector functions are controlled through interactions between the Fc region and two principal cell surface receptors FcεRI and CD23. The interaction with FcεRI is primarily responsible for allergic sensitization and the inflammatory response, while IgE binding to CD23 is involved in the regulation of IgE synthesis and allergen transcytosis. Here we present the crystal structure of a CD23/IgE-Fc complex and conduct isothermal titration calorimetric binding studies. Two lectin-like “head” domains of CD23 bind to IgE-Fc with affinities that differ by more than an order of magnitude, but the crystal structure reveals only one head bound to one of the two identical heavy-chains in the asymmetrically bent IgE-Fc. These results highlight the subtle interplay between receptor binding sites in IgE-Fc and their affinities, the understanding of which may be exploited for therapeutic intervention in allergic disease.

An estimated 40% of the population suffer from at least one form of allergy, with their prevalence continuing to soar worldwide¹. Allergic diseases, which include allergic rhinitis (‘hay fever’), respiratory allergy (e.g. atopic asthma) and food allergy, have a detrimental impact on a person’s quality of life^{2–5}, a nation’s economy^{6,7}, and in acute cases can be life threatening⁸.

The antibody immunoglobulin E (IgE) plays a central role in allergic disease mechanisms. For reasons poorly understood, IgE recognizes seemingly innocuous substances known as allergens *via* its Fab regions, whilst its effector functions are controlled through interactions of the Fc region with two principal cell surface receptors, FcεRI and CD23 (also known as FcεRII)^{9,10}. FcεRI, found on mast cells and basophils, binds IgE with high affinity (K_D of 0.01 to 0.1 nM)¹¹ and is responsible for allergic sensitization and the immediate hypersensitivity response in which minute amounts of allergen crosslink FcεRI-bound IgE triggering cellular degranulation, release of proinflammatory mediators such as histamine and the initiation of an allergic response¹².

The second receptor for IgE, CD23, is unique among Ig receptors in that it belongs to the C-type lectin-like superfamily¹³. Expressed in several hematopoietic cell types, including B cells and antigen-presenting cells, in its membrane-bound form CD23 consists of three C-type lectin ‘head’ domains connected to the membrane by a trimeric α -helical coiled-coil ‘stalk’¹⁴. A single head domain binds to human IgE-Fc with an affinity K_D of $\sim 1 \mu\text{M}$ ¹⁵, although avidity of the trimer^{16–19} and the presence of Ca^{2+} ^{15,20} can substantially enhance the interaction. The stalk region of CD23 is susceptible to attack by endogenous proteases such as ADAM10²¹, releasing soluble trimeric and monomeric forms of CD23. The house dust mite allergenic protease *Der p 1* generates a soluble monomeric form of CD23 consisting of just the lectin head domain, termed derCD23²².

The interaction of CD23 with IgE is involved in regulating many allergic processes²³. For example, when expressed on epithelial cells, CD23 is involved in the transportation of IgE-allergen immune complexes across the gut²⁴ and airways²⁵. On B cells that have switched to IgE production, the CD23-IgE interaction is critical to both the up- and down-regulation of IgE synthesis depending on the oligomeric form of CD23⁹. This interaction is also essential in the process of facilitated antigen presentation (FAP)⁹, in which IgE-allergen complexes are internalized by CD23 on antigen-presenting cells. The allergen is then proteolytically cleaved and its peptides

¹Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, Faculty of Biology, Medicine and Health, Manchester Academic Health Sciences Centre and Manchester Institute of Biotechnology, University of Manchester, 131 Princess Street, Manchester, M1 7DN, U.K. ²Randall Division of Cell and Molecular Biophysics, Faculty of Life Sciences & Medicine, King’s College London, New Hunt’s House, Guy’s Campus, London, SE1 1UL, U.K. ³Medical Research Council and Asthma UK Centre in Allergic Mechanisms of Asthma, London, U.K. [†]Present address: Department of Biochemistry, University of Oxford, South Parks Road, Oxford, OX1 3QU, U.K. [‡]Present address: Blizard Institute, Barts and The London Medical School, Queen Mary University of London, London E1 2AT, U.K. Correspondence and requests for materials should be addressed to B.D. (email: balvinder.dhaliwal@manchester.ac.uk) or B.J.S. (email: brian.sutton@kcl.ac.uk) or A.J.B. (email: andrew.beavil@kcl.ac.uk)




Figure 1. Structure of the 1:1 derCD23/IgE-Fc complex. The derCD23 (green C α traces with surfaces) binds to one heavy chain of IgE-Fc (red and blue) contacting the C ϵ 3 and C ϵ 4 domains. The C ϵ 2 domains are asymmetrically bent back onto one C ϵ 3 domain and make some additional contacts with derCD23. The carbohydrate is shown in all-atom representation (red and yellow, without surfaces) and can be seen behind the C ϵ 3 domain.

complexed with MHC class II molecules and recycled to the cell surface for Type 2 helper (T_H2) cell recognition and activation. The activated T_H2 cells secrete cytokines that ultimately lead to an increase in IgE synthesis and exacerbation of allergic inflammation^{26,27}. Recent evidence indicates that the CD23 surface density on B cells determines its Fc ϵ RI activity²⁸. The CD23-IgE interaction is thus involved in both the development and regulation of the allergic response, and indeed, molecules designed to target the interaction such as the anti-CD23 IgG antibody Lumiliximab and DARPins have been shown to inhibit IgE synthesis^{29,30}. A molecular understanding of the CD23-IgE interaction thus has implications for the control of allergic disease.

We previously reported the crystal structure of derCD23 in complex with Fc ϵ 3-4³¹, a subfragment of IgE-Fc consisting of a dimer of the C ϵ 3 and C ϵ 4 domains. One head domain was found to bind to each heavy chain of IgE, principally to the C ϵ 3 domains but with a contribution from C ϵ 4, explaining the known 2:1 stoichiometry of monomeric CD23 binding to both IgE-Fc and Fc ϵ 3-4^{19,32}. Although the binding sites for Fc ϵ RI and CD23 lie at opposite ends of the C ϵ 3 domain, we went on to demonstrate that the binding of the two receptors to IgE are allosterically linked and mutually exclusive, with conformational changes in Fc ϵ 3-4 upon CD23 binding precluding the binding of Fc ϵ RI and *vice versa*³¹.

Here we present the crystal structure of derCD23 in complex with the complete IgE-Fc region, including the C ϵ 2 domains, which in the free molecule fold back onto the Fc ϵ 3-4 region in an asymmetrical fashion³³. One derCD23 molecule is seen binding to one IgE-Fc chain, making contact with all three domains. Isothermal titration calorimetry (ITC) experiments confirm the 2:1 (derCD23:IgE-Fc) stoichiometry in solution and reveal that the two binding affinities differ by more than an order of magnitude; this is explained in terms of the crystal structure.

Results

Quaternary structure of the derCD23/IgE-Fc complex. The crystal structure contains two essentially identical complexes of derCD23/IgE-Fc within the asymmetric unit. Each complex consists of one IgE-Fc molecule bound to one molecule of derCD23 (Fig. 1). The derCD23 molecule interacts with the IgE-Fc chain onto which the C ϵ 2 domains are folded back. The 1:1 stoichiometry of the complex (*i.e.*, one IgE-Fc dimer bound to one molecule of derCD23) was unexpected, as it is well documented that one CD23 head domain binds to each heavy chain of IgE, whether the Fc ϵ 3-4 sub-fragment or the complete IgE-Fc^{15,19,31,32}.

The derCD23/IgE-Fc interface. The derCD23 molecule binds principally to the C ϵ 3 domain with lesser interactions with C ϵ 4 and the short C ϵ 3-C ϵ 4 linker region. The interface is predominantly hydrophilic, consisting of five salt bridges with hydrogen bonds, and six additional hydrogen bonds (Fig. 2). The majority of these interactions are similar to those observed at the interface of the derCD23/Fc ϵ 3-4 crystal structure³¹, as well as being consistent with NMR chemical shift perturbation studies and binding experiments using proteins with mutated interface residues³⁴. There is an additional interaction between Asp258 of CD23 and Arg440 in the C ϵ 3-C ϵ 4 linker, an interaction that had only previously been observed in the Ca²⁺-bound derCD23/Fc ϵ 3-4 crystal structure (Figure S1). In the absence of Ca²⁺ binding to CD23, as in the complex presented here, the loop including Asp258 is found to be partially flexible^{20,31,35,36}. Despite the close approach of derCD23 to the C ϵ 2 domain in the same heavy chain, just a single hydrogen bond is observed between His216 of derCD23 and Thr260 of C ϵ 2 (Figs 1 and 2).

The derCD23/IgE-Fc interface shows an even poorer surface complementarity³⁷ than that found in the derCD23/Fc ϵ 3-4 complex; the calculated S_c values³⁸ range between 0.49–0.51 for the former, compared to 0.63–0.71 for the latter (the range of values refers to the different molecular complexes in the asymmetric unit). The




Figure 2. Salt bridges and hydrogen bonds at the derCD23/IgE-Fc interface. The five H-bonds associated with salt bridges in both independent complexes are shown in red, and the six remaining H-bonds are shown in black.




Figure 3. Orthogonal views depicting conformational changes in IgE-Fc upon binding a single molecule of derCD23 (not shown). A substantial rigid body movement of the IgE-Fc Cε2 domains upon derCD23 binding is observed when compared to free IgE-Fc (yellow and green). In addition, the Cε3 domain in the derCD23-bound chain (red) adopts an extreme closed conformation, in contrast to the Cε3 domain in the free chain (blue), which adopts an extreme open conformation. The structures were superposed on their (Cε4)₂ domains.

buried surface area of the derCD23/IgE-Fc interface ranges from 965 to 1042 Å², with relative contributions from the Cε2 domain of 24%, the Cε3 domain 56%, the Cε3-Cε4 linker (Ser437 to Arg440) 10%, and the Cε4 domain 10%. The Cε2-Cε3 linker (Lys326 to Ser337) does not contribute to the interface. Relatively few well-defined residues are involved in the network of van der Waals contacts at the interface (Gly180, Trp184, Val185, Phe272 and Asp274 in derCD23; Phe321, Ser378, Lys380, Ile411 and Lys435 in IgE-Fc). Of these hydrophobic interactions, only one involves the Cε2 domain: Phe321 interacts with Gly180 of derCD23. No interactions are observed between derCD23 and the other IgE-Fc heavy chain.

Comparison with previous IgE-Fc structures. IgE-Fc strongly favors an asymmetrically bent structure in its free state³⁹, with the Cε2 domain pair folded back and making contact not only with one of the Cε3 domains but also Cε4³³. Previous modeling of the derCD23/IgE-Fc complex by docking the (Cε2)₂ domain pair from the free IgE-Fc structure onto the derCD23/Fcε3-4 complex had suggested that only minor rearrangements of a few N-terminal residues in the Cε2 domain were required to prevent a steric clash with derCD23³¹. However, the experimentally determined complex shows that upon binding of a single derCD23 molecule, the disulfide-linked (Cε2)₂ domain pair move as a rigid unit, swinging out 16° to accommodate the derCD23 molecule (Fig. 3 and Supplementary movie).

Experiment	K_{D1} (μM)	ΔH_1 (kcal/mol)	$-T\Delta S_1$ (kcal/mol)	K_{D2} (μM)	ΔH_2 (kcal/mol)	$-T\Delta S_2$ (kcal/mol)
derCD23 titrated into IgE-Fc	1.2 (± 0.3)	-7.2 (± 0.4)	-0.9	14.4 (± 3.0)	-3.7 (± 0.6)	-3.0

Table 1. Binding affinities of derCD23 to IgE-Fc studied by ITC.

The Fc ϵ 3-4 moiety of the IgE-Fc dimer in the complex also adopts a surprising conformation upon derCD23 binding. Flexibility in the C ϵ 3-C ϵ 4 interdomain angle has been well documented in both unliganded Fc ϵ 3-4^{40,41} and IgE-Fc³³, and in complex with sFc ϵ RI α ^{11,42} or derCD23^{20,31,36}. A comparison of all previous Fc ϵ 3-4 and IgE-Fc structures shows that the C ϵ 3-C ϵ 4 angle varies over a range of 25°, with IgE-Fc or Fc ϵ 3-4 in complex with sFc ϵ RI α adopting the most ‘open’ conformation for the C ϵ 3 domains, and Fc ϵ 3-4 in complex with derCD23 adopting the most ‘closed’ conformation. Examination of the C ϵ 3-C ϵ 4 interdomain angles of IgE-Fc in complex with derCD23 presented here reveals that the chain bound to derCD23 adopts a closed conformation. In contrast, the free IgE-Fc chain unexpectedly adopts a very open conformation; indeed, the C ϵ 3-C ϵ 4 interdomain angles of the two chains differ by 16° (Figure S2). Comparison of the two IgE-Fc chains in the complex with all other IgE-Fc and Fc ϵ 3-4 structures shows that the derCD23-bound IgE-Fc chain has the same closed C ϵ 3-C ϵ 4 interdomain angle previously observed in the Ca²⁺-free derCD23-Fc ϵ 3-4 complex crystallized in space group P1³⁶ (Figure S2). This closed conformation is clearly incompatible with Fc ϵ RI binding^{31,42}. The free IgE-Fc chain, on the other hand, adopts the most open conformation observed, the same conformation previously only seen in complex with sFc ϵ RI α ^{11,42}. Earlier structural studies had indicated that free IgE-Fc chains favoured an intermediate position between the open and closed conformations³³, however our results reveal that the free chain of IgE-Fc adopts an extremely open conformation when the other chain is bound to derCD23: clearly a conformational change involving a closing of the C ϵ 3-C ϵ 4 interdomain angle is required to bind a second derCD23 molecule (Figure S3).

The two CD23 binding sites on IgE have different affinities. The binding affinity of derCD23 to IgE-Fc was determined using ITC, by titrating derCD23 into a fixed concentration of IgE-Fc. The data fitted best to a two-site binding model (Figure S4). One binding site has an affinity of 1.2 μM (K_{D1} ; Table 1), while the other is over an order of magnitude (12-fold) weaker with an affinity of 14.4 μM (K_{D2}).

Discussion

The interaction of CD23 and IgE is critical in many allergic disease processes, including the regulation of IgE synthesis, facilitated allergen presentation and allergen transportation across the gut and airways. Here we present the crystal structure of a lectin-like head domain of CD23 in complex with the complete Fc receptor-binding region of IgE. The results of ITC experiments reported here confirm the 2:1 stoichiometry reported earlier^{19,32}, but now reveal that the affinities for the two sites, one in each heavy chain, are very different ($K_D = 1.2 \mu\text{M}$ and 14.4 μM), with distinct thermodynamic profiles (Table 1). This is consistent with results from an earlier sedimentation equilibrium study of derCD23 binding to Fc ϵ 3-4, which concluded that the two sites were thermodynamically distinct, with a greater enthalpy change for one site than the other, although the affinities were similar ($K_D \sim 10 \mu\text{M}$)³⁶. The work reported here investigates the effect upon CD23 binding of the C ϵ 2 domains in the complete IgE-Fc, which is known to have an acutely and asymmetrically bent structure in the free state³³.

The crystal structure of derCD23 bound to IgE-Fc unexpectedly revealed only a 1:1 complex, with the single derCD23 molecule bound principally to the heavy chain onto which the C ϵ 2 domains are folded back. The derCD23 molecule primarily contacts C ϵ 3 and, as in the complexes with Fc ϵ 3-4^{20,31,36}, also contacts C ϵ 4 and the C ϵ 3-C ϵ 4 linker region; in addition however, there are a few contacts with C ϵ 2 including a single hydrogen bond. The binding of this derCD23 molecule is also accompanied by a swinging out of the C ϵ 2 domain pair (an ‘unbending’ of the IgE-Fc) by 16°. Another aspect of the asymmetry of the IgE-Fc is the difference in the C ϵ 3-C ϵ 4 interdomain angle in the 1:1 complex: in the chain to which derCD23 is bound the angle is closed (as seen in the complexes with Fc ϵ 3-4), whereas in the other, free chain, the angle is very open and incompatible with derCD23 binding. This is in contrast to the earlier Fc ϵ 3-4 complexes in which two derCD23 molecules were bound symmetrically to the two chains, both of which adopted a closed interdomain conformation. Clearly further conformational changes must occur upon binding of a second derCD23 molecule to IgE-Fc.

Despite the undoubted structural asymmetry between the two derCD23 binding sites in IgE-Fc, we cannot definitively assign the stronger and weaker affinity values to these sites. However, it is reasonable to assume that the derCD23 molecule seen in the crystal structure corresponds to the higher affinity site; it also has contributions from C ϵ 2 which could not occur at the site in the other heavy chain. In any event, the binding of the first derCD23 molecule at least preserves, and perhaps even exacerbates, the asymmetry in IgE-Fc and in particular the C ϵ 3-C ϵ 4 interdomain angle; if the latter, then binding to the first site may influence the affinity of the second site (Supplementary Figure S2 and Supplementary Movie). If there is ‘cross-talk’ between the two sites, this is yet another example of communication between receptor binding sites in IgE-Fc, such as that for CD23 and Fc ϵ RI. We have previously reported that the binding sites of CD23 and Fc ϵ RI on IgE, although at opposite ends of the C ϵ 3 domain, are allosterically linked, whereby conformational changes in IgE upon CD23 binding preclude the binding of Fc ϵ RI and *vice versa*³¹. Moreover, Ca²⁺-binding to CD23 causes local allosteric conformational changes that increase its affinity for IgE²⁰.

Thus the CD23/IgE/Fc ϵ RI network of interactions involves at least two and perhaps three allosteric mechanisms within IgE-Fc. Together these permit an extraordinarily intricate level of control of protein-protein interactions regulating many aspects of the allergic response. Mutual exclusion of Fc ϵ RI and CD23 binding to IgE is essential to prevent mast cell and basophil activation by trimeric sCD23 in the absence of allergen³¹.

Conformational changes in CD23 on antigen-presenting cells, upon release of bound Ca^{2+} , has recently been proposed to be involved in the mechanism that releases IgE/allergen complexes and recycles receptor back to the plasma membrane⁴³. On B cells, cross-linking of membrane CD23 by soluble IgE, and/or the formation of extended arrays of mIgE cross-linked by soluble trimeric CD23 to create signaling platforms for the survival and differentiation of IgE-committed cells, have been proposed to control IgE homeostasis⁹. All of these interactions critically depend upon the concentrations and oligomeric states of CD23 and IgE and their mutual affinities. While this study extends our understanding of CD23 binding to IgE-Fc by including the C ϵ 2 domains and their associated conformational changes, it is important to consider the consequences of the presence of the IgE Fab arms, and also possible Fc glycosylation, for the interaction between the complete molecules. We have previously modelled the range of conformations available to the Fab arms in both free and Fc ϵ RI-bound IgE, and found that they are more restricted than in IgG, occupying volumes that are mutually exclusive and distinct from the space occupied by the Fc region^{44,45}. Thus the derCD23/IgE-Fc interaction that we observe in the crystal structure may indeed be extended to the whole IgE molecule. Regarding glycosylation, two sites on the surface of IgE-Fc (asparagine residues 265 and 371) have been shown to be glycosylated⁴⁶, and were mutated to glutamine in the construct used in this study to facilitate crystallization⁴⁷. However, these attachment sites are distant from the CD23 binding site, and therefore unlikely to affect the interaction.

Extrapolation of the results presented here to the binding of either one or two complete CD23 trimers to IgE is also important. We have previously reported that the two head domains that engage Fc ϵ 3-4 cannot come from the same trimer³¹, and this is equally true for IgE-Fc or the whole IgE molecule. However, taking all of the above structural considerations into account, it is also clear that two CD23 trimers can bind to one IgE molecule, or one CD23 trimer can cross-link two IgE molecules, either in the context of soluble trimeric CD23 binding to membrane IgE, or soluble IgE binding to membrane CD23, in relation to IgE regulation or allergen/IgE immune complex transcytosis. A detailed molecular understanding of the IgE/CD23 interaction may thus permit novel approaches to intervene therapeutically in allergic disease.

Methods

Protein purification. Recombinant human derCD23 (Ser156-Glu298) and IgE-Fc (Val224-Lys547, incorporating glycosylation double mutations N265Q and N371Q) proteins were expressed, refolded and purified according to protocols previously described^{15,31,32,47}.

Crystallization and data collection. The derCD23-IgE-Fc complex was crystallized by vapor diffusion. IgE-Fc was concentrated to 37 mg/mL, and derCD23 to 18 mg/mL, both in 25 mM Tris-HCl (pH 7.5), 20 mM NaCl, 0.05% (w/v) sodium azide, followed by mixing of 0.56 mM derCD23 (8.6 mg/mL) with 0.28 mM IgE-Fc (20 mg/mL), and diluting the complex with an equal volume (100 nL) of 15% (w/v) PEG 8,000, 0.1 M sodium citrate, 0.05 M ammonium sulfate. A single needle-like crystal suitable for data collection grew after one and a half months at 295 K. The crystal was flash-cooled (using 18% (w/v) PEG 8,000, 0.1 M sodium citrate, 0.05 M ammonium sulfate, 20% (v/v) glycerol as cryoprotectant) and stored in liquid nitrogen. Diffraction data were collected at 100 K at beamline I03, Diamond Light Source.

Structure solution and refinement. Indexing and integration of data were carried out with *MOSFLM*^{48,49}, and merging of data performed using *AIMLESS*⁵⁰. The derCD23-IgE-Fc complex was solved by molecular replacement using *PHASER*⁵¹; two copies of the Fc ϵ 3-4 dimer were identified in the asymmetric unit, using chains A and B from the derCD23-Fc ϵ 3-4 complex (PDB 4EZM)³¹. Subsequently, two sets of C ϵ 2 dimers (chains A and B, residues 235–325, PDB 2WQR)¹¹ were located. Finally, two Ca^{2+} -free derCD23 molecules (using PDB 4J6J)³⁵ were identified. Iterative cycles of refinement using *PHENIX*⁵², *REFMAC5*⁵³ and *BUSTER-TNT*⁵⁴ alternated with manual model building with *COOT*⁵⁵. The model was built into $2F_o - F_c$ composite omit, $2F_o - F_c$ and $F_o - F_c$ electron density maps to minimize bias. Disruption of some disulfide bonds due to radiation damage was observed in the derCD23/IgE-Fc complex structure. Carbohydrate atoms were subsequently incorporated into the structures. During refinement, tight NCS restraints were initially used; these were gradually relaxed and finally local structure similarity restraints⁵⁴ applied. TLS groups⁵⁶ were identified using the *TLSMD* Web server⁵⁷. Data processing and refinement statistics are provided in Supplementary Table S1. *PISA*⁵⁸ and *CONTACTS*⁵⁸ were used to analyze protein-protein interfaces, and *DynDom*⁵⁹ was used for conformational change analysis. All of the structural figures presented were generated using *PyMOL*⁶⁰.

Isothermal titration calorimetry. The derCD23 and IgE-Fc proteins were dialyzed overnight into 25 mM Tris-HCl (pH 7.4), 125 mM NaCl supplemented with 4 mM CaCl_2 ²⁰. The experiment was carried out at 25 °C in an iTC200 microcalorimeter (Microcal, GE Healthcare). To measure the affinity of derCD23 to IgE-Fc, 260 μM derCD23 was loaded in the syringe and titrated into 12.5 μM of IgE-Fc dimer within the sample cell. Data analysis was carried out using the Origin software supplied with the machine using a 2:1 binding model. Values in parentheses in Table 1 indicate the standard deviation.

References

- Pawankar, R., Walter Canonica, G., Holgate, S. T. & Lockey, R. F. World Allergy Organization. *White Book on Allergy 2011–2012 Executive Summary* (2011).
- DunnGalvin, A., Dubois, A. E., Flokstra-de Blok, B. M. & Hourihane, J. O. The effects of food allergy on quality of life. *Chem Immunol Allergy* **101**, 235–252, doi: 10.1159/000375106 (2015).
- Lifshitz, C. The impact of atopic dermatitis on quality of life. *Ann Nutr Metab* **66** Suppl 1, 34–40, doi: 10.1159/000370226 (2015).
- Meltzer, E. O. Allergic Rhinitis: Burden of Illness, Quality of Life, Comorbidities, and Control. *Immunol Allergy Clin North Am* **36**, 235–248, doi: 10.1016/j.jiac.2015.12.002 (2016).
- Sharma, H. P. & Herbert, L. J. Food allergy: psychosocial impact and public policy implications. *Chem Immunol Allergy* **101**, 221–226, doi: 10.1159/000373907 (2015).

6. Gupta, R. *et al.* The economic impact of childhood food allergy in the United States. *JAMA Pediatr* **167**, 1026–1031, doi: 10.1001/jamapediatrics.2013.2376 (2013).
7. Mir, E., Panjabi, C. & Shah, A. Impact of allergic rhinitis in school going children. *Asia Pacific allergy* **2**, 93–100, doi: 10.5415/apallergy.2012.2.2.93 (2012).
8. Bock, S. A., Munoz-Furlong, A. & Sampson, H. A. Fatalities due to anaphylactic reactions to foods. *J Allergy Clin Immunol* **107**, 191–193, doi: 10.1067/mai.2001.112031 (2001).
9. Gould, H. J. & Sutton, B. J. IgE in allergy and asthma today. *Nature reviews. Immunology* **8**, 205–217, doi: 10.1038/nri2273 (2008).
10. Sutton, B. J. & Davies, A. M. Structure and dynamics of IgE-receptor interactions: FcepsilonRI and CD23/FcepsilonRII. *Immunol Rev* **268**, 222–235, doi: 10.1111/imr.12340 (2015).
11. Holdom, M. D. *et al.* Conformational changes in IgE contribute to its uniquely slow dissociation rate from receptor FcεRI. *Nature structural & molecular biology* **18**, 571–576, doi: 10.1038/nsmb.2044 (2011).
12. Voehringer, D. Protective and pathological roles of mast cells and basophils. *Nature reviews. Immunology* **13**, 362–375, doi: 10.1038/nri3427 (2013).
13. Zelensky, A. N. & Greedy, J. E. The C-type lectin-like domain superfamily. *The FEBS journal* **272**, 6179–6217, doi: 10.1111/j.1742-4658.2005.05031.x (2005).
14. Beavil, A. J., Edmeades, R. L., Gould, H. J. & Sutton, B. J. Alpha-helical coiled-coil stalks in the low-affinity receptor for IgE (Fc epsilon RII/CD23) and related C-type lectins. *Proceedings of the National Academy of Sciences of the United States of America* **89**, 753–757 (1992).
15. Hibbert, R. G. *et al.* The structure of human CD23 and its interactions with IgE and CD21. *The Journal of experimental medicine* **202**, 751–760, doi: 10.1084/jem.20050811 (2005).
16. Chen, B. H. *et al.* Necessity of the stalk region for immunoglobulin E interaction with CD23. *Immunology* **107**, 373–381 (2002).
17. Dierks, S. E. *et al.* The oligomeric nature of the murine Fc epsilon RII/CD23. Implications for function. *J Immunol* **150**, 2372–2382 (1993).
18. Kelly, A. E., Chen, B. H., Woodward, E. C. & Conrad, D. H. Production of a chimeric form of CD23 that is oligomeric and blocks IgE binding to the Fc epsilonRI. *Journal of immunology* **161**, 6696–6704 (1998).
19. McCloskey, N. *et al.* Soluble CD23 monomers inhibit and oligomers stimulate IgE synthesis in human B cells. *The Journal of biological chemistry* **282**, 24083–24091, doi: 10.1074/jbc.M703195200 (2007).
20. Yuan, D. *et al.* Ca²⁺-dependent Structural Changes in the B-cell Receptor CD23 Increase Its Affinity for Human Immunoglobulin E. *The Journal of biological chemistry* **288**, 21667–21677, doi: 10.1074/jbc.M113.480657 (2013).
21. Weskamp, G. *et al.* ADAM10 is a principal 'shedase' of the low-affinity immunoglobulin E receptor CD23. *Nature immunology* **7**, 1293–1298, doi: 10.1038/nri1399 (2006).
22. Schulz, O. *et al.* Cleavage of the low-affinity receptor for human IgE (CD23) by a mite cysteine protease: nature of the cleaved fragment in relation to the structure and function of CD23. *Eur J Immunol* **27**, 584–588, doi: 10.1002/eji.1830270303 (1997).
23. Acharya, M. *et al.* CD23/FcepsilonRII: molecular multi-tasking. *Clin Exp Immunol* **162**, 12–23, doi: 10.1111/j.1365-2249.2010.04210.x (2010).
24. Tu, Y. *et al.* CD23-mediated IgE transport across human intestinal epithelium: inhibition by blocking sites of translation or binding. *Gastroenterology* **129**, 928–940, doi: 10.1053/j.gastro.2005.06.014 (2005).
25. Palaniyandi, S., Tomei, E., Li, Z., Conrad, D. H. & Zhu, X. CD23-dependent transcytosis of IgE and immune complex across the polarized human respiratory epithelial cells. *Journal of immunology* **186**, 3484–3496, doi: 10.4049/jimmunol.1002146 (2011).
26. Mudde, G. C., Reischul, I. G., Corvaia, N., Hren, A. & Poellbauer, E. M. Antigen presentation in allergic sensitization. *Immunol Cell Biol* **74**, 167–173, doi: 10.1038/icb.1996.23 (1996).
27. van der Heijden, F. L., Joost van Neerven, R. J., van Katwijk, M., Bos, J. D. & Kapsenberg, M. L. Serum-IgE-facilitated allergen presentation in atopic disease. *Journal of immunology* **150**, 3643–3650 (1993).
28. Selb, R. *et al.* CD23 surface density on B cells is associated with IgE levels and determines IgE-facilitated allergen uptake, as well as activation of allergen-specific T cells. *J Allergy Clin Immunol*, doi: 10.1016/j.jaci.2016.03.042 (2016).
29. Fellmann, M., Buschor, P., Rothlisberger, S., Zellweger, F. & Vogel, M. High affinity targeting of CD23 inhibits IgE synthesis in human B cells. *Immun Inflamm Dis* **3**, 339–349, doi: 10.1002/iid3.72 (2015).
30. Yabuuchi, S., Nakamura, T., Kloetzer, W. S. & Reff, M. E. Anti-CD23 monoclonal antibody inhibits germline Cepsilon transcription in B cells. *Int Immunopharmacol* **2**, 453–461 (2002).
31. Dhaliwal, B. *et al.* Crystal structure of IgE bound to its B-cell receptor CD23 reveals a mechanism of reciprocal allosteric inhibition with high affinity receptor FcepsilonRI. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 12686–12691, doi: 10.1073/pnas.1207278109 (2012).
32. Shi, J. *et al.* Interaction of the low-affinity receptor CD23/Fc epsilonRII lectin domain with the Fc epsilon3-4 fragment of human immunoglobulin E. *Biochemistry* **36**, 2112–2122, doi: 10.1021/bi961231e (1997).
33. Wan, T. *et al.* The crystal structure of IgE Fc reveals an asymmetrically bent conformation. *Nature immunology* **3**, 681–686, doi: 10.1038/nri811 (2002).
34. Borthakur, S. *et al.* Mapping of the CD23 binding site on IgE and allosteric control of the IgE-FcεRI interaction. *The Journal of biological chemistry*, doi: 10.1074/jbc.C112.397059 (2012).
35. Dhaliwal, B. *et al.* Conformational plasticity at the IgE-binding site of the B-cell receptor CD23. *Mol Immunol* **56**, 693–697, doi: 10.1016/j.molimm.2013.07.005 (2013).
36. Dhaliwal, B., Pang, M. O., Yuan, D., Beavil, A. J. & Sutton, B. J. A range of Cε3–Cε4 interdomain angles in IgE Fc accommodate binding to its receptor CD23. *Acta crystallographica. Section F, Structural biology communications* **70**, 305–309, doi: 10.1107/S2053230X14003355 (2014).
37. Lawrence, M. C. & Colman, P. M. Shape complementarity at protein/protein interfaces. *Journal of molecular biology* **234**, 946–950, doi: 10.1006/jmbi.1993.1648 (1993).
38. Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta crystallographica. Section D, Biological crystallography* **67**, 235–242, doi: 10.1107/S0907444910045749 (2011).
39. Beavil, A. J., Young, R. J., Sutton, B. J. & Perkins, S. J. Bent domain structure of recombinant human IgE-Fc in solution by X-ray and neutron scattering in conjunction with an automated curve fitting procedure. *Biochemistry* **34**, 14449–14461 (1995).
40. Wurzburg, B. A., Garman, S. C. & Jardetzky, T. S. Structure of the human IgE-Fc Cε3–Cε4 reveals conformational flexibility in the antibody effector domains. *Immunity* **13**, 375–385 (2000).
41. Wurzburg, B. A. & Jardetzky, T. S. Conformational flexibility in immunoglobulin E-Fc 3-4 revealed in multiple crystal forms. *Journal of molecular biology* **393**, 176–190, doi: 10.1016/j.jmb.2009.08.012 (2009).
42. Garman, S. C., Wurzburg, B. A., Tarchevskaya, S. S., Kinet, J. P. & Jardetzky, T. S. Structure of the Fc fragment of human IgE bound to its high-affinity receptor Fc epsilonRI alpha. *Nature* **406**, 259–266, doi: 10.1038/35018500 (2000).
43. Andersen, C. B. & Moestrup, S. K. How calcium makes endocytic receptors attractive. *Trends in biochemical sciences* **39**, 82–90, doi: 10.1016/j.tibs.2013.12.003 (2014).
44. Hunt, J. *et al.* A fluorescent biosensor reveals conformational changes in human immunoglobulin E Fc: implications for mechanisms of receptor binding, inhibition and allergen recognition. *The Journal of biological chemistry*, doi: 10.1074/jbc.M111.331967 (2012).
45. Drinkwater, N. *et al.* Human immunoglobulin E flexes between acutely bent and extended conformations. *Nature structural & molecular biology* **21**, 397–404, doi: 10.1038/nsmb.2795 (2014).

46. Plomp, R. *et al.* Site-specific N-glycosylation analysis of human immunoglobulin E. *J Proteome Res* **13**, 536–546, doi: 10.1021/pr400714w (2014).
47. Young, R. J. *et al.* Secretion of recombinant human IgE-Fc by mammalian cells and biological activity of glycosylation site mutants. *Protein engineering* **8**, 193–199 (1995).
48. Powell, H. R. The Rossmann Fourier autoindexing algorithm in MOSFLM. *Acta crystallographica. Section D, Biological crystallography* **55**, 1690–1695 (1999).
49. Battye, T. G., Kontogiannis, L., Johnson, O., Powell, H. R. & Leslie, A. G. iMOSFLM: a new graphical interface for diffraction-image processing with MOSFLM. *Acta crystallographica. Section D, Biological crystallography* **67**, 271–281, doi: 10.1107/S0907444910048675 (2011).
50. Evans, P. R. & Murshudov, G. N. How good are my data and what is the resolution? *Acta crystallographica. Section D, Biological crystallography* **69**, 1204–1214, doi: 10.1107/S0907444913000061 (2013).
51. McCoy, A. J. *et al.* Phaser crystallographic software. *J. Appl. Crystallogr.* **40**, 658–674, doi: 10.1107/S0021889807021206 (2007).
52. Adams, P. D. *et al.* The Phenix software for automated determination of macromolecular structures. *Methods* **55**, 94–106, doi: 10.1016/j.jymeth.2011.07.005 (2011).
53. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal structures. *Acta crystallographica. Section D, Biological crystallography* **67**, 355–367, doi: 10.1107/S0907444911001314 (2011).
54. Smart, O. S. *et al.* Exploiting structure similarity in refinement: automated NCS and target-structure restraints in BUSTER. *Acta crystallographica. Section D, Biological crystallography* **68**, 368–380, doi: 10.1107/S0907444911056058 (2012).
55. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta crystallographica. Section D, Biological crystallography* **66**, 486–501, doi: 10.1107/S0907444910007493 (2010).
56. Painter, J. & Merritt, E. A. Optimal description of a protein structure in terms of multiple groups undergoing TLS motion. *Acta crystallographica. Section D, Biological crystallography* **62**, 439–450, doi: 10.1107/S0907444906005270 (2006).
57. Painter, J. & Merritt, E. A. A molecular viewer for the analysis of TLS rigid-body motion in macromolecules. *Acta crystallographica. Section D, Biological crystallography* **61**, 465–471, doi: 10.1107/S0907444905001897 (2005).
58. Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline state. *Journal of molecular biology* **372**, 774–797, doi: 10.1016/j.jmb.2007.05.022 (2007).
59. Hayward, S. & Berendsen, H. J. Systematic analysis of domain motions in proteins from conformational change: new results on citrate synthase and T4 lysozyme. *Proteins* **30**, 144–154 (1998).
60. Schrödinger, L. The PyMOL Graphics System, Version 1.5.0 (2011).

Acknowledgements

This study was supported by The Wellcome Trust, Asthma UK, Medical Research Council (UK) (grant reference numbers: MRC G1100090; WT 076343 & 085944). We thank Diamond Light Source for access to beamline I03 (grant MX9495) that contributed to the results presented here.

Author Contributions

B.D., H.J.G., J.M.M., B.J.S. and A.J.B. conceived the project. B.D., M.O.Y.P., A.H.K. and L.K.J. designed and performed the experiments, and analyzed the data. B.D., J.M.M., B.J.S. and A.J.B. wrote the paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing Interests: The authors declare no competing financial interests.

How to cite this article: Dhaliwal, B. *et al.* IgE binds asymmetrically to its B cell receptor CD23. *Sci. Rep.* **7**, 45533; doi: 10.1038/srep45533 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017