

Review

Autophagic programmed cell death in *Drosophila*EH Baehrecke^{*,1}¹ Center for Biosystems Research, University of Maryland Biotechnology Institute, College Park, MD 20742, USA

* Corresponding author: EH Baehrecke E-mail: baehreck@umbi.umd.edu

Received 26.2.03; accepted 22.4.03

Edited by M Piacentini

Abstract

Autophagic programmed cell death occurs during the development of diverse animal groups, but the mechanisms that control this genetically regulated form of cell killing are poorly understood. Genetic studies of bulk protein degradation in yeast have provided important advances in our understanding of autophagy, and recent investigations of *Drosophila* autophagic cell death suggest that some of these mechanisms may be conserved. In *Drosophila*, several steroid-regulated genes that encode transcription regulators are required for autophagic cell death. These transcription regulators appear to activate a large number of genes that play a more direct role in cell killing, including genes that function in apoptosis such as caspases. While caspase function is required for autophagic cell death during *Drosophila* development, genes encoding proteins that are similar to the yeast autophagy regulators are also induced in dying salivary glands. Furthermore, numerous noncaspase proteases, cytoplasmic organizing factors, signaling molecules, and unknown factors are expressed in interesting patterns during autophagic cell death. This article reviews the current knowledge of the regulation of autophagic programmed cell death during development of *Drosophila*.

Cell Death and Differentiation (2003) 10, 940–945. doi:10.1038/sj.cdd.4401280

Keywords: autophagy; apoptosis; proteolysis; cytoplasmic degradation; *Drosophila*

Abbreviations: dark, fly Apaf-1 homolog ark; Nc, the caspase dronc

Introduction

Schweichel and Merker¹ recognized that at least three forms of cell death occur during development of animals based on the location and role of lysosomes. While nonlysosomal cell death has not been commonly observed, apoptosis and autophagic programmed cell death are prominent in developing animals of diverse taxa.² Apoptotic cells exhibit condensation of the nucleus and cytoplasm, fragmentation, and they are engulfed by phagocytes where the lysosome

completes degradation of the dying cell.³ Unlike apoptosis, cells that die by autophagic cell death are degraded with little or no help from phagocytes and utilize lysosomes within the dying cell for degradation. Defects in apoptosis have been associated with a variety of human disorders including cancer,⁴ and studies of the mechanisms underlying apoptosis have been the focus of cell death research.^{5,6} In contrast, little is known about the mechanisms that regulate autophagic cell death.

The degradation of cytoplasmic organelles and proteins is an important component of proper cell life and death. While the ubiquitin/proteasome pathway is used to degrade many short-lived regulatory proteins,⁷ long-lived proteins are targeted to the lysosome by five mechanisms including direct transport, vacuolar import and degradation, endocytosis, microautophagy, and macroautophagy.⁸ Macroautophagy (hereafter referred to as autophagy) is the best known mechanism for sequestration of long-lived proteins and cytoplasmic components into the lysosomal compartment of the cell for degradation. The first descriptions of lysosomal structures that contained cytoplasmic components such as mitochondria were observed in rat livers perfused with glucagon⁹ and rat hepatomas.¹⁰ These autophagic structures have since been described in organisms that are as different as yeast and humans. During autophagy, cytoplasm and organelles are sequestered in double-membrane vesicles known as autophagic vacuoles in organisms such as insects and mammals and as autophagosomes in yeast¹¹ (Figure 1). Although the origin of autophagic vacuole membranes are a subject of debate, evidence indicates that they are derived from the rough endoplasmic reticulum.¹² Autophagic vacuoles then dock against the lysosomal vacuole, and fusion of the outer autophagic vacuole membrane with the lysosomal membrane releases the inner membrane-bound cytoplasmic contents into the lysosome where degradation occurs.

Autophagic structures were first associated with developmental programmed cell death in insects^{13,14} and chickens,¹⁵ and it is now clear that autophagic cell death occurs during development of diverse organisms including humans.^{2,16–18} The recent association of autophagic cell death with models of tumorigenesis^{19–22} and neurodegenerative disorders^{23–26} has led to greater interest in the mechanisms that regulate this form of cell destruction.

Lessons from yeast

The molecular genetic basis of protein and organelle turnover by autophagy has been well characterized in the yeast *Saccharomyces cerevisiae*. In yeast, autophagy is induced under nutrient-limiting conditions. Genetic screens for yeast that either lacked accumulation of autophagic structures or exhibited defects in the degradation of cytoplasmic proteins under starvation conditions resulted in the identification of the *apg²⁷* and *aur²⁸* mutants. In addition, the isolation of altered




Figure 1 Diagram of the steps in autophagy. Autophagy is induced by starvation in yeast and cell death signals such as steroids during autophagic cell death in *Drosophila*. Induction results in the formation of a double-membraned autophagic vacuole that encloses cytoplasmic components including organelles such as mitochondria. The autophagic vacuole then docks against the lysosome, and the fusion of the outer autophagic vacuole membrane with the lysosome releases the inner membrane-bound cytoplasmic components into the lysosome where degradation occurs. Redrawn after Klionsky and Emr¹¹

cytoplasm to vacuole-targeting mutants (*cvt*) in screens of yeast for defects in proteolytic processing of aminopeptidase I²⁹ led to the recognition of the genetic overlap between the *apg*, *aut*, and *cvt* pathways.³⁰ In total, 19 *apg*, *aut*, and *cvt* genes have been identified that regulate the induction, formation, size, and breakdown steps in yeast autophagy (reviewed in Ohsumi⁸ and Kilonsky and Emr¹¹).

Genetic and biochemical studies of yeast indicate that the *apg*, *aut*, and *cvt* genes have distinct and related functions. While the role of some of these genes remains mysterious, two related ubiquitin-like systems are required for yeast autophagy. APG12 and AUT7/APG8/CVT5 encode proteins that show no apparent homology to ubiquitin, but both of these proteins appear to function in ubiquitin-like conjugation systems.^{31,32} APG12 and AUT7/APG8/CVT5 are both activated by the E1-like protein APG7/CVT2.^{33,34} APG12 is then conjugated to the E2-like protein APG10,³⁵ while AUT7/APG8/CVT5 becomes conjugated to the E2-like protein AUT1/APG3.³¹ APG12 is then conjugated to APG5³², and AUT7/APG8/CVT5 is attached to the lipid phosphatidylethanolamine.³¹ Mutations in either of these pathways prevent the proper formation of autophagic vacuoles,²⁷ and indicate that ubiquitin-like systems are involved in the bulk degradation of long-lived cytoplasmic proteins. While it is not clear if the genes that are required for yeast autophagy are also required for the formation of autophagic structures during programmed cell death, recent studies in *Drosophila* suggest similarity in the mechanisms of cytoplasmic degradation.³⁶

Activation of autophagic cell death in *Drosophila*

The steroid hormone 20-hydroxyecdysone (ecdysone) titer begins to rise 10 h after puparium formation, and reaches its peak level 12 h after puparium formation when it triggers the synchronous death of larval salivary gland cells.^{37,38} This cell death is preceded by markers of apoptosis including DNA fragmentation and nuclear acridine orange staining.³⁹ While these data suggest that salivary glands die by apoptosis, their morphology indicates that they die by autophagy (Figure 2). At 1 h after this peak in ecdysone titer (13 h after puparium

formation), salivary gland cells become round in shape, large cytoplasmic vacuoles appear to fragment, and a second class of vacuoles appears that is associated with the plasma membrane.⁴⁰ By 14 h after puparium formation, salivary glands possess autophagic vacuoles that contain cytoplasmic structures including mitochondria.⁴⁰ The cellular changes that occur following the rise in ecdysone titer are accompanied by changes in the tubulin and actin cytoskeleton, and accumulation of acid phosphatase activity.⁴¹ Salivary glands are destroyed 16-h following puparium formation. These changes in cell morphology indicate that the production and movement of vacuoles, reorganization of the cytoplasm and cell shape, and cytoplasmic degradation by proteases may all be involved in the autophagic destruction of salivary glands.

Drosophila salivary glands have served as a long-standing model for studies of steroid signaling.^{42–44} Ecdysone activates salivary gland cell death through a transcriptional regulatory hierarchy (Figure 3). The ecdysone receptor is encoded by the nuclear receptor genes *EcR* and *usp*.^{45–50} The ecdysone receptor complex, and the nuclear receptor competence factor β FTZ-F1, activate transcription of the early genes *BR-C*, *E74A*, and *E93*.^{51,52} Salivary glands are not destroyed in animals with mutations in β FTZ-F1, *BR-C*, *E74A*, and *E93*.^{52–55} While β FTZ-F1 and *E93* mutants exhibit defects in vacuolar changes during salivary gland autophagic cell death, *BR-C* and *E74A* mutant salivary glands progress to a very late stage in cell death even though complete destruction is prevented.⁴⁰ In addition, *E93* mutants exhibit similar defects in the formation of autophagic vacuoles in dying midgut cells,⁵⁶ indicating that *E93* is required for autophagy in dying *Drosophila* cells. *BR-C*, *E74A*, and *E93* all encode proteins that regulate transcription of target late genes,^{54,55,57,58} and these downstream genes appear to function more directly in programmed cell death.




Figure 2 Steroid regulation of autophagic programmed cell death in *Drosophila* salivary glands. At 10 h after puparium formation, the ecdysone titer starts to rise peaking 12 h after puparium formation. This pulse of steroid triggers the autophagic destruction of salivary glands, and this process is complete 16 h after puparium formation. Following the peak in ecdysone, cells become round in shape, large eosin-positive vacuoles (V) appear to fragment, and a new class of eosin-negative vacuoles (*) appear along the plasma membrane, and the nuclei (N) remains intact even though DNA fragmentation has taken place. At 14 hours after puparium formation, autophagic vacuoles containing mitochondria are present in the cytoplasm of salivary gland cells (arrows). Data reproduced from Lee and Baehrecke⁴⁰




Figure 3 Model for genetic regulation of autophagic programmed cell death in *Drosophila* salivary glands. At 10 h after puparium formation, the ecdysone titer rises and this steroid is bound by the receptor complex that is encoded by *EcR* and *usp*. The ecdysone receptor complex and β *FTZ-F1* activate the *BR-C*, *E74A*, and *E93* early genes. *BR-C*, *E74A*, and *E93* regulate genes involved in signaling, cytoplasmic reorganization, proteolysis, autophagy (*apg*-related genes), and apoptosis

Apoptosis genes function in *Drosophila* autophagic cell death

Several lines of evidence indicate that the core apoptosis machinery functions in autophagic cell death during *Drosophila* development. Autophagic cell death of *Drosophila* salivary glands and midguts is foreshadowed by markers of apoptosis including DNA fragmentation.^{39,40,56} In addition, destruction of salivary glands and midguts is prevented by expression of the caspase inhibitor p35.^{39,40,56} Significantly, the proapoptotic genes *rpr* and *hid* (*W*) increase in transcription following the rise in ecdysone that triggers salivary autophagic cell death, and this change is concurrent with a decrease in the inhibitor of apoptosis *diap2*.³⁹ At the same time, the fly *Apaf-1* homologue *ark* (*dark*, *dapaf1*, *hac1*), the caspases *dronc* (*Nc*), *drice* (*Ice*), and *dream* (*strica*), the Bcl-2 family member *buffy* (*dborg-2*), the CD36 realtive *crq*, and the DNase *rep4*, all exhibit dramatic increases in transcription just before salivary gland autophagic cell death.^{36,59,60}

Animals with mutations in the transcription regulators *BR-C*, *E74A*, and *E93* prevent proper transcription of apoptosis genes (Figure 3). Both *BR-C* and *E93* mutant salivary glands have dramatically reduced levels of *rpr*, *hid*, *dronc*, *dream*, and *crq*.^{36,59} While *buffy* and *drice* RNA levels are also decreased in *E93* mutant salivary glands, *drice* is not altered and *buffy* is ectopically expressed in *BR-C* mutants.³⁶ *E74A* mutants exhibit reduced levels of *hid*, *crq*, *buffy*, *drice*, and *dream*.^{36,59} Studies of apoptosis gene promoters indicate that *rpr* is a direct target of the ecdysone receptor⁵⁴ and that *dronc* is a direct target of *BR-C*.⁶¹ These data suggest that the ecdysone-regulated early genes *BR-C*, *E74A*, and *E93* have overlapping and distinct target apoptosis genes that they regulate. Significantly, it appears that proper transcription of apoptosis genes is essential for autophagic cell death, since animals with mutations in *BR-C*, *E74A*, and *E93* prevent destruction of salivary glands. This is intriguing, as studies of the human mammary carcinoma cell line MCF-7 indicate that caspases are not required for autophagic death of these cells.⁶²

Yeast autophagy genes appear to be conserved in higher animals, and are induced during autophagic programmed cell death

In all, 11 of the 19 genes that are required for yeast autophagy encode related genes in flies and humans (hereafter referred to as *apg*-related genes)^{16,63} (Table 1). Nine of these 11 *apg*-related genes are transcribed in dying *Drosophila* glands, and the seven genes that are most similar to *apg2* (*CG1241*), *apg3* (*CG6877*), *apg4* (*CG6194*), *apg5* (*CG1643*), *apg7* (*CG5489*), *apg9* (*CG3615*), and *aut10/cvt18* (*CG7986*) exhibit increased transcription following the rise in steroid hormone that triggers autophagic programmed cell death of this tissue.^{36,64} It is interesting that flies possess genes that are related to the yeast genes involved in both of the autophagy ubiquitin conjugation systems, and that they are also the genes that increase in transcription following the rise in steroid that triggers autophagic cell death. *CG1241* encodes a novel protein that is similar to APG2 that functions in the formation and completion of autophagic vacuoles.⁶⁵ APG2 interacts with the integral membrane protein APG9/AUT9/CVT7,⁶⁶ which is similar to the predicted protein encoded by *CG3615*. *CG7986* encodes a predicted protein that is most similar to AUT10/CVT18 which has WD40 domains, functions in the formation of autophagic vacuoles, and is required for proper localization of APG2.⁶⁷ *CG6194* encodes a predicted protein that is similar to the AUT2/APG4 cysteine protease, which interacts with and processes AUT7/APG8/CVT5, and is involved in the formation and completion of autophagic vacuoles.⁶⁸ AUT7/APG8/CVT5 is conjugated with the ubiquitin-E1-like enzyme APG7/CVT2,³¹ which is similar to the protein encoded by *CG5489*, indicating that one of the ubiquitin-like conjugation systems may function in salivary gland autophagy. AUT7/APG8/CVT5 is also conjugated with the ubiquitin-E2-like enzyme AUT1/APG3.³¹ AUT1/APG3 is similar to the predicted protein encoded by *CG6877*, suggesting the possibility that the second ubiquitin-like conjugation system functions in salivary gland autophagy. *CG1643* encodes a predicted protein that is similar to the novel protein APG5,⁶⁹ which also functions in the formation of

Table 1 Yeast autophagy proteins, their biochemical properties, and the *Drosophila* gene encoding the most similar predicted protein

Yeast protein	Biochemical properties	<i>Drosophila</i> gene
APG1/AUT3/CVT10	Serine/threonine kinase	CG8866
APG2/AUT8	Interacts with APG9	CG1241
AUT1/APG3	E2-like, conjugated with AUT7/APG8/CVT5	CG6877
AUT2/APG4	Cysteine protease, interacts with AUT7/APG8/CVT5	CG6194
APG5	Conjugated with APG12	CG1643
APG6	Interacts with APG14	CG5429
APG7/CVT2	E1-like, conjugated with APG12 or AUT7/APG8/CVT5	CG5489
AUT7/APG8/CVT5	Conjugated with APG7/CVT2, AUT1/APG3	CG32672
APG9/AUT9/CVT7	Integral membrane protein, interacts with APG2	CG3615
APG12	Conjugated with APG7/CVT2, APG10, APG5	CG10861
AUT10/CVT18	WD40 domains	CG7986

Drosophila melanogaster genes were identified by using BLAST searches of the National Center for Biotechnology Information database (<http://www.ncbi.nlm.nih.gov/>). BLAST scores lower than 10^{-5} were considered similar. Yeast genes that function in autophagy, but exhibit low similarity to predicted fly genes, are not included

autophagic vacuoles. APG5 is conjugated to APG12, and this complex is also activated by the ubiquitin-E1-like enzyme APG7/CVT2,³² further indicating that both ubiquitin-like conjugation systems that are used during yeast autophagy may also function in salivary gland cell death.

Mutations in the ecdysone-regulated transcription factor genes that prevent proper salivary gland cell death also inhibit proper transcription of the *apg*-related genes.³⁶ Animals with mutations in *BR-C* exhibit ectopic transcription of *CG6194* (*apg4*-like) and *CG1643* (*apg5*-like), and decreased transcription of *CG5489* (*apg7*-like) in salivary glands at the stage that they would normally die. While animals with mutations in *E74A* only exhibit decreased transcription of *CG6194*, salivary glands dissected from *E93* mutants have decreased levels of *CG1241* (*apg2*-like), *CG6194*, *CG1643*, and *CG5489*. Yeasts with mutations in *apg4*, *apg5*, and *apg7* are defective in autophagic vacuole formation.²⁷ Thus, it is striking that *E93* mutants exhibit decreased *CG6194*, *CG1643*, and *CG5489* RNA levels, since *E93* mutant salivary gland and midgut cells are defective in the formation of autophagic vacuoles.^{40,56}

Additional factors that are induced during *Drosophila* autophagic cell death

Genomewide studies of dying cells have identified numerous interesting genes that are induced just prior to autophagic programmed cell death (Figure 3). For example, several signaling molecules and transcription regulators increase in transcription in dying salivary glands. The *Drosophila* gene *CG8304* encodes a serine/threonine kinase that is most similar to the death-associated protein kinase (DAPK) in humans, and transcription of this gene increases 36-fold in dying salivary glands.³⁶ DAPK mediates membrane blebbing and formation of vesicles in dying human cell lines,⁷⁰ and suggests that *CG8304* may have a related function in *Drosophila*. Steroids usually function by activating transcription and, therefore, it is not surprising that transcription regulators are induced in dying salivary glands including the corepressors (*Smarter* and *CG4756*), the NFκB regulator *cactus*, the NFκB family member *dif*, several other genes encoding DNA-binding proteins (*bun*, *sox14*, *CG8319*), and components of transcription complexes (*trap95* and *hsf*).^{36,64}

Salivary gland cells exhibit dynamic changes in cell shape and vacuole localization just before autophagic cell destruction, suggesting that cytoplasmic reorganization and proteolysis are important for their death. Consistent with this notion, genes encoding motor proteins including *ctp*, *ck*, and *dIc90F*, and members of the Rho, Rac, and Rab families of small guanosine triphosphates (GTPases) exhibited increase in transcription just before salivary gland autophagic cell death.^{36,64} While cell remodeling is known to play an important role in phagocyte engulfment of apoptotic cells,⁷¹ the role of these factors in autophagic cell death is less clear. In addition to caspases, several cysteine, serine, and metalloproteases increase in transcription in dying salivary glands, and these changes in RNA level are complemented by decreased transcription of cysteine, serine, and metalloprotease inhibitors.^{36,64} Furthermore, upregulation of the matrix metalloprotease *mmp1* is abolished in mutants that prevent salivary gland cell death.³⁶ These proteases likely complement caspase function during autophagic cell death, as mutations in *mmp2* prevent proper destruction of *Drosophila* midguts,⁷² this tissue dies by autophagic cell death,⁵⁶ and inhibition of caspases does not completely prevent autophagic changes in dying cells.⁴⁰

The most important advances in science often come from the exploration of the unknown. Therefore, it is interesting that many genes encoding novel factors are induced in dying cells. We have identified genes that increase following both radiation-induced apoptosis and steroid-triggered autophagy. Very few common genes were identified using this approach, but the genes *CG10965*, *CG17323*, *CG7144*, *EG25E8.4*, and *CG5254* increased in RNA levels during both ecdysone- and radiation-triggered cell death.³⁶ Transcription of these genes was also altered in salivary glands of *BR-C*, *E74A*, and *E93* mutants, suggesting a possible functional role for these genes in autophagic programmed cell death.

Concluding Remarks and Future Directions

Studies of developing animals indicate that at least two forms of programmed cell death have been conserved during evolution.² Studies of apoptosis in cell lines indicate that the

biochemical mechanisms underlying programmed cell death are similar in different cell types. Dying cell morphology can be very different,⁷³ however, and identifying the mechanisms that regulate different types of physiological cell death will lead to a better understanding of how defects in this fundamental cellular process lead to aberrations in animal growth and development.

The steroid ecdysone triggers autophagic cell death during *Drosophila* development, but other factors must provide competence signals for cells to be capable of dying. These 'competence factors' may include cell age, status in the cell cycle, the presence of a transcriptional activator, absence of a transcriptional repressor, or the state of chromatin to name a few possibilities. Ultimately, this steroid signal activates a group of transcription regulators that include *BR-C*, *E74A*, and *E93*. These early genes appear to regulate a complex series of downstream cellular events that include cytoplasmic reorganization including cell shape and vacuole transport, proteolysis by caspase and apparently noncaspase proteases, genes related to yeast genes that function in autophagy, and many unknowns.

Many gaps exist in our knowledge of the mechanism underlying autophagic cell death, and this is an exciting time to ask critical questions about this understudied form of cell destruction. How many of the genes that are expressed in dying cells are required, and is it critical for these factors to integrate in a specific temporal manner? Do the *apg*-related genes serve the same purpose in yeast and higher animals? Are the noncaspase genes that are used during autophagy used during apoptosis? What is the clinical relevance of autophagic programmed cell death? Given our limited knowledge of the mechanisms underlying autophagic programmed cell death, it is clear that this fundamental cellular process will be a fruitful area of investigation for years to come.

Acknowledgements

I thank many colleagues and the members of the Baehrecke laboratory for helpful discussions, and Deborah Berry for comments on the manuscript. Work on this subject has been supported by NIH grant GM59136.

References

- Schweichel J-U and Merker H-J (1973) The morphology of various types of cell death in prenatal tissues. *Teratology* 7: 253–266
- Clarke PGH (1990) Developmental cell death: morphological diversity and multiple mechanisms. *Anat. Embryol.* 181: 195–213
- Kerr JF, Wyllie AH and Currie AR (1972) Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br. J. Cancer* 26: 239–257
- Johnstone RW, Ruefli AA and Lowe SW (2002) Apoptosis: a link between cancer genetics and chemotherapy. *Cell* 108: 153–164
- Hengartner MO (2000) The biochemistry of apoptosis. *Nature* 407: 770–776
- Shi Y (2002) Mechanisms of caspase activation and inhibition during apoptosis. *Mol. Cell* 9: 459–470
- Weissman AM (2001) Themes and variations on ubiquitylation. *Nat. Rev. Mol. Cell Biol.* 2: 169–178
- Ohsumi Y (2001) Molecular dissection of autophagy: two ubiquitin-like systems. *Nat. Rev. Mol. Cell Biol.* 2: 211–216
- Ashford TP and Porter KR (1962) Cytoplasmic components in hepatic cell lysosomes. *J. Cell. Biol.* 12: 198–202
- Essner E and Novikoff AB (1962) Cytological studies on two functional hepatomas, interrelations of endoplasmic reticulum, golgi apparatus, and lysosomes. *J. Cell. Biol.* 15: 289–312
- Klionsky DJ and Emr SD (2000) Autophagy as a regulated pathway of cellular degradation. *Science* 290: 1717–1721
- Dunn WAJ (1990) Studies on the mechanisms of autophagy: formation of the autophagic vacuole. *J. Cell. Biol.* 110: 1923–1933
- Scharrer B (1966) Ultrastructural study of the regressing prothoracic glands of blattarian insects. *Z. Zellerforsch.* 69: 1–21
- Schin KS and Clever U (1965) Lysosomal and free acid phosphatase in salivary glands of *Chironomus tentans*. *Science* 150: 1053–1055
- Scheib D (1965) Sur la regression du canal de de Müller male de l'embryon de poulet: localisation de la phosphatase acide au microscope electronique. *C. R. Acad. Sci. Hebd. Seances. Acad. Sci. D* 261: 5219–5221
- Baehrecke EH (2002) How death shapes life during development. *Nat. Rev. Mol. Cell Biol.* 3: 779–787
- Tanida I, Tanida-Miyake E, Komatsu M, Ueno T and Kominami E (2002) Human Apg3p/Aut1p homologue is an authentic E2 enzyme for multiple substrates, GATE-16, GABARAP, and MAP-LC3, and facilitates the conjugation of hApg12p to hApg5p. *J. Biol. Chem.* 277: 13739–13744
- Tanida I, Tanida-Miyake E, Ueno T and Kominami E (2001) The human homolog of *Saccharomyces cerevisiae* Apg7p is a protein-activating enzyme for multiple substrates including human Apg12p, GATE-16, GABARAP, and MAP-LC3. *J. Biol. Chem.* 276: 1701–1706
- Schulte-Hermann R, Bursch W, Grasl-Kraupp B, Marian B, Torok L, Kahl-Rainer P and Ellinger A (1997) Concepts of cell death and application to carcinogenesis. *Toxicol. Pathol.* 25: 89–93
- Bursch W, Ellinger A, Kienzl H, Torok L, Pandey S, Sikorska M, Walker R and Herrmann RS (1996) Active cell death induced by the anti-estrogens tamoxifen and ICI164384 in human mammary carcinoma cells (MCF-7) in culture: the role of autophagy. *Carcinogenesis* 17: 1595–1607
- Liang XH, Jackson S, Seaman M, Brown K, Kempkes B, Hibshoosh H and Levine B (1999) Induction of autophagy and inhibition of tumorigenesis by *beclin 1*. *Nature* 402: 672–676
- Jia L, Doumashkin RR, Allen PD, Gray AB, Newland AC and Kelsey SM (1997) Inhibition of autophagy abrogates tumour necrosis factor alpha induced apoptosis in human T-lymphoblastic leukaemic cells. *Br. J. Haematol.* 98: 673–685
- Kegel KB, Kim M, Sapp E, McIntyre C, Castano JG, Aronin N and DiFiglia M (2000) Huntingtin expression stimulates endosomal-lysosomal activity, endosome tubulation, and autophagy. *J. Neurosci.* 20: 7268–7278
- Yue Z, Horton A, Bravin M, DeJager PL and Selimi FNH (2002) A novel protein complex linking the delta 2 glutamate receptor and autophagy: implications for neurodegeneration in lurcher mice. *Neuron* 35: 921–933
- Stefanis L, Larsen KE, Rideout HJ, Sulzer D and Greene LA (2001) Expression of A53T mutant but not wild-type alpha-synuclein in PC12 cells induces alterations of the ubiquitin-dependent degradation system, loss of dopamine release, and autophagic cell death. *J. Neurosci.* 21: 9549–9560
- Mastroberardino PG, Iannicola C, Nardacci R, Bernassola F, De Laurenzi V, Melino G, Moreno S, Pavone F, Oliverio S, Fesus L and Piacentini M (2002) 'Tissue' transglutaminase ablation reduces neuronal death and prolongs survival in a mouse model of Huntington's disease. *Cell Death Differ.* 9: 873–880
- Tsukada M and Ohsumi Y (1993) Isolation and characterization of autophagy-defective mutants of *Saccharomyces cerevisiae*. *FEBS Lett.* 1–2: 169–174
- Thumm M, Egner R, Koch B, Schlumpberger M, Straub M, Veenhuis M and Wolf DH (1994) Isolation of autophagocytosis mutants of *Saccharomyces cerevisiae*. *FEBS Lett.* 349: 275–280
- Harding TM, Morano KA, Scott SV and Klionsky DJ (1995) Isolation and characterization of yeast mutants in the cytoplasm to vacuole protein targeting pathway. *J. Cell. Biol.* 131: 591–602
- Harding TM, Hefner-Gravink A, Thumm M and Klionsky DJ (1996) Genetic and phenotypic overlap between autophagy and the cytoplasm to vacuole protein. *J. Biol. Chem.* 271: 17621–17624
- Ichimura Y, Kirisako T, Takao T, Satomi Y, Shimonishi Y, Ishihara N, Mizushima N, Tanida I, Kominami E, Ohsumi M, Noda T and Ohsumi Y (2000) A ubiquitin-like system mediates protein lipidation. *Nature* 408: 488–492

32. Mizushima N, Noda T, Yoshimori T, Tanaka Y, Ishii T, George MD, Klionsky DJ, Ohsumi M and Ohsumi Y (1998) A protein conjugation system essential for autophagy. *Nature* 395: 395–398
33. Tanida I, Mizushima N, Kiyooka M, Ohsumi M, Ueno T, Ohsumi Y and Kominami E (1999) Apg7p/Cvt2p: a novel protein-activating enzyme essential for autophagy. *Mol. Biol. Cell* 10: 1367–1379
34. Kim J, Dalton VM, Eggerton KP, Scott SV and Klionsky DJ (1999) Apg7p/Cvt2p is required for the cytoplasm-to-vacuole targeting, macroautophagy, and peroxisome degradation pathways. *Mol. Biol. Cell* 10: 1337–1351
35. Shintani T, Mizushima N, Ogawa Y, Matsuura A, Noda T and Ohsumi Y (1999) Apg10p, a novel protein-conjugating enzyme essential for autophagy in yeast. *EMBO J.* 18: 5234–5241
36. Lee C-Y, Clough EA, Yellon P, Teslovich TM, Stephan DA and Baehrecke EH (2003) Genome-wide analyses of steroid- and radiation-triggered programmed cell death in *Drosophila*. *Curr. Biol.* 13: 350–357
37. Bodenstein D (1965) The postembryonic development of *Drosophila*. In *Biology of Drosophila* Demerec M (ed) New York: Hafner Publishing Co., pp. 275–367
38. Robertson CW (1936) The metamorphosis of *Drosophila melanogaster*, including an accurately timed account of the principal morphological changes. *J. Morphol.* 59: 351–399
39. Jiang C, Baehrecke EH and Thummel CS (1997) Steroid regulated programmed cell death during *Drosophila* metamorphosis. *Development* 124: 4673–4683
40. Lee CY and Baehrecke EH (2001) Steroid regulation of autophagic programmed cell death during development. *Development* 128: 1443–1455
41. Jochova J, Zakeri Z and Lockshin RA (1997) Rearrangement of the tubulin and actin cytoskeleton during programmed cell death in *Drosophila* salivary glands. *Cell Death Differ.* 4: 140–149
42. Becker HJ (1959) Die puffs der speicheldrüsenchromosomen von *Drosophila melanogaster* I. Beobachtungen zur verhalten des puffmusters in normalstamm und bei zwei mutanten, *giant* und *lethal-giant-larvae*. *Chromosoma* 10: 654–678
43. Ashburner M (1967) Patterns of puffing activity in the salivary gland chromosomes of *Drosophila*. I. Autosomal puffing patterns in a laboratory stock of *Drosophila melanogaster*. *Chromosoma* 21: 398–428
44. Andres AJ and Thummel CS (1992) Hormones, puffs, and flies: the molecular control of metamorphosis by ecdysone. *Trends Genet.* 8: 132–138
45. Koelle MR, Talbot WS, Segrevas WA, Bender MT, Cherbas P and Hogness DS (1991) The *Drosophila* EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily. *Cell* 67: 59–77
46. Shea MJ, King DL, Conboy MJ, Mariani BD and Kafatos FC (1990) Proteins that bind to *Drosophila* chorion cis-regulatory elements: a new C₂H₂ zinc finger protein and a C₂C₂ steroid receptor-like component. *Genes Dev.* 4: 1128–1140
47. Henrich VC, Sliter TJ, Lubahn DB, MacIntyre A and Gilbert LI (1990) A steroid/thyroid hormone receptor superfamily member in *Drosophila melanogaster* that shares extensive sequence similarity with a mammalian homologue. *Nucleic Acids Res.* 18: 4143–4148
48. Oro AE, McKeown M and Evans RM (1990) Relationship between the product of the *Drosophila ultraspiracle* locus and vertebrate retinoid X receptor. *Nature* 347: 298–301
49. Thomas HE, Stunnenberg HG and Stewart AF (1993) Heterodimerization of the *Drosophila* ecdysone receptor with retinoid X receptor and *ultraspiracle*. *Nature* 362: 471–475
50. Yao T-P, Segrevas WA, Oro AE, McKeown M and Evans RM (1992) *Drosophila ultraspiracle* modulates ecdysone receptor function via heterodimer formation. *Cell* 71: 63–72
51. Woodard CT, Baehrecke EH and Thummel CS (1994) A molecular mechanism for the stage-specificity of the *Drosophila* prepupal genetic response to ecdysone. *Cell* 79: 607–615
52. Broadus J, McCabe JR, Endrizzi B, Thummel CS and Woodard CT (1999) The *Drosophila* β FTZ-F1 orphan nuclear receptor provides competence for stage-specific responses to the steroid hormone ecdysone. *Mol. Cell* 3: 143–149
53. Restifo LL and White K (1992) Mutations in a steroid hormone-regulated gene disrupt the metamorphosis of internal tissues in *Drosophila*: salivary glands, muscle, and gut. *Wilhelm Roux's Arch. Dev. Biol.* 201: 221–234
54. Jiang C, Lamblin A-FJ, Steller H and Thummel CS (2000) A steroid-triggered transcriptional hierarchy controls salivary gland cell death during *Drosophila* metamorphosis. *Mol. Cell* 5: 445–455
55. Lee C-Y, Wendel DP, Reid P, Lam G, Thummel CS and Baehrecke EH (2000) *E93* directs steroid-triggered programmed cell death in *Drosophila*. *Mol. Cell* 6: 433–443
56. Lee C-Y, Cooksey BAK and Baehrecke EH (2002) Steroid regulation of midgut cell death during *Drosophila* development. *Dev. Biol.* 250: 101–111
57. Burtis KC, Thummel CS, Jones CW, Karim FD and Hogness DS (1990) The *Drosophila* 74EF early puff contains *E74*, a complex ecdysone-inducible gene that encodes two *ets*-related proteins. *Cell* 61: 85–99
58. DiBello PR, Withers DA, Bayer CA, Fristrom JW and Guild GM (1991) The *Drosophila Broad-Complex* encodes a family of related proteins containing zinc fingers. *Genetics* 129: 385–397
59. Lee C-Y, Simon CR, Woodard CT and Baehrecke EH (2002) Genetic mechanism for the stage- and tissue-specific regulation of steroid-triggered programmed cell death in *Drosophila*. *Dev. Biol.* 252: 138–148
60. Dorstyn L, Colussi PA, Quinn LM, Richardson H and Kumar S (1999) DRONC, an ecdysone-inducible *Drosophila* caspase. *Proc. Natl. Acad. Sci. USA* 96: 4307–4312
61. Kakourou D, Daish T, Martin D, Baehrecke EH and Kumar S (2002) Ecdysone-induced expression of the caspase DRONC during hormone-dependent programmed cell death in *Drosophila* is regulated by Broad-Complex. *J. Cell Biol.* 157: 985–995
62. Paglin S, Hollister T, Delohery T, Hackett N, McMahon M, Sphicas E, Domingo D and Yahalom J (2001) A novel response of cancer cells to radiation involves autophagy and formation of acidic vesicles. *Cancer Res.* 61: 439–444
63. Reggiori F and Klionsky DJ (2002) Autophagy in the eukaryotic cell. *Eukaryot. Cell* 1: 11–21
64. Gorski SM, Chittaranjan S, Pleasance ED, Freeman JD, Anderson CL, Varhol RJ, Coughlin SM, Zuyderduyn SD, Jones SJ and Marra MA (2003) A SAGE approach to discovery of genes involved in autophagic cell death. *Curr. Biol.* 13: 358–363
65. Wang CW, Kim J, Huang WP, Abeliovich H, Stromhaug PE, Dunn WAJ and Klionsky DJ (2001) Apg2 is a novel protein required for the cytoplasm to vacuole targeting, autophagy, and pexophagy pathways. *J. Biol. Chem.* 276: 30442–30451
66. Noda T, Kim J, Huang WP, Baba M, Tokunaga C, Ohsumi Y and Klionsky DJ (2000) Apg9p/Cvt7p is an integral membrane protein required for transport vesicle formation in the Cvt and autophagy pathways. *J. Cell Biol.* 148: 465–480
67. Guan J, Stromhaug PE, George MD, Habibzadegah-Tari P, Bevan A, Dunn WAJ and Klionsky DJ (2001) Cvt18/Gsa12 is required for cytoplasm-to-vacuole transport, pexophagy, and autophagy in *Saccharomyces cerevisiae* and *Pichia pastoris*. *Mol. Biol. Cell* 12: 3821–3838
68. Kirisako T, Ichimura Y, Okada H, Kabeya Y, Mizushima N, Yoshimori T, Ohsumi M, Takao T, Noda T and Ohsumi Y (2000) The reversible modification regulates the membrane-binding state of Apg8/Aut7 essential for autophagy and the cytoplasm to vacuole targeting pathway. *J. Cell. Biol.* 151: 263–276
69. Kametaka S, Matsuura A, Wada Y and Ohsumi Y (1996) Structural and functional analyses of APG5, a gene involved in autophagy in yeast. *Gene* 178: 139–143
70. Inbal B, Bialik S, Sabanay I, Shani G and Kimchi A (2002) DAP kinase and DRP-1 mediate membrane blebbing and the formation of autophagic vesicles during programmed cell death. *J. Cell Biol.* 157: 455–468
71. Hengartner MO (2001) Apoptosis: coralling the corpses. *Cell* 104: 325–328
72. Page-McCaw A, Serano J, Sante JM and Rubin GM (2003) *Drosophila* matrix metalloproteinases are required for tissue remodeling, but not embryonic development. *Dev. Cell* 4: 95–106
73. Leist M and Jätelä M (2001) Four deaths and a funeral: from caspases to alternative mechanisms. *Nat. Rev. Mol. Cell Biol.* 2: 1–10