

Correspondence

Cyclophilin A release as a biomarker of necrotic cell death

Cell Death and Differentiation (2010) 17, 1942–1943; doi:10.1038/cdd.2010.123; published online 8 October 2010

Dear Editor,

Necroptosis is a form of programmed necrotic cell death mediated by the activity of receptor-interacting protein 1 (RIP1) kinase.¹ This type of cell death is caspase-independent, is marked by early membrane permeabilization and does not exhibit any other hallmarks of apoptotic cell death such as phosphatidylserine externalization or DNA strand breaks.¹ Thus, the markers available to characterize apoptotic cell death cannot be used to analyze necroptosis. Only one marker of necrosis has been identified, and that is the release of the chromatin protein high-mobility group B1 (HMGB1).² However, we have found the release of HMGB1 to occur at a late stage of death. In this study, we have found that cyclophilin A (CypA), a cytosolic peptidyl-prolyl *cis-trans* isomerase, is released early in necroptosis. We propose that the release of CypA may be used as a biomarker for necroptosis and other cell death processes when the integrity of the plasma membrane is compromised.

As necroptotic cells exhibit early permeabilization of the plasma membrane,¹ we hypothesized that the release of intracellular protein(s) may be used as a biomarker for necroptosis. Treatment of L929 cells with zVAD.fmk induces necroptosis, which can be inhibited by treatment with the RIP1

kinase inhibitor, necrostatin-1 (Nec-1).¹ We observed the appearance of a strong band around 17 kDa by SDS-PAGE in the supernatant of zVAD-treated L929 cells at an early stage of cell death (Supplementary Figure 1, top panel). Mass spectrometry analysis identified this band as CypA, a soluble cytosolic protein that catalyzes the *cis-trans* isomerization of proline bonds. The release of CypA was further confirmed by western blot (Supplementary Figure 1, bottom panel). In contrast to the necrosis marker HMGB1, the timing of CypA release precedes or occurs around the same time as that of HMGB1 (Figure 1a). Nec-1 blocks the release of CypA into the supernatant (Figure 1b).

To determine whether the release of CypA from the cells was specific to necroptosis, we tested the supernatant of apoptotic cells. We were unable to detect CypA released from Jurkat cells treated with TNF α and CHX to induce apoptosis, even though intracellular CypA appears to be degraded at 24 h when cell death was completed (Figure 1c). However, FADD-deficient Jurkat cells, which undergo necroptosis in response to TNF α , did release CypA (Figure 1c). We also tested different inducers of apoptosis for the ability to release CypA. Treatment of doxorubicin and staurosporine caused a small amount of CypA




Figure 1 Cyclophilin A is released from necroptotic cells as a result of cytoplasmic membrane permeabilization. (a) L929 cells were treated with 100 μ M zVAD.fmk for indicated hours. The supernatants were collected, concentrated and analyzed by SDS-PAGE followed by western blotting using anti-CypA and anti-HMGB1. (b) L929 cells were treated with 100 μ M zVAD.fmk in the presence or absence of 30 μ M Nec-1. The cell lysate and supernatant were analyzed by western blotting using anti-CypA. Anti-tubulin was used as a control. Cell viability was determined by ATP assay and indicated as a bar graph below. (c) FADD-deficient Jurkat cells (left panel) or wt Jurkat cells (right panel) were treated with 10 ng/ml hTNF α to induce necroptosis, or 10 ng/ml hTNF α + 0.5 μ g/ml CHX to induce apoptosis, respectively. The cell lysate and supernatant were analyzed by western blotting using anti-CypA. The percentages of cell death were determined by ATP assay and indicated below

release at later stages of cell death, hours after caspase-3 activation (Supplementary Figures 2 and 3). Thus, CypA may also be released after permeabilization of the plasma membrane in the late phase of apoptotic cell death. However, as apoptotic cells are efficiently engulfed *in vivo*, the release of CypA by apoptotic cells may not occur *in vivo*.

To examine whether CypA could be released by permeabilization of the cells, we treated cells with the detergent digitonin, which led to cell membrane permeabilization. We found that digitonin-permeabilized cells also release CypA (Supplementary Figure 4). This may explain why CypA is released during necroptosis, which exhibit early permeabilization of the plasma membrane.¹

Elevated serum CypA levels have been found in inflammatory diseases such as rheumatoid arthritis and CypA is secreted *in vivo* by progressive solid tumors.^{3,4} CypA release from necrotic cells may have a role in the effects of necrosis on surrounding tissue. Necrosis is typically associated with inflammation. Necroptosis occurs in models of neuronal ischemia and traumatic brain injury.^{1,5} Treatment with Nec-1 significantly reduces cell death and inhibits the recruitment of neutrophils to the injured area.⁵ Extracellular CypA has been shown to induce a rapid inflammatory response upon injection *in vivo* and to recruit neutrophils. This recruitment is dependent on the CypA receptor, CD147 (EMMPRIN).³ Signaling of CypA through CD147 is proposed to contribute to the migration of neutrophils into the joints in rheumatoid arthritis and possibly in the destruction of bone and cartilage in the disease.⁶

We propose that CypA may be used as a biomarker for necroptosis. The release of CypA, in combination with Nec-1, a specific inhibitor of necroptosis, may be used as an assay for the activation of necroptosis, both *in vitro* and *in vivo*.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements. This work was supported in part by an NIH Director's Award and an R37 Merit Award (to JY) and an NIH F31Predoctoral Fellowship (to DC). Mass spectrometry analysis was conducted in the Taplin Mass Spectrometry Facility.

DE Christofferson¹ and J Yuan^{*,1}

¹ Department of Cell Biology, Harvard Medical School, 240 Longwood Avenue, Boston, MA 02115, USA

* Corresponding author: J Yuan, Department of Cell Biology, Harvard Medical School, 240 Longwood Avenue, Boston, MA 02115, USA.
Tel: +6 17 432 4187; Fax: +6 17 432 4177;
E-mail: jyuan@hms.harvard.edu

1. Degterev A *et al.* *Nat Chem Biol* 2005; **1**: 112–119.
2. Bianchi ME, Manfredi A. *Biochim Biophys Acta* 2004; **1677**: 181–186.
3. Arora K *et al.* *J Immunol* 2005; **175**: 517–522.
4. Huang CM *et al.* *Proteomics* 2006; **6**: 6107–6116.
5. You Z *et al.* *J Cereb Blood Flow Metab* 2008; **28**: 1564–1573.
6. Wang CH *et al.* *J Cell Mol Med* 2010; e-pub ahead of print.

Supplementary Information accompanies the paper on Cell Death and Differentiation website (<http://www.nature.com/cdd>)