

Tousled-like kinase mediated a new type of cell death pathway in *Drosophila*

Y Zhang^{1,2,3}, R Cai^{1,2,3}, R Zhou², Y Li^{*2} and L Liu^{*2}

Programmed cell death (PCD) has an important role in sculpting organisms during development. However, much remains to be learned about the molecular mechanism of PCD. We found that ectopic expression of *tousled-like kinase (tlk)* in *Drosophila* initiated a new type of cell death. Furthermore, the TLK-induced cell death is likely to be independent of the canonical caspase pathway and other known caspase-independent pathways. Genetically, *atg2* RNAi could rescue the TLK-induced cell death, and this function of *atg2* was likely distinct from its role in autophagy. In the developing retina, loss of *tlk* resulted in reduced PCD in the interommatidial cells (IOCs). Similarly, an increased number of IOCs was present in the *atg2* deletion mutant clones. However, double knockdown of *tlk* and *atg2* by RNAi did not have a synergistic effect. These results suggested that ATG2 may function downstream of TLK. In addition to a role in development, *tlk* and *atg2* RNAi could rescue calcium overload-induced cell death. Together, our results suggest that TLK mediates a new type of cell death pathway that occurs in both development and calcium cytotoxicity.

Cell Death and Differentiation (2016) 23, 146–157; doi:10.1038/cdd.2015.77; published online 19 June 2015

Cell death subtypes can be classified into uncontrollable accidental cell death and regulated cell death (RCD). As a further subtype of RCD, the cell death that occurs in development is referred as programmed cell death (PCD).¹ Although caspase-dependent apoptosis has crucial roles in development, other type(s) of PCD may exist.² The *Drosophila* eye is an elegant model system with which to study PCD in development;^{3,4} the patterning of the *Drosophila* eye is highly stereotypic and well characterized. The development of the fly retina begins in the eye disc of the third instar larvae, where the formation of ommatidium initiates from the differentiation of eight photoreceptor cells (R cells) followed by the recruitment of four cone cells. At the pupal stage, two primary pigment cells are recruited to surround the cone cells. Then, the interommatidial cells (IOCs) are chosen from a pool of undifferentiated cells and further refined into a highly stereotypical hexagonal lattice.³ Each hexagonal lattice contains 12 IOCs, including six secondary pigment cells at the edges, three tertiary pigment cells and three bristle cells at the vertices.^{5,6} The undetermined IOCs are then removed by apoptosis.⁷

It has been shown that intercellular communication has an essential role in regulating IOC apoptosis.⁸ The cone cells and primary pigment cells release survival ligands, such as Spitz, to promote the survival of IOCs, whereas IOCs release Delta to promote the death of their neighbors by activating the Notch pathway.^{2,8} Excessive IOCs are not the only cell type to be eliminated; the perimeter ommatidia are also trimmed during

development. This process is mediated by the secretion of a glycoprotein, Wingless, which promotes its own expression in the periphery of the eye and activates the caspase-dependent apoptosis pathway.⁶ The entire cell population of the perimeter ommatidia is eliminated, including the photoreceptor cells, cone cells, primary pigment cells and IOCs.⁶

Apoptosis is an important variant of PCD and is executed by caspases.¹ In *Drosophila*, these caspases include Dronc, a main initiator caspase, and DrICE and Dcp-1, which are activated by Dronc.⁹ The activities of caspases can be inhibited by two endogenous proteins, DIAP1 and DIAP2.¹⁰ Similarly, p35, a viral protein, is capable of inhibiting the caspase activity of DrICE and Dcp-1, resulting in the survival of extra IOCs and primary pigment cells in the adult fly retina.^{4,9} In *Drosophila*, the RHG (Reaper, Hid and Grim) family members are antagonists of DIAPs.¹⁰ In cells that are doomed to die, transcription of the RHG genes is increased.^{11–13} In addition, *Drosophila* p53 can promote apoptosis by acting together with the JNK signaling pathway to regulate the RHG proapoptotic machinery.^{14,15} Although deletion of the RHG genes blocks the majority of apoptosis, other PCD pathways likely exist during *Drosophila* eye development.²

In addition to apoptosis, other cell death pathways exist, although their roles in eye development are unclear. Ectopic expression of *eiger* (the fly homolog of mammalian TNF- α) induces cell death in the fly eyes. This type of apoptosis can be weakly inhibited by p35, but is strongly suppressed by the loss

¹State Key Laboratory of Membrane Biology, School of Life Sciences, Peking University, Beijing, China and ²Center of Stroke, Beijing Institute for Brain Disorder and Xuanwu Hospital, Capital Medical University, Beijing, China

*Corresponding author: Y Li or L Liu, Center of Stroke, Beijing Institute for Brain Disorder and Xuanwu Hospital, Capital Medical University, Beijing 100069, China. Tel: +86 10 62759691; Fax: +86 10 62756100; E-mail: yuhongli@pku.edu.cn or leiliu@ccmu.edu.cn

³These authors contributed equally to this work.

Abbreviations: TLK, tousled-like kinase; PCD, programmed cell death; IOCs, interommatidial cells; RCD, regulated cell death; RHG, reaper, hid and grim; AIF, apoptosis-inducing factor; JNK, Jun N-terminal kinase; MF, morphogenetic furrow; AO, acridine orange; APF, after pupa formation; pH3, anti-phospho-histone H3; TUNEL, TdT-mediated dUTP nick end labeling; PI, propidium iodide

Received 30.6.14; revised 21.4.15; accepted 05.5.15; Edited by E Baehrecke; published online 19.6.15

of JNK (Jun N-terminal kinase also called BSK) signaling, indicating that the Eiger/JNK-induced RCD is caspase-independent.^{16,17} Moreover, *Drosophila* AIF (apoptosis-inducing factor)-mediated cell death is also independent of the canonical caspase pathway.¹⁸ Autophagic cell death has been described to participate in *Drosophila* embryogenesis and is involved in the removal of the salivary gland and midgut tissues during *Drosophila* metamorphosis.^{19–21}

Beyond development, cell death has important roles in human diseases.²² For example, calcium overload is a pivotal stressor that induces cell death in many human diseases, such as stroke, traumatic brain injury, epilepsy, Alzheimer's disease and glaucoma.^{23–25} However, much remains to be learned regarding calcium-induced cell death pathways.²⁶

Here, we reported the discovery of a new type of TLK-induced PCD in *Drosophila* and delineated the function of TLK in both eye development and calcium-induced cell death.

Results

Overexpression of *tlk* induced cell death in *Drosophila* eyes. The adult *Drosophila* can survive without eyes.²⁷ Therefore, a genetic screen with the eye-specific promoter

*GMR-Gal4*²⁸ and various *UAS* lines is an elegant approach to discover the function of genes that cause *Drosophila* lethality. Here, the *Gal4/UAS* binary expression system is simplified as '>' throughout the text. We observed that the overexpression of *tlk* in the fly eyes (*GMR>tlk*) resulted in loss of pigmentation (Figure 1A a and b). To visualize each ommatidium, a membrane-tagged GFP transgene was added to the *GMR>tlk* flies (*GMR>mCD8-GFP/tlk*). This fly also showed the defective eye phenotype (Figure 1A c). To further characterize the cell death, we cross-sectioned the eyes to visualize the internal structure. In the *GMR-Gal4* control flies, each ommatidium displays a hexagonal profile with 7 R cells and accessory pigment cells²⁹ (Figure 1A d). In contrast, almost no intact ommatidia were visible and massive vacuoles were present throughout the eyes of *GMR>tlk* and *GMR>mCD8-GFP/tlk* flies (Figure 1A e and f). The defects could be suppressed by two independent *tlk RNAi* lines (Figure 1B). These RNAi lines target different regions of the *tlk* transcript and are designed to avoid off-target effect.³⁰ The quantitative RT-PCR verified that these RNAi lines indeed reduced the *tlk* transcripts (Supplementary Figure S1).

The absence of ommatidia in *GMR>tlk* may result from a failure to differentiate or enhanced cell death. To distinguish




Figure 1 Ectopic expression of *tlk* induced cell death in *Drosophila* eyes. (A) Overexpression of *tlk* induces eye defect. a, b and c are the images of the light microscope. d, e and f shows the toluidine blue-stained semi-thin sections of the eyes. The genotypes are shown on top of each micrograph. (B) The effect of two independent *tlk*^{RNAi} lines on the *GMR>tlk* flies. (C) Immunostaining of the eye disc from the 3rd instar larvae. The mCD8-GFP is a membrane-targeted GFP, and ELAV is a neuron marker. The enlarged images show that a cell cluster in b contains less R cells than a




Figure 2 Loss of *tlk* affects the developmental eye patterning. **(A)** Immunostaining of the 40 h APF retina with anti-Armadillo (red) or GFP (green). (a) The wild-type fly, *w¹¹¹⁸*. A hexagon is superimposed on the image to indicate the target area, which exemplifies the scoring method. Twenty-one cells lie within a target area of the wild-type fly, including 18 intact cells and six corner (labeled as *) counting as 3 cells. A representative cone cell, bristle cell, primary (1°), secondary (2°) and tertiary (3°) pigment cell are marked on the micrograph. (b) The *tlk* mutant clones are visualized by the absence of GFP. The wild-type clones and homozygous mutant clones are labeled with +/+ and -/-, respectively. (b') An enlarged view of the dashed white box in b and the extra IOCs are pointed by white arrows. **(B)** Quantification of the number of IOCs in **A**. In all figures, at least 30 hexagonal target areas were scored from three to six flies. **(C)** Effect of the *tlk* mutant (*tlk^{(1)G0054}*) on the development of the cone cells. An enlarged view is shown on the right panel. The wild-type clones always have four cone cells, whereas ~20% mutant clones showed either increased (with five cone cells) or decreased (with three cone cells) number of cone cells, as labeled by the asterisks. **(D)** Effect of *tlk* RNAi on the eye development. a and b as a control, the *white^{RNAi}* could effectively reduce the pigmentation of the eye compared with its promoter line (*GMR>Gal4*). (c and d) The scan electron microscope (S.E.M.) images of the eyes. (e and f) The anti-Armadillo staining. The enlarged views are shown on the right panels (e' and f'), and the extra IOCs are pointed by yellow arrows. **(E)** Quantification of the number of IOCs in **D**. **(F)** Effect of *tlk* RNAi on the development of perimeter ommatidia. (a and b) The images of S.E.M. (c and d) Immunostaining of the 40 h APF retina with anti-Armadillo (red). The arrows point to the extra small perimeter ommatidia. **(G)** Effect of *tlk* RNAi on cell proliferation. Immunostaining of the eye disc from the 3rd instar larvae with anti-phospho-pH3 (pH3). The arrow heads point to the morphogenic furrow. The number of pH3-positive cells in the posterior region of the eye disc is counted. The statistic results from five eye discs are shown under each condition. **(H)** AO staining of retina. Twenty areas in each eye disc were scored, and three to five eye discs were examined. The statistic results from five eye discs are shown under each condition. **(I)** Genetic interaction between *tlk* and *DIAP1*. **(J)** Quantification of IOCs




Figure 2 (Continued)

these possibilities, we examined eye development in the 3rd instar larvae. At this stage, the morphogenetic furrow (MF) marks the forward edge of differentiation, and the cell clusters at different stages of differentiation are aligned posterior to the MF, with the more mature cell clusters located at a further posterior region.²⁹ Some cells in the posterior region undergo a secondary division wave. Eventually, clusters of five R cells are formed, followed by recruitment of more R cells and other cell types at a later developmental stage.³¹ Consistent with the expression of the *GMR* promoter in the posterior region,²⁸ R cells were present in the eye disc of the *GMR>tlk* flies (R cells were labeled with ELAV) (Figure 1C a and b), suggesting that differentiation was not significantly affected. However, the number of R cells in some clusters was reduced (Figure 1C b).

Because almost all R cells were missing in the adult stage of *GMR>tlk* (Figure 1A e and f), the R cells likely started to die at the late larval stage. Therefore, overexpression of *tlk* promotes cell death instead of affecting differentiation.

TLK regulated PCD during eye development. When raised at 25 °C, each ommatidium adopts a rigid hexagonal pattern in the wild-type flies at 40 h APF (after pupa formation), and anti-Armadillo staining can visualize each cell in the developing retina.³² To quantify the number of IOCs in the developing retina, a hexagonal target area is defined by connecting the center of six ommatidia surrounding a central ommatidium (Figure 2A a).³³

To test whether loss-of-function *tlk* may promote cell survival in the developing retina, we studied a *tlk* mutant. As a member of the conserved serine/threonine kinase family, TLK regulates cell cycle progression and chromatin assembly.³⁴ A P-element-mediated loss-of-function mutant, *tlk^{l(1)G0054}*, has been well characterized, and it induces DNA fragmentation, cell death and pupal lethality.³⁴ To determine the effect of this mutant on the *Drosophila* eye, we performed mosaic analysis. In the pupal retina of the mosaics, the wild-type clones were labeled with two copies of GFP, whereas the homozygous mutant *tlk (tlk^{l(1)G0054}/tlk^{l(1)G0054})* cells displayed no GFP labeling (Figure 2A b). Strikingly, the *tlk* homozygous mutant clones showed a significant increase in the number of IOCs (Figures 2A b' and 2B). In addition, ~20% of the ommatidia developed more or fewer cone cells (3 or 5 compared with 4 in the wild-type) in the *tlk* homozygous mutant clones (Figure 2C, the asterisks labeled ommatidia). This result indicated that the cone cell division may be affected in the *tlk* homozygous mutant. Because cone cells can affect the recruitment and survival of IOCs,^{2,8} the defects in the cone cells may complicate the interpretation of TLK function in the IOCs. To determine the function of TLK at a later developmental stage, we examined the *tlk* RNAi lines driven by *GMR-Gal4*. A genetically matched line, *white^{RNAi}*, was used as the control. The *white^{RNAi}* line was functional because it can efficiently reduce the pigmentation of *GMR-Gal4* (Figure 2D a and b). In the *GMR>tlk^{RNAi}* adult eyes, the shape of the ommatidia were irregular, indicating defects in the patterning of the eyes (Figure 2D c and d). To quantify the defect, we examined the pupal retina at 40 h APF. The result showed that extra IOCs (Figures 2D e, f and 2E) and perimeter ommatidia (Figure 2F) were present. Therefore, extra IOCs survived with the loss of *tlk*.

To determine whether the extra IOCs resulted from reduced cell death or increased proliferation, the eye discs were stained with anti-phospho-histone H3 (pH3), a marker of cell proliferation.³⁵ The result showed that the pH3 staining pattern was unaltered in the 3rd instar larval eye disc of the *GMR>tlk^{RNAi}* lines (Figure 2G), indicating that the extra number of IOCs and perimeter ommatidia were not due to increased cell proliferation. To further test whether PCD was reduced in the *GMR>tlk^{RNAi}* lines, we stained the pupal retina (at 35 h APF) with acridine orange (AO), a marker of cell death.³⁶ The result showed reduced AO staining in the retina of the *GMR>tlk^{RNAi}* flies, compared with the control retina (Figure 2H). Therefore, TLK mediated PCD during eye patterning.

The canonical caspase pathway is the main executioner of PCD in eye patterning.² To test the genetic interaction between TLK and the caspase pathway, we examined the effect of overexpression of DIAP1 on IOCs. DIAP1 blocks the executioner caspases, including DrICE and Dcp-1.⁹ The result showed that the tested fly (*GMR>DIAP1/tlk^{RNAi}*) eyes exhibited even more IOCs than either *GMR>tlk^{RNAi}* or *GMR-DIAP1* alone (Figure 2I and J), suggesting that TLK mediated a new type of PCD that is independent of the apoptotic pathway.

Characterization of TLK-induced cell death. To study TLK-induced cell death, we stained the eye discs of the 3rd

instar larvae with AO (Figure 3A a–c). *GMR-hid* was used as a positive control (Figure 3A c). Interestingly, the AO staining intensity was even stronger in the *GMR>tlk* flies than the *GMR>hid* flies, suggesting that ectopic expression of *tlk* can strongly induce cell death. Further, we examined DNA fragmentation by TdT-mediated dUTP nick end labeling (TUNEL) (Figure 3A d–f). The results demonstrated intensely labeled TUNEL signals (Figure 3A e). Because TUNEL can also label necrotic cells, in which the membrane integrity is disrupted,³⁷ we tested whether the membrane was intact by PI (propidium iodide) staining (Figure 3B a–c). The results showed that the PI signal was negative in the *GMR>tlk* flies (Figure 3B b). As a positive control, the necrosis model of *sev>GlutR1^{Lc}*³⁷ was PI positive (Figure 3B c). These results suggested that overexpression of TLK was sufficient to induce cell death.

Next, we asked whether the apoptotic pathway was involved in the cell death of the *GMR>tlk* flies. The results showed that inhibition of the caspase pathway by overexpression of *p35*, *DIAP1* and *DIAP2* had no effect on the eye defect in the *GMR>tlk* flies (Figure 4A). As positive controls, these lines suppressed apoptosis in the *GMR-hid* fly (Supplementary Figure S2). Moreover, we examined an *in vivo* sensor of caspase activation, Apoliner, which comprises two fluorophores: enhanced green fluorescent protein (eGFP) and monomeric red fluorescent protein (mRFP).³⁸ These fluorescent proteins are linked by a peptide sequence containing a caspase-sensitive cleavage site. Upon caspase activation, the cleavage of Apoliner allows the eGFP to translocate into the nucleus, whereas the mRFP remains on the cell membrane.³⁸ As a positive control, the mRFP was presented on the cell membrane in the posterior region of the eye disc in the *GMR-hid* flies (Figure 4B b). In contrast, the eye disc of *GMR>tlk* flies showed no enhanced presence of the mRFP on the cell membrane (Figure 4B c). In addition, we performed immunostaining using the cleaved Dcp-1 antibody (Figure 4B d–f). The results showed that there was no significant activation of the caspase activity in the *GMR>tlk* flies compared with the control flies (Figure 4B f). Together, these results suggest that the TLK-induced cell death was independent of caspase function.

JNK is known as a mediator of caspase-independent cell death in *Drosophila*.^{16,18} We observed that inhibition of the JNK pathway by overexpression of a dominant-negative *Bsk* (*Bsk^{DN}*) did not suppress the eye defect of *GMR>tlk* flies (Figure 4C). Consistently, the JNK pathway was not activated in the eye disc of *GMR>tlk* flies as determined by *puc-lacZ*, an *in vivo* reporter of JNK activation³⁹ (Figure 4D a–b). In contrast, the positive control of *GMR>eiger¹⁷* showed robust induction of the JNK signal (Figure 4D c). AIF is a mediator of caspase-independent cell death in *Drosophila*.¹⁸ We found that loss-of-function of *AIF* had no effect on the cell death in *GMR>tlk* flies; a similar effect was observed with the *p53* mutant or overexpression of ROS chelating enzymes (catalase and GTPx-1) (Figure 4E).

TLK has an important role in chromatin assembly, including chromosome segregation, replication, transcription and DNA repair.⁴⁰ In *Drosophila*, TLK phosphorylates ASF1 and causes nuclear division arrest and pupal lethality.³⁴ Our result showed that *asf1^{RNAi}* had no effect on the TLK-induced cell death




Figure 3 Cell death induced by ectopic expression of *tlk* in *Drosophila* eye. (A) AO and TUNEL staining of the eye disc from the 3rd instar larvae. *GMR-hid* is shown as a positive control. (a–c) The AO staining. (a'–c') The merged image of bright field and AO staining. (d–f) TUNEL staining. The integrative density (Intden) of the posterior region in each eye disc was obtained by ImageJ. Five eye discs were examined for each genotype. (B) PI staining of the eye disc. (a–c) PI staining. (a'–c') Merged image of bright field with PI staining. The *sev>GlutR1^{Lc}* is shown as a positive control




Figure 4 Characterization of the TLK-induced cell death. **(A)** Effect of inhibition of caspase pathway on the *GMR > tlk*. **(B)** Caspase activity in the larval eye disc. (a, b and c) Alteration of apoliner fluorescence. (d, e and f) Immunostaining with anti-cleaved Dcp-1. The number of positive dots in the posterior region of the eye disc is counted. *GMR-hid* is shown as a positive control. The statistic results from five eye discs are shown under each condition. d', e' and f' are merged images of DAPI and cleaved Dcp-1 staining. **(C)** Effect of inhibition JNK signaling pathway on *GMR > tlk*. **(D)** JNK activity in the larval eye disc. *puc-lacZ* is an *in vivo* reporter of JNK activation that can be detected by the X-gal staining. The *GMR > eiger* is shown as a positive control. **(E)** Effect of several genes related to cell death or chromatin assembly on *GMR > tlk* eye defects

(Figure 4E), suggesting that the function of TLK on cell death was likely different from its role in chromatin organization. Together, these results suggested that TLK may induce a new cell death pathway.

ATG2 functioned downstream of the TLK-mediated PCD.

To understand the molecular mechanism behind TLK-induced cell death, we screened ~1000 TRiP RNAi lines using the *GMR> tdk* flies. These RNAi lines are inserted into a defined genomic locus, thus eliminating the variation of positional effects compared with random P-element insertions.³⁰ We identified nine suppressor and enhancer lines (Table 1). Most of the suppressors are components of the mediator family, which affected the transcription of *tdk*. In contrast, *atg2* RNAi did not affect the transcription of *tdk* and was further studied. The *atg2^{RNAi}* showed a strong rescue effect on the *GMR> tdk* flies (Figure 5A a and b). Moreover, *atg2^{RNAi}* reduced AO and TUNEL staining in the *tdk*-overexpressing eye discs (Figure 5B), suggesting that cell death was suppressed. Interestingly, the number of IOCs was also increased at 40h APF in the retina of the *GMR> atg2^{RNAi}* flies (Figure 5C a). The efficiency of the *atg2^{RNAi}* was confirmed (Supplementary Figure S3). Using the CRISPR-associated single-guide RNA system (Cas9/sgRNA),⁴¹ we generated a large deletion (2164 bp) in the exon region of the *atg2* gene. The homozygous deletion was pupal lethal. Therefore, we generated mosaics in the fly eyes. Similar to the RNAi effect, the clones of the homozygous deletion displayed a significant increase in the number of IOCs (Figure 5C b). In addition, knocking down both *tdk* and *atg2* by RNAi resulted in a similar number of IOCs as the *GMR> tdk^{RNAi}* alone (Figure 5C c), suggesting that ATG2 acts in the same pathway with TLK. Moreover, *atg2^{RNAi}* had a synergistic effect with *GMR-DIAP1*, similar to *tdk^{RNAi}* (Figure 5C d). The statistical results are presented (Figure 5D). Because ATG2 is known to be involved in autophagy,¹⁹ we asked whether autophagy was activated. There was no LysoTracker staining in the larval eye disc of the *GMR> tdk* flies (Supplementary Figure S4 a and b). As a positive control, the eye disc of *sev> GluR1^{Lc}* showed intense LysoTracker staining (Supplementary Figure S4 c), as reported previously.³⁷ We also tested several autophagy-related genes by RNAi, including *atg1*, *atg4*, *atg8a*, *atg9* or *atg16*. The result showed that none of the genes rescued the eye defect of the *GMR> tdk* flies (Supplementary Figure S5). Together, these results suggested that ATG2 might function downstream of TLK.

To explore the mechanism of TLK-induced cell death, we stained the eye disc of *GMR> tdk-flag* with a nuclear membrane marker (NPC), anti-Flag and DAPI. In the *GMR> tdk-flag* cells, TLK-Flag was stained in the nucleus (Supplementary Figure S6a and b), as reported.³⁴ Meanwhile, the nuclear membranes were damaged or had disappeared in some of the *tdk*-expressing cells (Supplementary Figure S6b). Because TLK is a nuclear protein, it was likely that the cell death caused dysfunction of the nucleus.

TLK mediated mild calcium overload-induced cell death.

Our data suggested that TLK may induce a new type of PCD in *Drosophila* eye development. Does TLK also function in pathological conditions? Previously, we were interested in calcium overload-induced cell death and generated a transgenic *Drosophila* line to express *GluR1^{Lc}* (rat glutamate receptor 1 lurcher mutant),³⁷ a constitutively open cation channel *in vitro*.⁴² Another transgenic line that we generated is *hs-GluR1^{Lc}*, in which *GluR1^{Lc}* is driven directly by a heat shock (*hs*) promoter. These flies have defects in the eyes when raised at 25 °C (Figure 6A a and d). To further assess cell death, the fly eyes were sectioned (Figure 6A b, c, e and f). We observed that some ommatidia disappeared and were accompanied by the presence of large vacuoles throughout the eyes in the 1-day-old *hs-GluR1^{Lc}* flies (Figure 6A e), and no cellular structure was observed in the 10-day-old *hs-GluR1^{Lc}* flies (Figure 6A f). These results suggested that mild calcium overload induced cell death in the *Drosophila* eyes. To quantify calcium overload, we measured the intracellular calcium levels in the 3rd instar larval eye disc using Fura-2. Indeed, the calcium levels were increased by ~20% in the *hs-GluR1^{Lc}* flies compared with the control flies (Figure 6B).

Next, we investigated the cell death pathways involved in the *hs-GluR1^{Lc}* flies. The results showed that *tdk^{RNAi}* strongly rescued the eye defect of the *hs-GluR1^{Lc}* flies (Figure 6C b and c). For the other pathways, suppression of the caspase-mediated pathway had a partial rescue effect, whereas the other pathways, including AIF, JNK and autophagy, showed no effect (Supplementary Figure S7). In addition, *atg2^{RNAi}* showed a partial rescue of the *hs-GluR1^{Lc}* eyes (Figure 6C d and h). Although TLK may regulate transcription,⁴³ the transcription of *GluR1^{Lc}* was not affected by the *tdk^{RNAi}* lines (Figure 6D). Therefore, the TLK-mediated cell death was active in the mild calcium overload condition in the fly eyes.

Table 1 Genetic modifiers of cell death in *GMR> tdk*

CG number	Gene symbol	Gene function	Effect
CG13867	MED8	RNA polymerase II transcription cofactor activity	Rescue
CG9936	skd/MED13	RNA polymerase II transcription cofactor activity	Rescue
CG5546	MED19	RNA polymerase II transcription cofactor activity	Rescue
CG17397	MED21	RNA polymerase II transcription cofactor activity	Rescue
CG5121	MED28	RNA polymerase II transcription cofactor activity	Rescue
CG17183	MED30	RNA polymerase II transcription cofactor activity	Rescue
CG1241	Atg2	Autophagy	Rescue
CG43665	eIF-2γ	GTPase activity	Enhance
CG6779	RpS3	Structural constituent of ribosome	Enhance




Figure 6 The *hs-GluR1^{Lc}* model and the effect of *tlk^{RNAi}*. (A) The pigmentation defect in the eye of *hs-GluR1^{Lc}*. Light images and toluidine blue-stained semi-thin sections are shown. (B) Intracellular calcium level in the larval eye disc measured by Fura-2. Trial $n = 3$. Six larvae were examined for each trial. The open bar indicates the control, whereas the black bar indicates statistically significant difference. * represents t -test, $P < 0.05$. (C) Effect of *tlk^{RNAi}* and *atg2^{RNAi}* on the *hs-GluR1^{Lc}* eye defect. The upper panels are the light images, and the lower panel show the toluidine blue-stained semi-thin sections. (D) Effect of *tlk^{RNAi}* on *GluR1^{Lc}* transcription in the *hs-GluR1^{Lc}* flies. Trial $n = 6$

In theory, *tlk^{RNAi}* might promote a reduction of PCD by enhancing survival signals, such as EGF and Ras, or suppressing death signals, such as Notch.^{2,8} Because these pathways partially converge onto the apoptotic pathway to regulate eye patterning,^{44–46} the independence of *tlk^{RNAi}* with respect to DIAP1 function suggests that this possibility is less likely. Moreover, we find that TLK-induced cell death cannot be rescued by most of the known cell death pathways, including caspase, autophagy, AIF, JNK, ROS and p53. Therefore, we propose that TLK regulates a new type of PCD.

It is not clear how TLK causes cell death. Interestingly, the nuclear membrane seems severely disrupted in TLK-overexpressing cells. Because TLK is located in the nucleus, it may phosphorylate nuclear substrates and promote the disruption of the nuclear membrane. On the other hand, ATG2 is known to function in the formation of the autophagosome membrane,⁴⁷ but may also function in the maintenance of the nuclear membrane. This hypothesis requires further investigation.

In addition to regulating developmental PCD, TLK may also be involved in disease. Our data suggest that the cell death

observed in the *hs-GluR1^{Lc}* fly eye is partially regulated by the apoptotic pathway. However, this calcium-induced cell death was strongly suppressed by the *tlk^{RNAi}* lines, indicating that TLK may function in calcium overload-induced cell death. Mild calcium overload has been implicated in human diseases, such as Alzheimer's disease and glaucoma.^{25,48} The potential function of TLK in these diseases requires further study. In addition, how TLK may be activated *in vivo* remains unclear.

In conclusion, we have identified a potential novel cell death pathway that functions in both eye development and calcium cytotoxicity. Strikingly, TLK is both necessary and sufficient for this variant of cell death.

Materials and Methods

***Drosophila* stocks and maintenance.** *Drosophila* stocks were maintained at 25 °C on standard fly food. The following fly strains were kindly provided by different labs: *puc^{ES9}-lacZ* (Dr. Tian Xu); *AIF^{KO}* (Dr. Josef Penninger); *UAS-IAP2* (Dr. Pascal Meier); *UAS-catalase* and *UAS-GTPx-I* (Dr. Utpal Banerjee); *UAS-eiger* (Dr. Lei Xue); *GMR-hid, GMR-Gal4* (Dr. Andreas Bergmann); *p53/Tb* (Dr. John Abrams) TRIP RNAi stocks were obtained from the Tsinghua *Drosophila* stock center: *tlk^{RNAi}* (RNAi-1: THU0178, RNAi-2: THU1326); *Atg2^{RNAi}* (THU3698). All the other lines were obtained from the Bloomington Stock Center, including *tlk^{(1)G0054}* (BL11593); *UAS-Apoliner* (BL32122). For the RNAi lines, *y¹ v¹*; *P{CaryP}attP2* (BL36303) and *white^{RNAi}* (BL33623) were used as the genetic background matched controls.

Generation of a transgenic fly line. The *GluR1^{Lc}* cDNA was digested from pCAGGS-mGluR1^{Lc} construct (a gift from Dr. Michisuke Yuzaki) and subcloned into the *pCaSpeR-hs/act* vector. The *hs-GluR1^{Lc}* transgenic fly was generated in a *w¹¹¹⁸* background using the P-element-mediated transformation.

Generation of *atg2* deletion. The *atg2* deletion mutant was generated using the Cas9/sgRNA system. To generate gene deletion, two sg-RNAs were designed to target the *atg2* locus. The sg-RNAs sequences that flank a 2164 bp exon region of the *atg2* gene are the following: g1, 5'-GGATGGCTACCACATTCGACCGG-3'; g2, 5'-GGATGTCTCCGATGACCGAGGG-3'. The deletion of 2164 bp was confirmed by PCR sequencing.

Histology of adult eyes. The adult fly heads were removed from the body and fixed in the fixative containing 4% paraformaldehyde and 2.5% glutaraldehyde in 0.1 M PB, post-fixed in 1% osmium tetroxide, dehydrated in an ethanol series followed by exchange with propylene oxide. Then, the heads were infiltrated in a mixture of propylene oxide and Spurr's medium, and finally imbedded in Spurr's medium as previously described.⁴⁹ The eyes were semi-thin sectioned to 1.5 μm using an ultramicrotome (UC7; Leica, Tokyo, Japan) and stained with toluidine blue.

Histology of larval tissue. AO, immunostaining, PI, lacZ and lysotracker staining were performed as described.^{36,37,50,51} For TUNEL staining, the procedures were followed by the manual of the manufacture (*In Situ* Cell Death TMR, Product No. 12156792910, Roche, Indianapolis, IN, USA), except an additional treatment of protease K before the enzymatic reaction. The quantification analysis was performed using ImageJ.

Antibodies used include anti-ELAV (9F8A9, DSHB, Iowa, IA, USA; 1:1000), anti-Armadillo (N27A1, DSHB, 1:2), anti-phospho-Histone-H3 (06-570, Millipore, Billerica, MA, USA; 1:1000), and anti-GFP (A-11122, Thermo Fisher, Rockford, IL, USA; 1:1000), anti-cleaved *Drosophila* Dcp-1 (Asp216) (#9578, CST, Danvers, MA, USA; 1:100), anti-Flag (#2368, CST; 1:1000), anti-Nuclear Protein Complex (NPC) (ab24609, Abcam, Cambridge, MA, USA; 1:1000).

Fura-2 measurement. The eye discs of the 3rd instar larvae were dissected and incubated with 5 μM Fura-2 AM (F-1221, Invitrogen Molecular Probe, Eugene, OR, USA) in Schneider's medium at 25 °C for 30 min in the dark. With the excitation wavelengths at 340 and 380 nm, we recorded two emission intensity values at 510 nm, respectively. The ratio of emission intensity at 510 nm excited by 340 or 380 nm was calculated as the relative calcium concentration.

qRT-PCR method. Total RNA was extracted by Trizol reagent (Invitrogen, Eugene, OR, USA) followed by DNase I treatment based on the manufacturer's standard protocol, then the purity and integrity of total RNA was determined by 1% agarose gel electrophoresis. The concentration of total RNA was measured by Nanodrop. Two microgram mRNA was reverse transcribed into cDNA library by oligo-dT primer using RevertAid First Strand cDNA Synthesis Kit (Thermo scientific) based on the manufacturer's standard protocol. The final volume of q-PCR reaction was 25 μl using Platinum SYBRGreen qPCR SuperMix-UDG Kit (Invitrogen) containing 1 μl diluted cDNA sample (1:3). The q-PCR was performed in triplicate using 7500 real time PCR system (Thermo Fisher). The quantification of target gene was conducted by ΔΔCt method.⁵² Primers used are as following:

GAPDH F: 5'-CGCAGCGCCATTCTCTCTA-3';

R: 5'-GACTGCCGCTTTTTCCTTTTC-3'

tlk F: 5'-GGGCGGGAACCTACTGGA-3';

R: 5'-TTTTCGGGGATTTTTGC-3'.

atg2 F: 5'-CCAACGCCTATACCATAGTGAGAGA-3'

R: 5'-TCTGGTCGTCTCCGTGAT-3'

Scanning electron microscopy. Adult flies were anaesthetized to death by chloroform, mounted on stages and then observed using a scanning electron microscope TM-1000 (HITACHI, Tokyo, Japan).

IOP cell counts. At least 30 hexagonal target areas were scored from three to six different flies for each experiment as reported by others.³²

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. This work is supported by the Chinese Ministry of Science and Technology (2013CB530700) and National Natural Science Foundation of China (NSFC31171324) to L. L.

- Galluzzi L, Bravo-San Pedro JM, Vitale I, Aaronson SA, Abrams JM, Adam D et al. Essential versus accessory aspects of cell death: recommendations of the NCCD 2015. *Cell Death Differ* 2015; **22**: 58–73.
- Brachmann CB, Cagan RL. Patterning the fly eye: the role of apoptosis. *Trends Genet* 2003; **19**: 91–96.
- Cagan RL, Ready DF. The emergence of order in the *Drosophila* pupal retina. *Dev Biol* 1989; **136**: 346–362.
- Hay BA, Wolff T, Rubin GM. Expression of baculovirus P35 prevents cell death in *Drosophila*. *Development* 1994; **120**: 2121–2129.
- Copeland JM, Bosdet I, Freeman JD, Guo M, Gorski SM, Hay BA. echinus, required for interommatidial cell sorting and cell death in the *Drosophila* pupal retina, encodes a protein with homology to ubiquitin-specific proteases. *BMC Dev Biol* 2007; **7**: 82.
- Lin HV, Rogulja A, Cadigan KM. Wingless eliminates ommatidia from the edge of the developing eye through activation of apoptosis. *Development* 2004; **131**: 2409–2418.
- Cordero J, Jassim O, Bao S, Cagan R. A role for wingless in an early pupal cell death event that contributes to patterning the *Drosophila* eye. *Mech Dev* 2004; **121**: 1523–1530.
- Miller DT, Cagan RL. Local induction of patterning and programmed cell death in the developing *Drosophila* retina. *Development* 1998; **125**: 2327–2335.
- Hay BA, Guo M. Caspase-dependent cell death in *Drosophila*. *Annu Rev Cell Dev Biol* 2006; **22**: 623–650.
- Steller H. Regulation of apoptosis in *Drosophila*. *Cell Death Differ* 2008; **15**: 1132–1138.
- White K, Grether ME, Abrams JM, Young L, Farrell K, Steller H. Genetic control of programmed cell death in *Drosophila*. *Science* 1994; **264**: 677–683.
- Chen P, Nordstrom W, Gish B, Abrams JM. grim, a novel cell death gene in *Drosophila*. *Genes Dev* 1996; **10**: 1773–1782.
- Grether ME, Abrams JM, Agapite J, White K, Steller H. The head involution defective gene of *Drosophila melanogaster* functions in programmed cell death. *Genes Dev* 1995; **9**: 1694–1708.
- Shevkov E, Morata G. A dp53/JNK-dependant feedback amplification loop is essential for the apoptotic response to stress in *Drosophila*. *Cell Death Differ* 2012; **19**: 451–460.
- Brodsky MH, Nordstrom W, Tsang G, Kwan E, Rubin GM, Abrams JM. *Drosophila* p53 binds a damage response element at the reaper locus. *Cell* 2000; **101**: 103–113.
- Igaki T, Kanda H, Yamamoto-Goto Y, Kanuka H, Kurana E, Aigaki T et al. Eiger, a TNF superfamily ligand that triggers the *Drosophila* JNK pathway. *EMBO J* 2002; **21**: 3009–3018.
- Moreno E, Yan M, Basler K. Evolution of TNF signaling mechanisms: JNK-dependent apoptosis triggered by Eiger, the *Drosophila* homolog of the TNF superfamily. *Curr Biol* 2002; **12**: 1263–1268.

18. Joza N, Galindo K, Pospisilik JA, Benit P, Rangachari M, Kanitz EE *et al*. The molecular archaeology of a mitochondrial death effector: AIF in *Drosophila*. *Cell Death Differ* 2008; **15**: 1009–1018.
19. Berry DL, Baehrecke EH. Growth arrest and autophagy are required for salivary gland cell degradation in *Drosophila*. *Cell* 2007; **131**: 1137–1148.
20. Denton D, Shrivage B, Simin R, Mills K, Berry DL, Baehrecke EH *et al*. Autophagy, not apoptosis, is essential for midgut cell death in *Drosophila*. *Curr Biol* 2009; **19**: 1741–1746.
21. Denton D, Nicolson S, Kumar S. Cell death by autophagy: facts and apparent artefacts. *Cell Death Differ* 2012; **19**: 87–95.
22. Fuchs Y, Steller H. Programmed cell death in animal development and disease. *Cell* 2011; **147**: 742–758.
23. Syntichaki P, Tavernarakis N. The biochemistry of neuronal necrosis: rogue biology? *Nat Rev Neurosci* 2003; **4**: 672–684.
24. Yuan J, Lipinski M, Degterev A. Diversity in the mechanisms of neuronal cell death. *Neuron* 2003; **40**: 401–413.
25. Crish SD, Calkins DJ. Neurodegeneration in glaucoma: progression and calcium-dependent intracellular mechanisms. *Neuroscience* 2011; **176**: 1–11.
26. Zhivotovsky B, Orrenius S. Calcium and cell death mechanisms: a perspective from the cell death community. *Cell Calcium* 2011; **50**: 211–221.
27. Jenny A. Preparation of adult *Drosophila* eyes for thin sectioning and microscopic analysis. *J Vis Exp* 2011; **54**: 2959.
28. Liu H, Ma C, Moses K. Identification and functional characterization of conserved promoter elements from glass: a retinal development gene of *Drosophila*. *Mech Dev* 1996; **56**: 73–82.
29. Charlton-Perkins M, Cook TA. Building a fly eye: terminal differentiation events of the retina, corneal lens, and pigmented epithelia. *Curr Top Dev Biol* 2010; **93**: 129–173.
30. Ni JQ, Zhou R, Czech B, Liu LP, Holderbaum L, Yang-Zhou D *et al*. A genome-scale shRNA resource for transgenic RNAi in *Drosophila*. *Nat Methods* 2011; **8**: 405–407.
31. de Nooij JC, Hariharan IK. Uncoupling cell fate determination from patterned cell division in the *Drosophila* eye. *Science* 1995; **270**: 983–985.
32. Mendes CS, Arama E, Brown S, Scherr H, Srivastava M, Bergmann A *et al*. Cytochrome c-d regulates developmental apoptosis in the *Drosophila* retina. *EMBO Rep* 2006; **7**: 933–939.
33. Johnson RI, Cagan RL. A quantitative method to analyze *Drosophila* pupal eye patterning. *PLoS One* 2009; **4**: e7008.
34. Carrera P, Moshkin YM, Gronke S, Sillje HH, Nigg EA, Jackle H *et al*. Tousled-like kinase functions with the chromatin assembly pathway regulating nuclear divisions. *Genes Dev* 2003; **17**: 2578–2590.
35. Hendzel MJ, Wei Y, Mancini MA, Van Hooser A, Ranalli T, Brinkley BR *et al*. Mitosis-specific phosphorylation of histone H3 initiates primarily within pericentromeric heterochromatin during G2 and spreads in an ordered fashion coincident with mitotic chromosome condensation. *Chromosoma* 1997; **106**: 348–360.
36. Arama E, Steller H. Detection of apoptosis by terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling and acridine orange in *Drosophila* embryos and adult male gonads. *Nat Protoc* 2006; **1**: 1725–1731.
37. Yang Y, Hou L, Li Y, Ni J, Liu L. Neuronal necrosis and spreading death in a *Drosophila* genetic model. *Cell Death Dis* 2013; **4**: e723.
38. Bardet PL, Kolahgar G, Mynett A, Miguel-Aliaga I, Briscoe J, Meier P *et al*. A fluorescent reporter of caspase activity for live imaging. *Proc Natl Acad Sci USA* 2008; **105**: 13901–13905.
39. Adachi-Yamada T, Fujimura-Kamada K, Nishida Y, Matsumoto K. Distortion of proximodistal information causes JNK-dependent apoptosis in *Drosophila* wing. *Nature* 1999; **400**: 166–169.
40. De Benedetti A. The tousled-like kinases as guardians of genome integrity. *ISRN Mol Biol* 2012; **2012**: 627596.
41. Ren X, Sun J, Housden BE, Hu Y, Roesel C, Lin S *et al*. Optimized gene editing technology for *Drosophila melanogaster* using germ line-specific Cas9. *Proc Natl Acad Sci USA* 2013; **110**: 19012–19017.
42. Kohda K, Wang Y, Yuzaki M. Mutation of a glutamate receptor motif reveals its role in gating and delta2 receptor channel properties. *Nat Neurosci* 2000; **3**: 315–322.
43. Han Z, Saam JR, Adams HP, Mango SE, Schumacher JM. The *C. elegans* Tousled-like kinase (TLK-1) has an essential role in transcription. *Curr Biol* 2003; **13**: 1921–1929.
44. Yu SY, Yoo SJ, Yang L, Zapata C, Srinivasan A, Hay BA *et al*. A pathway of signals regulating effector and initiator caspases in the developing *Drosophila* eye. *Development* 2002; **129**: 3269–3278.
45. Kurada P, White K. Ras promotes cell survival in *Drosophila* by downregulating hid expression. *Cell* 1998; **95**: 319–329.
46. Bergmann A, Agapite J, McCall K, Steller H. The *Drosophila* gene hid is a direct molecular target of Ras-dependent survival signaling. *Cell* 1998; **95**: 331–341.
47. Velikkakath AK, Nishimura T, Oita E, Ishihara N, Mizushima N. Mammalian Atg2 proteins are essential for autophagosome formation and important for regulation of size and distribution of lipid droplets. *Mol Biol Cell* 2012; **23**: 896–909.
48. Berridge MJ. Calcium hypothesis of Alzheimer's disease. *Pflugers Arch* 2010; **459**: 441–449.
49. Porter JA, Hicks JL, Williams DS, Montell C. Differential localizations of and requirements for the two *Drosophila* ninaC kinase/myosins in photoreceptor cells. *J Cell Biol* 1992; **116**: 683–693.
50. Sweeney ST, Hidalgo A, de Belle JS, Keshishian H. X-gal staining of the central nervous system in adult *Drosophila*. *Cold Spring Harb Protoc* 2012; **2012**: 239–241.
51. Wu JS, Luo L. A protocol for dissecting *Drosophila melanogaster* brains for live imaging or immunostaining. *Nat Protoc* 2006; **1**: 2110–2115.
52. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M *et al*. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 2009; **55**: 611–622.

Supplementary Information accompanies this paper on Cell Death and Differentiation website (<http://www.nature.com/cdd>)