

ARTICLE

Genomic screening identifies novel linkages and provides further evidence for a role of *MYH9* in nonsyndromic cleft lip and palate

Brett T Chiquet^{1,2}, Syed S Hashmi^{1,3}, Robin Henry¹, Amber Burt⁴, John B Mulliken⁵, Samuel Stal⁶, Molly Bray⁷, Susan H Blanton⁴ and Jacqueline T Hecht^{*1}

¹Department of Pediatrics, University of Texas Medical School at Houston, Houston, TX, USA; ²University of Texas Dental Branch at Houston, Houston, TX, USA; ³University of Texas School of Public Health, Houston, TX, USA; ⁴Miami Institute for Human Genetics, University of Miami Miller School of Medicine, Miami, FL, USA; ⁵Department of Plastic Surgery, Children's Hospital, Boston, MA, USA; ⁶Division of Plastic Surgery, Texas Children's Hospital, Houston, TX, USA; ⁷Department of Pediatrics, Baylor College of Medicine, Houston, TX, USA

Nonsyndromic cleft lip with or without cleft palate (NSCLP) is a common birth anomaly that requires prolonged multidisciplinary rehabilitation. Although variation in several genes has been identified as contributing to NSCLP, most of the genetic susceptibility loci have yet to be defined. To identify additional contributory genes, a high-throughput genomic scan was performed using the Illumina Linkage IVb Panel platform. We genotyped 6008 SNPs in nine non-Hispanic white NSCLP multiplex families and a single large African-American NSCLP multiplex family. Fourteen chromosomal regions were identified with $LOD > 1.5$, including six regions not previously reported. Analysis of the data from the African-American and non-Hispanic white families revealed two likely chromosomal regions: 8q21.3–24.12 and 22q12.2–12.3 with LOD scores of 2.98 and 2.66, respectively. On the basis of biological function, syndecan 2 (*SDC2*) and growth differentiation factor 6 (*GDF6*) in 8q21.3–24.12 and myosin heavy-chain 9, non-muscle (*MYH9*) in 22q12.2–12.3 were selected as candidate genes. Association analyses from these genes yielded marginally significant *P*-values for SNPs in *SDC2* and *GDF6* ($0.01 \leq P < 0.05$). Evidence for an altered transmission was found for four *MYH9* SNPs ($P < 0.01$). SNP rs1002246 exhibited altered transmission by all analytic methods. However, analysis of two SNP *MYH9* haplotypes did not identify a single high-risk haplotype. Our results confirm a previous report that 8q21.3–24.12 may harbor a clefting gene and identify 22q12.2–12.3 as a new candidate region that contains *MYH9*. Most importantly, we confirm the previous report of an association with *MYH9*.

European Journal of Human Genetics (2009) 17, 195–204; doi:10.1038/ejhg.2008.149; published online 20 August 2008

Keywords: cleft lip and palate; genome scan; *SDC2*; *GDF6*; *MYH9*

*Correspondence: Dr JT Hecht, Department of Pediatrics, University of Texas Houston Medical School, 6431 Fannin St MSB 3.136, PO Box 20708, Houston, TX 77225, USA.

Tel: 1 713 500 5764; Fax: 1 713 500 5689;

E-mail: jacqueline.t.hecht@uth.tmc.edu

Received 4 April 2008; revised 15 July 2008; accepted 16 July 2008; published online 20 August 2008

Introduction

Nonsyndromic cleft lip with or without cleft palate (NSCLP) is a common congenital anomaly with a prevalence of 1/500–1/1400 live births and varies with ethnicity.^{1–3} Non-Hispanic white and Hispanic populations have a higher prevalence than African-American populations but a lower prevalence than Asian populations.^{1,3,4} NSCLP is considered to be a multifactorial disorder, that is, caused by both genes and environmental

exposures, neither of which has been fully defined. Anywhere from 2 to 14 genes have been proposed to contribute to the NSCLP phenotype and association and linkage studies have identified some of these putative genes.^{5–7}

Genome scans have proven to be a useful tool for identifying NSCLP candidate chromosomal regions. Seven NSCLP genome scans have been reported involving populations from East Asia, India, Europe, Syria and Turkey.^{8–14} In addition, a meta-analysis was performed on data generated from 13 NSCLP studies, including 5 of the aforementioned studies and 8 additional unpublished data sets.¹⁵ The individual genomic scans have identified 3–20 candidate regions (Supplementary Table 1), and 1p31–36, 2q32–37, 6q21–27 and 16q21–24 were common to at least four of the eight studies listed in Supplementary Table 1.^{8–15}

To confirm previously reported chromosomal regions and to identify new chromosomal regions likely to contain NSCLP genes, we subjected 10 multiplex NSCLP families to genomic screening utilizing the Illumina Linkage IVb 6K SNP Linkage Panel (Illumina Inc., San Diego, CA, USA). Of these, nine families are from our well-characterized non-Hispanic white data set and one family was African American. We identified one new region from each ethnic group with an LOD >2.5; one from chromosome 8q21.3–24.12 (2.98; African American) and one from chromosome 22q12.2–q12.3 (2.66; non-Hispanic white). Two candidate genes from the chromosome 8 region and one from the chromosome 22 region were further evaluated.

Inspection of the region(s) with the highest LOD score from each ethnic group identified three genes of interest: growth differentiation factor 6 (*GDF6*) and syndecan 2 (*SDC2*) in 8q21.3–24.12 and myosin heavy chain 9 (*MYH9*) in 22q12.2–12.3.¹⁶ *SDC2* and *GDF6* on chromosome 8 were considered to be good candidate genes because of their known biological function. *SDC2* is involved in cell signaling, migration and cell–matrix interactions and interacts with members of the transforming growth factor- β family.^{17–19} *GDF6* is a member of the bone morphogenetic protein group that forms part of the transforming growth factor- β superfamily and is expressed in the craniofacial region during embryogenesis^{20–24} *MYH9* on chromosome 22 is of particular interest because it is highly expressed in the palatal shelves before fusion.²⁵ Moreover, in a recent study of NSCLP in a group of Italians, an association was identified with an SNP in *MYH9*.²⁶ In this study, we identified SNPs spanning the candidate genes and evaluated them for evidence and/or association.

Materials and methods

Data set

The data set consists of 123 families with a history of NSCLP. Of these, there are 81 extended multiplex families (65 non-Hispanic white, 15 Hispanic and 1 African

American) and 42 multiplex trios/duos (23 non-Hispanic white and 19 Hispanic) with a positive history of NSCLP. In addition, there are 316 simplex trios/duos (226 non-Hispanic white, 82 Hispanic and 8 African American) with no family history of NSCLP. These families, methods of ascertainment and inclusion criteria have been reported earlier.²⁷ Blood or saliva samples were collected after obtaining informed consent. DNA was extracted from blood using Roche DNA Isolation Kit for Mammalian Blood (Roche, Basel, Switzerland) or from saliva using Oragene Purifier (DNA Genotek Inc., Ottawa, ON, Canada) following the manufacturer's protocol.

Genome scan

Ten multiplex families were subjected to a genome scan using the 6K Illumina Linkage IVb mapping panel. These families were selected from our data set on the basis of the number of affected individuals and availability of DNA. Nine families were non-Hispanic white and one (F1100) was African American (Figure 1). Under our dominant model, four of the non-Hispanic white families, 100, 1000, 1200 and 3000, can yield maximum LOD scores greater than 1 (1.6, 1.5, 1.1 and 1.2, respectively). A maximum LOD score of 2.98 is possible for the African-American family. Among the non-Hispanic white families, there were 30 affected individuals of 67 individuals. In family 100 (F100), there were two siblings (III-4 and III-9) who were reported to have a notched gum in childhood, a finding that could indicate a microform cleft lip. However, these diagnoses could not be clinically confirmed. Therefore, the linkage analysis was performed twice: first, assuming these individuals to be affected and then coding them as unknown with regard to the cleft phenotype. There were 27 genotyped individuals in the African-American family, 7 of whom were affected.

The Illumina Linkage IVb mapping panel consisting of six 008 SNPs was used on a BeadStation system (Illumina Inc.). Allele detection and genotype calling were performed using the BeadStudio software (Illumina Inc.).

Candidate genes

Thirty-seven SNPs spanning the three candidate genes identified in the genome scan (*GDF6* – 6; *SDC2* – 21; and *MYH9* – 10) were subsequently typed in the entire NSCLP data set with the exclusion of the African-American families; selection criteria for these SNPs were discussed earlier.²⁷ *GDF6* and *SDC2* SNPs were chosen with minor allele frequency >0.3, preferentially selecting coding and TagSNPs to obtain maximum coverage of these genes. Six of the *MYH9* SNPs were chosen on the basis of a previous publication;²⁶ the remaining four SNPs in the *MYH9* gene were selected using the above criteria. TaqMan Assays (Applied Biosystems; Foster City, CA, USA) were used for genotyping; alleles were detected using the ABI 7900HT Sequence Detection System. All genotyping results were




Figure 1 Families included in genome scan. Family 1100 is African American; all the remaining families are non-Hispanic white. *DNA included in the genome scan; arrow denotes proband. Filled symbols denote affected individuals; half-filled symbols denote notched gum; empty symbols denote unaffected individual.

imported into Progeny Lab (South Bend, Indiana, USA), and PedCheck was run on all SNPs to identify Mendelian inconsistencies.²⁸

Analysis

Results of genotyping from the genome scan were subjected to parametric and non-parametric linkage analyses using MERLIN.²⁹ Linkage parameters were as pre-

viously calculated, that is, dominant model with penetrances of 0.24 in females and 0.32 in males and a phenocopy rate of 0.001.³⁰ An r^2 of 0.19 was the cutoff value for clustering SNPs to remove the effect of linkage disequilibrium. Family 1100 was analyzed separately because of its different race/ethnicity. Analysis of the candidate gene SNPs was restricted to our Hispanic and non-Hispanic white families, as the number of families of

other ethnicities was small. For analyses of the candidate gene SNPs, probands/families were stratified by ethnicity alone or by ethnicity and family history. In addition, the Hispanic and non-Hispanic white families were analyzed together for the *MYH9* SNPs, as the allele frequencies and LD patterns did not differ between the two ethnicities. Allele frequencies and the Hardy–Weinberg equilibrium (HWE) were calculated using SAS (v9.1). Pairwise linkage disequilibrium values (D' and r^2) were calculated using GOLD.³¹ To extract the maximum amount of information from the data set, multiple approaches for assessing linkage and/or association were used. Parametric and non-parametric linkage analyses were performed as discussed above. The pedigree disequilibrium test (PDT), Geno-PDT (G-PDT) and association in the presence of linkage (APL) test were used to evaluate evidence for association.^{32–34} This panel of analytic tools was chosen because each has different strengths and assumptions. PDT is an extension of the transmission disequilibrium test and allows for the incorporation of extended pedigrees in the analysis of allelic association. The G-PDT examines the association between marker genotypes and disease, whereas APL allows for missing parental genotypes. Each program may be more powerful for certain genetic models and for different pedigree structures. In addition, APL can be used to examine multi-marker haplotypes; it was used to look for the overtransmission of two-marker haplotypes within genes; all possible pairwise combinations of SNPs were included.

Results

Genome scan

Ten families were subjected to the Illumina Linkage IVb SNP genome-wide panel. Nine non-Hispanic white families generated 11 chromosomal regions with an LOD > 1.5 with parametric and/or non-parametric analysis (Table 1).

The highest LOD score for either methodology was two SNPs in the chromosome 22q12.2–q12.3 region (rs762883 and rs9862). Three of the families, 100, 1000 and 3000, had maximum LOD score > 1.0 in the region (1.05, 1.11 and 1.12, respectively), although not for the same set of SNPs (data not shown). Three chromosomal regions with an LOD > 1.5 were identified in the large African-American family (Table 2). The highest LOD score (2.98) was found in an SNP in chromosome 8q21.3–24.12. This is the maximum possible LOD score for this family. There was no overlap in the regions identified in these two groups. The multipoint graphs for chromosomes with LOD scores > 1.5 are shown in Supplementary Figures 2 and 3.

Candidate genes

All 37 SNPs from the three candidate genes, *MYH9*, *GDF6* and *SDC2*, were in HWE. The allele frequencies of most SNPs in *GDF6* and *SDC2* were significantly different between the non-Hispanic whites and Hispanics, even after Bonferroni correction ($P=0.001$) (Table 3A). Therefore, the data were stratified by ethnicity for the analysis of these genes. In contrast, the differences in the *MYH9* SNPs did not meet this criterion (Table 3B). For this reason, *MYH9* SNPs were analyzed with the two ethnicities combined.

Two-point parametric and non-parametric analyses in the combined data set found evidence for linkage to a single SNP, rs1002246, in *MYH9* (HLOD = 1.58 and $P=0.0006$, respectively) (Table 4). There was no evidence for linkage by multipoint analysis (data not shown). However, all three methods of association analysis identified evidence for altered transmission of this same SNP, rs1002246 (Table 5). In addition, PDT detected evidence for an altered transmission of three additional SNPs. When the data were stratified by family history, a marginally significant association was still present for the SNP,

Table 1 Chromosomal regions with LOD > 1.5 in nine non-Hispanic white families

Chromosome	Region ^a	dbSNP	cM	Maximum LOD score			
				Parametric (hLOD)		Non-parametric (LOD)	
				Aff ^b	Unk ^c	Aff ^b	Unk ^c
3	3p23	rs342758	58.08	0.35	0.85	0.92	1.64
3	3p21.1	rs11235	71.53	0.87	1.58	0.49	1.03
3	3p13–14.1 ^a	rs736333	94.05	0.24	0.67	0.65	1.53
5	5p13.3 ^a	rs1966983	51.74	0.94	1.64	0.72	1.19
5	5q35.2	rs1875189	192.51	0.57	0.53	1.59	1.04
6	6q27	rs727811	177.17	0.85	1.66	0.28	0.45
7	7q36.2–36.3 ^a	rs7795368	182.8	1.7	2.59	0.57	1.16
9	9q22.2–22.32 ^a	rs1048510	97.53	1.78	1.27	0.96	0.82
9	9q33.2–33.3	rs1984001	132.41	1.46	0.95	1.87	1.73
22	22q11.23–12.1	rs763281	21.41	1.75	1.22	1.46	0.99
22	22q12.2–12.3 ^a	rs762883	36.94	2.66	2.12	2.49	2.17

^aNovel region identified.

^bAff, Analysis of F100 individuals with notched gums classified as affected.

^cUnk, Analysis of F100 individuals with notched gums classified as unknown.

Table 2 Chromosomal regions with LOD > 1.5 in the African-American family

Chromosome	Region ^a	dbSNP	cM	Maximum LOD score	
				Parametric (hLOD)	Non-parametric (LOD)
2	2p22 ^a	rs1446297	62.97	2.27	1.07
3	3p26	rs11130180	15.16	2	0.71
8	8q21.3–24.12	rs536161	97.66	2.98	1.47

^aIndicates novel region identified.

Table 3A Allele frequencies of chromosome 8 dbSNPs

Gene	dbSNP ^a	Base pair	Location	MAF ^b	HCF ^{c,d}
<i>SDC2</i>	rs2452993	97 574 572	Upstream	0.42	0.59***
<i>SDC2</i>	rs2437772	97 582 153	Intron 1	0.31	0.27
<i>SDC2</i>	rs2437780	97 588 852	Intron 1	0.30	0.26
<i>SDC2</i>	rs2575738	97 599 578	Intron 1	0.29	0.20*
<i>SDC2</i>	rs7003874	97 607 128	Intron 1	0.46	0.35*
<i>SDC2</i>	rs874643	97 611 319	Intron 1	0.19	0.40***
<i>SDC2</i>	rs2440681	97 613 757	Intron 1	0.35	0.22***
<i>SDC2</i>	rs2582831	97 620 954	Intron 1	0.28	0.44***
<i>SDC2</i>	rs2582839	97 627 443	Intron 1	0.29	0.46***
<i>SDC2</i>	rs1444573	97 633 837	Intron 1	0.37	0.58***
<i>SDC2</i>	rs1984456	97 636 282	Intron 1	0.35	0.48***
<i>SDC2</i>	rs2582846	97 642 858	Intron 1	0.20	0.43***
<i>SDC2</i>	rs12156240	97 646 383	Intron 1	0.49	0.69***
<i>SDC2</i>	rs2575702	97 654 007	Intron 1	0.34	0.46**
<i>SDC2</i>	rs2651472	97 665 849	Intron 1	0.49	0.58*
<i>SDC2</i>	rs2028086	97 673 354	Intron 1	0.24	0.14**
<i>SDC2</i>	rs1836910	97 680 704	Intron 2	0.26	0.17*
<i>SDC2</i>	rs1042381	97 683 837	Exon 3	0.18	0.14
<i>SDC2</i>	rs2651454	97 690 146	Intron 4	0.23	0.38***
<i>SDC2</i>	rs2651458	97 694 626	Downstream	0.23	0.38***
<i>SDC2</i>	rs714046	97 700 749	Downstream	0.19	0.2
<i>GDF6</i>	rs2247792	97 219 221	Upstream	0.41	0.21***
<i>GDF6</i>	rs2440199	97 225 587	Exon 1	0.41	0.63***
<i>GDF6</i>	rs2255182	97 234 080	Intron 1	0.49	0.28***
<i>GDF6</i>	rs2514527	97 238 503	Intron 1	0.38	0.64***
<i>GDF6</i>	rs2514531	97 243 999	Downstream	0.45	0.26***
<i>GDF6</i>	rs714552	97 252 618	Downstream	0.40	0.61***

^aBold dbSNPs are also genotyped in genome scan.

^bMinor allele frequency of non-Hispanic white data set.

^cCorresponding frequency in Hispanic of non-Hispanic white minor allele.

^dP-value less than *0.05, **0.001 and ***0.0001.

rs1002246, in the simplex data set ($P=0.03$). Inspection of all two-SNP haplotypes found that most haplotypes constructed with rs1002246 and containing the 'A' allele were overtransmitted, but generally did not reach significance, even when stratified by ethnicity and/or family history (data not shown). The minor allele frequency (ie, frequency of the 'A' allele) was 0.34 in the non-Hispanic whites. Haplotypes with the 'G' rs1002246 allele were either undertransmitted or demonstrated the expected transmission.

There was no evidence for linkage to SNPs in *SDC2*. Marginal significance for altered transmission was detected in both the non-Hispanic white and Hispanic samples (Table 6A and B and Supplementary Tables 2 and 3). In the simplex non-Hispanic white sample, rs1042381

Table 3B Allele frequencies of chromosome 22 dbSNPs

Gene	dbSNP	Base pair	Location	MAF ^a	HCF ^{b,c}
<i>MYH9</i>	rs7078	35 007 860	Downstream	0.30	0.25
<i>MYH9</i>	rs4821478	35 020 059	Intron 28	0.35	0.35
<i>MYH9</i>	rs2009930	35 029 252	Intron 19	0.35	0.37
<i>MYH9</i>	rs2239783	35 035 074	Intron 15	0.26	0.27
<i>MYH9</i>	rs3752462	35 040 129	Intron 13	0.36	0.37
<i>MYH9</i>	rs1002246	35 044 605	Intron 10	0.34	0.33
<i>MYH9</i>	rs2071731	35 048 804	Intron 5	0.37	0.34
<i>MYH9</i>	rs713659	35 058 638	Intron 3	0.42	0.37
<i>MYH9</i>	rs713839	35 063 884	Intron 3	0.38	0.35
<i>MYH9</i>	rs739097	35 076 025	Intron 1	0.44	0.36

^aMinor allele frequency of non-Hispanic white data set.

^bCorresponding frequency in Hispanic of non-Hispanic white minor allele.

^cP-value less than *0.05, **0.001 and ***0.0001.

was marginally significant (Table 6A). In the Hispanic sample, significant association was detected in the entire data set for rs198456, in the multiplex families for rs2437772 and rs2437780, and for the simplex families for rs2582831 and rs2582846 (Table 6B). Two *SDC2* haplotypes demonstrated excess transmission in the non-Hispanic white group and three in the Hispanic group (Table 7).

No evidence for linkage to SNPs in *GDF6* was detected. Again, altered transmission was only marginally significant for two SNPs: rs2255182 in the complete non-Hispanic white sample and both the multiplex and simplex subgroups ($P=0.04$, $P=0.02$ and $P=0.04$, respectively) and rs2247792 in the non-Hispanic white multiplex families ($P=0.03$) (Table 6A). Overtransmission of one haplotype involving rs2440199 and rs2514527 was observed in the Hispanic sample ($P=0.008$) (Table 7).

Discussion

This study was undertaken to identify additional genes that contribute to the NSCLP phenotype. One African-American and nine non-Hispanic white families were analyzed by a 6K genome scan. Fourteen regions were identified with LOD scores > 1.5 and four regions had LOD scores > 2 (2p22, 7q36.2–36.3, 8q21.3–24.12 and 22q12.1-q12.3) (Tables 1 and 2). Of these, only the 8q21.3–24.12 region had previously been identified by genomic scanning (Supplementary Table 1).^{11,15} The region from each

Table 4 Parametric and non-parametric linkage analysis for *MYH9*

<i>dbSNP</i>	<i>cM</i>	<i>Base pair</i>	<i>Parametric</i>		<i>Non-parametric</i>	
			<i>hLOD</i>	α	<i>LOD</i>	<i>P-value</i> ^a
rs7078	35.008	35 007 860	0.018	0.083	0.01	0.4
rs4821478	35.02	35 020 059	0.033	0.109	0.02	0.4
rs2009930	35.029	35 029 252	0	0	-0.07	0.7
rs2239783	35.035	35 035 074	0.174	0.213	0	0.5
rs3752462	35.04	35 040 129	0.108	0.205	0.12	0.2
rs1002246	35.045	35 044 605	1.578	0.694	2.26	0.0006
rs2071731	35.049	35 048 804	0	0	0	0.5
rs713659	35.059	35 058 638	0	0	0	0.5
rs713839	35.064	35 063 884	0	0.005	0	0.5
rs739097	35.076	35 076 025	0	0	-0.01	0.6

^a*P*-value <0.05 bolded.**Table 5** Results for *MYH9* from current and previously reported study

<i>dbSNP</i>	<i>Location</i>	<i>Martinelli et al</i>		<i>Entire sample</i>				<i>Current data</i> ^a			<i>Simplex families</i>		
		<i>Italian sample</i>	<i>TDT</i> ^b	<i>PDT</i>	<i>G-PDT</i>	<i>APL</i>	<i>PDT</i>	<i>G-PDT</i>	<i>APL</i>	<i>PDT</i>	<i>G-PDT</i>	<i>APL</i>	
rs5995288	Intron 1	0.82	—	—	—	—	—	—	—	—	—	—	
rs739097	Intron 1	0.5	0.17	0.395	0.142	0.617	0.862	0.948	0.114	0.295	0.067		
rs713839	Intron 3	—	0.067	0.191	0.11	0.452	0.558	0.596	0.043	0.151	0.094		
rs713659	Intron 3	—	0.211	0.397	0.098	0.756	0.894	0.545	0.091	0.17	0.014		
rs2071731	Intron 5	0.88	0.12	0.186	0.165	0.607	0.663	0.456	0.065	0.144	0.026		
rs1002246	Intron 10	1	0.02	0.043	0.009	0.056	0.149	0.127	0.185	0.262	0.033		
rs3752462	Intron 13	0.02	0.035	0.117	0.602	0.114	0.216	0.662	0.162	0.382	0.353		
rs2239783	Intron 15	—	0.681	0.834	0.369	0.478	0.726	0.143	0.86	0.61	0.857		
rs2009930	Intron 19	0.4	0.039	0.131	0.177	0.195	0.386	0.718	0.087	0.257	0.065		
rs4821478	Intron 28	—	0.023	0.077	0.381	0.053	0.163	0.899	0.223	0.465	0.328		
rs2269529	Exon 34	0.69	—	—	—	—	—	—	—	—	—		
rs7078	Downstream	0.34	0.836	0.832	0.345	0.564	0.712	0.137	0.766	0.879	0.770		

—, not genotyped.

^a*P*-values have not been corrected for multiple testing.^b*P*-value <0.05 bolded.**Table 6** Association results for *SDC2* and *GDF6* on chromosome 8

<i>Gene</i>	<i>dbSNP</i>	<i>All NHW</i> ^{a,b}			<i>Simplex NHW</i>			<i>Multiplex NHW</i>		
		<i>PDT</i>	<i>G-PDT</i>	<i>APL</i>	<i>PDT</i>	<i>G-PDT</i>	<i>APL</i>	<i>PDT</i>	<i>G-PDT</i>	<i>APL</i>
<i>(A) Non-Hispanic white population</i> ^c										
<i>SDC2</i>	rs1042381	0.796	0.822	0.632	0.046	0.029	0.289	0.123	0.075	0.349
<i>GDF6</i>	rs2247792	0.804	0.223	0.149	0.804	0.223	0.147	0.631	0.013	0.244
<i>GDF6</i>	rs2440199	0.955	0.348	0.597	0.955	0.348	0.587	0.449	0.05	0.887
<i>GDF6</i>	rs2255182	0.582	0.036	0.764	0.582	0.036	0.763	0.679	0.019	0.678
<i>Gene</i>	<i>dbSNP</i>	<i>PDT</i>	<i>All</i> ^{a,b} <i>G-PDT</i>	<i>APL</i>	<i>PDT</i>	<i>Simplex</i> <i>G-PDT</i>	<i>APL</i>	<i>PDT</i>	<i>Multiplex</i> <i>G-PDT</i>	<i>APL</i>
<i>(B) Hispanic population</i> ^c										
<i>SDC2</i>	rs2437772	0.522	0.168	0.819	0.336	0.487	0.716	1	0.035	0.915
<i>SDC2</i>	rs2437780	0.796	0.139	0.95	0.578	0.839	0.975	0.858	0.04	0.959
<i>SDC2</i>	rs2582831	0.655	0.89	0.147	1	0.793	0.04	0.564	0.668	0.878
<i>SDC2</i>	rs1984456	0.753	0.939	0.037	0.706	0.837	0.154	0.9	0.974	0.086
<i>SDC2</i>	rs2582846	0.274	0.594	0.058	0.132	0.281	0.036	0.715	0.804	0.548

^a*P*-value <0.05 bolded.^b*P*-values have not been corrected for multiple testing.^cResults for SNPs with at least one test statistic *P*<0.05.

Table 7 Overtransmitted haplotypes for *SDC2* and *GDF6*

Gene	dbSNP1	dbSNP2	Ethnicity	P-value	Haplotype
<i>SDC2</i>	rs2452993	rs7003874	NHW	0.01	1-2
<i>SDC2</i>	rs2452993	rs12156240	NHW	0.003	2-1
<i>SDC2</i>	rs2575738	rs2651472	H	0.04	2-1
<i>SDC2</i>	rs2575738	rs1836910	H	0.04	1-2
<i>SDC2</i>	rs2582846	rs12156240	H	0.016	2-1
<i>GDF6</i>	rs2440199	rs2514527	H	0.008	2-1

NHW, non-Hispanic white; H, Hispanic.

ethnicity that resulted in the highest LOD score was subjected to candidate gene analysis.

Eleven regions on six chromosomes with LOD scores > 1.5 were identified in the non-Hispanic white families (Table 2). The highest LOD score was found in chromosomal region 22q12.2–12.3 and the total linkage area spans over 20 cM. This region contains the *MYH9* gene, which is expressed in the developing palate and was found to be associated with NSCLP in an Italian population.^{25,26} To further evaluate the role of *MYH9* in NSCLP, 10 intragenic and flanking SNPs, which included six of the SNPs genotyped in the Italian families, were interrogated in our NSCLP sample.

Linkage was found between a single SNP, rs1002246, in the *MYH9* gene and our entire data set (Table 5), consistent with the prior identification of this region in our genome scan. PDT, G-PDT and APL analyses, all identified this SNP in our entire NSCLP sample. rs1002246 differs from the SNP found in the Italian NSCLP sample (Table 5). rs1002246 is in intron 10 of the *MYH9* gene but this is not in a region associated with intron–exon splicing. Although intronic SNPs do not typically alter protein structure, an association with intronic variants has been reported for a number of complex diseases.^{27,35–40} rs1002246 is located in a region of reduced LD that can complicate identifying at-risk haplotypes (Supplementary Table 3). In fact, we were unable to identify a high-risk haplotype. Interestingly, rs3752462, the SNP identified by the Italian group, was only marginally significant in both data sets (Table 5). Our finding of linkage by parametric and non-parametric analyses and association strongly suggests that *MYH9* may play a causal role in NSCLP. Nevertheless, we cannot conclusively exclude the possibility that another gene in the 22q12.2–12.3 chromosomal region is contributing to the positive linkage and association. We are continuing to evaluate this region.

This is the first NSCLP genomic scan to include an African-American family. NSCLP is relatively rare in African Americans with a birth prevalence of 1/2500, when compared with 1/1400 in non-Hispanic whites and Hispanics in Texas.³ Although earlier genome scans have

been performed on non-Hispanic white, Indian, Chinese and Syrian populations, it is a rarity to find a large African-American family with segregating NSCLP^{8–14} Using this family, three chromosomal regions, 2p22, 3p26 and 8q21.3–24.12, were identified with LOD scores > 1.5, of which only 2p22 is new (Table 2).

The most significant linkage (LOD = 2.98) was to 8q21.3–24.12 region, the same region identified by the meta-analysis, providing further support that this region contains an NSCLP gene.¹⁵ *SDC2* and *GDF6* were found in this region and, because of their known biological function, they were considered to be good candidate genes. Nevertheless, linkage was not detected between either of these genes and NSCLP in our data set. Marginal evidence for association ($0.01 < P < 0.05$) was detected for SNPs in both of these genes in the non-Hispanic white and Hispanic samples. In *SDC2*, SNPs rs2437772 and rs2437780 were both significant in the multiplex non-Hispanic white data set using the G-PDT; however, neither PDT nor APL detected a significant association. Therefore, these results are marginal at best. Haplotype analysis revealed altered transmission for two *SDC2* haplotypes ($P < 0.01$). However, given the large number of haplotypes tested, this does not provide strong evidence for a role of *SDC2* in NSCLP.

G-PDT analysis of SNPs in *GDF6* found borderline evidence for association in the non-Hispanic white data set; PDT and APL tests were negative. rs2255182, in intron 1, was significant in the total non-Hispanic white data set as well as when the data were stratified by family history (Table 3A). In addition, rs2247792, 22.5 kb upstream of the *GDF6* gene, was associated in the multiplex non-Hispanic white data set. Interestingly, analysis identified a significant haplotype in the Hispanic data set, but not in the non-Hispanic white sample (Table 7). Although association analyses of the SNPs in *SDC2* and *GDF6* showed some evidence for altered transmission, the level did not correspond to the LOD score detected in this region. This may be the result of linkage without an association. An alternative explanation is that this region was identified in an African-American family, and genes in this region do not play a significant role in the non-Hispanic white and Hispanic populations. NSCLP displays genetic heterogeneity with a variety of genes playing an etiologic role, and it is likely that these genes vary with ethnicity.⁴¹ However, because this region was also identified in other ethnicities, additional testing is underway to interrogate this region in the entire data set. Of particular interest is the Frizzled 6 (*Fzd6*) gene that is located in this region. *Fzd6* is a Wnt receptor whose expression is upregulated during osteogenic differentiation of mesenchymal stem cells.⁴² *Fzd6* functions in the noncanonical Wnt pathway to decrease the amount of binding of TCF/LEF and TCF- β -catenin to target DNA.⁴³ We have recently identified an association between several Wnt genes and NSCLP,⁴⁴ and studies are

underway to evaluate *Fzd6* for association with NSCLP and interaction with Wnt genes.

Regions 2p22 and 3p26 were also identified in the African-American family; these are new avenues for discovery of NSCLP genes. 2p22 is a novel chromosomal region, whereas 3p26 was previously reported in a genome scan of Chinese multiplex families.⁹ Further examination of these regions needs to be undertaken to identify novel NSCLP genes.

In addition to 22q12.2–q12.3 (which contains *MYH9*), 10 regions were identified in the non-Hispanic white families with LOD scores greater than 1.5 but less than 2. Four of these regions, 3p14.1–p13, 5p13.3, 7q36.2–q36.3 and 9q22.22–q22.32, were novel; however, and there are no obvious candidate genes in two of these regions (3p14.1–p13 and 5p13.3). The remaining six regions have been reported earlier in genomic scans.^{8–11,13–15}

The 7q36.2–36.3 region is novel and contains sonic hedgehog homolog (*SHH*) and engrailed homeobox 2 (*EN2*), two known NSCLP candidate genes. *SHH* plays a critical role in the development and patterning of the craniofacial processes that give rise to the developing lip and palate.⁴⁵ Mutations in the human *SHH* gene cause holoprosencephaly-3, which can be associated with cleft lip and palate (OMIM: #142945),⁴⁶ *SHH* null mice can have cleft palate.⁴⁷ However, single-strand conformational polymorphism analysis of the *SHH* gene in NSCLP patients has not identified any disease-causing mutations.⁴⁸ *EN2* plays a role in the central nervous system development and has also been implicated in autism spectrum disorders (OMIM: #608636).⁴⁹ Previous RFLP testing found no association between the *EN2* gene and NSCLP.⁵⁰ Although no positive findings have been found, these remain candidate genes of interest.

Two regions on chromosome 9, 9q22.2–22.32 and 9q33.2–33.3, were identified in this genomic scan. The 9q22.2–22.32 region is novel and contains both the BarH-like homeobox 1 (*Barx1*) and patched homolog 1 (*PTCH1*). Mouse *BARX1* is expressed in the first and second branchial arches that form the craniofacial processes.⁵¹ The role of *Barx1* in NSCLP has not been evaluated. *PTCH1*, an SHH receptor, in combination with *BARX1*, plays a role in mid-face structure formation.⁵² Mutations in *PTCH1* cause basal cell nevus syndrome (BCNS; OMIM: #109400) and 5% of BCNS patients have a cleft lip and palate.⁵³ Interestingly, forkhead box E1 (*FOXE1*), which plays a role in thyroid morphogenesis, is located just outside this region in 9q22.33.⁵⁴ The *FOXE1* region has been identified in the NSCLP genome scans.⁵⁵ Mutations in *FOXE1* cause the Bamforth–Lazarus syndrome of which cleft palate is an associated finding (OMIM: #241850). However, sequencing of the *FOXE1* gene in the NSCLP probands identified two missense mutations, both of which were predicted to be benign.⁵⁶

There are some limitations to the study. NSCLP is clearly heterogeneous, and among the non-Hispanic white families, none have the potential to yield an LOD score >1.6, making it difficult to obtain conclusive evidence for linkage. The one family that can yield a nearly significant LOD score is African American; this racial group has a significantly lower risk of NSCLP than non-Hispanic whites and Hispanics and very likely has a different etiology. In addition, the sample for the association studies is not large enough to detect SNPs with minor effects. Nonetheless, we have identified new regions of interest through linkage analyses as well as candidate genes through our association studies, both of which warrant further evaluation.

Here, we report six new regions (2p22, 3p13–14.1, 5p13.3, 7q36.2–36.3, 9q22.2–22.32 and 22q12.2–12.3) that can be interrogated for NSCLP genes. We identified eight regions that were previously implicated in other genomic scans in a variety of populations. The strongest linkages were found for the 8q21.3–24.12 and 22q12.2–12.3 regions, with the *MYH9* (chromosome 22) gene exhibiting the strongest evidence for a causal role. This gene should be further evaluated in other populations for its possible role in NSCLP. The results of this study are important as they provide additional regions to search for new NSCLP genes and confirm the findings of earlier genomic scans. Our findings also demonstrate that large multiplex families with complex disorders can be successfully used in genome mapping strategies.

Acknowledgements

We thank Maria Elena Serna for expert clinical co-ordination and for managing the databases. We also thank Jacqueline Bui and Priscilla Roche for their technical assistance, Yuki Bradford for data analysis at the Center for Human Genetics Research at Vanderbilt University and all of the NSCLP families who kindly participated in this study. This study was approved by the Committee for the Protection of Human Subjects of the University of Texas Health Science Center at Houston (HSC-MS-03-090). This work was funded by grants from the National Institutes of Health (R01-DE011931 to JTH, T32-DE015355-05 trainee BTC).

Conflict of interest

The authors state no conflict of interest.

References

- 1 Gorlin RJ, Cohen MM, Hennekam RCM: *Syndromes of the Head and Neck*. New York: Oxford University Press, 2001.
- 2 Wyszynski D: *Cleft Lip and Palate: From Origin to Treatment*. Oxford: Oxford University Press, 2002.
- 3 Hashmi SS, Waller DK, Langlois P, Canfield M, Hecht JT: Prevalence of nonsyndromic oral clefts in Texas: 1995–1999. *Am J Med Genet A* 2005; **134**: 368–372.
- 4 Canfield MA, Honein MA, Yuskiv N *et al*: National estimates and race/ethnic-specific variation of selected birth defects in the

- United States, 1999–2001. *Birth Defects Res A Clin Mol Teratol* 2006; **76**: 747–756.
- 5 Schliekelman P, Slatkin M: Multiplex relative risk and estimation of the number of loci underlying an inherited disease. *Am J Hum Genet* 2002; **71**: 1369–1385.
- 6 Lidral AC, Moreno LM: Progress toward discerning the genetics of cleft lip. *Curr Opin Pediatr* 2005; **17**: 731–739.
- 7 Carinci F, Scapoli L, Palmieri A, Zollino I, Pezzetti F: Human genetic factors in nonsyndromic cleft lip and palate: an update. *Int J Pediatr Otorhinolaryngol* 2007; **71**: 1509–1519.
- 8 Field LL, Ray AK, Cooper ME, Goldstein T, Shaw DF, Marazita ML: Genome scan for loci involved in nonsyndromic cleft lip with or without cleft palate in families from West Bengal, India. *Am J Med Genet A* 2004; **130**: 265–271.
- 9 Marazita ML, Field LL, Cooper ME et al: Genome scan for loci involved in cleft lip with or without cleft palate in Chinese multiplex families. *Am J Hum Genet* 2002; **71**: 349–364.
- 10 Marazita ML, Field LL, Tuncbilek G, Cooper ME, Goldstein T, Gursu KG: Genome-scan for loci involved in cleft lip with or without cleft palate in consanguineous families from Turkey. *Am J Med Genet A* 2004; **126**: 111–122.
- 11 Prescott NJ, Lees MM, Winter RM, Malcolm S: Identification of susceptibility loci for nonsyndromic cleft lip with or without cleft palate in a two stage genome scan of affected sib-pairs. *Hum Genet* 2000; **106**: 345–350.
- 12 Radhakrishna U, Ratnamala U, Gaines M et al: Genomewide scan for nonsyndromic cleft lip and palate in multigenerational Indian families reveals significant evidence of linkage at 13q33.1–34. *Am J Hum Genet* 2006; **79**: 580–585.
- 13 Riley BM, Schultz RE, Cooper ME et al: A genome-wide linkage scan for cleft lip and cleft palate identifies a novel locus on 8p11–23. *Am J Med Genet A* 2007; **143**: 846–852.
- 14 Wyszynski DF, Albacha-Hejazi H, Aldirani M et al: A genome-wide scan for loci predisposing to non-syndromic cleft lip with or without cleft palate in two large Syrian families. *Am J Med Genet* 2003; **123A**: 140–147.
- 15 Marazita ML, Murray JC, Lidral AC et al: Meta-analysis of 13 genome scans reveals multiple cleft lip/palate genes with novel loci on 9q21 and 2q32–35. *Am J Hum Genet* 2004; **75**: 161–173.
- 16 Kent WJ, Sugnet CW, Furey TS et al: The human genome browser at UCSC. *Genome Res* 2002; **12**: 996–1006.
- 17 Chen L, Klass C, Woods A: Syndecan-2 regulates transforming growth factor-beta signaling. *J Biol Chem* 2004; **279**: 15715–15718.
- 18 Essner JJ, Chen E, Ekker SC: Syndecan-2. *Int J Biochem Cell Biol* 2006; **38**: 152–156.
- 19 Oh ES, Couchman JR: Syndecans-2 and -4; close cousins, but not identical twins. *Mol Cells* 2004; **17**: 181–187.
- 20 Settle Jr SH, Rountree RB, Sinha A, Thacker A, Higgins K, Kingsley DM: Multiple joint and skeletal patterning defects caused by single and double mutations in the mouse Gdf6 and Gdf5 genes. *Dev Biol* 2003; **254**: 116–130.
- 21 Sena K, Morotome Y, Baba O, Terashima T, Takano Y, Ishikawa I: Gene expression of growth differentiation factors in the developing periodontium of rat molars. *J Dent Res* 2003; **82**: 166–171.
- 22 Portnoy ME, McDermott KJ, Antonellis A et al: Detection of potential GDF6 regulatory elements by multispecies sequence comparisons and identification of a skeletal joint enhancer. *Genomics* 2005; **86**: 295–305.
- 23 Hanel ML, Hensey C: Eye and neural defects associated with loss of GDF6. *BMC Dev Biol* 2006; **6**: 43.
- 24 Portnoy ME, McDermott KJ, Antonellis A et al: Detection of potential GDF6 regulatory elements by multispecies sequence comparisons and identification of a skeletal joint enhancer. *Genomics* 2005; **86**: 295–305.
- 25 Marigo V, Nigro A, Pecci A et al: Correlation between the clinical phenotype of MYH9-related disease and tissue distribution of class II nonmuscle myosin heavy chains. *Genomics* 2004; **83**: 1125–1133.
- 26 Martinelli M, Di Stazio M, Scapoli L et al: Cleft lip with or without cleft palate: implication of the heavy chain of non-muscle myosin IIA. *J Med Genet* 2007; **44**: 387–392.
- 27 Chiquet BT, Lidral AC, Stal S et al: CRISPLD2: A novel NSCLP candidate gene. *Hum Mol Genet* 2007; **16**: 2241–2248.
- 28 O'Connell JR, Weeks DE: The VITESSE algorithm for rapid exact multilocus linkage analysis via genotype set-recoding and fuzzy inheritance. *Nat Genet* 1995; **11**: 402–408.
- 29 Abecasis GR, Cherny SS, Cookson WO, Cardon LR: Merlin – rapid analysis of dense genetic maps using sparse gene flow trees.[see comment]. *Nat Genet* 2002; **30**: 97–101.
- 30 Blanton SH, Bertin T, Patel S, Stal S, Mulliken JB, Hecht JT: Nonsyndromic cleft lip and palate: four chromosomal regions of interest. *Am J Med Genet A* 2004; **125**: 28–37.
- 31 Abecasis GR, Cookson WO: GOLD – graphical overview of linkage disequilibrium. *Bioinformatics* 2000; **16**: 182–183.
- 32 Martin ER, Bass MP, Gilbert JR, Pericak-Vance MA, Hauser ER: Genotype-based association test for general pedigrees: the genotype-PDT. *Genet Epidemiol* 2003; **25**: 203–213.
- 33 Martin ER, Monks SA, Warren LL, Kaplan NL: A test for linkage and association in general pedigrees: the pedigree disequilibrium test. *Am J Hum Genet* 2000; **67**: 146–154.
- 34 Chung RH, Hauser ER, Martin ER: The APL test: extension to general nuclear families and haplotypes and examination of its robustness. *Hum Hered* 2006; **61**: 189–199.
- 35 Blanton SH, Cortez A, Stal S, Mulliken JB, Finnell RH, Hecht JT: Variation in IRF6 contributes to nonsyndromic cleft lip and palate. *Am J Med Genet A* 2005; **137**: 259–262.
- 36 Li M, Atmaca-Sonmez P, Othman M et al: CFH haplotypes without the Y402H coding variant show strong association with susceptibility to age-related macular degeneration. *Nat Genet* 2006; **38**: 1049–1054.
- 37 Maller J, George S, Purcell S et al: Common variation in three genes, including a noncoding variant in CFH, strongly influences risk of age-related macular degeneration. *Nat Genet* 2006; **38**: 1055–1059.
- 38 McWhinney SR, Boru G, Binkley PK et al: Intronic single nucleotide polymorphisms in the RET protooncogene are associated with a subset of apparently sporadic pheochromocytoma and may modulate age of onset. *J Clin Endocrinol Metab* 2003; **88**: 4911–4916.
- 39 Scapoli L, Palmieri A, Martinelli M et al: Strong evidence of linkage disequilibrium between polymorphisms at the IRF6 locus and nonsyndromic cleft lip with or without cleft palate, in an Italian population. *Am J Hum Genet* 2005; **76**: 180–183.
- 40 Zucchero TM, Cooper ME, Maher BS et al: Interferon regulatory factor 6 (IRF6) gene variants and the risk of isolated cleft lip or palate. *N Engl J Med* 2004; **351**: 769–780.
- 41 Reich DE, Goldstein DB: Detecting association in a case-control study while correcting for population stratification.[see comment]. *Genet Epidemiol* 2001; **20**: 4–16.
- 42 Boland GM, Perkins G, Hall DJ, Tuan RS: Wnt 3a promotes proliferation and suppresses osteogenic differentiation of adult human mesenchymal stem cells. *J Cell Biochem* 2004; **93**: 1210–1230.
- 43 Wang Y, Guo N, Nathans J: The role of Frizzled3 and Frizzled6 in neural tube closure and in the planar polarity of inner-ear sensory hair cells. *J Neurosci* 2006; **26**: 2147–2156.
- 44 Chiquet BT, Blanton SH, Burt A et al: Variation in WNT genes is associated with non-syndromic cleft lip with or without cleft palate. *Hum Mol Genet* 2008; **17**: 2212–2218.
- 45 Hu D, Helms JA: The role of sonic hedgehog in normal and abnormal craniofacial morphogenesis. *Development* 1999; **126**: 4873–4884.
- 46 Verlinsky Y, Rechitsky S, Verlinsky O et al: Preimplantation diagnosis for sonic hedgehog mutation causing familial holoprosencephaly. *N Engl J Med* 2003; **348**: 1449–1454.
- 47 Rice R, Spencer-Dene B, Connor EC et al: Disruption of Fgf10/Fgfr2b-coordinated epithelial-mesenchymal interactions causes cleft palate. *J Clin Invest* 2004; **113**: 1692–1700.

- 48 Orioli IM, Vieira AR, Castilla EE, Ming JE, Muenke M: Mutational analysis of the Sonic Hedgehog gene in 220 newborns with oral clefts in a South American (ECLAMC) population. *Am J Med Genet* 2002; **108**: 12–15.
- 49 Benayed R, Gharani N, Rossman I *et al*: Support for the homeobox transcription factor gene ENGRAILED 2 as an autism spectrum disorder susceptibility locus. *Am J Hum Genet* 2005; **77**: 851–868.
- 50 Chenevix-Trench G, Jones K, Green AC, Duffy DL, Martin NG: Cleft lip with or without cleft palate: associations with transforming growth factor alpha and retinoic acid receptor loci. *Am J Hum Genet* 1992; **51**: 1377–1385.
- 51 Tissier-Seta JP, Mucchielli ML, Mark M, Mattei MG, Goridis C, Brunet JF: Barx1, a new mouse homeodomain transcription factor expressed in cranio-facial ectomesenchyme and the stomach. *Mech Dev* 1995; **51**: 3–15.
- 52 Cox TC: Taking it to the max: the genetic and developmental mechanisms coordinating midfacial morphogenesis and dysmorphology. *Clin Genet* 2004; **65**: 163–176.
- 53 Hahn H, Wicking C, Zaphiropoulos PG *et al*: Mutations of the human homolog of *Drosophila* patched in the nevoid basal cell carcinoma syndrome. *Cell* 1996; **85**: 841–851.
- 54 Trueba SS, Auge J, Mattei G *et al*: PAX8, TITF1, and FOXE1 gene expression patterns during human development: new insights into human thyroid development and thyroid dysgenesis-associated malformations. *J Clin Endocrinol Metab* 2005; **90**: 455–462.
- 55 Machida J, Moreno LM, Mansilla MA, Bullard SB *et al*: The role of FOXE1 in the etiology of cleft lip, 2005. 57th Annual Meeting of the American Society of Human Genetics San Diego, CA, 470p.
- 56 Vieira AR, Avila JR, Daack-Hirsch S *et al*: Medical sequencing of candidate genes for nonsyndromic cleft lip and palate. *PLoS Genet* 2005; **1**: e64.

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)