

ARTICLE

Genomic analysis of Meckel–Gruber syndrome in Arabs reveals marked genetic heterogeneity and novel candidate genes

Ranad Shaheen¹, Eissa Faqeih², Muneera J Alshammari^{1,3}, Abdulrahman Swaid⁴, Lihadh Al-Gazali⁵, Elham Mardawi⁶, Shinu Ansari¹, Sameera Sogaty⁷, Mohammed Z Seidahmed⁸, Muhammed I AlMotairi¹, Chantal Farra⁹, Wesam Kurdi¹⁰, Shatha Al-Rasheed⁴ and Fowzan S Alkuraya^{*,1,3,11}

Meckel–Gruber syndrome (MKS, OMIM #249000) is a multiple congenital malformation syndrome that represents the severe end of the ciliopathy phenotypic spectrum. Despite the relatively common occurrence of this syndrome among Arabs, little is known about its genetic architecture in this population. This is a series of 18 Arab families with MKS, who were evaluated clinically and studied using autozygome-guided mutation analysis and exome sequencing. We show that autozygome-guided candidate gene analysis identified the underlying mutation in the majority ($n = 12$, 71%). Exome sequencing revealed a likely pathogenic mutation in three novel candidate MKS disease genes. These include *C5orf42*, Ellis–van-Creveld disease gene *EVC2* and *SEC8* (also known as *EXOC4*), which encodes an exocyst protein with an established role in ciliogenesis. This is the largest and most comprehensive genomic study on MKS in Arabs and the results, in addition to revealing genetic and allelic heterogeneity, suggest that previously reported disease genes and the novel candidates uncovered by this study account for the overwhelming majority of MKS patients in our population.

European Journal of Human Genetics (2013) **21**, 762–768; doi:10.1038/ejhg.2012.254; published online 21 November 2012

Keywords: autozygome; ciliopathy; encephalocele; EVC2; EXOC4

INTRODUCTION

Meckel–Gruber syndrome (MKS, OMIM #249000) is a multiple congenital malformation syndrome characterized mainly by early lethality, occipital encephalocele, dysplastic kidneys, and polydactyly, but the phenotype also includes liver fibrosis, microphthalmia, cleft palate, and a wide-array of associated brain anomalies.¹ MKS tends to be a rare disease in most populations (<1:20 000) with the notable exception of Finland where a birth prevalence of 1:9000 was estimated, and Arabia where Teebi and Teebi reported a much higher frequency of 1:3500.^{2,3}

Our understanding of MKS pathogenesis has improved greatly over the past few years since it was first revealed that *MKS1*, the first gene found to be mutated in MKS, encodes a ciliary protein.⁴ Subsequently, nine additional genes have been identified, all similarly encoding ciliary proteins (*TMEM216* (MKS2), *TMEM67* (MKS3), *CEP290* (MKS4), *RPGRIP1L* (MKS5), *CC2D2A* (MKS6), *NPHP3* (MKS7), *TCTN2* (MKS8), *B9D1* (MKS9) and *B9D2* (MKS10)).^{4–14} The finding that defective ciliary biology is the core molecular pathology of MKS made it possible to dissect the pathogenesis of each of its manifestations. For instance, the primary cilium plays a critical role in SHH signaling that controls anterior–

posterior and dorsal–ventral patterning of the developing limb buds and neural tube, respectively, thus explaining the polydactyly and neural tube defects that characterize MKS at the molecular level. Similarly, the third classical feature of cystic dysplasia of the kidneys is likely caused by abnormal patterning secondary to defective urine flow-induced signaling that is normally transduced by the cilia on the luminal surface of renal epithelium.

Defective ciliary biology is not unique to MKS; an expansive group of disorders commonly known as ciliopathies share this fundamental pathological feature.¹⁵ In fact, mutations in MKS genes have been reported to cause other ciliopathies, for example, *MKS1* and Bardet–Biedl syndrome,¹⁶ *TMEM216* and Joubert syndrome,¹⁷ *TMEM67* and Joubert syndrome and nephrophthisis,^{18,19} *CEP290* in non-syndromic retinal dystrophy, Senior–Loken syndrome, nephronophthisis, Joubert syndrome, and Bardet–Biedl syndrome,²⁰ *RPGRIP1L* and Joubert syndrome,²¹ and *CC2D2A* and Joubert syndrome.²² The factors that determine the ultimate clinical phenotype are not completely understood but there is growing evidence that ciliopathies represent a spectrum of clinical severity that correlates to some extent with the severity of the ciliary defect. Consistent with MKS being at the severe end of the clinical spectrum, most causative mutations are truncating

¹Department of Genetics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia; ²Department of Pediatrics, King Fahad Medical Complex, Riyadh, Saudi Arabia; ³Department of Pediatrics, King Khalid University Hospital and College of Medicine, King Saud University, Riyadh, Saudi Arabia; ⁴Department of Pediatrics, King Abdulaziz Medical City, Riyadh, Saudi Arabia; ⁵Department of Pediatrics, Faculty of Medicine and Health Sciences, United Arab Emirates University, Al-Ain, UAE; ⁶Department of Obstetrics and Gynecology, Security Forces Hospital, Riyadh, Saudi Arabia; ⁷Department of Medical Genetic, Jeddah King Fahad General Hospital, Jeddah, Saudi Arabia; ⁸Department of Pediatrics, Security Forces Hospital, Riyadh, Saudi Arabia; ⁹Department of Pediatrics, American University of Beirut, Beirut, Lebanon; ¹⁰Department of Obstetrics and Gynecology, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia; ¹¹Department of Anatomy and Cell Biology, College of Medicine, Alfaisal University, Riyadh, Saudi Arabia

*Correspondence: Professor FS Alkuraya, Developmental Genetics Unit, King Faisal Specialist Hospital and Research Center, MBC-03 PO Box 3354, Riyadh 11211, Saudi Arabia. Tel: +966 1 442 7875; Fax: +966 1 442 4585; E-mail: falkuraya@kfshrc.edu.sa

Received 21 June 2012; revised 18 September 2012; accepted 25 September 2012; published online 21 November 2012

in nature while hypomorphic mutations in the same genes cause less severe phenotypes.²³

The genetic heterogeneity of MKS has yet to be fully captured. Indeed, a recent study showed that only half of the cases can be traced to a mutation in any of the known genes (seven were included in the analysis, so the additional contribution of *TCTN2* could not be inferred).¹⁴ The autosomal recessive nature of MKS, the highly consanguineous nature of the Arab population where MKS is particularly prevalent, and the success of exome sequencing in other recessive disorders encouraged us to undertake a comprehensive genomic study to fully delineate the genetic architecture of MKS in our population by combining the power of autozygome and exome analysis. In this largest genomic MKS case series in Arabs to date, we show that autozygome-guided candidate gene analysis successfully identified the underlying mutation in >70% of the cases. In the remaining cases, we used exome sequencing followed by autozygome filtration to identify three novel candidate genes, one of which was very recently reported to be also mutated in Joubert syndrome. Assuming the candidate MKS genes that we report here are replicated by future studies, it appears that most of the genetic heterogeneity of MKS, at least in Arabs, has been captured.

MATERIALS AND METHODS

Human subjects

For case definition, we used a relaxed definition of occipital encephalocele as a must-have feature plus any combination of cleft palate, dysplastic kidneys, liver fibrosis, polydactyly, and early lethality. All cases were fully evaluated by an experienced dysmorphologist and/or neonatologist. Eligible cases were recruited after obtaining IRB-approved informed written consent from the parents (KFHSRC RAC#2080006). Blood in EDTA and, when possible, PAXGene tubes was collected for DNA and RNA extraction, respectively.

Autozygome analysis

DNA samples from both affected and unaffected family members were genotyped on Affymetrix Chip platform as per the manufacturer's protocol (Affymetrix, Santa Clara, CA, USA). Runs of homozygosity (ROH) >2 Mb and that span 107 SNPs were used as surrogates of autozygosity given the consanguineous nature of the families using autoSNPa (<http://dna.leeds.ac.uk/autosnpa/>). The entire set of autozygosity blocks (autozygome) was determined for each patient. The shared haplotype between parents as well as relatives with similarly affected children was determined by using IBDelphi (<http://dna.leeds.ac.uk/ibdelfi/>).

Exome sequencing and analysis

Exome capture was performed using TruSeq Exome Enrichment kit (Illumina, San Diego, CA, USA) following the manufacturer's protocol. Samples were prepared as an Illumina sequencing library, and in the second step, the sequencing libraries were enriched for the desired target using the Illumina Exome Enrichment protocol. The captured libraries were sequenced using Illumina HiSeq 2000 Sequencer. The reads were mapped against UCSC hg19 (<http://genome.ucsc.edu/>) by BWA (<http://bio-bwa.sourceforge.net/>). The SNPs and Indels were detected by SAMTOOLS (<http://samtools.sourceforge.net/>). The resulting variants were filtered by only considering homozygous novel changes (absent in dbSNP, 1000 Genomes and 200 in-house Saudi exomes) within the autozygome followed by *in silico* analysis for potential pathogenicity. Variants that survived these filters were then Sanger sequenced in at least 192 normal Saudi controls, that is, 384 chromosomes.

Workflow

As summarized in Figure 1, cases were first subjected to autozygome-guided mutation analysis of known MKS genes. If negative, cases were exome sequenced. Exomes were first examined for potential compound heterozygous mutations in known MKS genes since these will be missed by the autozygome




Figure 1 Workflow of the current study.

approach. If negative, we proceeded with the filtration scheme outlined above (see Exome Sequencing).

RESULTS

Human subjects

In total, 18 families that met the inclusion criteria were recruited, all consanguineous. All pedigrees are shown in Supplementary Figure S1 and representative clinical pictures for some of the MKS patients are shown in Figure 2. Table 1 summarizes their clinical features. One family was initially recruited despite lack of additional features of MKS besides occipital encephalocele because there were no ultrasonographic features suggestive of another diagnosis and the fetus was unavailable for clinical assessment after delivery. This case was later found to have two base pair deletion in *POMT1* gene (NM_007171.3:c.2179_2180delTC), so it was excluded and is not considered further in this study although it serves as a reminder that encephalocele is a birth defect with a broad differential diagnosis.²⁴ As expected, our relaxed clinical definition resulted in the recruitment of two cases (MKS_F1 and MKS_F15) that do not meet the classical minimal definition of early lethality, encephalocele, and cystic/dysplastic kidneys. However, we opted to retain these cases in our molecular analysis to explore the extent of variable expressivity in MKS especially when these cases were subsequently found to harbor mutations in genes recently reported to cause Joubert syndrome (see below).

Autozygome-guided mutational analysis

Consistent with all families being consanguineous, all index cases had evidence of autozygosity and the percentage of their genomes represented by the autozygome ranged 2.5–18.4% (Supplementary Table S1). At the time of the study, 10 MKS disease genes were identified (*TMEM67*, *TMEM216*, *CEP290*, *CC2D2A*, *NPHP3*, *RPGRIP1L*, *TCTN2*, *B9D1*, and *B9D2*). Gene(s) that overlap with autozygome were Sanger sequenced and this led to the identification of a likely pathogenic homozygous mutation in 9 of 17 families (Figure 1). Supplementary Figure S2 shows both the multi-species alignment of orthologs as well as the PolyPhen/SIFT scores for each of




Figure 2 (a–c) Facial features of some MKS patients MKS_F1, F2, and F8, respectively, showing: microcephaly, sloping forehead, hypertelorism, micrognathia, potter-like facies, and severe occipital encephalocele (black arrows). (d, e) Representative hand and foot images showing postaxial polydactyly in MKS_F2. (f, g) Antenatal ultrasound for MKS_F12 patient showed encephalocele (white arrow) and enlarged polycystic kidney (dashes arrows).

the novel missense variants. For three families (MKS_F4, MKS_F11, and MKS_F13), we had no access to the index so we used IBDelphi to infer the shared ancestral haplotype between the parents which helped us identify the following pathogenic mutations: c.1506-2A>G in *TCTN2* (NM_024809.3), c.1855_1858del in *CEP290* (NM_025114) and c.613C>T; p.R205* in *CEP290* (NM_025114.3), respectively. Thus autozygome-guided mutational analysis and analysis to the shared haplotype between parents led to the successful identification of the underlying mutation in 71% of cases (Table 2; Figure 3).

Autozygome-filtered exome sequencing

Five families in which autozygome-guided mutation analysis failed to identify the causative mutation in our initial list of 10 genes were exome sequenced (Figure 1). Summary of the depth of coverage and the amount of exome covered are shown in Supplementary Table S2. Variants were excluded when present homozygous in Saudi controls and heterozygous at a frequency > 1% ie present twice in 200 exomes. The filtration steps used in interpreting the exome results are summarized in Supplementary Table S3 and the novel variants that survived the various filters are listed in Table 3. MKS_F1 was found to harbor a novel homozygous mutation that abolished a consensus donor site in *TMEM237* (c.869 + 1G>A) and RT-PCR confirmed the resulting exonic skipping, which predicts in-frame deletion of > 60 amino acids p(Ap.Met227_Arg291del). Mutations in this gene have recently been reported to cause a phenotypic spectrum between Joubert syndrome and MKS.²⁵ MKS_F15 was found to harbor the same homozygous truncating mutation we recently reported in a gene that causes Joubert syndrome (*C5orf42*).^{26,27} Two exomes (MKS_16 and MKS_2) highlighted novel MKS candidate genes: *EXOC4* and *EVC2*, respectively, whereas exome sequencing of MKS_F18 failed to

generate any compelling candidate variants after applying the various filters (Figure 1; Supplementary Table S4).

DISCUSSION

Although many MKS genes were mapped using consanguineous Arab families, the mutation distribution of these genes in Arabs is unknown but deserves investigation to inform clinical sequencing and genetic counseling. Furthermore, the success we have had in combining autozygome and exome analysis in delineating the genetic architecture of other genetically heterogeneous disorders, for example, Osteogenesis imperfecta, retinal dystrophy, cataract, mitochondrial diseases, Bardet–Biedl syndrome, and Joubert syndrome^{26,28–31} (and Abu Safieh *et al* (2012)³², Shaheen *et al* (2012)³³), encouraged us to use a similar approach on MKS to not only determine the mutation distribution in known MKS disease genes but to also potentially identify novel candidate disease genes.

To address the first aim, that is, determination of the mutation distribution in known MKS genes, we took advantage of the consanguineous nature of our study population to trace causative mutations by virtue of the autozygosity signature that typically characterizes identical-by-descent recessive mutations. Indeed, we show that autozygome-guided mutational analysis has efficiently identified the causative variant in each family in which MKS is caused by a mutation in a known gene. Importantly, even though the mutations in *TMEM237* and *C5orf42* were identified by exome sequencing (these were unknown to cause MKS at the time of the analysis), both could have easily been identified using the same approach further increasing the yield of this approach to 88% (15 out of 17) in the setting of MKS. This is consistent with the trend we and others have shown in terms of the power of this approach in the

Table 1 Summary of clinical description of MKS individuals (classical features are underlined>)

Patient features	MKS_F1	MKS_F2	MKS_F3	MKS_F4	MKS_F5	MKS_F6	MKS_F7	MKS_F8	MKS_F9	MKS_F10	MKS_F11	MKS_F12	MKS_F13	MKS_F14	MKS_F15	MKS_F16	MKS_F17	MKS_F18
Gene	TMEM237	EVC2	CC2D2A	TCTN2	CC2D2A	TMEM67	MKS1	CC2D2A	TCTN2	CC2D2A	CC2D2A	CC2D2A	CEP290	CC2D2A	C5orf42	EXOC4	MKS1	
<u>Early death</u>	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<u>Microcephaly</u>	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<u>Variable prenatal growth deficiency</u>	?	-	+	?	+	+	?	?	?	+	+	+	+	?	-	?	?	-
<u>Head and neck</u>																		
<u>Potter-like facies</u>	-	+	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Sloping forehead</u>	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Microphthalmia</u>	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Hypertelorism</u>	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Low-set ears</u>	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Cleft lip</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Cleft palate</u>	-	-	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Macrostomia</u>	?	+	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Lobulated tongue</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Natal teeth</u>	?	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Short neck</u>	+	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Webbed neck</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>CNS</u>																		
<u>Occipital encephalocele</u>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<u>Anencephaly</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Hydrocephalus</u>	+	-	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Dandy-Walker malformation</u>	+	?	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Arnold-Chiari malformation</u>	-	?	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Cerebral hypoplasia</u>	+	?	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>CVS</u>																		
<u>Patent ductus arteriosus</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Lung</u>																		
<u>Pulmonary hypoplasia</u>	-	?	+	?	+	+	?	?	?	?	?	?	?	?	?	?	?	?
<u>Urinary system</u>																		
<u>Polycystic kidneys</u>	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<u>Ambiguous genitalia</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Extremities</u>																		
<u>Syndactyly</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Polydactyly</u>	-	+	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Bowed long bones</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Talipes</u>	-	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
<u>Oligohydramnios</u>	?	?	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<u>Prenatal diagnosis by ultrasound</u>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

Table 2 Summary of autozygome-guided mutation analysis

Family ID	No. of affected used for analysis	MKS genes within autozygome or within shared haplotype of the parents		Sequence analysis results			Reference
MKS_F1 ^a	1	<i>TMEM237</i>	<i>TMEM237</i>	NM_001044385.2	c.869+1G>A	p.Met227_Arg291del	Current study
MKS_F2 ^a	1	<i>TMEM67</i>	<i>EVC2</i>	NM_147127.4	c.3870_3893dup	p.Lys1293_Lys1300dup	Current study
MKS_F3	1	<i>CC2D2A</i>	<i>CC2D2A</i>	NM_001080522.2	c.3084delG	p.Lys1029Argfs ^b 3	12
MKS_F4 ^b	–	<i>CEP290, TCTN2, TMEM237</i>	<i>TCTN2</i>	NM_024809.3	c.1506-2A>G		13
MKS_F5	1	<i>CC2D2A</i>	<i>CC2D2A</i>	NM_001080522.2	c.3084delG	p.Lys1029Argfs ^b 3	12
MKS_F6	1	<i>TMEM67</i>	<i>TMEM67</i>	NM_153704.5	c.2306del T	p.Leu769Tyrfs ^b 4	Current study
MKS_F7	1	<i>NPHP3, MKS1, CC2D2A</i>	<i>MKS1</i>	NM_017777.3	c.1126dupA	p.Thr376Asnfs ^b 3	Current study
MKS_F8	1	<i>CC2D2A</i>	<i>CC2D2A</i>	M_001080522.2	c.4531T>C	p.Trp1511Arg	Current study
MKS_F9	1	<i>TCTN2</i>	<i>TCTN2</i>	NM_024809.3	c.1506-2A>G		13
MKS_F10	1	<i>CC2D2A</i>	<i>CC2D2A</i>	NM_001080522.2	c.3084delG	p.Lys1029Argfs ^b 3	12
MKS_F11 ^b	–	<i>TMEM67, TCTN2, CEP290</i>	<i>CEP290</i>	NM_025114.3	c.1860_1863delAAGA	p.Arg621Ilefs ^b 2	Current study
MKS_F12	1	<i>TMEM67, CC2D2A</i>	<i>CC2D2A</i>	NM_001080522.2	c.3084delG	p.Lys1029Argfs ^b 3	12
MKS_F13 ^b	–	<i>CEP290, TMEM67, MKS1, TCTN2</i>	<i>CEP290</i>	NM_025114.3	c.613C>T	p.R205 ^b	9
MKS_F14	1	<i>B9D1, TMEM216, CC2D2A, NPHP3</i>	<i>CC2D2A</i>	NM_001080522.2	c.4531T>C	p.Trp1511Arg	Current study
MKS_F15 ^a	1	<i>CEP290</i>	<i>C5orf42</i>	NM_023073.3	c.7988_7989delGA	p.G2663Afs ^b 40	Current study
MKS_F16 ^a	1	–	<i>EXOC4</i>	NM_021807.3	c.1733A>G	p.Gln578Arg	Current study
MKS_F17 ^c	–	–	<i>MKS1</i>	NM_017777.3	c.1126dupA	p.Thr376Asnfs ^b 3	Current study
MKS_F18	1	<i>C5orf42, B9D1</i>	–	–	–	–	–

^aUncovered using exome capture.^bAnalysis was done on parents using IBDelphi.^cAnalysis was done by direct sequencing.

setting of genetically heterogeneous autosomal recessive diseases.^{28,29,34}

In order to explore the potential of revealing novel disease genes, all cases in which autozygome-guided mutational analysis was negative were exome sequenced. As we have shown previously, autozygome served as an extremely powerful filter of the resulting variants.^{35,36} In Family MKS_F1, two variants survived filtration but *TMEM237* was recently described as a novel Joubert gene.²⁵ Some of the original reported cases had occipital encephalocele and cystic kidney so it seems likely that our mutation is the causal variant.

In Family MKS_F15, only two variants survived the various filters (*C5orf42*, NM_023073.3; c.7988_7989delGA and *FAM48A*, NM_001014286; c.236A>G). We could not determine which of the two is likely to be causal because while the former is truncating in nature, the latter missense mutation affected a gene that is clearly linked to neural tube defects in mice.³⁷ During the preparation of this manuscript, we and others identified several truncating mutations in *C5orf42* in the context of Joubert syndrome.^{26,27} Since MKS is a more severe phenotype, it is tempting to speculate that the additional allele in *FAM48A* may have influenced the severity of the phenotype. However, we caution against the overinterpretation of the presence of the *FAM48A* allele because occipital encephalocele/meningocele has been previously reported in the setting of *C5orf42* mutations.^{26,27} Indeed, we note that case MKS_F15, while meeting our operational clinical definition of MKS, did not have polydactyly or renal involvement so it can be argued that even if *C5orf42* is indeed an MKS disease gene, it causes atypical forms of the disease. It is worth highlighting that the only two cases in our series that did not meet the classical definition of MKS are those with mutations in *TMEM237* and *C5orf42*, which have been shown to cause both classical Joubert syndrome as well as cases that are Meckel-like as we explained above. Only through analysis of the full extent of *C5orf42* and *TMEM237* contribution to MKS using large case series will it be apparent if classical MKS can also be caused by mutations in these genes.

Families MKS_F2 and MKS_F16 deserve special emphasis. In Family MKS_F2, the child had all the classical features of MKS and yet his exome sequencing revealed a novel variant in *EVC2*, a known

disease gene for Ellis-van-Creveld syndrome. Although the mutation is in-frame, it inserts eight amino acids p.Lys1293_Lys1300dup. Interestingly, mutations in *EVC2* have been shown to modulate SHH signaling. Furthermore, *EVC2* has been found recently to localize at the basal body of the primary cilium so it is tempting to speculate that this may be the link between *EVC2* and MKS.^{38–40} Thus, our results are build on recent data to suggest that *EVC* is a ciliopathy. However, we note that lack of *EVC* findings on skeletal survey and fetal echocardiography in this case strongly argue that this is a bona fide MKS phenotype rather than a simple clinical overlap between *EVC* and MKS as was once reported.⁴¹ In Family MKS_F16, the variant we identified in *EXOC4* (also known as *SEC8*) is particularly interesting. The gene encodes one of the several exocyst proteins that are recruited to the basal body of budding cilia along with the members of disheveled family of proteins in order to dock the basal body to the membrane and allow the nascent cilia to form.^{42,43} Knockdown of another exocyst gene *SEC10* was found to impair ciliogenesis, raising the possibility that defective *SEC8* may lead a similar cellular phenotype. Furthermore, another exocyst gene *EXOC8* was recently found mutated in a patient with Joubert syndrome.⁴⁴ Of note, the variant we identified is conserved down to Fugu and zebrafish.

One of the cases (MKS_F18) was not solved by Exome sequencing. In our experience, only around 70% of autosomal recessive cases are solved by this techniques and this could be explained by various reasons. First, the mutation may not be in the exon or exon/intron boundary. Second, the mutation may be in an exon that has not been annotated and thus not included in the capture design. Third, the mutation may belong to a class that is particularly difficult to assay using next-generation sequencing (eg, genomic rearrangements, repeats, etc). Fourth, the mutation may simply have been skipped by either the capture or sequencing reaction.

Of the three proposed novel MKS candidate genes, *C5orf49* has been independently confirmed through recent reports of Joubert syndrome patients with biallelic mutations in these genes.^{26,27} *EVC2* and *EXOC4*, however, will have to await future confirmation as bona fide MKS loci. If independently confirmed, our study shows that, at




Figure 3 Sequence chromatograms of novel mutations listed in Table 2. Mutation sites are denoted with asterisks or lines.

Table 3 Variants that survived filtration in each exome

Family	Name of the gene	Reference sequence	Nucleotide level	Protein level	Comments
MKS_F1	<i>TMEM237</i>	NM_001044385.2	c.869 + 1G>A	p.Met227_Arg291del	<i>TMEM237</i> was recently described as a novel Joubert gene
	<i>NCOA6</i>	NM_014071.3	c.822_830del	p.Gln283_Gln285del	
MKS_F2	<i>EVC2</i>	NM_147127.4	c.3870_3893dup	p.Lys1293_Lys1300dup	<i>EVC2</i> has been found to localize at the basal body of the primary cilium and modulate SHH signaling
MKS_F15	<i>C5orf42</i>	NM_023073.3	c.7988_7989delGA	p.Gly2663Alafs*40	<i>C5orf42</i> was recently described as a novel Joubert gene
	<i>FAM48A</i>	NM_001014286.2	c.236A>G	p.Asn79Ser	<i>FAM48A</i> is linked to neural tube defect in mice
MKS_F16	<i>TXNDC15^a</i>	NM_024715.3	c.673_687del	p.Ser225_his229del	<i>TXNDC15</i> : Two stop gain mutations are reported in the Exome Variant Server
	<i>ETV1^a</i>	NM_001163149.1	c.1234C>G	p.Gln412Glu	<i>ETV1</i> : Has incompatible mouse model
	<i>EXOC4^a</i>	NM_021807.3	c.1733A>G	p.Gln578Arg	<i>EXOC4</i> : Encodes exocyst proteins that are recruited to the basal body of budding cilia in order to dock the basal body to the membrane and allow the nascent cilia to form. There are no reported truncating mutation in the Exome Variant Server

^aPresent in the shared haplotype between the two parents who have affected children with MKS within the same family.

least in Arabs, most of the genetic heterogeneity of this disorder has been captured and that the contribution of future MKS loci is likely to be minimal. One previous study showed that seven MKS genes (*TMEM67*, *TMEM216*, *CEP290*, *CC2D2A*, *B9D1*, and *RPGRIP1L*) were found to be biallelically mutated in 54% of their cohort, a higher contribution for these particular genes than what we observed in our study sample (32%). Thus, it seems possible that the higher percentage of mutation-positive MKS cases (94% by including

TCTN2, *TMEM237*, *C5orf42*, and if *EVC2* and *EXOC4* are confirmed by future reports) may potentially be relevant to other populations as well.

In summary, we show in the largest genomic study on an MKS cohort the mutation distribution among Arabs where the disease is particularly prevalent. This genomic approach revealed interesting novel candidates that may potentially bring, along with recently described MKS disease genes, the quest to unravel the genetics of

MKS to its final stages making it possible for the overwhelming majority of these patients to receive genetic counseling that is informed by molecular confirmation.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the study families for their enthusiastic participation. We thank Dr Mohammed Al-Balawi for his help in extracting and storing DNA samples. We also thank the Genomic and Sequencing Core Facilities at KFSHRC for their technical help. This study was funded in part by KACST Grant 09-MED941-20 (FSA) and DHFMR Collaborative Research Grant (FSA).

Author contributions: Ranad Shaheen and Fowzan S Alkuraya: collected and analyzed the data and wrote the manuscript. Eissa Faqeih, Muneera J Alshammari, Abdulrahman Swaid, Lihadh Al-Gazali, Elham Mardawi, Shinu Ansari, Mohammed Z Seidahmed, Muhammed Mutairi, Chantal Farra, Wesam Kurdi, and Shatha Al-Rasheed: collected and analyzed the data.

- Seller MJ: Phenotypic variation in Meckel syndrome. *Clin Genet* 1981; **20**: 74–77.
- Teebi AS, Teebi SA: Genetic diversity among the Arabs. *Community Genet* 2005; **8**: 21–26.
- Salonen R, Norio R: The Meckel syndrome in Finland: epidemiologic and genetic aspects. *Am J Med Genet* 1984; **18**: 691–698.
- Kyttala M, Tallila J, Salonen R *et al*: MKS1, encoding a component of the flagellar apparatus basal body proteome, is mutated in Meckel syndrome. *Nat Genet* 2006; **38**: 155–157.
- Consugar MB, Kubly VJ, Lager DJ *et al*: Molecular diagnostics of Meckel-Gruber syndrome highlights phenotypic differences between MKS1 and MKS3. *Hum Genet* 2007; **121**: 591–599.
- Dawe HR, Smith UM, Cullinane AR *et al*: The Meckel-Gruber Syndrome proteins MKS1 and meckelin interact and are required for primary cilium formation. *Hum Mol Genet* 2007; **16**: 173–186.
- Valente EM, Logan CV, Mougou-Zerelli S *et al*: Mutations in TMEM216 perturb ciliogenesis and cause Joubert, Meckel and related syndromes. *Nat Genet* 2010; **42**: 619–625.
- Smith UM, Consugar M, Tee LJ *et al*: The transmembrane protein meckelin (MKS3) is mutated in Meckel-Gruber syndrome and the wpk rat. *Nat Genet* 2006; **38**: 191–196.
- Baala L, Audollent S, Martinovic J *et al*: Pleiotropic effects of CEP290 (NPHP6) mutations extend to Meckel syndrome. *Am J Hum Genet* 2007; **81**: 170–179.
- Frank V, den Hollander AI, Bruchle NO *et al*: Mutations of the CEP290 gene encoding a centrosomal protein cause Meckel-Gruber syndrome. *Hum Mutat* 2008; **29**: 45–52.
- Delous M, Baala L, Salomon R *et al*: The ciliary gene RPGRIP1L is mutated in cerebello-oculo-renal syndrome (Joubert syndrome type B) and Meckel syndrome. *Nat Genet* 2007; **39**: 875–881.
- Tallila J, Jakkula E, Peltonen L, Salonen R, Kestila M: Identification of CC2D2A as a Meckel syndrome gene adds an important piece to the ciliopathy puzzle. *Am J Hum Genet* 2008; **82**: 1361–1367.
- Shaheen R, Faqeih E, Seidahmed MZ *et al*: A TCTN2 mutation defines a novel Meckel Gruber syndrome locus. *Hum Mutat* 2011; **32**: 573–578.
- Hopp K, Heyer CM, Hommerding CJ *et al*: B9D1 is revealed as a novel Meckel syndrome (MKS) gene by targeted exon-enriched next-generation sequencing and deletion analysis. *Hum Mol Genet* 2011; **20**: 2524–2534.
- Nigg EA, Raff JW: Centrioles, centrosomes, and cilia in health and disease. *Cell* 2009; **139**: 663–678.
- Leitch CC, Zaghoul NA, Davis EE *et al*: Hypomorphic mutations in syndromic encephalocoele genes are associated with Bardet-Biedl syndrome. *Nat Genet* 2008; **40**: 443–448.
- Edvardson S, Shaag A, Zenvirt S *et al*: Joubert syndrome 2 (JBTS2) in Ashkenazi Jews is associated with a TMEM216 mutation. *Am J Hum Genet* 2010; **86**: 93–97.
- Baala L, Romano S, Khaddour R *et al*: The Meckel-Gruber syndrome gene, MKS3, is mutated in Joubert syndrome. *Am J Hum Genet* 2007; **80**: 186–194.
- Otto EA, Tory K, Attanasio M *et al*: Hypomorphic mutations in meckelin (MKS3/TMEM67) cause nephronophthisis with liver fibrosis (NPHP11). *J Med Genet* 2009; **46**: 663–670.
- Coppieters F, Lefever S, Leroy BP, De Baere E: CEP290, a gene with many faces: mutation overview and presentation of CEP290base. *Hum Mutat* 2010; **31**: 1097–1108.
- Arts HH, Doherty D, van Beersum SE *et al*: Mutations in the gene encoding the basal body protein RPGRIP1L, a nephrocystin-4 interactor, cause Joubert syndrome. *Nat Genet* 2007; **39**: 882–888.
- Gorden NT, Arts HH, Parisi MA *et al*: CC2D2A is mutated in Joubert syndrome and interacts with the ciliopathy-associated basal body protein CEP290. *Am J Hum Genet* 2008; **83**: 559–571.
- Salonen R, Kestila M, Bergmann C: Clinical utility gene card for: Meckel syndrome. *Eur J Hum Genet* 2011; **19**: e-pub ahead of print 2 February 2011; doi:10.1038/ejhg.2010.255.
- Cohen Jr. MM, Lemire RJ: Syndromes with cephaloceles. *Teratology* 1982; **25**: 161–172.
- Huang L, Szymanska K, Jensen VL *et al*: TMEM237 is mutated in individuals with a Joubert syndrome related disorder and expands the role of the TMEM family at the ciliary transition zone. *Am J Hum Genet* 2011; **89**: 713–730.
- Alazami AM, Alshammari MJ, Salih MA *et al*: Molecular characterization of Joubert syndrome in Saudi Arabia. *Hum Mutat* 2012; **33**: 1423–1428.
- Srouf M, Schwartzentruber J, Hamdan FF *et al*: Mutations in C5ORF42 cause Joubert syndrome in the French Canadian population. *Am J Hum Genet* 2012; **90**: 693–700.
- Shamseldin HE, Alshammari M, Al-Sheddi T *et al*: Genomic analysis of mitochondrial diseases in a consanguineous population reveals novel candidate disease genes. *J Med Genet* 2012; **49**: 234–241.
- Abu Safieh L, Aldahmesh MA, Shamseldin H *et al*: Clinical and molecular characterisation of Bardet-Biedl syndrome in consanguineous populations: the power of homozygosity mapping. *J Med Genet* 2010; **47**: 236–241.
- Abu-Safieh L, Al-Anazi S, Al-Abdi L *et al*: In search of triallelism in Bardet-Biedl syndrome. *EJHG* 2012; **20**: 420–427.
- Aldahmesh MA, Khan AO, Mohamed JY *et al*: Genomic analysis of pediatric cataract in Saudi Arabia reveals novel candidate disease genes. *Genet Med* 2012.
- Abu-Safieh L, Alrashed M, Anazi S *et al*: Autozygome-guided exome sequencing in retinal dystrophy patients reveals pathogenetic mutations and novel candidate disease genes. *Genome Res* 2012; e-pub ahead of print 26 October 2012.
- Shaheen R, Alazami AM, Alshammari MJ *et al*: Study of autosomal recessive osteogenesis imperfecta in Arabia reveals a novel locus defined by TMEM38B mutation. *J Med Genet* 2012; **49**: 630–635.
- Shaheen R, Al-Dirbashi OY, Al-Hassnan ZN *et al*: Clinical, biochemical and molecular characterization of peroxisomal diseases in Arabs. *Clin Genet* 2011; **79**: 60–70.
- Shaheen R, Faqeih E, Sunker A *et al*: Recessive mutations in DOCK6, encoding the guanidine nucleotide exchange factor DOCK6, lead to abnormal actin cytoskeleton organization and Adams-Oliver syndrome. *Am J Hum Genet* 2011; **89**: 328–333.
- Aldahmesh MA, Mohamed JY, Alkuraya HS *et al*: Recessive mutations in ELOVL4 cause ichthyosis, intellectual disability, and spastic quadriplegia. *Am J Hum Genet* 2011; **89**: 745–750.
- Zohn IE, Li Y, Skolnik EY, Anderson KV, Han J, Niswander L: p38 and a p38-interacting protein are critical for downregulation of E-cadherin during mouse gastrulation. *Cell* 2006; **125**: 957–969.
- Blair HJ, Tompson S, Liu YN *et al*: Evc2 is a positive modulator of Hedgehog signalling that interacts with Evc at the cilia membrane and is also found in the nucleus. *BMC Biol* 2011; **9**: 14.
- Valencia M, Lapunzina P, Lim D *et al*: Widening the mutation spectrum of EVC and EVC2: ectopic expression of Weyer variants in NIH 3T3 fibroblasts disrupts Hedgehog signaling. *Hum Mutat* 2009; **30**: 1667–1675.
- Ruiz-Perez VL, Goodship JA: Ellis-van Creveld syndrome and Weyers acrodermal dysostosis are caused by cilia-mediated diminished response to hedgehog ligands. *Am J Med Genet C Semin Med Genet* 2009; **151C**: 341–351.
- Kemperdick H, Ammermann M, Janssen F, Lange H, Moubayed P: [The differential diagnosis of the Meckel syndrome and the Ellis-van-Creveld syndrome with encephalocoele (author's transl)]. *Klin Padiatr* 1975; **187**: 87–93.
- Ganner A, Lienkamp S, Schafer T *et al*: Regulation of ciliary polarity by the APC/C. *Proc Natl Acad Sci USA* 2009; **106**: 17799–17804.
- Park TJ, Mitchell BJ, Abitua PB, Kintner C, Wallingford JB: Dishevelled controls apical docking and planar polarization of basal bodies in ciliated epithelial cells. *Nat Genet* 2008; **40**: 871–879.
- Dixon-Salazar TJ, Silhavy JL, Udpa N *et al*: Exome sequencing can improve diagnosis and alter patient management. *Sci Transl Med* 2012; **4**: 138ra178.

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)