

Endocytic pathways regulate Toll-like receptor 4 signaling and link innate and adaptive immunity

Harald Husebye^{1,5}, Øyvind Halaas^{1,5},
Harald Stenmark^{1,3}, Gro Tunheim²,
Øystein Sandanger¹, Bjarne Bogen²,
Andreas Brech³, Eicke Latz⁴
and Terje Espevik^{1,*}

¹Institute of Cancer Research and Molecular Medicine, The Norwegian University of Science and Technology, Trondheim, Norway, ²Institute of Immunology, Rikshospitalet University Hospital, Oslo, Norway, ³Department of Biochemistry, The Norwegian Radiumhospital, Oslo, Norway and ⁴Division of Infectious Diseases and Immunology, University of Massachusetts Medical School, Worcester, MA, USA

Immune responses are initiated when molecules of microbial origin are sensed by the Toll-like receptors (TLRs). We now report the identification of essential molecular components for the trafficking of the lipopolysaccharide (LPS) receptor complex. LPS was endocytosed by a receptor-mediated mechanism dependent on dynamin and clathrin and colocalized with TLR4 on early/sorting endosomes. TLR4 was ubiquitinated and associated with the ubiquitin-binding endosomal sorting protein hepatocyte growth factor-regulated tyrosine kinase substrate, Hrs. Inhibition of endocytosis and endosomal sorting increased LPS signaling. Finally, the LPS receptor complex was sorted to late endosomes/lysosomes for degradation and loading of associated antigens onto HLA class II molecules for presentation to CD4⁺ T cells. Our results show that endosomal trafficking of the LPS receptor complex is essential for signal termination and LPS-associated antigen presentation, thus controlling both innate and adaptive immunity through TLR4.

The EMBO Journal (2006) 25, 683–692. doi:10.1038/sj.emboj.7600991; Published online 9 February 2006

Subject Categories: membranes & transport; immunology

Keywords: antigen presentation; endocytosis; sorting; Toll-like receptor (TLR)

Introduction

The Toll-like receptors (TLRs) are key initiators of innate and adaptive immune responses through production of proinflammatory cytokines and chemokines, upregulation of costimulatory molecules and activation of antigen presentation (Akira and Takeda, 2004). The signaling receptor for lipopolysaccharide (LPS) is TLR4/MD-2 receiving LPS from CD14. MD-2 is a small, secreted glycoprotein that associates with

*Corresponding author. Institute of Cancer Research and Molecular Medicine, Norwegian University of Science and Technology, 7489 Trondheim, Norway. Tel.: +47 7359 8668; Fax: +47 7359 8801; E-mail: terje.espevik@ntnu.no

⁵These authors contributed equally to this work

Received: 26 August 2005; accepted: 13 January 2006; published online: 9 February 2006

TLR4 in a gp96-dependent manner in the ER. MD-2 is necessary both for translocation of TLR4 from Golgi to the surface (Nagai *et al*, 2002) and LPS-induced activation (Shimazu *et al*, 1999). LPS delivered by CD14 to TLR4/MD-2 initiates a signaling cascade through the Toll/interleukin-1 receptor (TIR)-adaptors MyD88, TIRAP (also known as Mal), TRIF and TRAM, which eventually leads to nuclear translocation of the proinflammatory transcription factor NFκB (Akira and Takeda, 2004). After LPS has engaged CD14, LPS is transported into Golgi-like structures together with TLR4 (Thieblemont and Wright, 1999; Latz *et al*, 2002).

Excessive host responses towards LPS may lead to life-threatening complications such as septic shock, multiorgan failure and death (Waage *et al*, 1989). Thus, it is instrumental that LPS-induced responses are finely tuned. Part of the desensitization to LPS in human monocytes is caused by downregulation of surface TLR4/MD-2 (Kitchens *et al*, 1998). Furthermore, several negative regulators of TLR4 signaling have been reported, such as IRAK-M, short MyD88, SOCS-1, SIGIRR, ST-2, Tollip, A20, Triad3A (Liew *et al*, 2005), and recently DAP12 (Hamerman *et al*, 2005). Some of these negative regulators interfere with signaling molecules, whereas others, like Triad3A (Chuang and Ulevitch, 2004) and A20 (Boone *et al*, 2004), promote ubiquitin-dependent degradation of TLRs or signaling components, respectively. Little is known about the mechanisms of the endosomal degradation pathway of the TLRs, and if this pathway serves important functions for TLR4 and associated LPS co-receptors.

We undertook this study to delineate the mechanism by which the activated receptor complex is trafficked after engaging LPS. We have characterized the molecular mechanisms of endocytic trafficking of the LPS receptor complex and found important functional consequences of the lysosomal degradation pathway, both for the termination of signaling and the presentation of associated antigens to the CD4⁺ T helper cells.

Results

LPS downregulates plasma membrane TLR4 and both appear on endosomes

During the early hours of incubation with LPS, a gradual decrease in plasma membrane TLR4 was observed in monocytes (Figure 1A). This occurred concomitantly with formation of endosome-like structures containing LPS^{Cy5} (Figure 1B, upper panel). At later time points, LPS accumulated in the perinuclear area (Figure 1B, lower panels). For the detection of TLR4, a TLR4 antibody (anti-TLR4^{A546}) was added simultaneously with fluorescent LPS (LPS^{OG}) to monocytes. Colocalization of TLR4 and LPS was observed in microdomains on the endosomal limiting membrane 1 h after stimulation (Figure 1C).

We went on to study the localization and transport of TLR4 fused to YFP (TLR4^{YFP}) in live HEK293 cells expressing the co-receptors MD-2 and CD14 using confocal microscopy and




Figure 1 The LPS receptor complex is endocytosed and appears on endosomal structures in human monocytes. **(A)** Flow cytometry showing downregulation of surface TLR4. The surface TLR4 level was monitored in monocytes before and after LPS (2 ng/ml) stimulation using the TLR4 antibody HTA125 and the mouse IgG2a as an isotype control. **(B)** Confocal images of monocytes stimulated with Cy5-labeled LPS (LPS^{CY5}). The monocytes were incubated with LPS^{CY5} (250 ng/ml) and monitored over a period of 5 h. Arrows indicate LPS-positive endosomal structures. **(C)** Confocal images of monocytes stimulated with OregonGreen-labeled LPS (LPS^{OG}) and coincubated with Alexa546-labeled HTA125 (anti-TLR4^{A546}). Images were captured 1 h after addition of anti-TLR4^{A546} (2 µg/ml) and LPS^{OG} (600 ng/ml). No uptake was observed for the Alexa546-labeled mouse isotype control IgG2a (2 µg/ml) (data not shown). Bar = 5 µm.

LPS^{CY5} for stimulation. In unstimulated cells, TLR4^{YFP} was found in the endoplasmic reticulum, at the plasma membrane, as a large pool in the Golgi area and sporadically on endosomal structures (Figure 2A). Following LPS stimulation, endosomes showing LPS and TLR4 colocalization appeared within 15 min, suggesting active endocytosis (Figure 2A and B). Addition of LPS increased the number of TLR4-containing endosomes seven-fold within the first 40 min of stimulation and this increase was dependent on the co-receptor MD-2 (Figure 2B and C). Although we found endosomes containing TLR4 and LPS in the absence of MD-2, the presence of MD-2 enhanced the process, and also promoted aggregation of TLR4 in patches on the endosomes, as seen in Figure 1C and upper right panel of Figure 2A. CD14 greatly increased both binding to the surface and uptake of LPS, as has previously been reported (Kitchens *et al*, 1998; Latz *et al*, 2002).

LPS and TLR4 are trafficked to early/sorting endosomes

We further characterized the compartments into which LPS and TLR4 were sorted. Since LPS binding to the surface was only detected in cells expressing CD14, we subsequently used LPS binding as a marker for cells that had been successfully transfected with the CD14 plasmid. To characterize the TLR4-positive endosomes, cells were transfected with plasmids encoding CD14 and a CFP chimera containing tandem

FYVE domains from the mouse Hrs (2 × FYVE^{CFP}). This construct binds phosphatidylinositol-3-phosphate (PI3P) that is highly enriched on early/sorting endosomes (Gaullier *et al*, 1998; Simonsen *et al*, 1998; Gillooly *et al*, 2000). Within the first hour of LPS stimulation, most endosomes containing TLR4 and LPS were FYVE positive (Figure 2D). Endosomes containing TLR4 and LPS, and identified by 2 × FYVE^{CFP}, were also positive for transferrin (Figure 2E, Supplementary Figure 2). Transferrin is endocytosed in a clathrin-dependent manner and sorted on early/sorting endosomes. In cells stimulated for longer periods, TLR4 and LPS appeared on endosomes being FYVE negative (data not shown). TLR4 showed extensive colocalization with LPS in all compartments. The localization of TLR4 to early endosomes was also confirmed by staining fixed cells with an antibody to the early endosomal marker EEA1 (data not shown) or by cotransfecting the cells with a truncated but functional EEA1 (Stenmark *et al*, 1996) fused to CFP (EEA1^{CFP}). In unstimulated cells, TLR4 was present on small EEA1^{CFP}-positive endosomes that grew significantly larger in size after LPS stimulation (Supplementary Figure 1A). These EEA1-positive endosomes, like the FYVE-positive endosomes, contained a substantial amount of LPS and TLR4 on the limiting membrane. In addition, images were obtained demonstrating apparent fusion events between LPS-positive endosomes (Supplementary Figure 1B and Supplementary Movie). Taken together, our data show that LPS and TLR4 are trafficked to early/sorting endosomes and that LPS-induced signaling increases the size of these endosomes probably by endosome–endosome fusions.

LPS is endocytosed by a receptor-mediated mechanism involving dynamin and clathrin

Receptor-mediated endocytosis is often classified by being clathrin-dependent (through coated pits as seen with transferrin) or clathrin-independent (through caveoli or lipid rafts as seen for many glycosylphosphatidylinositol (GPI)-anchored proteins). Both mechanisms are dependent on dynamin, whereas (macro)pinocytosis is dependent on actin (Nichols, 2003). Dynamin proteins are GTPases that are essential for budding of vesicles from the plasma membrane and from recycling endosomes. Whereas transmembrane proteins contain sorting information in the cytosolic domain, GPI-anchored proteins (like CD14) have, in general, been shown to either be coendocytosed with the transmembrane proteins or pinocytosed together with fluid phase contents (Mayor and Riezman, 2004). We next investigated whether LPS endocytosis was dependent on dynamin using constructs of dynamin II wild type (WT) and dominant negative dynamin II (Dyn K44A) (Damke *et al*, 1994). Cells expressing TLR4^{YFP}/MD-2 were cotransfected with CD14 and dynamin WT or Dyn K44A, and investigated by confocal microscopy (Figure 3A). Fluorescent transferrin and LPS were added together for selection of cells with a dynamin knock down phenotype. LPS and transferrin uptake were not affected in the dynamin WT transfectants (Figure 3A, left panel), whereas in the Dyn K44A transfectants (Figure 3A, right panel) a clear accumulation of transferrin on the plasma membrane was observed and, as a consequence, reduced transferrin uptake (Damke *et al*, 1994). In DynK44A transfectants, total LPS uptake was reduced to about 18% of the uptake in dynamin WT transfectants (Figure 3B), supporting




Figure 2 LPS is internalized on early endosomes containing TLR4. HEK293 cells expressing TLR4^{YFP}/MD-2 or TLR4^{YFP} only were transfected with CD14 and stimulated by adding LPS^{Cy5} (250 ng/ml). (A) Confocal images before (left) and 30 min after (right) the addition of LPS^{Cy5}. (B) Estimation of the number TLR4^{YFP}-containing endosomes (>0.5 μm) before and after stimulation. The TLR4^{YFP}-containing endosomes were counted in living cells without stimulation and in the period 10–40 min after LPS^{Cy5} stimulation. The number of cells analyzed were $n = 52$ cells for each situation. (C) Comparison of the number of TLR4^{YFP}-containing endosomes (>0.5 μm) with and without coexpression of MD-2. The LPS-positive endosomes with TLR4 colocalization was quantified as above in CD14-transfected cells expressing TLR4^{YFP}/MD-2 ($n = 85$) or cells expressing TLR4^{YFP} ($n = 103$). (D) Confocal images of cells showing colocalization of TLR4 and LPS on early/sorting endosomes. HEK293 cells expressing TLR4^{YFP}/MD-2 cotransfected with CD14 and the marker for early/sorting endosomes, 2 × FYVE^{CFP}, 45 min after the addition of LPS^{Cy5}. (E) Confocal image of a cell transfected as in (D) and simultaneously given Alexa⁵⁴⁶-labeled transferrin Tf^{A546} (5 μg/ml) and LPS^{Cy5}. Full-sized images of each of the four channels of the overlaid image are shown in Supplementary Figure 2. All the images presented in this figure represent optical sections of 1.3 μm and one individual scan per cell was used for the counting of TLR4-positive endosomes. Bar = 5 μm.

that LPS is endocytosed by a receptor-mediated vesicular mechanism.

Clathrin is a scaffold protein involved in assembly of selected transmembrane proteins in invaginations called clathrin-coated pits. We knocked down clathrin heavy chain by short interfering (si) RNA technology (Motley *et al*, 2003) in HEK293 cells expressing TLR4^{YFP}/MD-2 including a non-interfering scrambled RNA duplex (Bache *et al*, 2003) as a control. Following siRNA treatment, the cells were transfected with CD14 and the uptake and distribution of LPS and transferrin were investigated (Figure 3C). The introduction of control siRNA did not affect LPS and transferrin uptake (Figure 3C, left panel), whereas clathrin siRNA resulted in significant accumulation of both transferrin and LPS at the plasma membrane in these cells (Figure 3C, right panel). Cells showing accumulation of transferrin at the plasma membrane were scored as clathrin knock down transfectants. In these cells, LPS did not appear on endosomes during the first 40 min of stimulation (Figure 3C and D). At later time points, LPS appeared on large endosomes also positive for transferrin (data not shown), suggesting an additional but slower clathrin-independent uptake mechanism.

As we found siRNA against clathrin to be very effective in inhibiting both transferrin and LPS uptake, our results show clathrin to be an essential component for the early endocytosis of the LPS receptor complex.

TLR4 is ubiquitinated and associates with Hrs

The targeting of transmembrane proteins to lysosomes is determined by tyrosine phosphorylation and di-leucine motifs of which there are candidates in TLR4. Tyrosine-based motifs (YXXΦ) are found in positions 587 (YDAF), 622 (YRDF) and 707 (YLEW), whereas a di-leucine motif ([DE]XXXL[L]) is found in position 697 (ELYRLL), both recognized by adaptor protein (AP) complexes involved in the initial phase of endocytosis (Bonifacino and Traub, 2003). A major additional mechanism for endocytosis and lysosomal targeting of transmembrane proteins is by the covalent attachment of one or more ubiquitins to lysines to the cytosolic domain (Hicke, 2001; Raiborg *et al*, 2003). A recent report showed that the E3 ubiquitin ligase Triad3A was involved in degradation of TLRs (except TLR2) (Chuang and Ulevitch, 2004). Thus, we wanted to address if LPS induced ubiquitination of TLR4 and if this had any effect on endocytosis and degradation of




Figure 3 Endocytosis of the LPS receptor complex is dependent on dynamin and clathrin. (A) Confocal images of dynamin WT (left) and Dyn K44A (right)-transfected cells. HEK293 cells expressing TLR4^{YFP}/MD-2 were transfected with CD14 together with dynamin WT or Dyn K44A and left for 6 days before analysis. The cells were simultaneously given LPS^{Cy5} (250 ng/ml) and Alexa⁵⁴⁶-labeled transferrin (Tf^{A546}) (5 μg/ml) and imaged after 50 min. (B) Comparison of the total LPS uptake in dynamin WT ($n = 46$)- and Dyn K44A ($n = 52$)-transfected cells in fixed cells 75 min after LPS^{Cy5} stimulation. The LPS uptake was quantified using the LSM510 software by measuring total intracellular fluorescence in manually drawn regions of interest and the number of cells analyzed were $n = 46$ for the dynamin WT and $n = 52$ for the Dyn K44A transfectants. (C) Confocal images of control siRNA-treated cells (left) and clathrin siRNA-treated cells (right) following LPS^{Cy5} stimulation. HEK293 cells expressing TLR4^{YFP}/MD-2 were treated with control or clathrin siRNA for 48 h followed by a 24 h transfection with CD14. The cells were simultaneously given LPS^{Cy5} (250 ng/ml) and Tf^{A546} (5 μg/ml) and imaged after 35 min. (D) Comparison of the number of LPS-positive endosomes ($> 0.5 \mu\text{m}$) per cell in cells transfected with control siRNA and clathrin siRNA 10–40 min after LPS^{Cy5} stimulation. The number of cells monitored was $n = 33$ (control siRNA) and $n = 25$ (clathrin siRNA). Clathrin- and control siRNA-treated HEK293 cells transfected with CD14 alone were included for comparison ($n = 40$ and 16, respectively). Bar = 10 μm.

TLR4. We found that TLR4 ubiquitination was both constitutive and enhanced by LPS in HEK293 cells expressing TLR4^{YFP}/MD-2 (Figure 4A). In monocytes, a weak increase in ubiquitination of TLR4 was observed at 1 h of LPS stimulation, followed by a reduction after 3 h (Figure 4B). TLR4 was ubiquitinated in cells transfected with *c-myc* Ub wild type (UbRGG) but not in cells transfected with a *c-myc* nonfunctional Ub (UbR) lacking the two C-terminal glycines required for covalent conjugation to proteins (data not shown) (Stang *et al*, 2004). There was, however, no effect of UbR on uptake of LPS within the first hour of stimulation (data not shown), suggesting that LPS uptake is Ub-independent. Ubiquitin ligases are recruited by their SH2 domains to tyrosine-phosphorylated targets, and we found that TLR4 was indeed

tyrosine phosphorylated in response to LPS (data not shown). These data suggest that although TLR4 is ubiquitinated as a response of LPS stimulation, ubiquitination is not required for the initial endocytosis of the LPS receptor complex.

Hrs is located to clathrin-coated microdomains of the early/sorting endosomal limiting membrane where it is involved in the recognition and targeting of ubiquitinated protein cargo to the lysosomal degradation pathway by promoting translocation of the target proteins to the lumen of endosomes forming multivesicular bodies (MVB) (Gruenberg and Stenmark, 2004). Hrs has recently also been shown to associate with phagosomes and aid in fusion with lysosomes (Vieira *et al*, 2004). We investigated the involvement of Hrs in TLR4 trafficking in monocytes and in HEK293-TLR4^{YFP}/MD-2 cells transiently transfected with *c-myc*-tagged Hrs. In monocytes Hrs constitutively associated with TLR4; however, the association increased transiently after 1 h of LPS stimulation (Figure 4B). Control IgG did not co-precipitate TLR4 (Figure 4B) or Hrs (data not shown). The TLR4 immunoprecipitated with Hrs was slightly larger and appeared more smeared than the mature form of TLR4 (Figure 4B, middle), possibly because Hrs interacts mainly with multiple ubiquitins on the intracellular domain of TLR4. Also in HEK293 cells overexpressing *c-myc*-Hrs, we found that TLR4 co-precipitated with Hrs (Figure 4C) in an LPS-inducible manner, whereas control IgG did not (control C1, Figure 4C). The upper band in Figure 4C represents the *c-myc*-tagged Hrs and the lower band is the endogenous Hrs as shown by comigration with the endogenous Hrs from nontransfected HEK293 cells (control C2, Figure 4C). Importantly, a moderate overexpression of *c-myc* Hrs (here estimated to be less than three times the endogenous level) lead to a rapid decrease in the amount of total cellular TLR4 after addition of LPS, whereas the amounts of Hrs and tubulin were unaffected (Figure 4C). This result is in accordance with a recent paper (Scoles *et al*, 2005), which demonstrated that overexpression of Hrs (e.g. less than 10 times the endogenous level) results in a rapid decrease in the EGFR levels upon stimulation with EGF. Furthermore, TLR4^{CFP} and Hrs^{YFP} were colocalized on discrete microdomains on the endosomal limiting membrane following LPS stimulation (Figure 4D).

TLR4 is trafficked to lysosomes for degradation

Next, we investigated the effect of chloroquine and lactacystin on TLR4 levels following LPS stimulation in HEK293 cells expressing TLR4^{YFP}/MD-2 and transfected with CD14. Chloroquine inhibits maturation of early endosomes to lysosomes by inhibiting acidification and thus preventing lysosomal degradation (Mellman *et al*, 1986). Lactacystin is an irreversible proteasome inhibitor. Previous studies have shown that cellular TLR4 is present in two different sizes, the larger is a 130 kDa heavily glycosylated, MD-2-dependent, surface-translocated form and the smaller is a 110 kDa partially glycosylated form found predominantly in Golgi (Nagai *et al*, 2002; Ohnishi *et al*, 2003). LPS seemed to increase the total cellular TLR4 after 10 h which may be due to increased transcription of the CMV-promoted TLR4-YFP constructs by LPS (Lee *et al*, 2004). However, 1 h of LPS stimulation did not increase the TLR4 level. Treatment with chloroquine resulted in a significant cellular accumulation of both forms of TLR4 as early as 1 h after LPS (Figure 5A). Interestingly, lactacystin




Figure 4 LPS promotes TLR4 ubiquitination and the association of TLR4 with Hrs. **(A)** HEK293 cells expressing TLR4^{YFP}/MD-2 were stimulated with LPS (1 µg/ml) for the indicated time periods. TLR4^{YFP} was immunoprecipitated with a polyclonal anti-GFP antibody and blotted for ubiquitin (FK2 antibody) and GFP (monoclonal GFP antibody). **(B)** Monocytes were either left untreated or stimulated with LPS (1 µg/ml) for the indicated time periods. Cellular lysates were made and TLR4 immunoprecipitated with a polyclonal TLR4 antibody and blotted for ubiquitin (left), TLR4 (middle) and Hrs (lower left) or the reverse immunoprecipitation of Hrs or control IgG blotted for TLR4 (upper right). The Hrs in lysate is shown in the lower right panel. **(C)** HEK293 cells expressing TLR4^{YFP}/MD-2 were transfected with *c-myc* Hrs for 24 h before LPS stimulation for the indicated periods. Cellular lysates were prepared and immunoprecipitated with anti-GFP (TLR4^{YFP}) or control IgG (C1) and blotted for Hrs. Control C2 represents Hrs immunoprecipitated from nontransfected HEK293 cells showing the size of endogenous Hrs. Overexpression of Hrs resulted in an LPS-induced reduction in TLR4 levels in lysates, whereas Hrs and tubulin levels are unaffected (lower three panels). **(D)** Confocal images of HEK293 cells expressing TLR4^{CFP}/MD-2 transfected with CD14 and Hrs^{YFP} following 3 h of LPS (250 ng/ml) stimulation. Bar = 5 µm.

reported to block Triad3A-mediated degradation of TLR9 (Chuang and Ulevitch, 2004) did not cause the same accumulation of TLR4^{YFP} in LPS-stimulated cells (Figure 5A). LY294002 (LY) inhibits the PI3-kinase enzymes responsible for the formation of the cellular PI3P controlling endosome fusions (Simonsen *et al*, 1998) and the formation of intraluminal vesicles in late endosomes/MVBs (Futter *et al*, 2001). PI3-kinase activity is also required for the recruitment of Hrs to endosomal membranes (Komada and Soriano, 1999; Urbe *et al*, 2000). Cells treated with LY and stimulated with LPS showed a dramatic increase in the level of TLR4^{YFP} after 10 h of LPS stimulation compared to LPS-stimulated cells treated with control vehicle (Figure 5B). Altogether, these data suggest that endosomal sorting and acidification are required for degradation of TLR4.

Furthermore, we investigated the involvement of Hrs in TLR4 degradation by knocking down the endogenous level of Hrs by siRNA technology (Bache *et al*, 2003). Hrs siRNA-treated HEK293 cells expressing TLR4^{YFP}/MD-2 and transfected with CD14 showed an increased levels of both the large fully glycosylated and the partially glycosylated TLR4^{YFP} form compared to the cells treated with control siRNA (Figure 5C, upper panel). Our results suggest that ubiquitinated TLR4 associates with Hrs directing the activated TLR4 to lysosomes for degradation. This result is in

contrast to TLR9 that has been reported to be degraded by proteasomes after ubiquitination by the Triad3A ubiquitin ligase (Chuang and Ulevitch, 2004).

Inhibition of the endosomal pathway increases LPS-induced NFκB activation

After describing mechanisms of the TLR4 sorting through the endosomal pathway, we investigated the role of this pathway on LPS-induced NFκB activation. By comparing the effect of wild-type versus dominant negative constructs for the trafficking components dynamin II (dynamin WT versus Dyn K44A) and ubiquitin (UbrGG versus Ubr), we found that the reduced endocytic activity caused by Dyn K44A, or saturation of Ub-interacting proteins with the nonfunctional Ubr, caused increased NFκB activation compared to overexpression of the WT constructs (Figure 5D and Supplementary Figure 3). This result strongly supports that signaling is initiated at the plasma membrane (Ahmad-Nejad *et al*, 2002; Latz *et al*, 2002), and that endocytosis and lysosomal targeting of TLR4 limit NFκB activation. Additional arguments for this conclusion was also obtained in knock down experiments of Hrs with siRNA showing a 70% increase in LPS-induced NFκB activation compared to control (Figure 5E and Supplementary Figure 3). Furthermore, by knocking down an essential component (HCRP1) of the endosomal




Figure 5 Lysosomal degradation of TLR4 limits signaling. (A) HEK293 cells expressing TLR4^{YFP}/MD-2 were transfected with CD14 and treated with chloroquine (10 μM) or lactacystin (10 μM) for 30 min prior to LPS (1 μg/ml) stimulation for the indicated time periods. Cellular lysates were made and TLR4^{YFP} was detected using a polyclonal anti-GFP antibody and reblotted for α-tubulin as a control for equal loading. The samples displayed represent sections of the original blot. (B) HEK293 cells expressing TLR4^{YFP}/MD-2 were transfected with CD14 and treated with 10 μM LY or control vehicle (EtOH) for 30 min and stimulated with LPS (1 μg/ml) for 10 h. Cellular lysates were made and TLR4^{YFP} was detected using a polyclonal anti-GFP antibody and reblotted for α-tubulin as a control for equal loading. (C) HEK293 cells expressing TLR4^{YFP}/MD-2 were treated with control of Hrs siRNA and transfected with CD14. Cellular lysates were made and blotted for GFP (TLR4^{YFP}), Hrs, and α-tubulin for the examination of TLR4^{YFP} and Hrs levels. The samples displayed represent sections of the original blot. (D) Effect of dynamin and ubiquitination on LPS-induced NFκB activation through TLR4. HEK293 cells expressing TLR4 were transfected with MD-2, CD14 and Elam-luc together with WT constructs (dynamin and UbRGG) or a dominant negative version of dynamin (Dyn K44A) or an inhibitory version of ubiquitin (UbR, lacking the two C-terminal glycines necessary for conjugation to proteins but able to occupy ubiquitin-binding pockets). The cells were stimulated for 6 h with LPS (10 ng/ml) and assayed for NFκB-induced luciferase expression. (E) Effect of Hrs and ESCRT-I on LPS-induced NFκB activation through TLR4. HEK293 cells expressing TLR4^{YFP}/MD-2 were treated with control siRNA or siRNA directed towards Hrs or HCRP1 (component of ESCRT-I) and cotransfected with CD14 and Elam-luc as described in the Materials and methods section and assayed for NFκB-induced luciferase expression.

sorting complexes required for transport-I (ESCRT-I) with siRNA (Bache *et al*, 2004), the LPS (10 ng/ml)-induced NFκB activation increased by 245% compared to the siRNA control (Figure 5E). These data suggest that dynamin, ubiquitin, Hrs and HCRP1 (i.e. ESCRT-I) are involved in desensitizing the cell after LPS activation, most likely through endocytosis and lysosomal degradation of the activated LPS receptor complex.

Antigens associated with the LPS receptor complex are presented on HLA class II to CD4⁺ T helper cells

Time-lapse series of LPS-stimulated monocytes showed that LPS overlapped with dextran in dynamic tubular structures protruding from the perinuclear area after 3–5 h of incubation (data not shown). These protrusions were directed towards the plasma membrane as has been described for MHC class II containing lysosome-like tubulae after LPS stimulation in dendritic cells (Boes *et al*, 2002), a phenotype not observed in HEK293 cells. By counterstaining HEK293 cells expressing TLR4^{YFP}/MD-2 transfected with CD14 with an antibody against the late endosomal/lysosomal marker protein LAMP-1, we observed that TLR4^{YFP} was present in the lumen of late endosomes/MVBs following LPS stimulation (Figure 6A). Treatment of stimulated live cells with LysoTracker Red showed that TLR4^{YFP} and LPS^{CY5} colocalized at the limiting membrane and in the lumen of nonacidic endosomes (Figure 6B). Intraluminal TLR4^{YFP} was lost fol-

lowing endosomal acidification, whereas LPS^{CY5} was still detectable. Since TLR4 and LPS/CD14 are directed through the endocytic compartments to lysosomes for degradation, associated antigens may be presented on HLA class II to CD4⁺ T cells. We performed antigen presentation assays using mouse antibodies to TLR4 (HTA125), CD14 (5C5) and MD-2 (IIC2) and proliferation of a CD4⁺ T cell specific for the mouse κ-light chain-derived peptide C_κ^{40–48} presented on HLA-DR4 (Schjetne *et al*, 2002) in an LPS-free system. This system is designed for functional testing of the class II presentation pathway by providing a read-out for lysosomal antigen processing, loading and presentation in primary human cells. We found efficient presentation of the antibody-derived peptide (Figure 6C) suggesting that TLR4, MD-2 and CD14 were directed to a functional class II loading compartment, most commonly ascribed to MVB (Peters *et al*, 1991). This result implies that the LPS receptor complex, TLR4/MD-2/CD14, not only signals the presence of microbes but also assists in directing the endocytosis of antigens on class II molecules for presentation to T helper cells.

Discussion

Molecular mechanisms involved in sorting of the LPS receptor complex

The TLR4 agonist, LPS, is a potent immune stimulator which can cause sepsis. TLR4 signaling is under tight regulatory




Figure 6 The LPS receptor complex is trafficked to the class II loading compartment. (A) Confocal images showing TLR4^{YFP} in the lumen of LAMP-1-positive endosomes. HEK293 cells expressing TLR4^{YFP}/MD-2 transfected with CD14 were treated for 2 h with LPS (1 μg/ml) and fixed before intracellular staining for LAMP1. Arrows indicate LAMP-1-positive endosomes showing intraluminal TLR4^{YFP}. (B) Confocal images showing TLR4^{YFP} in the lumen of nonacidic endosomes. HEK293 cells expressing TLR4^{YFP}/MD-2 transfected with CD14 were stimulated with LPS (1 μg/ml) for 3 h and incubated with 10 nM LysoTracker Red (C). Antibodies towards TLR4 (HTA125), CD14 (5C5) or MD-2 (IIC1) or isotype-matched controls were added to human HLA-matched monocytes together with cloned DR4-restricted CD4⁺ T cells specific for mouse C_K⁴⁰⁻⁴⁸. T-cell proliferation was measured after 48 h by incorporation of [³H] TdR. Data points are mean of duplicates from a representative experiment. Isotype-matched monoclonal antibodies were used as controls. Bar = 5 μm.

control at multiple levels and several inhibitors of TLR4 signaling have now been described. Internalization of receptors regulates signal transduction (Di Guglielmo *et al*, 2003); however, little attention has been devoted to mechanisms and functional consequences of the endocytic uptake of the LPS receptor complex. In this study, we have characterized the uptake of LPS and the sorting of the LPS receptor complex from the plasma membrane to the late endosomes/lysosomes and explored functional consequences of this endosomal pathway.

Internalization of membrane receptors occurs both through clathrin- and caveolae/raft-mediated pathways. The EGF receptor is internalized within minutes after ligand addition by a clathrin-mediated pathway (Le and Wrana, 2005), whereas the IL-2 receptor appears to be taken up by a lipid raft-dependent manner (Lamaze *et al*, 2001). Previous reports have shown that CD14 is located in lipid rafts and that a fraction of TLR4 moves into the lipid raft domain after LPS stimulation, a result suggesting a clathrin-independent uptake of the LPS receptor complex (Triantafilou *et al*, 2004). Furthermore, Kitchens *et al* (1998) reported that the CD14-mediated uptake of LPS occurs by both clathrin-dependent

and -independent pathways. Knocking down clathrin by siRNA results in a significant inhibition of EGF uptake at low EGF doses and less effective at higher EGF doses (Sigismund *et al*, 2005). Our results using siRNA for clathrin clearly demonstrate that the early phase (up to 40 min) of LPS internalization is predominantly clathrin-dependent, whereas clathrin-independent pathways operate at later time points.

Monoubiquitination serves an important role in sorting membrane proteins through the endosomal pathway (Raiborg *et al*, 2003), whereas polyubiquitination is the recognition signal for proteasomal degradation (Marmor and Yarden, 2004). As mentioned above, Triad3A has been reported to be an E3 ubiquitin ligase that promotes proteasomal degradation of TLR9 (Chuang and Ulevitch, 2004). Triad3A-mediated ubiquitination does not seem to play a role in the lysosomal degradation pathway as the effect of this ligase on TLR9 degradation is dependent on proteasomal, but not lysosomal activity (Chuang and Ulevitch, 2004). We observed increased ubiquitination of TLR4 upon LPS activation both in monocytes and in genetically engineered cells. Hrs has an essential role in sorting ubiquitinated proteins into clathrin-coated microdomains of early endosomes (Raiborg

et al, 2002) and Hrs has interacting domains for PI3P (the FYVE domain), clathrin and ubiquitin (Bache *et al*, 2003). We provide evidence for Hrs interaction with TLR4 on endosomes and that this interaction was increased upon LPS activation. Thus, Hrs seems to play an important role in the initial steps of TLR4 sorting to the endosomal pathway. A part of the LPS receptor complex also traffics from the plasma membrane to the Golgi (Latz *et al*, 2002), and this pathway may either go directly from the plasma membrane or through the early/sorting endosomes. Whether Hrs also plays a role in TLR4 trafficking to the Golgi apparatus is not known. Another protein, Tollip, has been shown to inhibit LPS-induced NF κ B activation (Zhang and Ghosh, 2002) by a mechanism requiring localization to endosomes or activated plasma membranes and through binding of ubiquitinated cargo and recruitment to clathrin-coated microdomains (Kato *et al*, 2004), not unlike Hrs. It is thus likely that redundancy exists in the machinery that ensures that inflammatory signals are shut off to avoid hyperinflammation and the accompanied adverse effects.

Endosomal trafficking of the LPS receptor complex leading to lysosomal degradation and signal termination was consistently demonstrated in the HEK293-TLR4 cells. Our results suggest that a similar pathway of LPS receptor endocytosis and degradation also operates in monocytes. Firstly, LPS induced a clear downregulation of TLR4 on the plasma membrane with subsequent localization of LPS and TLR4 to endosomes. Secondly, ubiquitination of TLR4 was constitutive, weakly increased after 1 h of LPS stimulation and markedly decreased after 3 h. Finally, TLR4 associated with Hrs and this association was transiently increased at 1 h of LPS stimulation. Direct evidence for LPS-induced lysosomal degradation of TLR4 in monocytes was difficult to obtain due to low TLR4 signals in immunoblots and probably a high turnover rate for TLR4. However, antibodies towards TLR4, MD-2 and CD14 were degraded in MVBs/lysosomes since an antibody-derived peptide was loaded onto HLA class II (Figure 6C). Thus, it is likely that the pathway of endosomal degradation of TLR4 that we demonstrated in the HEK293-TLR4 cells also, at least to some extent, takes place in monocytes.

Inhibition of endocytosis and endosomal sorting increase LPS-induced proinflammatory signaling

Our studies provide evidence that sorting the LPS receptor complex through the endosomal pathway has important functional consequences. We observed a striking increase in LPS signaling when proteins involved in endosomal trafficking and maturation were inhibited. By reducing the amount of functional ESCRT-I, which associates with Hrs and is involved in the sorting of ubiquitinated proteins into multivesicular bodies (Bache *et al*, 2002), a more than two-fold increase in the LPS signaling was observed. Inhibition of ubiquitination and depletion of Hrs or ESCRT-I resulted in increased LPS signaling. Furthermore, overexpression of Hrs lead to a significant increase in LPS-induced TLR4 degradation. From these results it is apparent that LPS signaling starts at the plasma membrane; however, signaling is likely to continue along the endosomal pathway until an Hrs- and ESCRT-I-mediated translocation of TLR4 to the luminal side of endosomes has taken place. This view is also supported by confocal imaging showing the presence of the signaling

adapters MyD88 and TRAM at the plasma membrane and on endosomes in LPS-stimulated cells (H Husebye and T Espevik, unpublished results). Signaling on endosomes is also reported for TGF- β which is taken up by a clathrin-dependent mechanism and trafficked into early endosomes where the signaling component SARA is enriched (Di Guglielmo *et al*, 2003).

The LPS receptor complex links innate and adaptive immunity

A very important consequence of the trafficking of the LPS receptor complex described in this paper is that antigens associated with the LPS receptor complex are directly transported to the antigen presentation machinery for presentation to CD4⁺ T helper cells. The loading of antigens on class II molecules occurs by fusion of limiting and intraluminal membranes in MVBs (Peters *et al*, 1991), resulting in exchange of the class II inhibitory peptide (CLIP) with lysosomal peptides derived from processed exogenous proteins. This is also highly relevant for other suggested ligands of TLR4, like heat shock proteins, viral proteins (Akira and Takeda, 2004) and cotrafficked antigens, by providing a direct link between activation of innate and acquired immune cells. This may be exploited in design of synthetic vaccines aimed at the CD4⁺ T-helper cell population, as has been shown for conjugated complexes of lipoproteins and immunogenic peptides acting through TLR2 (Jackson *et al*, 2004).

Altogether, our results strongly support that distinct components of the endocytic pathway inhibit inflammatory signals induced by TLR4 and that long-lived information is simultaneously passed on to the adaptive branch of the immune system.

Materials and methods

Reagents

LPS (0111:B4) from *Escherichia coli* was purchased from Invivogen. Cy5- and OregonGreen (OG) labeling of LPS was performed as previously described (Latz *et al*, 2002). Alexa labeling of antibodies was performed according to the manufacturer's protocol (Invitrogen). Alexa⁵⁴⁶-labeled transferrin and LysoTracker Red were purchased from Invitrogen. LY294002 and lactacystin were purchased from Calbiochem. Chloroquine was purchased from Sigma. Mouse antibodies used: anti-GFP (JL-8) (Clontech), anti-TLR4 (HTA125) was kindly provided by Dr Kensuke Miyake (Saga Medical School, Japan) (Yang *et al*, 2000), anti-c-myc (9E10) (BD Bioscience), anti-ubiquitin-protein conjugates (FK2) (Affiniti Research), anti-LAMP1 (R&D Systems), anti-CD14 (5C5) (Lien *et al*, 1998) and anti-MD-2 (IIC1). The mouse monoclonal antibody IIC1 (IgM) detects MD-2 in CHO and HEK293 cells expressing TLR4/MD-2 as well as on the surface of human monocytes (T Espevik and L Ryan, unpublished results). Rabbit polyclonal antibodies used: anti-TLR4 (eBioscience), anti-GFP (Clontech) and anti-Hrs (Raiborg *et al*, 2001). The following mammalian expression vectors were used: pcDNA3 (Invitrogen), CD14 (Latz *et al*, 2002), c-myc-Hrs, c-myc-ubiquitin wild-type UbrGG (Stang *et al*, 2004) and c-myc-ubiquitin DN UbR (Stang *et al*, 2004) were in pcDNA3. Dynamin-II wild type and Dynamin-II K44A in pcDNA3 were kindly provided by Dr Sandy Schmid (Scripps, USA). MD-2 in pEF-BOS was kindly provided by Dr Miyake (Yang *et al*, 2000). Hrs^{YFP} was in pEYFP-C1 (Clontech). The ECFP-2 \times FYVE construct was made by substituting GFP from EGFP-2 \times FYVE (Gillooly *et al*, 2000) with CFP using pECFP (Clontech) as a template and PCR primers 5'-CCA AGT ACG CCC CCT ATT GA-3' and 5'-ATT TAA GCT TGT ACA GCT CGT CCA TGC-3'. The CFP encoding fragment was inserted as an *NheI/XhoI* fragment.

Cells and cell lines

HEK293 cell lines that stably expressed human TLR4 (Yang *et al*, 2000), TLR4^{YFP}, TLR4^{YFP}/MD-2 (Latz *et al*, 2002) or TLR4^{CFP}/MD-2 were used and cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum (FCS) and 0.5 mg/ml G418 and transfected using GeneJuiceTM transfection reagent (Novagene) according to the manufacturer's protocol. Human monocytes were isolated by plastic adherence from Lymphoprep (Axis-Shield) separated buffycoats (The Blood Bank, St Olavs Hospital, Trondheim, Norway) and grown in RPMI/10% pooled A + serum (The Blood Bank). LPS was sonicated for 5 min and preincubated in serum-containing medium at 37°C for 5 min before being added to cells. For confocal imaging, the cells were seeded on 35 mm glass bottom γ -irradiated tissue cell dishes (MatTek Corporation) and buffered with 25 mM HEPES buffer just before stimulation. LPS^{CYS} (250 ng/ml) or LPS^{OG} (600 ng/ml) was used in LPS uptake studies.

Flow cytometry

Adherent monocytes were incubated with or without LPS (100 ng/ml) in medium/10% A + serum for 2 h, detached with EDTA and stained with 10 μ g/ml anti-TLR4 (HTA125) or mouse IgG2a for 30 min on ice, washed and incubated with 10 μ g/ml secondary anti-mouse Ig-FITC (Becton Dickinson) for 30 min on ice and washed and analyzed by flow cytometry (Coulter).

Short interfering RNA treatment

siRNA duplexes targeting the coding region of clathrin heavy chain (Motley *et al*, 2003), the coding region of Hrs (Bache *et al*, 2003) and the coding region of the ESCRT-I component HCRP1 (Bache *et al*, 2004) were used. A scrambled RNA duplex (Bache *et al*, 2003) was used as a noninterfering control. All siRNA duplexes used were synthesized by Dharmacon. HEK293 cells and cells expressing TLR4^{YFP}/MD-2 were transfected with 20 nM siRNA for 48 h using Oligofectamine (Invitrogen) according to the manufacturer's protocol. For the silencing of Hrs and HCRP1, the cells were replated and subjected to a second round of siRNA treatment. For the introduction of plasmid DNA into the siRNA-transfected cells, the cells were replated and transfected for 24 h using the GeneJuice transfection reagent.

Western blotting and immunoprecipitation

HEK293 cells expressing TLR4^{YFP} were plated on 35 mm dishes and transfected 24–48 h before two washes in PBS and lysed in 100–300 μ l of lysis buffer (20 mM Tris-HCl, 1 mM EDTA, 1 mM EGTA, 137 mM NaCl, 1% Triton X-100, 1 mM sodium deoxycholate, 10% glycerol, 1 mM Na₃VO₄, 50 mM NaF and Complete protease inhibitor (Roche)). For Western blotting, the samples were denatured in 1 \times NuPAGE LDS sample buffer supplemented with 25 mM DTT for 10 min at 70°C. Electrophoresis was performed using NuPAGE novex 7% Tris-Acetate polyacrylamide or 10% Bis-Tris polyacrylamide gels (Invitrogen) and the samples blotted onto 0.45 μ m nitrocellulose filters (Biorad laboratories) and developed

References

- Ahmad-Nejad P, Hacker H, Rutz M, Bauer S, Vabulas RM, Wagner H (2002) Bacterial CpG-DNA and lipopolysaccharides activate Toll-like receptors at distinct cellular compartments. *Eur J Immunol* **32**: 1958–1968
- Akira S, Takeda K (2004) Toll-like receptor signalling. *Nat Rev Immunol* **4**: 499–511
- Bache KG, Raiborg C, Mehlum A, Madhus IH, Stenmark H (2002) Phosphorylation of Hrs downstream of the epidermal growth factor receptor. *Eur J Biochem* **269**: 3881–3887
- Bache KG, Raiborg C, Mehlum A, Stenmark H (2003) STAM and Hrs are subunits of a multivalent ubiquitin-binding complex on early endosomes. *J Biol Chem* **278**: 12513–12521
- Bache KG, Slagsvold T, Cabezas A, Rosendal KR, Raiborg C, Stenmark H (2004) The growth-regulatory protein HCRP1/hVps37A is a subunit of mammalian ESCRT-I and mediates receptor down-regulation. *Mol Biol Cell* **15**: 4337–4346
- Boes M, Cerny J, Massol R, Op den Brouw M, Kirchhausen T, Chen J, Ploegh HL (2002) T-cell engagement of dendritic cells rapidly rearranges MHC class II transport. *Nature* **418**: 983–988

using ECL (Amersham Biosciences). For the analysis of immune complexes, lysates were precleared for 1 h with protein G Sepharose and subsequently incubated for 2 h with the appropriate antibody (2–3 μ g/ml). Immune complexes were harvested with protein G Sepharose for 2 h before four washes in lysis buffer. Elution of the immune complexes was performed at 95°C for 10 min in 2 \times LDS sample buffer supplemented with 50 mM DTT.

Confocal imaging

Images of live cells were captured at 37°C using an Axiovert 100-M microscope with a heated stage equipped with a Zeiss LSM 510 META scanning unit and a 1.4 NA \times 63 plan apochromat objective. For intracellular staining, the cells were permeabilized in 0.05% saponin, fixed in 3% paraformaldehyde and incubated with the appropriate antibody as described previously (Simonsen *et al*, 1998). For LPS uptake studies in monocytes, fluorescently labeled LPS was added to the cells alone or simultaneously with 2 μ g/ml Alexa⁵⁴⁶-labeled anti-TLR4 (HTA125) or 2 μ g/ml Alexa546-labeled isotype control. Only ring formed endosomes positive for TLR4 and/or LPS on the limiting membrane (minimal cutoff size 0.5 μ m) were counted. The total intracellular LPS fluorescence in Figure 3B was quantified by manually drawing a region of interest that covered the cytoplasm and using the LSM510 software to calculate total fluorescence.

Antigen presentation

The human CD4⁺ Th1 cell clone (T18) specific for mouse antibody constant κ -chain-derived peptide C κ ^{40–48} and restricted by HLA-DR4 (DRA1, B1*0401) was used to measure T helper-cell responses (Schjetne *et al*, 2003). Antigen-presenting cells used were HLA-matched adherent peripheral blood monocytes (CD14⁺). Tissue culture medium was RPMI 1640 with glucose and L-glutamine (Gibco, Paisley, Scotland) supplemented with pooled 10% human serum (HS) from blood donors.

Luciferase assay

NF κ B activation was determined by an NF κ B luciferase reporter assay as described (Latz *et al*, 2002).

Supplementary data

Supplementary data are available at *The EMBO Journal* Online.

Acknowledgements

We thank Liv Ryan for technical assistance. Professor Oddmund Bakke, University of Oslo, generously provided the EEA1^{CFP} construct. This work was supported by The Norwegian Research Council, The Norwegian Cancer Society and the Commission of the European Communities, LSMH-CT-2004-512093, AMIS. Andreas Brech is the recipient of a career scientist fellowship from the FUGE program of the Research Council of Norway. The authors have no conflicting financial interests.

- Bonifacino JS, Traub LM (2003) Signals for sorting of transmembrane proteins to endosomes and lysosomes 2. *Annu Rev Biochem* **72**: 395–447
- Boone DL, Turer EE, Lee EG, Ahmad RC, Wheeler MT, Tsui C, Hurley P, Chien M, Chai S, Hitotsumatsu O, McNally E, Pickart C, Ma A (2004) The ubiquitin-modifying enzyme A20 is required for termination of Toll-like receptor responses. *Nat Immunol* **5**: 1052–1060
- Chuang TH, Ulevitch RJ (2004) Triad3A, an E3 ubiquitin-protein ligase regulating Toll-like receptors. *Nat Immunol* **5**: 495–502
- Damke H, Baba T, Warnock DE, Schmid SL (1994) Induction of mutant dynamin specifically blocks endocytic coated vesicle formation. *J Cell Biol* **127**: 915–934
- Di Guglielmo GM, Le RC, Goodfellow AF, Wrana JL (2003) Distinct endocytic pathways regulate TGF-beta receptor signalling and turnover. *Nat Cell Biol* **5**: 410–421
- Futter CE, Collinson LM, Backer JM, Hopkins CR (2001) Human VPS34 is required for internal vesicle formation within multivesicular endosomes. *J Cell Biol* **155**: 1251–1264

- Gaullier JM, Simonsen A, D'Arrigo A, Bremnes B, Stenmark H, Aasland R (1998) FYVE fingers bind Ptdins(3)P. *Nature* **394**: 432–433
- Gillooly DJ, Morrow IC, Lindsay M, Gould R, Bryant NJ, Gaullier JM, Parton RG, Stenmark H (2000) Localization of phosphatidylinositol 3-phosphate in yeast and mammalian cells. *EMBO J* **19**: 4577–4588
- Gruenberg J, Stenmark H (2004) The biogenesis of multivesicular endosomes. *Nat Rev Mol Cell Biol* **5**: 317–323
- Hamerman JA, Tchao NK, Lowell CA, Lanier LL (2005) Enhanced Toll-like receptor responses in the absence of signaling adaptor DAP12. *Nat Immunol* **6**: 579–586
- Hicke L (2001) Protein regulation by monoubiquitin. *Nat Rev Mol Cell Biol* **2**: 195–201
- Jackson DC, Lau YF, Le T, Suhrbier A, Deliyannis G, Cheers C, Smith C, Zeng W, Brown LE (2004) A totally synthetic vaccine of generic structure that targets Toll-like receptor 2 on dendritic cells and promotes antibody or cytotoxic T cell responses. *Proc Natl Acad Sci USA* **101**: 15440–15445
- Katoh Y, Shiba Y, Mitsuhashi H, Yanagida Y, Takatsu H, Nakayama K (2004) Tollip and Tom1 form a complex and recruit ubiquitin-conjugated proteins onto early endosomes. *J Biol Chem* **279**: 24435–24443
- Kitchens RL, Wang PY, Munford RS (1998) Bacterial lipopolysaccharide can enter monocytes via two CD14-dependent pathways. *J Immunol* **161**: 5534–5545
- Komada M, Soriano P (1999) Hrs, a FYVE finger protein localized to early endosomes, is implicated in vesicular traffic and required for ventral folding morphogenesis. *Genes Dev* **13**: 1475–1485
- Lamaze C, Dujeancourt A, Baba T, Lo CG, Benmerah A, Utry-Varsat A (2001) Interleukin 2 receptors and detergent-resistant membrane domains define a clathrin-independent endocytic pathway. *Mol Cell* **7**: 661–671
- Latz E, Visintin A, Lien E, Fitzgerald KA, Monks BG, Kurt-Jones EA, Golenbock DT, Espevik T (2002) Lipopolysaccharide rapidly traffics to and from the Golgi apparatus with the toll-like receptor 4-MD-2-CD14 complex in a process that is distinct from the initiation of signal transduction. *J Biol Chem* **277**: 47834–47843
- Le RC, Wrana JL (2005) Clathrin- and non-clathrin-mediated endocytic regulation of cell signalling. *Nat Rev Mol Cell Biol* **6**: 112–126
- Lee Y, Sohn WJ, Kim DS, Kwon HJ (2004) NF-kappaB- and c-Jun-dependent regulation of human cytomegalovirus immediate-early gene enhancer/promoter in response to lipopolysaccharide and bacterial CpG-oligodeoxynucleotides in macrophage cell line RAW 264.7. *Eur J Biochem* **271**: 1094–1105
- Lien E, Aukrust P, Sundan A, Muller F, Froland SS, Espevik T (1998) Elevated levels of serum-soluble CD14 in human immunodeficiency virus type 1 (HIV-1) infection: correlation to disease progression and clinical events. *Blood* **92**: 2084–2092
- Liew FY, Xu D, Brint EK, O'Neill LAJ (2005) Negative regulation of toll-like receptor-mediated immune responses. *Nat Rev Immunol* **5**: 446–458
- Marmor MD, Yarden Y (2004) Role of protein ubiquitylation in regulating endocytosis of receptor tyrosine kinases. *Oncogene* **23**: 2057–2070
- Mayor S, Riezman H (2004) Sorting GPI-anchored proteins. *Nat Rev Mol Cell Biol* **5**: 110–120
- Mellman I, Fuchs R, Helenius A (1986) Acidification of the endocytic and exocytic pathways. *Annu Rev Biochem* **55**: 663–700
- Motley A, Bright NA, Seaman MNJ, Robinson MS (2003) Clathrin-mediated endocytosis in AP-2-depleted cells. *J Cell Biol* **162**: 909–918
- Nagai Y, Akashi S, Nagafuku M, Ogata M, Iwakura Y, Akira S, Kitamura T, Kosugi A, Kimoto M, Miyake K (2002) Essential role of MD-2 in LPS responsiveness and TLR4 distribution. *Nat Immunol* **3**: 667–672
- Nichols B (2003) Caveosomes and endocytosis of lipid rafts. *J Cell Sci* **116**: 4707–4714
- Ohnishi T, Muroi M, Tanamoto K (2003) MD-2 is necessary for the Toll-like receptor 4 protein to undergo glycosylation essential for its translocation to the cell surface. *Clin Diagn Lab Immunol* **10**: 405–410
- Peters PJ, Neefjes JJ, Oorschot V, Ploegh HL, Geuze HJ (1991) Segregation of Mhc class-II molecules from Mhc class-I molecules in the Golgi-complex for transport to lysosomal compartments. *Nature* **349**: 669–676
- Raiborg C, Bache KG, Gillooly DJ, Madhus IH, Stang E, Stenmark H (2002) Hrs sorts ubiquitinated proteins into clathrin-coated microdomains of early endosomes. *Nat Cell Biol* **4**: 394–398
- Raiborg C, Bache KG, Mehlum A, Stang E, Stenmark H (2001) Hrs recruits clathrin to early endosomes. *EMBO J* **20**: 5008–5021
- Raiborg C, Rusten TE, Stenmark H (2003) Protein sorting into multivesicular endosomes. *Curr Opin Cell Biol* **15**: 446–455
- Schjetne KW, Thompson KM, Aarvak T, Fleckenstein B, Sollid LM, Bogen B (2002) A mouse C{kappa}-specific T cell clone indicates that DC-SIGN is an efficient target for antibody-mediated delivery of T cell epitopes for MHC class II presentation. *Int Immunol* **14**: 1423–1430
- Schjetne KW, Thompson KM, Nilsen N, Flo TH, Fleckenstein B, Iversen JG, Espevik T, Bogen B (2003) Cutting edge: link between innate and adaptive immunity: Toll-like receptor 2 internalizes antigen for presentation to CD4+ T cells and could be an efficient vaccine target. *J Immunol* **171**: 32–36
- Scoles DR, Qin Y, Nguyen V, Gutmann DH, Pulst SM (2005) HRS inhibits EGF receptor signaling in the RT4 rat schwannoma cell line. *Biochem Biophys Res Commun* **335**: 385–392
- Shimazu R, Akashi S, Ogata H, Nagai Y, Fukudome K, Miyake K, Kimoto M (1999) MD-2, a molecule that confers lipopolysaccharide responsiveness on Toll-like receptor 4. *J Exp Med* **189**: 1777–1782
- Sigmund S, Woelk T, Puri C, Maspero E, Tacchetti C, Transidico P, Di Fiore PP, Polo S (2005) Clathrin-independent endocytosis of ubiquitinated cargos. *Proc Natl Acad Sci USA* **102**: 2760–2765
- Simonsen A, Lippe R, Christoforidis S, Gaullier JM, Brech A, Callaghan J, Toh BH, Murphy C, Zerial M, Stenmark H (1998) EEA1 links PI(3)K function to Rab5 regulation of endosome fusion. *Nature* **394**: 494–498
- Stang E, Blystad FD, Kazazic M, Bertelsen V, Brodahl T, Raiborg C, Stenmark H, Madhus IH (2004) Cbl-dependent ubiquitination is required for progression of EGF receptors into clathrin-coated pits. *Mol Biol Cell* **15**: 3591–3604
- Stenmark H, Aasland R, Toh BH, D'Arrigo A (1996) Endosomal localization of the autoantigen EEA1 is mediated by a zinc-binding FYVE finger. *J Biol Chem* **271**: 24048–24054
- Thieblemont N, Wright SD (1999) Transport of bacterial lipopolysaccharide to the Golgi apparatus. *J Exp Med* **190**: 523–534
- Triantafilou M, Morath S, Mackie A, Hartung T, Triantafilou K (2004) Lateral diffusion of Toll-like receptors reveals that they are transiently confined within lipid rafts on the plasma membrane. *J Cell Sci* **117**: 4007–4014
- Urbe S, Mills IG, Stenmark H, Kitamura N, Clague MJ (2000) Endosomal localization and receptor dynamics determine tyrosine phosphorylation of hepatocyte growth factor-regulated tyrosine kinase substrate. *Mol Cell Biol* **20**: 7685–7692
- Vieira OV, Harrison RE, Scott CC, Stenmark H, Alexander D, Liu J, Gruenberg J, Schreiber AD, Grinstein S (2004) Acquisition of Hrs, an essential component of phagosomal maturation, is impaired by mycobacteria. *Mol Cell Biol* **24**: 4593–4604
- Waage A, Brandtzaeg P, Halstensen A, Kierulf P, Espevik T (1989) The complex pattern of cytokines in serum from patients with meningococcal septic shock. Association between interleukin 6, interleukin 1, and fatal outcome. *J Exp Med* **169**: 333–338
- Yang H, Young DW, Gusovsky F, Chow JC (2000) Cellular events mediated by lipopolysaccharide-stimulated Toll-like receptor 4. MD-2 is required for activation of mitogen-activated protein kinase and Elk-1. *J Biol Chem* **275**: 20861–20866
- Zhang G, Ghosh S (2002) Negative regulation of toll-like receptor-mediated signaling by Tollip. *J Biol Chem* **277**: 7059–7065