

## REVIEW

# Genes without frontiers?

D Bensasson<sup>1</sup>, JL Boore<sup>1</sup> and KM Nielsen<sup>2,3</sup>

<sup>1</sup>Evolutionary Genomics Department, DOE Joint Genome Institute, 2800 Mitchell Drive, Walnut Creek, CA 94598, USA; <sup>2</sup>Department of Pharmacy, University of Tromsø, N9037 Tromsø, Norway; <sup>3</sup>The Norwegian Institute for Gene Ecology, Tromsø, Norway

For bacteria, the primary genetic barrier against the genetic exchange of DNA that is not self-transmissible is dissimilarity in the bacterial DNA sequences concerned. Genetic exchange by homologous recombination is frequent among close bacterial relatives and recent experiments have shown that it can enable the uptake of closely linked nonhomologous foreign DNA. Artificial vectors are mosaics of mobile DNA elements from free-living bacterial isolates and so bear a residual similarity to their ubiquitous natural progenitors. This homology is tightly linked to the multitude of different DNA sequences that are inserted into synthetic vectors. Can homology between vector and bacterial DNA enable the

uptake of these foreign DNA inserts? In this review we investigate pUC18 as an example of an artificial vector and consider whether its homology to broad host-range antibiotic resistance transposons and plasmid origins of replication could enable the uptake of insert DNA in the light of studies of homology-facilitated foreign DNA uptake. We also discuss the disposal of recombinant DNA, its persistence in the environment and whether homologies to pUC18 may exist in naturally competent bacteria. Most DNA that is inserted into the cloning site of artificial vectors would be of little use to a bacterium, but perhaps not all.

*Heredity* (2004) 92, 483–489. doi:10.1038/sj.hdy.6800451

**Keywords:** horizontal gene transfer; biosafety; natural transformation

## Introduction

Gene flow across distantly related bacterial groups (horizontal gene transfer) is a major feature of bacterial evolution (Maynard Smith *et al*, 1991; Campbell, 2000; Ochman *et al*, 2000; Gogarten *et al*, 2002). This evolution need not be slow. The intense selection pressure imposed on microbial communities by worldwide antibiotic use reveals that new multiresistance plasmids can arise from diverse origins and spread in less than five decades (Hartl and Dykhuizen, 1984; Davies, 1994). In this case, the antibiotic resistance genes that spread so rapidly were mostly carried by mobile genetic elements.

Genetic exchange of DNA that is not self-transmissible can occur within a similar timeframe (Maynard Smith *et al*, 1991; Bowler *et al*, 1994), but such genetic exchanges occur by homologous recombination and so are usually restricted to very closely related bacteria. Thus the primary genetic barrier against genetic exchange of DNA that is not self-transmissible is dissimilarity in the bacterial DNA sequences concerned (DuBose *et al*, 1988; Rayssiguier *et al*, 1989; Maynard Smith *et al*, 1991; Vulic *et al*, 1997; Feil *et al*, 2001). The rate of homologous recombination is high enough to obliterate the phylogenetic signal among some close bacterial relatives (Feil *et al*, 2001), and for the nucleotide sites of some species, homologous recombination is a far more likely prospect than a point substitution (Guttman and Dykhuizen, 1994; Feil *et al*, 1999, 2000, 2001). Recent experiments show that close DNA sequence similarity can also facilitate the

stable acquisition of linked nonhomologous foreign DNA sequences (Nielsen *et al*, 2000a, b; de Vries and Wackernagel, 2002; Kay *et al*, 2002; Prudhomme *et al*, 2002).

The synthetic vectors that are heavily used in industry and research are mosaics of mobile DNA elements from natural bacterial isolates (Pouwels *et al*, 1988). Care was taken to render these vectors nontransmissible by removal of the genes necessary for their mobility (Berg *et al*, 1975; NIH guidelines for research involving recombinant DNA molecules: Appendix I, April 2002). Owing to their natural origins, artificial vectors still bear a residual similarity to DNA of naturally occurring bacteria. If this residual similarity were to enable recombination between bacterial DNA and only the parts of the artificial vectors that are homologous, there would most likely be no discernable effect on the bacterium involved. However, a plethora of different DNA sequences is inserted into artificial vectors and this raises the question whether homology between vector and bacterial DNA can facilitate the uptake of these foreign DNA inserts. In this review, we address this question by considering the extent of vector similarity to natural bacterial sequences, the requirements of homology-mediated foreign DNA uptake, the persistence of plasmid DNA and its bacterial uptake in the environment, and the possible effects of natural selection.

## The natural homologies of artificial vectors

How similar are the sequences of artificial vector DNA to those of naturally occurring bacterial DNA? We use the relatively small cloning vector pUC18 (a pBR322 derivative) to address this question, because it is one of the simplest synthetic vectors, is commonly used, and is

similar in sequence and organisation to many other artificial vectors.

The 2686 bp plasmid vector pUC18 is a mosaic of three different naturally occurring DNA sequences: an origin of replication and its flanking DNA from the *Escherichia coli* ColE1 plasmid, an antibiotic resistance gene and its flanking DNA from the transposon Tn3 of a naturally occurring resistance plasmid, and part of a  $\beta$ -d-galactosidase gene (LacZ) and flanking DNA from *E. coli* chromosomal DNA that may have originated from transposon Tn951 (Cornelis *et al*, 1978). The remaining 55 bp originate from the mostly synthetic multiple cloning site (MCS, or polylinker region originally developed in the bacteriophage M13) into which recombinant DNA is inserted (Figure 1) (Pouwels *et al*, 1988). The 'M13' universal primers that are generally used for PCR of recombinant DNA bind to sites that are actually homologous only to the  $\beta$ -d-galactosidase gene.

The largest component of pUC18 is the Tn3-like region containing the 860 bp TEM-1  $\beta$ -lactamase gene (*bla*<sub>TEM-1</sub>), which confers resistance to the antibiotic ampicillin, and other penicillins ( $\beta$ -lactam antibiotics). The Tn3-like region of pUC18 (1209 bp) is almost identical (97.5–99.8% similarity) to all reported natural TEM-type DNA sequences (Table 1). These include the TEM-type  $\beta$ -lactamases that are resistant to the new generation of synthetic  $\beta$ -lactams and those that are resistant to inhibitors manufactured to suppress  $\beta$ -lactamase activity. These highly related TEM-type sequences have spread, on Tn3-like transposons, to an extremely broad range of bacterial taxa. The TEM-type sequences deposited in GenBank alone (Table 1) represent species of commensal

and pathogenic bacteria of the Enterobacteriaceae, other Proteobacteria such as *Pseudomonas aeruginosa*, *Stenotrophomonas maltophilia*, *Neisseria meningitidis*, *Neisseria gonorrhoeae* (Table 1) and the phylogenetically distant gum-dwelling *Capnocytophaga ochracea* of the Flavobacteriia (Rosenau *et al*, 2000). TEM-type  $\beta$ -lactamases are carried by plasmid-borne transposons, to which pUC18 has at least 99% similarity, including Tn1, Tn2, Tn3, Tn801, Tn841 and TnSF1 (Table 1).


The broad distribution of these sequences is less surprising considering that  $\beta$ -lactams are the most commonly prescribed antibiotics worldwide, and that the single most common form of genetic resistance to these antibiotics is TEM-type  $\beta$ -lactamase production (Therrien and Levesque, 2000). Early work on antibiotic resistance has shown that TEM-type  $\beta$ -lactamases occur in a broad range of bacterial species, in many different countries, and on many different conjugative resistance plasmids, whose incompatibility groups include FI, FII, I $\alpha$ , N, A-C complex, H, L, M, P, S, T, W, X, and Y (Matthew and Hedges, 1976). The Tn3-like region of pUC18, and most ampicillin-resistant vectors, is derived from one of the first antibiotic resistance factors to be isolated in 1963 (Datta and Kontomichalou, 1965; Sutcliffe, 1978), and it is not surprising that one of the first natural isolates of an antibiotic resistance gene represents one of the most ubiquitous types.

The pUC18 ColE1-like replication region has at least 83% sequence identity to the replication origins of many naturally occurring, small, transmissible plasmids of the Enterobacteriaceae (Table 1). In some cases similarities flank the pUC18 cloning site, such that they also flank any DNA that is inserted into the vector. This is also the case for the  $\beta$ -d-galactosidase (LacZ) region (Table 1 and Figure 1).

The cloning vector pUC18 is not unusual among synthetic vectors. Most of the artificial vector sequences in the GenBank vector database share pUC18's ColE1-like origin of replication (671 of 911 sequences), and its Tn3-like region conferring ampicillin resistance (648 of 911). Other genes commonly used as selective markers for vectors are also derived from mobile, naturally occurring bacterial DNA sequences. For example, neomycin resistance (*nptII*) is conferred by a region of transposon Tn5, chloramphenicol resistance is from transposon Tn9, and kanamycin resistance (*nptI*) is from transposon Tn903 (Pouwels *et al*, 1988).

## A hitchhiker's guide to homology

How much homologous DNA sequence is required for the bacterial acceptance of how much foreign DNA? In a recent experiment, de Vries and Wackernagel (2002) grew Gram-negative naturally competent bacteria (*Acinetobacter* sp. BD413) in the presence of linear naked DNA. As little as 183 bp of homology between the linear naked DNA and a plasmid in the bacteria was enough to facilitate detectable levels of foreign DNA uptake. Further experiments, which used 1 kb of homology, showed that this homology enabled the frequent bacterial acquisition of a working bleomycin resistance gene that itself had no homology to the plasmid. This 1 kb of homology frequently anchored the stable integration of as much as 2.6 kb of foreign DNA and about 1 kb on average (de Vries and Wackernagel, 2002).



**Figure 1** The natural origins of plasmid vector pUC18. These were summarised using the details given in the GenBank entries of pUC18 (Accession: L08752), cloning vector pBR322 (J01749), cloning vector M13mp18 (X02513), the *E. coli* lactose operon (J01636), transposon Tn3 (V00613), the ColE1 plasmid (NC\_001371) and references therein, and by aligning these sequences with pUC18, using the NCBI 'BLAST 2 sequences' facility at <[www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)>.

**Table 1** Examples of homology between naturally occurring bacterial DNA and pUC18

DNA isolated from	Homology length (bp)	%S <sup>a</sup>	Description	Natural mobile element	Comments	Accession
Flavobacteria <i>Capnocytophaga ochracea</i> strain CIP105321	869	99.2	<i>bla</i> <sub>TEM-17</sub>	Uncharacterised plasmid	9 kb plasmid	Y14574
Proteobacteria Betaproteobacteria Neisseriales <i>Neisseria meningitidis</i> strains MC9690-129,-130	1209	99.4	<i>tnpR</i> , promoter, <i>bla</i> <sub>TEM-1</sub> , repeat	Transposon Tn3 on plasmid pAB6	5.6 kb plasmid has a mobilization and a replication protein	AF126482
<i>Neisseria gonorrhoeae</i> strain GC-182	1209	99.4	<i>tnpR</i> , promoter, <i>bla</i> , repeat	Transposon Tn2 on plasmid pJD4	7.4 kb plasmid has a mobilization and a replication protein	NC_002098
Gammaproteobacteria Pseudomonales <i>Pseudomonas aeruginosa</i> unknown strain	1209	99.3	<i>tnpR</i> , promoter, <i>bla</i> , repeat	Transposon Tn1 on plasmid RK2, and RP1	60 kb conjugative plasmid, broad host-range IncP-alpha	NC_001621
unknown strain	1209	99.2	<i>tnpR</i> , promoter, <i>bla</i> <sub>TEM-21</sub> , repeat	Transposon Tn801	Chromosomal, linked to IS6100 element	AF466526
unknown strain	>1062	99.8	<i>tnpR</i> , promoter, <i>bla</i>	Transposon Tn3	Broad host-range transposon	X50604
	>1062	99.4	<i>tnpR</i> , promoter, <i>bla</i>	Transposon Tn2	Broad host-range transposon	X50607
	>1062	99.2	<i>tnpR</i> , promoter, <i>bla</i>	Transposon Tn1	Broad host-range transposon	X50606
Xanthomonadales <i>Stenotrophomonas maltophilia</i> strain J6751a, UK	>1071	98.8	<i>tnpR</i> , promoter, <i>bla</i> <sub>TEM-2</sub>	Tn1-like transposon	Mobilizes to broad host-range conjugative plasmids	AJ251946
Enterobacteriales <i>Escherichia coli</i> many different strains strain SE53	861–1209	98.7–99.1	<i>tnpR</i> , promoter, <i>bla</i> , repeat	Tn1, 2 or 3-like transposons	>16 different TEM-type β-lactamases	See below <sup>b</sup>
strain K12 P678-54	672 <sup>c</sup>	86 <sup>d</sup>	ColE1-like replication origin	Plasmid pUD2380	8.5 kb mobilizable resistance plasmid	AJ008006
<i>Klebsiella pneumoniae</i> many different strains	629 <sup>c</sup>	87 <sup>d</sup>	ColE1-like replication origin	Plasmid pCloDF13	10 kb colicinogenic plasmid	NC_002119
	855–1209	99	<i>tnpR</i> , promoter, <i>bla</i> , repeat	Tn1, 2 or 3-like transposons, Tn1 is on multiresistance conjugative plasmid	9 different TEM-type β-lactamases.	See below <sup>c</sup>
<i>Klebsiella oxytoca</i> strain ROM	>1082	99.3	<i>tnpR</i> , promoter, <i>bla</i> <sub>TEM-12</sub> , repeat	Tn841 transposon on plasmid pOZ201	100 kb conjugative plasmid	M88143
strain Kox443	>862	99.1	<i>tnpR</i> , promoter, <i>bla</i> <sub>TEM-59</sub> , repeat	Conjugative plasmid	50 kb conjugative plasmid	AF062386
strain 26W	>1005	99.1	<i>tnpR</i> , promoter, part of <i>bla</i> <sub>TEM-10</sub>			U09188
strain KH11	401 <sup>c</sup>	83	ColE1-like replication origin	Plasmid pTKH11	8 kb cryptic resistance plasmid	Y17716
strain NBL63	625 <sup>c</sup>	88	ColE1-like replication origin	Plasmid pNBL63	7 kb cryptic resistance plasmid	Y17846
<i>Shigella flexneri</i> strains K24, K25, Hong Kong strain SH595	>1067	99.3	<i>tnpR</i> , promoter, <i>bla</i>	Tn2 on conjugative plasmid	120 kb broad host-range multiresistance plasmid	U48775
<i>Salmonella enteritidis</i> strain YMC4199 serovar Saintpaul, Korea	1209	99.5	<i>tnpR</i> , promoter, <i>bla</i> , repeat	Transposon TnSF1	Tn-21-like transposon	AF188331
	>836	99	<i>bla</i> <sub>TEM-52</sub>	Plasmid pYMC4199		AF126444
<i>Salmonella typhimurium</i> unknown strain	1209 & 545	99 and 95	<i>tnpR</i> , promoter, <i>bla</i> , repeat, ColE1-like replication origin	Plasmid pNTP16	8.3 kb mobilizable multi-resistance plasmid	NC_002090

Table 1 (continued)

DNA isolated from	Homology length (bp)	%S <sup>a</sup>	Description	Natural mobile element	Comments	Accession
<i>Salmonella enterica</i> serovar Borreze	469 <sup>c</sup> & 137	87 and 92	ColE1-like replication origin	Plasmid pWQ799	6.9 kb mobilizable plasmid carrying genes for O:54 antigen expression	L39794
<i>Salmonella panama</i> unknown strain	684 <sup>c</sup>	94	ColE1-like replication origin	Plasmid RSF1030	8.3 kb resistance plasmid carrying Tn2	J01784
<i>Yersinia pestis</i> strain CO-92 Biovar Orientalis	972 <sup>c</sup>	95–98	ColE1-like replication origin	Plasmid pPCP1	9.6 kb virulence plasmid essential for plague transmission by flea bite	AL109969
<i>Yersinia enterocolitica</i> strain 29807, biogroup 1A	447 <sup>c</sup>	90	ColE1-like replication origin	Plasmid p29807	2.7 kb cryptic plasmid	AJ32618
strain 842 (O:8,19)	> 118 & > 160 & 38	96.6, 96.3, 100	LacI, flank & β-d-galactosidase and Tn3 inverted repeat	Transposon Tn951 on 50 kb conjugative plasmid pGC1	16.6 kb lac transposon	M82979, M25019
<i>Proteus mirabilis</i> unknown strain	> 1084	99.4	<i>tnpR</i> , promoter, <i>bla</i> <sub>IRT-18</sub>			AJ012256
strain H223b, Tunisia	> 885	99.3	<i>bla</i> <sub>TEM-1</sub>	Plasmid pCMY-4		Y18200
strain FI-14, Italy	> 1008	98.8	<i>bla</i> <sub>TEM-72</sub>	Plasmid pPM14	40 kb plasmid	AF157413
<i>Morganella morgani</i> strain FFLM15	> 1095	98.8	<i>tnpR</i> , promoter, <i>bla</i> <sub>TEM-10</sub>	Conjugative plasmid	50 kb conjugative plasmid	AF093512
strain Mm126	> 919	98.9	<i>bla</i> <sub>TEM-21</sub>	Plasmid pMm126	40 kb conjugative multiresistance plasmid	AF052748
strain FI-13, Italy	> 1008	98.8	<i>bla</i> <sub>TEM-72</sub>	Plasmid pMN13	40 kb plasmid	AF157553
<i>Serratia marcescens</i> strain S5 (epidemic)	> 1016	97.5	<i>bla</i> <sub>TEM-AQ</sub>	Plasmid pSM5	54 kb plasmid	X97254
unknown strain	220	100	<i>tnpR</i> , promoter, part of <i>bla</i>	part of Tn3 on plasmid pAZ007	Multidrug resistance plasmid	M23634
<i>Pantoea agglomerans</i> from a kitchen sink	556 <sup>c</sup>	89 <sup>d</sup>	ColE1-like replication origin	Plasmid pPIGDM1	2.5 kb plasmid	AF014880
<i>Providencia stuartii</i> strain VR1, Italy	997	98.2	<i>bla</i> <sub>TEM-60</sub>	Plasmid pVR-1	20 kb plasmid	AF047171
<i>Edwardsiella ictaluri</i> unknown strain	467 <sup>c</sup>	84 <sup>d</sup>	ColE1-like replication origin	Plasmid pEI1	4.8 kb plasmid	NC_002497

<sup>a</sup>% similarity to pUC18.<sup>b</sup>V00613, AF188200, X57972, U37195, Y17584, AF190692-5, and AB049569.<sup>c</sup>BLAST alignment was gapped.<sup>d</sup>Weighted mean of sections aligned by BLAST.<sup>e</sup>X64523, Y17581-3, L19940, U95363, Y13612, AF104441, AJ239002, X65252,3,7.

pUC18 (Accession: L08752) was used to BLAST completed microbial genomes and all DNA sequences with the organism name 'plasmids'. NCBI sequence entries containing keywords such as 'bla NOT vector' were also compared to pUC18 using pairwise BLAST. To avoid contaminating results from accidental inclusions of vector sequences in GenBank sequence entries, results were excluded if they fulfilled several criteria: (i) they were not diverged from pUC18; (ii) if homology was at the start or the end of a sequence entry; (iii) if this homology was not a subset of a longer region of homology to the sequence of the naturally occurring Tn3 transposon (Accession: V00613), or ColE1 (Accession: NC\_001371); (iv) if reading of publications associated with the DNA sequence revealed that a synthetic vector with homology to pUC18 had been used to generate the DNA sequence.

Prudhomme *et al* (2002) present very similar findings for *Streptococcus pneumoniae*, a naturally competent Gram positive bacterium, where less than 1 kb of one-sided homology is sufficient for the chromosomal uptake of over 2 kb of foreign DNA. In the absence of DNA homology no stable integration of the naked DNA with bacterial DNA could be detected, but when homology to bacterial DNA flanks both sides of the foreign DNA the rate of uptake is higher than in the case of one-sided homology (de Vries and Wackernagel, 2002; Prudhomme *et al*, 2002) and close to the rate at which naked DNA is able to enter the bacterium (Nielsen *et al*, 2000a, b; de Vries *et al*, 2001). Prudhomme *et al* also show that a single homologous region (942 bp) facilitates the integration of whole plasmids into bacterial chromosomes at a 100-fold greater rate than foreign DNA uptake from the same plasmid when linearised.

Several studies estimate the degree of similarity required for homologous recombination in different types of bacteria. The frequency of recombination of DNA molecules in bacteria, such as the enterobacteria (Vulic *et al*, 1997, 1999), *Bacillus* (Majewski and Cohan, 1998), and *Streptococcus* (Majewski *et al*, 2000), increases exponentially as the nucleotide similarity of the participating DNA molecules increases. Most wild-type bacteria abort effective recombination with DNA that is over 1–2% diverged from their own; however, the constraints on sequence similarity for homologous recombination can be relaxed to ~20% divergence in certain 'mutator' individuals (Rayssiguier *et al*, 1989; Vulic *et al*, 1997, 1999; Majewski *et al*, 2000), and others predict recombination between molecules that are up to 30% diverged (Townsend *et al*, 2003). Mutator individuals make up 1–2% of natural *E. coli* and *Salmonella* populations (LeClerc *et al*, 1996; Matic *et al*, 1997), though one study of patients with cystic fibrosis, found that, on average, 20% of the pathogenic isolates of *P. aeruginosa* examined showed mutator phenotypes (Oliver *et al*, 2000). Homologous recombination in mutators could explain the finding that *S. pneumoniae* strains became resistant to antibiotics in the last 50 years through changes to their penicillin-binding protein genes by at least two independent recombination events, involving DNA sequences that were up to 20% divergent (Maynard Smith *et al*, 1991).

Most of the homologies described in Table 1 are between pUC18 and Tn3-like transposons. Most of these are less than 1% diverged from naturally occurring bacterial DNA sequences and span more than 1 kb. If artificial vectors and their inserts were available to bacteria, such sequence similarity would seem sufficient to facilitate the bacterial uptake of insert DNA. The sequences with which pUC18 has Tn3-like homology are on broad host-range conjugative plasmids and are associated with broad host-range mobile elements (Table 1). If new DNA sequences could be associated with such mobile elements, this may increase the chance of it spreading to other bacteria. The homology of pUC18 to plasmids with ColE1-like origins of replication is also sufficiently long for the uptake of inserted DNA sequences, but natural and artificial sequences are up to 17% diverged so homology-facilitated uptake of foreign DNA would seem likely only in mutators. Homologies of ColE1-like origins of replication are generally associated with small nonconjugative plasmids (Table 1) so even if recombination did occur with these, it

probably would not significantly increase the chances of spread to other bacterial species.

## Bacterial competence and naked DNA

Are artificial vectors and their inserts available to free-living bacteria? If DNA is associated with the appropriate mobile elements in living bacteria, genetic exchange can proceed by conjugation or transduction, and thus reach a broad range of bacteria at a potentially high rate. Artificial cloning vectors are not mobile, however, and the laboratory microorganisms that propagate them are killed prior to their environmental release. Recombinant DNA will not be propagated in the environment unless free-living bacteria pick it up by natural transformation. Given that most of the mobile elements with which artificial vectors have homology are the very ones that carried antibiotic resistance genes across bacterial phyla, the product of an initial natural transformation event could then spread in a relatively unimpeded way.

Some bacteria, for example *S. pneumoniae*, *Acinetobacter* sp. Strain BD13, *Pseudomonas stutzeri*, *N. gonorrhoeae* (reviewed in Lorenz and Wackernagel, 1994), are naturally competent, that is they are able to accept naked DNA from the environment into their cells. The various experiments that show how homology facilitates natural transformation, also illustrate how readily these bacterial species can absorb naked DNA (Nielsen *et al*, 2000a, b; de Vries *et al*, 2001; de Vries and Wackernagel, 2002). It is likely that some naturally competent bacteria carry elements homologous to artificial vectors because of the broad host-range of the elements to which pUC18 is homologous (Table 1; Matthew and Hedges, 1976).

There is tight homology between pUC18 and mobile elements in *Neisseria* species (Table 1), which are naturally competent. However, *Neisseria* (and *Haemophilus*) only take up DNA that is connected to specific uptake sequences (Solomon and Grossman, 1996). In *Neisseria* the specific uptake sequence required is 10 nucleotides long and there is no exact match to this in the pUC18 DNA sequence. In other species, such as *Acinetobacter* and *P. stutzeri*, DNA uptake is not sequence specific.

Several genes resembling those that encode the cellular machinery required for natural transformation exist in bacterial species that are not known for their natural competence, for example in *E. coli*, *Salmonella typhi*, *Klebsiella pneumoniae*, and *P. aeruginosa* (Finkel and Kolter, 2001). Some species are also known to be naturally competent in particular situations (Lorenz and Wackernagel, 1994; Demaneche *et al*, 2001b) and so could potentially absorb artificial DNA if it is available in their environment. For example, *E. coli* may become competent in mineral water, when hungry, when in human food, and when in soil that is struck by lightning (Bauer *et al*, 1999; Finkel and Kolter, 2001; Woegerbauer *et al*, 2002; Demaneche *et al*, 2001a), and *Pseudomonas fluorescens* is competent in soil but not *in vitro* (Demaneche *et al*, 2001b).

Could free-living bacteria be exposed to recombinant DNA in the environment? DNA can persist in the environment for thousands of years (Austin *et al*, 1997), though probably not in a form where the genes it may encode are readily utilizable. Experiments so far have demonstrated the transformation and unharmed activity

of genes after a few days in soil (Nielsen *et al*, 1997; Sikorski *et al*, 1998).

There are no recommendations or guidelines on the disposal of recombinant DNA, only on the disposal of living genetically modified organisms (e.g. NIH guidelines for research involving recombinant DNA molecules (2002): Sections I-B and III-F-1, and Appendices G and K). Recombinant DNA is most likely poured down the sink in large and diverse quantities and genes can stay intact even when heated or chemically treated along with the killed microbial biomass from which they came (Andersen *et al*, 2001).

## Natural selection

Most of the DNA that is inserted into artificial vectors would probably not benefit free-living bacteria. Without a phenotypic benefit to recipient bacteria, it is unlikely that non-homologous DNA picked up from artificial vectors would spread or persist (Campbell, 2000; Berg and Kurland, 2002), and it is unclear why such acquisitions would be of any concern. Given the large-scale genome sequencing of deadly microorganisms or the scope of research into virulence, however, there are almost certainly DNA inserts released in artificial vectors that would have a chance of persistence and would be of concern if acquired by some bacteria.

Even genetic variants that do not confer a beneficial trait sometimes spread among bacteria if linked to an advantageous locus (Gutman and Dykhuizen, 1994). The plasmids and transposons to which artificial vectors have homologies generally confer a whole suite of potentially beneficial traits, for example, multiple drug resistance or pathogenic characteristics (Table 1). Close linkage to such traits increases the chances of natural recombinants persisting in the environment even if newly acquired foreign DNA does not confer an immediate selective advantage on recipient bacteria.

## Conclusions

Many factors influence the risk that the natural homology of artificial vectors could facilitate the bacterial uptake of recombinant DNA inserts. As several of these factors are notoriously difficult to quantify in an open environment, the degree to which this risk is realistic remains unknown. The processes involved occur at low frequencies, but could act on a large number of different molecules, bacteria and potential environmental situations. Given the mobility of the elements to which artificial vectors have homology, the acquisition of an undesirable trait that is beneficial to bacteria need only happen once for potentially far-reaching consequences. Most DNA that is inserted into artificial vectors would most likely pose no risk to human health or the environment even if acquired by free-living bacteria.

## Acknowledgements

Many thanks for helpful comments and criticisms on various incarnations of the manuscript to Daniel L Hartl, M Pilar Francino, Richard H Thomas, Casey M Bergman, and anonymous reviewers. KMN was supported by The Research Council of Norway (MU: biodiversity program) and the Norwegian Institute for Gene Ecology; DB was

partly supported by a Royal Society – Fulbright fellowship. Part of this work was performed under the auspices of the US Department of Energy, Office of Biological and Environmental Research, by the University of California, Lawrence Berkeley National Laboratory under contract No. DE-AC0376SF00098.

## References

- Andersen J, Schafer T, Jorgensen P, Moller S (2001). Using inactivated microbial biomass as fertilizer: the fate of antibiotic resistance genes in the environment. *Res Microbiol* **152**: 823–833.
- Austin JJ, Smith AB, Thomas RH (1997). Palaeontology in a molecular world: the search for authentic ancient DNA. *Trends Ecol Evol* **12**: 303–306.
- Bauer F, Hertel C, Hammes WP (1999). Transformation of *Escherichia coli* in foodstuffs. *Syst Appl Microbiol* **22**: 161–168.
- Berg OG, Kurland CG (2002). Evolution of microbial genomes: sequence acquisition and loss. *Mol Biol Evol* **19**: 2265–2276.
- Berg P, Baltimore D, Brenner S, Roblin RO, Singer MF (1975). Summary statement of the Asilomar conference on recombinant DNA molecules. *Proc Nat Acad Sci USA* **72**: 1981–1984.
- Bowler LD, Zhang Q-Y, Riou J-Y, Spratt BG (1994). Interspecies recombination between the *penA* genes of *Neisseria meningitidis* and commensal *Neisseria* species during the emergence of penicillin resistance in *N. meningitidis*: natural events and laboratory simulation. *J. Bacteriol.* **176**: 333–337.
- Campbell AM (2000). Lateral gene transfer in prokaryotes. *Theor Pop Biol* **57**: 71–77.
- Cornelis G, Ghosal D, Saedler H (1978). Tn951: a new transposon carrying a lactose operon. *Mol Gen Genet* **160**: 215–224.
- Datta N, Kontomichalou P (1965). Penicillinase synthesis controlled by infectious *R* factors in Enterobacteriaceae. *Nature* **208**: 239–241.
- Davies J (1994). Inactivation of antibiotics and the dissemination of resistance genes. *Science* **264**: 375–382.
- de Vries J, Meier P, Wackernagel W (2001). The natural transformation of the soil bacteria *Pseudomonas stutzeri* and *Acinetobacter* sp. by transgenic plant DNA strictly depends on homologous sequences in the recipient cells. *FEMS Microbiol. Lett.* **195**: 211–215.
- de Vries J, Wackernagel W (2002). Integration of foreign DNA during natural transformation of *Acinetobacter* sp. by homology-facilitated illegitimate recombination. *Proc. Nat. Acad. Sci. USA* **99**: 2094–2099.
- Demaneche S, Bertolla F, Buret F, Nalin R, Sailland A, Auriol P *et al* (2001a). Laboratory-scale evidence for lightning-mediated gene transfer in soil. *Appl Environ Microbiol* **67**: 3440–3444.
- Demaneche S, Kay E, Gourbiere F, Simonet P (2001b). Natural transformation of *Pseudomonas fluorescens* and *Agrobacterium tumefaciens* in soil. *Appl Environ Microbiol* **67**: 2617–2621.
- DuBose RF, Dykhuizen DE, Hartl DL (1988). Genetic exchange among natural isolates of bacteria: recombination within the *phoA* gene of *Escherichia coli*. *Proc Nat Acad Sci USA* **85**: 7036–7040.
- Feil EJ, Holmes EC, Bessen DE, Chan M-S, Day NPJ, Enright MC *et al* (2001). Recombination within natural populations of pathogenic bacteria: short-term empirical estimates and long-term phylogenetic consequences. *Proc Nat Acad Sci USA* **98**: 182–187.
- Feil EJ, Maiden CJ, Achtman M, Spratt BG (1999). The relative contributions of recombination and mutation to the divergence of clones of *Neisseria meningitidis*. *Mol Biol Evol* **16**: 1496–1502.
- Feil EJ, Maynard Smith J, Enright MC, Spratt BG (2000). Estimating recombinational parameters in *Streptococcus pneu-*

- moniae* from multilocus sequence typing data. *Genetics* **154**: 1439–1450.
- Finkel SE, Kolter R (2001). DNA as a nutrient: novel role for bacterial competence gene homologs. *J Bacteriol* **183**: 6288–6293.
- Gogarten JP, Doolittle WF, Lawrence JG (2002). Prokaryotic evolution in the light of gene transfer. *Mol Biol Evol* **19**: 2226–2238.
- Guttman DS, Dykhuizen DE (1994). Clonal divergence in *Escherichia coli* as a result of recombination, not mutation. *Science* **266**: 1380–1383.
- Hartl DL, Dykhuizen DE (1984). The population genetics of *Escherichia coli*. *Annu Rev Genet* **18**: 31–68.
- Kay E, Vogel TM, Bertolla F, Nalin R, Simonet P (2002). In situ transfer of antibiotic resistance genes from transgenic (transplastomic) tobacco plants to bacteria. *Appl Environ Microbiol* **68**: 3345–3351.
- LeClerc JE, Li B, Payne WL, Cebula TA (1996). High mutation frequencies among *Escherichia coli* and *Salmonella* pathogens. *Science* **274**: 1208–1211.
- Lorenz MG, Wackernagel W (1994). Bacterial gene transfer by natural genetic transformation in the environment. *Microbiol Rev* **58**: 563–602.
- Majewski J, Cohan FM (1998). The effect of mismatch repair and heteroduplex formation on sexual isolation in *Bacillus*. *Genetics* **148**: 13–18.
- Majewski J, Zawadzki P, Pickerill P, Cohan FM, Dowson CG (2000). Barriers to genetic exchange between bacterial species: *Streptococcus pneumoniae* transformation. *J Bacteriol* **182**: 1016–1023.
- Matic I, Radman M, Taddei F, Picard F, Doit C, Bingen E et al (1997). Highly variable mutation rates in commensal and pathogenic *Escherichia coli*. *Science* **277**: 1833–1834.
- Matthew M, Hedges RW (1976). Analytical isoelectric focusing of R Factor-determined beta-lactamases: correlation with plasmid compatibility. *J Bacteriol* **125**: 713–718.
- Maynard Smith J, Dowson CG, Spratt BG (1991). Localized sex in bacteria. *Nature* **349**: 29–31.
- Nielsen KM, Smalla K, van Elsas J (2000a). Natural transformation of *Acinetobacter* sp. strain BD413 with cell lysates of *Acinetobacter* sp., *Pseudomonas fluorescens*, and *Burkholderia cepacia* in soil microcosms. *Appl Environ Microbiol* **66**: 206–212.
- Nielsen KM, van Elsas JD, Smalla K (2000b). Transformation of *Acinetobacter* sp. BD413(pFGΔntpII) with transgenic plant DNA in soil microcosms and effects of kanamycin on selection of transformants. *Appl Environ Microbiol* **66**: 1237–1242.
- Nielsen KM, van Weerelt MDM, Berg TN, Bones AM, Hagler AN, van Elsas JD (1997). Natural transformation and availability of transforming DNA to *Acinetobacter calcoaceticus* in soil microcosms. *Appl Environ Microbiol* **63**: 1945–1952.
- NIH guidelines for research involving recombinant DNA molecules (April 2002). <http://www4.od.nih.gov/oba/rac/guidelines/guidelines.html>.
- Ochman H, Lawrence JG, Groisman EA (2000). Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**: 299–304.
- Oliver A, Canton R, Campo P, Baquero F, Blasquez J (2000). High frequency of hypermutable *Pseudomonas aeruginosa* in cystic fibrosis lung infection. *Science* **288**: 1251–1253.
- Pouwels PH, Enger-Valk BE, Brammar WJ (1988). *Cloning Vectors*. Elsevier: Amsterdam.
- Prudhomme M, Libante V, Claverys J-P (2002). Homologous recombination at the border: insertion–deletions and the trapping of foreign DNA in *Streptococcus pneumoniae*. *Proc Natl Acad Sci USA* **99**: 2100–2105.
- Rayssiguier C, Thaler DS, Radman M (1989). The barrier to recombination between *Escherichia coli* and *Salmonella typhimurium* is disrupted in mismatch repair mutants. *Nature* **342**: 396–401.
- Rosenau A, Cattier B, Gousset N, Harriau P, Philippon A, Quentin R (2000). *Capnocytophaga ochracea*: characterization of a plasmid-encoded extended-spectrum TEM-17 beta-lactamase in the phylum *Flavobacter-Bacteroides*. *Antimicrob Agents Chemother* **44**: 760–762.
- Sikorski J, Graupner S, Lorenz MG, Wackernagel W (1998). Natural genetic transformation of *Pseudomonas stutzeri* in a non-sterile soil. *Microbiology* **144**: 569–576.
- Solomon JM, Grossman AD (1996). Who's competent and when: regulation of natural genetic competence in bacteria. *Trends Genet* **12**: 150–155.
- Sutcliffe JG (1978). Nucleotide sequence of the ampicillin resistance gene of *Escherichia coli* plasmid pBR322. *Proc Natl Acad Sci USA* **75**: 3737–3741.
- Therrien C, Levesque RC (2000). Molecular basis of antibiotic resistance and beta-lactamase inhibition by mechanism-based inactivators: perspectives and future directions. *FEMS Microbiol Rev* **24**: 251–262.
- Townsend JP, Nielsen KM, Fisher DS, Hartl DL (2003). Horizontal acquisition of divergent chromosomal DNA in bacteria: effects of mutator phenotypes. *Genetics* **164**: 13–21.
- Vulic M, Dionisio F, Taddel F, Radman M (1997). Molecular keys to speciation: DNA polymorphism and the control of genetic exchange in enterobacteria. *Proc Natl Acad Sci USA* **94**: 9763–9767.
- Vulic M, Lenski RE, Radman M (1999). Mutation, recombination, and incipient speciation of bacteria in the laboratory. *Proc Natl Acad Sci USA* **96**: 7348–7351.
- Woegerbauer M, Jenni B, Thalhammer F, Graninger W, Burgmann H (2002). Natural genetic transformation of clinical isolates of *Escherichia coli* in urine and water. *Appl Environ Microbiol* **68**: 440–443.