

SHORT COMMUNICATION

Interaction of *Salmonella enterica* with basil and other salad leaves

Cedric N Berger^{1,5}, Robert K Shaw^{2,5}, Derek J Brown³, Henry Mather³, Simon Clare⁴, Gordon Dougan⁴, Mark J Pallen² and Gad Frankel¹

¹Department of Life Science, Division of Cell and Molecular Biology, Imperial College London, London, UK;

²Division of Immunity and Infection, School of Medicine, University of Birmingham, Birmingham, UK;

³Scottish Salmonella Reference Laboratory, Stobhill Hospital, Glasgow, Scotland and ⁴Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, UK

Contaminated salad leaves have emerged as important vehicles for the transmission of enteric pathogens to humans. A recent outbreak of *Salmonella enterica* serovar Senftenberg (*S. Senftenberg*) in the United Kingdom has been traced to the consumption of contaminated basil. Using the outbreak strain of *S. Senftenberg*, we found that it binds to basil, lettuce, rocket and spinach leaves showing a pattern of diffuse adhesion. Flagella were seen linking *S. Senftenberg* to the leaf epidermis, and the deletion of *fljC* (encoding phase-1 flagella) resulted in a significantly reduced level of adhesion. In contrast, although flagella linking *S. enterica* serovar Typhimurium to the basil leaf epidermis were widespread, deletion of *fljC* did not affect leaf attachment levels. These results implicate the role of flagella in *Salmonella* leaf attachment and suggest that different *Salmonella* serovars use strain-specific mechanisms to attach to salad leaves.

The ISME Journal (2009) 3, 261–265; doi:10.1038/ismej.2008.95; published online 2 October 2008

Subject Category: microbe–microbe and microbe–host interactions

Keywords: *Salmonella enterica*; flagella; salad leaves

Salmonella enterica has long been recognized as an important source of food-borne infections and has commonly been associated with the consumption of contaminated poultry and poultry products (Olsen *et al.*, 1992; Baggesen and Wegener, 1994). The global increase in human infections with serovar Enteritidis observed in the late 1980s and early 1990s (Rodrigue *et al.*, 1990) was almost entirely attributable to the presence of this organism within the poultry production industry. However, implementation of control measures including vaccination has resulted in recent years in a reduction in cases of human salmonellosis associated with the consumption of poultry and egg products (Kessel *et al.*, 2001).

In recent years, there has been an increase in consumer demand for fruit, salad and vegetables. Such ready-to-eat foods require little or no cooking before consumption. In a study of general outbreaks of infectious intestinal diseases in England and Wales, carried out between 1992 and 2006, 23% of

outbreaks were of food-borne origin and 4% of food-borne outbreaks were linked to prepared salad (reviewed in Little and Gillespie, 2008). *Salmonella* was the most commonly identified pathogen acquired from fresh produce in the United States, being isolated in 48% of cases between 1973 and 1997 (Sivapalasingham *et al.*, 2004).

The newsworthy international outbreaks of *S. enterica* that have been linked to ready-to-eat plant produce include a Scandinavian/UK outbreak of *S. enterica* serovar Thompson associated with consumption of rucola lettuce (Nygård *et al.*, 2008), a Danish outbreak of *S. enterica* serovar Anatum infection linked to imported basil (Pakalniskiene *et al.*, 2006), a tomato-associated outbreak of *S. enterica* serovar Braenderup infection in the United States (Gupta *et al.*, 2007), an outbreak of *S. enterica* serovar Typhimurium (*S. Typhimurium*) DT204b in several European countries associated with the consumption of lettuce (Crook *et al.*, 2003) and an outbreak of *S. enterica* serovar Senftenberg (*S. Senftenberg*) infection associated with imported Israeli basil affecting the United Kingdom, Denmark, the Netherlands and the United States (Pezzoli *et al.*, 2007).

A recent study has shown differential binding of *S. enterica* to different lettuce cultivars (Klerks *et al.*, 2007). Adhesion of *Salmonella* to alfalfa sprouts was shown to be associated with thin aggregative

Correspondence: G Frankel, Department of Life Science, Division of Cell and Molecular Biology, Flowers Building, Imperial College, London SW7 2AZ, UK.

E-mail: g.frankel@imperial.ac.uk

⁵These authors contributed equally to this work.

Received 10 July 2008; revised 1 September 2008; accepted 2 September 2008; published online 2 October 2008

fimbriae (Tafi), the extracellular cellulose matrix involved in biofilm formation and the O-antigen capsule (Barak *et al.*, 2007). The aim of this study was to investigate the mechanism used by *S. enterica* to attach to salad leaves.

Leaf adhesion assays were initially performed using the UK basil outbreak *S. Senftenberg* strain 070885. Leaves were inoculated and processed as we described before for *Escherichia coli* O157 (Shaw *et al.*, 2008). Briefly, freshly excised leaves of 10 mm width from supermarket-sourced basil, lettuce, rocket and spinach plants were trimmed, affixed to petri dish base and immersed in 4 ml of bacterial cultures or bacterial culture pellets re-suspended in H₂O (OD₆₀₀ = 1.00, equivalent to $6.5 \pm 0.7 \times 10^8$ colony-forming unit (CFU) ml⁻¹) and incubated statically at 20 or 37 °C for 1 h; non-adherent bacteria were removed by washing. Inoculated leaves were processed for scanning electron microscopy (SEM) and immunofluorescence as described (Shaw *et al.*, 2008).

Diffusely adherent *S. Senftenberg*, covering large areas of basil (Figure 1a), lettuce, rocket and spinach (data not shown) leaf surfaces, were observed by SEM and immunofluorescence. Similar attachment levels and patterns were seen after incubations at 20 or 37 °C. No adherent epiphyte microorganisms were seen on unwashed or mock-inoculated leaves (data not shown). High-resolution SEM revealed peritrichous filaments linking *S. Senftenberg* to the leaf epidermis (Figure 1c). We used monoclonal JIM5 pectin antibodies (Plant Probes) (together with anti-mouse Alexa Fluor 594), anti-FliC_{g,s,t} antiserum (together with goat anti-rabbit Alexa Fluor 488) and immunofluorescence (Shaw *et al.*, 2008) to determine if the filaments are flagella. The depicted figures, representative of randomly selected fields, show abundance of flagella (green) linking *S. Senftenberg* (blue) to the leaf epidermis (red; Figure 1d).

To determine the role flagella play in leaf attachment, we deleted the phase-1 flagella *fliC* gene from *S. Senftenberg* 070885. Deletion of *fliC* and insertion of a Kn cassette were performed using the lambda red system (Datsenko and Wanner, 2000), generating strain ICC301. Primer pair Flic/Kn Forward (5'-ATGCCACAAGTCATTAATACCAACAGCCTCTCGCTGATCACTCAAATAATATCTGTGTAGGCTGGAGCTGCTTCG) and Flic/Kn Reverse (5'-TTAACGCAGTAAAGAGAGGACGTTTTGCGGAACCTGGTTCGCCTGCGCCAGCATATGAATATCCTCCTTAG) was used to amplify the kanamycin cassette in pKD4. Control *fliC* Forward and Reverse primers (5'-gtgccgatacaagggttacggtgag and 5'-GGAATTA AAAAAGGCAGCTTTCGCTG, respectively) and SEM verified the deletion and lack of flagellar expression (data not shown). Basil leaves were inoculated with *fliC* mutant bacterial cultures (OD₆₀₀ = 1.00, equivalent to $6.5 \pm 0.7 \times 10^8$ CFU ml⁻¹) and incubated statically at 20 or 37 °C for 1 h as above. The number of leaf-adherent bacteria was quantified by excising

30-mm² washed leaf disks into sterile water. The leaf disks were homogenized and serial dilutions were plated onto MacConkey agar plates. This revealed that at both 20 and 37 °C, wild-type *S. Senftenberg* adhered at significantly higher levels than the Δ *fliC* mutant (Figure 1e); the low-level-attached Δ *fliC* mutant bacteria formed aggregates on the leaf surface (Figure 1b). No bacterial growth was seen after plating gently washed and homogenized uninfected leaf extracts (data not shown).

We next investigated the ability of other *Salmonella* serovars (OD₆₀₀ = 1.00, equivalent to $6.5 \pm 0.7 \times 10^8$ CFU ml⁻¹) to bind basil leaves as above. Using SEM we found that *S. Typhimurium* (strain SL1344) (Figure 2a) and *S. enterica* serovar Enteritidis strain PT4 (data not shown) adhered efficiently through long filamentous structures (Figure 2b), whereas no adhesion was seen, after scanning 1.2 mm² (in two independent experiments), on leaves inoculated with *S. enterica* serovar Arizona (strain SARC5), *S. enterica* serovar Heidelberg (strain SARB23) and *S. enterica* serovar Agona (strain SARB1) (data not shown). The *S. Typhimurium* filaments were identified as phase-1 flagella by staining with mouse monoclonal anti-FliC_i antibodies and goat anti-mouse Alexa Fluor 488 (Figure 2c). No phase-1 flagella were seen by immunofluorescence on the *S. Typhimurium* Δ *fliC* mutant (Figure 2d). In contrast to *S. Senftenberg*, no significant difference was observed in adhesion levels (leaves inoculated with $6.5 \pm 0.7 \times 10^8$ CFU ml⁻¹) between the wild-type strain and *S. Typhimurium* Δ *fliC* mutant when incubations were carried out at either 20 or 37 °C (Figures 2d and e). These results suggest the existence of serovar-specific attachment mechanisms, which is in agreement with a report by Klerks *et al.* (2007).

Concluding remarks

Although the mechanisms used by enteric pathogens to colonize the mammalian gut mucosa have been extensively studied, those involved in attachment to vegetables and salad leaves are not well known. In this study, we have shown that the outbreak *S. Senftenberg* isolate attached to a variety of salad leaves (including basil, lettuce rocket and spinach) and that flagella played a major role in bacterial leaf interaction. In contrast, although abundant flagella were seen linking *S. Typhimurium* to basil leaf surface, deletion of the gene for the phase-1 flagellin *fliC* had no measurable effect on the level of leaf association.

The incidence of human infection by enteric bacteria through the consumption of contaminated salad leaves has increased in the last few years (Little and Gillespie, 2008). Leaf contamination can occur during crop growth (for example, through contaminated water, wild or domesticated animals, flies or birds), harvest, distribution, processing and




Figure 1 Attachment of *Salmonella enterica* serovar Senftenberg to basil leaves. Wild-type *S. Senftenberg* attaches in a diffuse pattern (**a**) whereas clumps of *S. Senftenberg* $\Delta fliC$ were seen at a low frequency (**b**). Peritrichous flagella-like structures bound laterally to the leaf surface were observed linking *S. Senftenberg* to the leaf epidermis (**c**). These structures were shown to be flagella (green) (arrows and inset) using FliC antibodies (**d**). *S. Senftenberg* and the leaf epidermis are shown in blue and red, respectively. Flagella-like structures were not seen on *S. Senftenberg* $\Delta fliC$ (**b**). Bars = 10 μ m (**a**, **b** and **d**), 1 μ m (**c**) and 0.5 μ m (**b** and **d**, inset). Quantification of leaf attachment by wild-type and *S. Senftenberg* $\Delta fliC$ to basil leaves at 20 or 37 °C (**e**). Deletion of *fliC* resulted in a significant reduction in leaf attachment. Results are presented as mean \pm s.d. Significant differences are indicated by asterisks (* P < 0.05).




Figure 2 Attachment of *Salmonella enterica* serovar Typhimurium to basil leaves. Wild-type (a) and $\Delta fliC$ *S. Typhimurium* (d) attach to leaves in a diffuse pattern. Peritrichous flagella-like structures bound laterally to the leaf surface were observed linking *S. Typhimurium* to the leaf epidermis (b). These structures were shown to be flagella (green) (arrows and inset) using FliC antibodies (c). *S. Typhimurium* and the leaf epidermis are shown in blue and red, respectively. No FliC staining was seen on *S. Typhimurium* $\Delta fliC$ (d). Bars = 10 μ m (a, c and d), 0.1 μ m (b) and 0.2 μ m (c, inset). Quantification of leaf attachment of wild-type and *S. Typhimurium* $\Delta fliC$ to basil leaves at 20 or 37 °C (e). Deletion of *fliC* did not affect leaf attachment levels. Results are presented as mean \pm s.d.

packing or cooking. A better understanding of the mechanism involved in the attachment of *S. enterica* to salad leaves would be useful in developing interventions to minimize contamination and transmission and in the development of accurate risk assessments.

Acknowledgements

We thank Dr Jay Hinton, IFR, Norwich for *S. Typhimurium* Δ *fliC* and Ms Florencia Minuzzi for her help in leaf adhesion assays. This study was supported by grants from the MRC and BBSRC.

References

- Baggesen DL, Wegener HC. (1994). Phage types of *Salmonella enterica* ssp. *enterica* serovar Typhimurium isolated from production animals and humans in Denmark. *Acta Vet Scand* **35**: 349–354.
- Barak JD, Jahn CE, Gibson DL, Charkowski AO. (2007). The role of cellulose and O-antigen capsule in the colonization of plants by *Salmonella enterica*. *Mol Plant Microbe Interact* **20**: 1083–1091.
- Crook PD, Aguilera JF, Threlfall EJ, O'Brien SJ, Sigmundsdóttir G, Wilson D *et al.* (2003). A European outbreak of *Salmonella enterica* serotype Typhimurium definitive phage type 204b in 2000. *Clin Microbiol Infect Dis* **9**: 839–845.
- Datsenko KA, Wanner BL. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K12 using PCR products. *Proc Natl Acad Sci USA* **97**: 6640–6645.
- Gupta SK, Nalluswami K, Snider C, Perch M, Balasegaram M, Burmeister D *et al.* (2007). Outbreak of *Salmonella* Braenderup infections associated with Roma tomatoes, northeastern United States, 2004: a useful method for subtyping exposures in field investigations. *Epidemiol Infect* **135**: 1165–1173.
- Kessel AS, Gillespie IA, O'Brien SJ, Adak GK, Humphrey TJ, Ward LR. (2001). General outbreaks of infectious intestinal disease linked with poultry, England and Wales, 1992–1999. *Commun Dis Public Health* **4**: 171–177.
- Klerks MM, Franz E, van Gent-Pelzer M, Zijlstra C, van Bruggen AH. (2007). Differential interaction of *Salmonella enterica* serovars with lettuce cultivars and plant–microbe factors influencing the colonization efficiency. *ISME J* **1**: 620–631.
- Little CI, Gillespie IA. (2008). Prepared salads and public health. *J Appl Microbiol*; e-pub ahead of print 4 April, doi:10.1111/j.1365-2672.2008.03801.x.
- Nygård K, Lassen J, Vold L, Andersson Y, Fisher I, Löfdahl S *et al.* (2008). Outbreak of *Salmonella* Thompson infections linked to imported rucola lettuce. *Foodborne Pathog Dis* **5**: 165–173.
- Olsen JE, Sørensen M, Brown DJ, Gaarslev K, Bisgaard M. (1992). Plasmid profiles as an epidemiological marker in *Salmonella enterica* serovar Berta infections. Comparison of isolates obtained from humans and poultry. *APMIS* **100**: 221–228.
- Pakalniskiene J, Falkenhorst G, Lisby M, Madsen SB, Olsen KE, Nielsen EM *et al.* (2006). A foodborne outbreak of enterotoxigenic *E. coli* and *Salmonella* Anatum infection after a high-school dinner in Denmark, November 2004. *Epidemiol Infect* **6**: 1–6.
- Pezzoli L, Elson R, Little C, Fisher I, Yip H, Peters T *et al.* (2007). International outbreak of *Salmonella* Senftenberg. *Euro Surveill* **12**: E070613–E070614.
- Rodrigue DC, Tauxe RV, Rowe B. (1990). International increase in *Salmonella* Enteritidis: a new pandemic? *Epidemiol Infect* **105**: 21–27.
- Shaw RK, Berger CN, Feys B, Knutton S, Pallen MJ, Frankel G. (2008). Enterohemorrhagic *Escherichia coli* exploits EspA filaments for attachment to salad leaves. *Appl Environ Microbiol* **74**: 2908–2914.
- Sivapalasingham S, Friedman CR, Cohen L, Tauxe RV. (2004). Fresh produce: a growing cause of outbreaks of foodborne illness in the United States, 1973 through 1997. *J Food Protect* **67**: 2342–2353.